



LJMU Research Online

Alwan, N, Moss, SL, Elliot-Sale, KJ, Davies, IG and Enright, KJ

A Narrative Review on Female Physique Athletes: The Physiological and Psychological Implications of Weight Management Practices

<http://researchonline.ljmu.ac.uk/id/eprint/10842/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Alwan, N, Moss, SL, Elliot-Sale, KJ, Davies, IG and Enright, KJ (2019) A Narrative Review on Female Physique Athletes: The Physiological and Psychological Implications of Weight Management Practices. International Journal of Sport Nutrition and Exercise Metabolism. ISSN 1526-484X

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

1 A Narrative Review on Female Physique Athletes: The Physiological
2 and Psychological Implications of Weight Management Practices

3 Nura Alwan¹, Samantha L. Moss², Kirsty J. Elliot-Sale³, Ian G.
4 Davies¹, and Kevin Enright¹

5

6 ¹School of Sport Studies, Leisure and Nutrition, Liverpool John
7 Moores University, UK.

8 ²Department of Sport and Exercise Sciences, University of Chester,
9 UK.

10 ³Musculoskeletal Physiology Research Group, Sport, Health and
11 Performance Enhancement Research Centre, Nottingham Trent
12 University, UK.

13

14 **Running head:** Health considerations in female physique athletes

15

16 **Corresponding author:**

17 Nura Alwan

18 ¹School of Sport Studies, Leisure and Nutrition,

19 Liverpool John Moores University,

20 Barkhill Road,

21 Liverpool,

22 L17 6BD

23 United Kingdom

24 Email: N.alwan@2016.ljmu.ac.uk

25

26 **Abstract**

27 Physique competitions are events in which aesthetic appearance and
28 posing ability are valued above physical performance. Female
29 physique athletes are required to possess high lean body mass and
30 extremely low fat mass in competition. As such, extended periods of
31 reduced energy intake and intensive training regimens are utilised with
32 acute weight loss practices at the end of the pre-competition phase.
33 This represents an increased risk for chronic low energy availability
34 and associated symptoms of *Relative Energy Deficiency in Sport*,
35 compromising both psychological and physiological health. Available
36 literature suggests that a large proportion of female physique athletes
37 report menstrual irregularities (*e.g.*, amenorrhea and oligomenorrhea),
38 which are unlikely to normalise immediately post-
39 competition. Furthermore, the tendency to reduce intakes of numerous
40 essential micronutrients is prominent among those using restrictive
41 eating patterns. Following competition reduced resting metabolic rate,
42 and hyperphagia, are also a concern for these female athletes, which
43 can result in frequent weight cycling, distorted body image and
44 disordered eating/eating disorders. Overall, female physique athletes
45 are an understudied population and the need for more robust studies to
46 detect low energy availability and associated health effects is
47 warranted. This narrative review aims to define the natural female
48 physique athlete, explore some of the physiological and psychological
49 implications of weight management practices experienced by female
50 physique athletes and propose future research directions.

51

52 **Keywords**

53 Fat loss, low energy availability, physique events, body composition,

54 nutrition

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73 **Background**

74 Physique competitions are events in which competitors are judged on
75 aesthetic appearance rather than on physical performance. Natural
76 (*i.e.*, drug-free) physique competitions have evolved dramatically in
77 recent years, with a growth in organisations, contests and classes
78 (Halliday et al., 2016). The International Federation of Body Building
79 and Fitness (IFBB) hosts over 2,000 competitions annually, in 196
80 affiliated countries. Approximately 1,300 female and male athletes
81 competed at the World Fitness Championships in 2017 (Rowbottom,
82 2017), and this number is anticipated to increase, with around 1,000
83 new members joining the sport each year (Parish et al., 2010).

84 Female physique (FP) athletes have aspirations of achieving a lean and
85 muscular body composition for competitive success (Halliday et al.,
86 2016). Preparing for a natural physique competition provides a myriad
87 of health benefits including improvement in cardiovascular status
88 (Kistler et al., 2014; Robinson et al., 2015), muscle strength
89 (Campbell et al., 2018), increasing feelings of accomplishment, and
90 transient improvements in self-esteem (Aspridis et al., 2014; Baghurst
91 et al., 2014; Probert et al., 2007). Despite these positive outcomes,
92 numerous unfavorable effects also exist, including, but not limited to:
93 diminished levels of reproductive hormones (Hulmi et al., 2016) and
94 symptoms of disordered eating and eating disorders (DE/ED)
95 (Walberg and Johnston, 1991). Available research on FP athletes
96 reveals prolonged periods of sustained energy restriction and intensive
97 training regimens in an attempt to acquire and maintain a lean body
98 composition, indicating an increased risk of low energy availability
99 (LEA) and its associated effects (Fagerberg, 2017). For a thorough

100 understanding of the existence, aetiologies and clinical consequences
101 of LEA, readers are directed to the review by Loucks et al. (2011).

102 Prolonged periods of LEA with or without disordered eating,
103 menstrual dysfunction and low bone mineral density is termed the
104 Female Athlete Triad (Triad), representing a medical condition
105 observed in females who perform high levels of physical activity
106 (Manore, 2007). In order to describe a wide range of physiological,
107 psychological and performance-related impairments associated with
108 LEA, the International Olympic Committee introduced the concept of
109 Relative Energy Deficiency in Sport (RED-S) in 2014 (Mountjoy et
110 al., 2014). Considering the health risks of RED-S, and the increasing
111 participation of females in physique events, the purpose of this
112 narrative review was three-fold: 1. to define the natural female
113 physique athlete; 2. to explore the physiological and psychological
114 implications of the weight management practices experienced by the
115 natural FP athlete; 3. to address future research directions.

116

117

118

119

120

121

122

123

124 **Literature Search**

125 A literature search was conducted using databases: PubMed, Web of
126 Science, Google Scholar, and SPORTDiscus (via EBSCO) up to 10th
127 September 2018. Despite slight variation in the terminology used for
128 ‘physique athlete’ in the literature, synonyms were included in the
129 search strategy. Various combinations of the following search terms
130 were used, for the search: ‘physique athlete’ OR ‘fitness competitor’
131 OR ‘bodybuilding’ OR ‘competitive body-builder’ OR ‘figure athlete’
132 AND (contest or competition OR dieting OR dietary intake or nutrition
133 OR macronutrient OR micronutrient OR training OR body
134 composition OR peak week OR practices OR weight loss OR weight
135 regain). Several other search terms associated with health outcomes
136 included: ‘physique athlete’ OR ‘fitness competitor’ OR
137 ‘bodybuilding’ OR ‘competitive body-builder’ OR ‘figure athlete’
138 AND (energy availability, menstrual cycle OR bone, OR eating OR
139 body image). Any additional articles relevant to the scope of this
140 narrative review were obtained through PubMed via the function
141 “similar articles” or from the reference lists of the included studies.

142 Criteria for inclusion were: *i*) studies published in English language
143 and in peer-reviewed articles within the past 30 years (*i.e.*, theses or
144 conference abstracts were not eligible), *ii*) studies involving human
145 participants, *iii*) studies with participants who were specifically
146 engaging or been engaged in physique competitions, across any
147 category (*i.e.*, bikini fitness, wellness fitness, and figure), *iv*) studies
148 using female participants, or studies using both female and male
149 participants, and *v*) studies investigating at least one of the following:
150 body composition, nutritional intake, micronutrients, training

151 strategies, psychology, menstrual cycle, hormonal markers, bone
152 mineral density, energy availability, and weight loss/management
153 practices). Exclusion criteria were studies that reported use of
154 performance-enhancing drugs, and only male participants.

155

156 **Definition of the natural female physique athlete**

157 Benjamin and Joseph Weider established the first organisation which
158 specialised solely in bodybuilding events, known as the IFBB (Vallet,
159 2017). To date, the IFBB is one of the most influential amateur sports
160 organisations in the bodybuilding sphere and is an official signatory of
161 the World Anti-Doping Code where athletes participate in random
162 drug testing programs, such as urinalysis and polygraph tests for
163 prohibited substances (IFBB, 2014).

164 Whilst bodybuilding is traditionally a male dominated sport, the
165 growth of female competitors has increased significantly in recent
166 times (Spendlove et al., 2015). This growth in popularity is largely due
167 to the introduction of new female-specific physique categories (*e.g.*,
168 Fitness, Body Fitness and Bikini Fitness) since 1995 (Spendlove et al.,
169 2015; Tajrobehkar, 2016). As these new categories allowed ‘smaller’
170 competitors to enter the sport, and reduced the emphasis on muscle
171 mass, they have encouraged healthier practices, indirectly attracting
172 more women from mainstream society than in previous decades
173 (Tajrobehkar, 2016).

174 Female physique athletes are assessed on aesthetic appearance and
175 posing ability whereby high lean body mass (LBM) and low fat mass
176 (FM) are key markers of performance (Kleiner et al., 1994).

177 Competitions involve comparison rounds; wherein athletes are
178 instructed to perform poses, and a final round; in which top ranked
179 athletes perform an individual posing routine (Steele et al., 2018). The
180 intricate scoring system assesses athlete features, such as symmetry,
181 muscularity, size and presentation (*i.e.*, personal confidence, facial
182 beauty, and skin condition) (Choi, 2003; Obel, 1996). Unlike other
183 weight-restricted sports (*e.g.*, male bodybuilding, wrestling and
184 boxing), in which weight categories are utilised, FP athletes are
185 allocated to categories based on their subjective assessment of the
186 amount of LBM and FM, and are then further sub-classified by height
187 (Fry et al., 1991). At one end of the continuum (*i.e.*, bikini fitness),
188 athletes typically have less LBM and higher FM, whilst at the other
189 end (*i.e.*, physique), athletes are diametrically opposed with high LBM
190 and a corresponding low FM (Fig.1).

191

192 **[Insert Figure 1 near here]**

193

194 **Body composition in competition**

195 Typically, an annual schedule for the physique athlete is divided into
196 an off-season phase and a pre-competition phase (Hackett et al., 2013).
197 Within the off-season phase, physique athletes manipulate resistance
198 training variables including volume, intensity and frequency for the
199 purpose of gaining LBM (Spendlove et al., 2015). This period can last
200 years and is characterised by a positive energy balance, in conjunction
201 with a high protein intake to stimulate muscle anabolism (Phillips,
202 2004; Campbell et al., 2018). In the pre-competition phase, the

203 majority of athletes attempt to reduce body fat levels and preserve
204 LBM using a combination of rigorous resistance and aerobic training,
205 while manipulating their nutritional intake to achieve a negative
206 energy balance (Hackett et al., 2013; Petrizzo et al., 2017). The pre-
207 competition phase lasts between 12 and 24 weeks (Mitchell et al.,
208 2018) and athletes are likely to compete between two to three times
209 per year (Chappell et al., 2018). Usually, the pre-competition phase is
210 followed by a recovery phase (a transition to off-season), during which
211 athletes increase their total energy intake and decrease their total
212 training load (Hulmi et al., 2016). Previous research reports the
213 magnitude of weight loss is in the range of 6-10 kg over a 18-24 week
214 period (Table 1). This suggests that FP athletes pursue a gradual
215 approach to weight loss (~ 0.4 kg per week), similar to male
216 bodybuilding and physique athletes (~ 0.6-0.8 kg per week) (Chappell
217 et al., 2018; Kistler et al., 2014; Robinson et al., 2015; Rossow et al.,
218 2013). In the end stages of the pre-competition phase, FP athletes
219 achieve 8.6 - 16% body fat (Hulmi et al., 2016; Rohrig et al., 2017;
220 Tinsley et al., 2018; Trexler et al., 2017), which is exceptionally lower
221 than the recommended values for female athletes (Sundgot-Borgen
222 and Garthe, 2011).

223

224 **Strategies to manipulate body composition during competition** 225 **week**

226 Whilst FP athletes employ a gradual approach to fat loss, acute weight
227 loss practices occur during the competition week (Helms et al., 2014).
228 Peer-reviewed articles suggest fluid, salt, and carbohydrate
229 manipulation is commonly practiced to reduce body water content in

230 order to enhance muscle definition on competition day (Mitchell et al.,
231 2017; Shephard, 1994). Nearly one-half of twenty-two FP athletes
232 practiced water manipulations (36 %), whereas more than two-quarters
233 practiced carbohydrate manipulations (77 %) (Chappell and Simper,
234 2018). Water loading, followed by water restriction is allegedly used
235 to modify renal hormones and encourage urination beyond the period
236 of increased fluid intake, resulting in reduced body water (Helms et
237 al., 2014; Mitchell et al., 2017). The physiological effects of water
238 loading have only been investigated in male combat sport athletes with
239 a purpose of making-weight (Crighton et al., 2016; Reale et al., 2018),
240 as opposed to physique athletes trying to enhance their aesthetic
241 appearance. The acute weight loss experienced early in competition
242 week (~7-5 days prior to competition) is likely to be mediated by
243 glycogen depletion prior to a carbohydrate loading protocol (Chappell
244 and Simper, 2018). Female physique athletes reduce their
245 carbohydrate intake from 4.1- 4.5 $\text{g}\cdot\text{kg}^{-1}\cdot\text{d}^{-1}$ before entering the pre-
246 competition phase, to 1.2 - 2.7 $\text{g}\cdot\text{kg}^{-1}\cdot\text{d}^{-1}$ at the end stages of pre-
247 competition phase (Halliday et al., 2016; Rohrig et al., 2017). In one
248 case, daily carbohydrate intake was reduced to ~ 0.3 $\text{g}\cdot\text{kg}\cdot\text{d}^{-1}$, three
249 days prior to competition (Tinsley et al., 2018). From the available
250 evidence, it appears that during the pre-competition phase, FP athletes
251 fall considerably below the carbohydrate recommendations for
252 moderate volume training (5-7 $\text{g}\cdot\text{kg}^{-1}\cdot\text{d}^{-1}$) (Manore, 2002). Addressing
253 the distribution of carbohydrate intake throughout the day and in
254 relation to training, could provide further insights into the strategies
255 used to optimise body composition (Slater and Phillips, 2011).

256 Based on limited data, the efficacy and safety of competition week
257 strategies in physique events are still unknown, but might be
258 detrimental to athlete health (Chappell and Simper, 2018; Helms et al.,
259 2014) by increasing the risks associated with hyponatremia and
260 glycogen depletion (Slater and Phillips, 2011).

261

262 **Health implications for the female physique athlete**

263 Physique athletes typically reduce their total energy intake to induce
264 gradual weight loss over a prolonged period of time, and progress
265 towards acute weight loss methods, such as restrictive diets (energy
266 availability [EA] $< 30 \text{ kcal} \cdot \text{kg}^{-1} \text{ FFM} \cdot \text{d}^{-1}$, where FFM = fat free mass),
267 in the latter stages of the pre-competition phase (Sundgot-Borgen et
268 al., 2013; Fagerberg et al., 2017). As such, FP athletes face major
269 health-related challenges in an attempt to reach and maintain a lean
270 body composition.

271

272 *Reduced energy availability in female physique athletes*

273 Current literature on FP athletes has documented prolonged periods of
274 LEA, specifically during the pre-competition phases. Halliday and
275 colleagues (Halliday et al., 2016) showed that during a 20-week pre-
276 competition phase, the estimated mean EA was categorised as low in
277 the initial ($27.9 \text{ kcal} \cdot \text{kg}^{-1} \text{ FFM} \cdot \text{d}^{-1}$) and latter ($23.3 \text{ kcal} \cdot \text{kg}^{-1} \text{ FFM} \cdot \text{d}^{-1}$)
278 stages of the phase, respectively. In this study (Halliday et al., 2016),
279 total energy intake and exercise energy expenditure were self-reported
280 and reproductive function was not measured. Similarly, Tinsley et al.
281 (2018) documented caloric intakes of between 18.2 and $31.1 \text{ kcal} \cdot \text{kg}^{-1}$

282 FFM·d⁻¹ in a FP athlete (during two different pre-competition phases)
283 indicating extreme caloric restriction (Manore, 2002). Although EA
284 was not objectively quantified, the authors estimated that the athlete
285 fell below the threshold of EA for the maintenance of normal
286 physiological function based on total energy intake and body
287 composition data. Self-report research designs are not uncommon in
288 the literature on physique athletes and, as such should be interpreted
289 with caution (Fagerberg, 2017). Therefore, EA data in FP athletes
290 remains questionable considering the lack of sensitive and relevant
291 screening tools (Heikura et al., 2018). Nonetheless, aforementioned
292 studies highlight that FP athletes may induce sub-optimal EA and
293 shows the importance for future studies on this topic to utilise more
294 robust measures of total energy intake and exercise energy expenditure
295 in order to accurately evaluate EA (Elliott-Sale et al., 2018; Fagerberg,
296 2017).

297

298 *Nutrient deficiency*

299 Bodybuilding diets are traditionally characterised as restrictive and
300 monotonous, as they often limit food variability (Kleiner et al., 1994).
301 As a consequence, compromised micronutrient status is often observed
302 in the pre-competition phase among FP athletes (Slater and Phillips,
303 2011). Calcium, iron, zinc and sodium intakes have been shown to
304 decrease significantly, to less than two-thirds (~ 67%) of the
305 Recommended Daily Allowance (RDA) (Newton et al., 1993;
306 Walberg-Rankin and Gwazdauskas, 1993) in the absence of dietary
307 supplements during the pre-competition phase. These results may be
308 attributed to restricted energy intake combined with the elimination of
309 sodium and dairy products from the diet (Steen, 1991). Considering

310 that weight loss trends/dietary fads typically change over time, it is
311 worth noting that the applicability of the aforementioned studies might
312 be limited (Spendlove et al., 2015).

313 More recently, Ismaeel et al. (2017) showed that FP athletes who used
314 extreme restrictive eating patterns consumed significantly less protein
315 (123.3 ± 22.9 g *cf.* 64.8 ± 16.2 g, $p = 0.02$), sodium ($4,059.6 \pm 397.0$
316 mg *cf.* $2,635.9 \pm 1,028.3$ mg, $p = 0.03$), vitamin E (9.9 ± 2.1 mg *cf.*
317 5.8 ± 1.2 mg, $p = 0.03$) and vitamin C (169.5 ± 47.4 mg *cf.* $65.5 \pm$
318 26.5 mg, $p = 0.02$) than athletes who permitted dietary flexibility
319 (Ismaeel et al., 2017). These differences may be caused by the large
320 variation in total energy intake ($1,964.9 \pm 258.9$ kcal·d⁻¹ *cf.* $1,454.7 \pm$
321 541.4 kcal·d⁻¹) consumed by each group. While the study (Ismaeel et
322 al., 2017) included dietary supplements in the micronutrient analysis,
323 it did not specify whether individuals were in the pre-competition or
324 off-season phase. Nevertheless, these results identify potential risks
325 for deficiencies in essential nutrients for FP athletes, thereby
326 suppressing the immune function and causing increased susceptibility
327 to illnesses and infections, especially for those engaging in restrictive
328 eating patterns (Sundgot-Borgen and Garthe, 2011). As the majority
329 of studies assessing micronutrient status have also used self-report
330 methods (Ismaeel et al., 2017; Kleiner et al., 1994; Newton et al., 1993;
331 Walberg-Rankin and Gwazdauskas, 1993; Walberg and Johnston,
332 1991), it is prudent that future measures are clarified using biomarkers
333 in blood or urine samples.

334

335

336 *Menstrual irregularities, endocrine effects and bone health in female*
337 *physique athletes*

338 Many active women with LEA develop various forms of reproductive
339 dysfunction, including oligomenorrhea, amenorrhea and luteal phase
340 defects (Manore, 2002). Low energy availability causes alterations in
341 the hypothalamic-pituitary-ovarian axis, namely diminished secretion
342 of luteinizing hormone and follicle stimulating-hormone, which
343 subsequently reduces oestrogen production. The final consequence is
344 typically described as functional hypothalamic amenorrhea (West,
345 1998). Previous research has shown that 82-86% of females (non-
346 contraceptive users) who entered at least one physique competition
347 were either oligomenorrheic or amenorrheic (Walberg-Rankin and
348 Gwazdauskas, 1993; Walberg and Johnston, 1991). Similarly, case
349 studies have also observed amenorrhea (Hulmi et al., 2016; Petrizzo et
350 al., 2017; Rohrig et al., 2017), with some reporting delays in
351 menstruation of up to 71 weeks post-competition (Halliday et al.,
352 2016; Kleiner et al., 1994; Kleiner et al., 1990).

353 Changes to reproductive and metabolic hormones in FP athletes have
354 been observed in the pre-competition phase, including decreases in
355 oestradiol, testosterone, thyroid stimulating hormone, triiodothyronine
356 (T3) and leptin (Table 1). These hormones were normalised within 4 -
357 16 weeks post-competition, when supported by an increased intake of
358 protein ($\sim 2. \text{g} \cdot \text{kg}^{-1} \cdot \text{d}^{-1}$) and greater EA (Hulmi et al., 2016; Trexler et
359 al., 2017) with the exception of serum T3 and testosterone (Hulmi et
360 al., 2016), which were only partially recovered 12-16 weeks after
361 competition. As such, the suppression of these key metabolic
362 hormones persist further into the recovery phase, possibly due to the

363 effects of dropping below the EA threshold regardless of altered
364 exercise regimen, as previously described by Loucks and Heath
365 (1994). More longitudinal data is required on endocrine and metabolic
366 function beyond the 16 weeks post-competition to better understand
367 the time-course for full restoration.

368 Regular menstrual cycles are often used as a surrogate marker of long-
369 term LEA; however, the use of hormonal contraceptives may
370 obfuscate this relationship (Heikura et al., 2018). Hormonal
371 contraceptives provide negative feedback to the hypothalamus and
372 pituitary glands, leading to suppression of follicle stimulating-
373 hormone, luteinizing hormone and gonadotropin-releasing hormone,
374 and continuous down-regulation of endogenous oestrogen and
375 progesterone (Elliott-Sale et al., 2013). Previous data in FP athletes
376 have failed to investigate female sex hormones (*i.e.*, oestrogen and
377 progesterone) (Trexler et al., 2017), did not include hormonal
378 contraceptive users (Halliday et al., 2016; Rohrig et al., 2017; Tinsley
379 et al., 2018) or grouped all oral contraceptive users together, making
380 the interpretation difficult (Elliott-Sale et al., 2013). Considering the
381 high prevalence of hormonal contraceptive use (Hulmi et al., 2016),
382 there is great concern that FP athletes, who are experiencing chronic
383 LEA, are going undetected, as hormonal contraceptive use maintains
384 regular menstrual cycles. To this end, there is a need for studies to
385 determine whether the FP athletes, who are using hormonal
386 contraceptives, are at increased risk of endocrine dysfunction.

387 Although it is not unusual for bone mineral density to be compromised
388 during calorie restriction and reduced body mass, it is possible that the
389 minimal changes observed in bone mineral density (1.062-1.204g.cm³)

390 (Van der Ploeg et al., 2001; Hulmi et al., 2016; Petrizzo et al., 2017)
391 is explained by the high-impact and weight-bearing activities
392 performed in their training regimens (Zanker et al., 2004). As a result,
393 this may have served to retain bone-mineral density compartment
394 (Layne & Nelson, 1999).

395

396 *Weight cycling*

397 Female physique athletes often experience rapid weight gain following
398 competitions (Andersen et al., 1995; Walberg-Rankin and
399 Gwazdauskas, 1993) with one study reporting uncontrollable binge
400 eating behaviour, reflecting a hyperphagic effect to intensive weight
401 loss protocols (Trexler et al., 2017). This practice is commonly known
402 as ‘weight cycling’ (*i.e.*, repeated cycles of weight loss and regain).
403 Previous research has reported unfavorable metabolic parameters
404 including a decline in resting metabolic rate (RMR) (reduced between
405 154.7 and 226 kcal) (Rohrig et al., 2017; Tinsley et al., 2018) during
406 pre-competition phase and weight regain of up to 8.6 kg at 4 weeks
407 post-competition refeeding in females (Walberg-Rankin and
408 Gwazdauskas, 1993). The RMR suppression is possibly induced by
409 the dietary restriction during weight loss resulting in alterations in
410 leptin levels, thyroid status and sympathetic nervous system activity
411 (Stiegler and Cunliffe, 2006). Conversely, recent case studies have
412 shown that some FP athletes use a “reverse dieting” technique, in order
413 to avoid those implications (Trexler et al., 2014). This strategy requires
414 athletes to slowly increase their energy intake in an effort to limit any
415 rapid increases in FM, and to prevent reductions in RMR (Trexler et
416 al., 2014). However, the effort to “reverse” (*i.e.*, slowly increase)

417 energy intake requires considerable discipline to curb with the
418 increases in appetite sensations (Greenway, 2015), and therefore the
419 authors speculate, whether such a strategy is achievable. Future
420 research on “reverse dieting” technique in the recovery phase is
421 warranted.

422

423 *Disordered eating /Eating Disorders behaviours*

424 Considering that appearance is a major criterion to judge performance
425 of FP athletes, the increased risk of DE/ED in this population is
426 perhaps unsurprising. Important risk and trigger factors of poor eating
427 habits in FP athletes may include the focus on aesthetic appearance as
428 the primary performance marker in competition (Sundgot-Borgen and
429 Torstveit, 2004), the peer/media pressure which can elicit body
430 dissatisfaction (Hausenblas et al., 2013) and the influences from
431 coaches with inadequate nutrition knowledge (Sundgot-Borgen,
432 1994). There is also evidence that FP athletes are particularly
433 vulnerable to DE/ED and body image dissatisfaction because of the
434 preoccupation with being muscular and lean (Devrim et al., 2018).

435 For example, a cross-sectional study by Walberg and Johnston (1991)
436 compared 12 aspiring and retired FP athletes with 103 recreational
437 weight-lifters on the Eating Disorder Inventory. Results revealed that
438 FP athletes had significantly greater food obsessions (67%),
439 uncontrolled urges to eat (58%) and felt more terrified of becoming fat
440 (58%; all $p < 0.05$). The use of laxatives, for weight loss, (17% *cf.*
441 15%) and binge eating (50% *cf.* 62%) were similar between the groups.

442 In another study, Andersen et al. (1998) reported that ten out of twenty-
443 six FP athletes experienced binge eating episodes in the recovery
444 phase, and eighteen out of twenty-six FP athletes displayed body and
445 weight dissatisfaction, reiterating that there is a high risk of eating and
446 body image-related problems within the sport (Pope et al., 1997).
447 Nevertheless, the small sample size and the lack of any comparative
448 group analysis by Andersen et al. (1998) somewhat limits the
449 interpretation. To the authors' knowledge, no quantitative data
450 examining disordered eating behaviours exists for a large cohort of
451 natural FP athletes.

452 Furthermore, it is difficult to capture sensitive data using questionnaire
453 methods concerning mental health and well-being without a
454 confirmatory interview (Andersen et al., 1998). Athletes may be
455 anxious of revealing inappropriate eating practices in fear of being
456 negatively judged, which could prevent honest disclosure.
457 Nevertheless, there is a plausible link between participation in
458 physique sports and DE behaviours. Further research is warranted to
459 explore the psychopathological and behavioural outcomes in these
460 athletes. Understanding the experiences and perceptions of weight
461 management and eating behaviours across the pre-competition,
462 recovery and off-season phases might be of particular importance.
463 Using validated screening tools to detect DE and EDs and follow-up
464 interviews will allow researchers to collect comprehensive data that
465 could inform practice.

466

467 **Conclusions and future research**

468 The ultimate determinant of competitive success in physique events is
469 a high degree of muscularity and minimal levels of body fat. As such,
470 physique athletes engage in both prolonged energy restriction and
471 intensive training regimens in order to meet these demands. Some FP
472 athletes may be vulnerable to chronic LEA and associated
473 physiological and psychological health effects, even during the
474 recovery phase. Despite an increased participation in physique events,
475 there is paucity in the literature on FP athletes. Future research should
476 therefore:

- 477 *i)* identify the weight loss strategies and DE/ED behaviours
478 of FP athletes, in order to determine the risks of LEA in
479 this population;
- 480 *ii)* explore such strategies using a qualitative approach, to
481 enable FP athletes to express and elaborate on their
482 experiences of weight management, eating behaviours
483 and psycho-physiological health implications;
- 484 *iii)* investigate endocrine and micronutrient changes in FP
485 athletes using objective biomarkers, to assess whether
486 these individuals are in chronic states of LEA throughout
487 the season;
- 488 *iv)* develop effective, safe and evidence-based nutritional
489 recovery guidelines to minimise any long-term health
490 implications.

491

492 **Practical Application Statement**

493 At present, it is difficult to draw upon practical applications from the
494 existing literature. FP athletes are an understudied population, and
495 methodological limitations exist. A primary issue is that the majority
496 of cited reports are case studies, or observational studies with small
497 sample sizes, which may be insufficient for drawing definite
498 conclusions on the possible physiological and psychological health
499 implications among natural FP athletes. More research will have a
500 valuable impact upon the advice and strategies provided by coaches
501 and sport science/health professionals who work with these athletes.

502 It is worth noting that many female athletes are reluctant to discuss
503 their competition strategies and health histories with health or sport
504 science professionals (Manore, 2002), making this population difficult
505 to research (Aspridis et al., 2014), and may explain the small sample
506 sizes reported by previous studies (Halliday et al., 2016; Ismaeel et al.,
507 2017; Petrizzo et al., 2017). Therefore, it is imperative that both
508 coaches and sport science/health professionals working in the field
509 build trusting relationships with physique athletes and respect their
510 desires to be lean, with a view to achieve an optimum body
511 composition and health outcomes through a collaborative relationship.

512

513 **Novelty statement**

514 This is the first review to summarise the common physiological and
515 psychological health implications among female physique athletes.

516

517 **Acknowledgements**

518 The authors would like to thank Dr Jim McVeigh, Director of Public
519 Health at Liverpool John Moors University for his valuable input to
520 this review.

521

522 **Author's contributions**

523 The lead author on this paper was NA. SM, KE-S, ID and KE
524 participated in the drafting of the manuscript. All authors read and
525 approved the final manuscript.

526

527 **Declaration of funding sources**

528 No funding was received for the preparation of this manuscript

529

530 **Conflicts of interest**

531 The authors declare that they have no conflicts of interest

532

533

534

535

536

537

538

539

540

541

542

543

544

545

546

547

548

549

550

551

552 **References**

553

554 American Psychiatric, A., American Psychiatric, A., and Force, D. S.
 555 M. T. (2013). Feeding and eating disorders. In *Diagnostic and*
 556 *statistical manual of mental disorders* (5th ed., pp. 329-354).
 557 Washington, DC: Author.

558

559 Andersen, R. E., Barlett, S. J., Morgan, G. D., and Brownell, K. D.
 560 (1995). Weight loss, psychological, and nutritional patterns in
 561 competitive male body builders. *International Journal of*
 562 *Eating Disorders*, 18(1), 49-57.

563

564 Andersen, R. E., Brownell, K. D., Morgan, G. D., and Bartlett, S. J.
 565 (1998). Weight loss, psychological, and nutritional patterns in
 566 competitive female bodybuilders. *Eating Disorders*, 6(2),
 567 159-167.

568

569 Aspridis, A., O'Halloran, P., and Liamputtong, P. (2014). Female
 570 Bodybuilding: Perceived Social and Psychological Effects of
 571 Participating in the Figure Class. *Women in Sport and*
 572 *Physical Activity Journal*, 22(1), 24-29.

573

574 Baghurst, T., Parish, A., and Denny, G. (2014). Why Women Become
 575 Competitive Amateur Bodybuilders. *Women in Sport and*
 576 *Physical Activity Journal*, 22(1), 5-9.

577

578

579 Campbell, B. I., Aguilar, D., Conlin, L., Vargas, A., Schoenfeld, B. J.,
 580 Corson, A., and Couvillion, K. (2018). Effects of High Versus
 581 Low Protein Intake on Body Composition and Maximal
 582 Strength in Aspiring Female Physique Athletes Engaging in
 583 an 8-Week Resistance Training Program. *International*
 584 *Journal of Sport Nutrition and Exercise Metabolism*, 1-6.

585

586 IFBB Elite Pro Categories. (2017). Retrieved from
 587 <https://eliteproifbb.com/our-disciplines/categories/>.

588

589 Chappell, A., and Simper, T. (2018). Nutritional Peak Week and
 590 Competition Day Strategies of Competitive Natural
 591 Bodybuilders. *Sports*, 6(4), 126.

592

593 Chappell, A. J., Simper, T., and Barker, M. E. (2018). Nutritional
 594 strategies of high level natural bodybuilders during
 595 competition preparation. *Journal of the International Society*
 596 *of Sports Nutrition*, 15, 4.

597

598 Choi, P. Y. L. (2003). Muscle matters: maintaining visible differences
 599 between women and men. *Sexualities, Evolution and Gender*,
 600 5(2), 71-81.

601

602 Crighton, B., Close, G. L., and Morton, J. P. (2016). Alarming weight
 603 cutting behaviours in mixed martial arts: a cause for concern
 604 and a call for action. *British Journal of Sports Medicine*, 50(8),
 605 446-447.

606

- 607 Devrim, A., Bilgic, P and Hongo, N. (2018). Is there any relationship
608 between body image perception, eating disorders, and muscle
609 dysmorphic disorders in male bodybuilders? *American Journal*
610 *of Men's Health*, 12(5), 1746-1758
611
612
- 613 Elliott-Sale, K. J., Smith, S., Bacon, J., Clayton, D., McPhilimey, M.,
614 Goutianos, G., and Sale, C. (2013). Examining the role of oral
615 contraceptive users as an experimental and/or control group in
616 athletic performance studies. *Contraception*, 88(3), 408-412.
617
- 618 Elliott-Sale, K. J., Tenforde, A. S., Parziale, A. L., Holtzman, B., and
619 Ackerman, K. E. (2018). Endocrine Effects of Relative Energy
620 Deficiency in Sport. *International Journal of Sport Nutrition*
621 *and Exercise Metabolism*, 28, 335-349.
622
- 623 Fagerberg, P. (2017). Negative Consequences of Low Energy
624 Availability in Natural Male Bodybuilding: A Review.
625 *International Journal of Sport Nutrition and Exercise*
626 *Metabolism*, 0(0), 1-18.
627
- 628 Fry, A. C., Ryan, A. J., Schwab, R. J., Powell, D. R., and Kraemer, W.
629 J. (1991). Anthropometric characteristics as discriminators of
630 body-building success. *Journal of Sports Sciences*, 9(1), 23-
631 32.
632
- 633 Greenway, F. L. (2015). Physiological adaptations to weight loss and
634 factors favouring weight regain. *International Journal Of*
635 *Obesity*, 39, 1188.
636
- 637 Hackett, D. A., Johnson, N. A., and Chow, C. M. (2013). Training
638 practices and ergogenic aids used by male bodybuilders.
639 *Journal of Strength and Conditioning Research*, 27(6), 1609-
640 1617.
641
- 642 Halliday, T. M., Loenneke, J. P., and Davy, B. M. (2016). Dietary
643 Intake, Body Composition, and Menstrual Cycle Changes
644 during Competition Preparation and Recovery in a Drug-Free
645 Figure Competitor: A Case Study. *Nutrients*, 8(11), 740.
646
- 647 Hausenblas, H. A., Campbell, A., Menzel, J. E., Doughty, J., Levin,
648 M. and Thompson, J. K. (2013). Media effects of experimental
649 presentation of the ideal physique on eating disorder
650 symptoms: A meta-analysis of laboratory studies. *Clinical*
651 *Psychology Review*, 33(1), 168-181.
652
- 653 Heikura, I. A., Uusitalo, A. L. T., Stellingwerff, T., Bergland, D.,
654 Mero, A. A., and Burke, L. M. (2018). Low Energy
655 Availability Is Difficult to Assess but Outcomes Have Large
656 Impact on Bone Injury Rates in Elite Distance Athletes.
657 *International Journal of Sport Nutrition and Exercise*
658 *Metabolism*, 28(4), 403-411.
659
- 660 Helms, E. R., Aragon, A.A., Fitschen PJ. (2014). Evidence-based
661 recommendations for natural bodybuilding contest

- 662 preparation: nutrition and supplementation. *Journal of the*
 663 *International Society of Sports Nutrition*, 12(11), 20.
 664
- 665 Hulmi, J. J., Isola, V., Suonpaa, M., Jarvinen, N. J., Kokkonen, M.,
 666 Wennerstrom, A., and Hakkinen, K. (2016). The Effects of
 667 Intensive Weight Reduction on Body Composition and Serum
 668 Hormones in Female Fitness Competitors. *Frontiers in*
 669 *Physiology*, 7, 689.
 670
- 671 IFBB. (2014). IFBB anti-doping area. Retrieved from
 672 <https://ifbb.com/wada-anti-doping-rules/>.
 673
- 674 Ismaeel, A., Weems, S., and Willoughby, D. (2017). A Comparison of
 675 the Nutrient Intakes of Macronutrient-based Dieting and Strict
 676 Dieting Bodybuilders. *International Journal of Sport*
 677 *Nutrition and Exercise Metabolism*, 28(4), 502-508.
 678
- 679
- 680 Kistler, B. M., Fitschen, P. J., Ranadive, S. M., Fernhall, B., and
 681 Wilund, K. R. (2014). Case study: Natural bodybuilding
 682 contest preparation. *International Journal of Sport Nutrition*
 683 *and Exercise Metabolism*, 24(6), 694-700.
 684
- 685 Kleiner, S., Bazzarre, TL., and Ainsworth, B. E. (1994). Nutritional
 686 status of nationally ranked elite bodybuilders. *International*
 687 *Journal of Sport Nutrition.*, 4(1), 54-69.
 688
- 689 Kleiner, S. M., Bazzarre, T. L., and Litchford, M. D. (1990). Metabolic
 690 profiles, diet, and health practices of championship male and
 691 female bodybuilders. *Journal of the American Dietetic*
 692 *Association*, 90(7), 962-967.
 693
- 694 Layne, J. E., and Nelson, M. E. (1999). The effects of progressive
 695 resistance training on bone density: a review. *Medicine and*
 696 *Science in Sports and Exercise*, 31(1), 25-30.
 697
- 698 Loucks, A. B., and Heath, E. M. (1994). Induction of low-T3
 699 syndrome in exercising women occurs at a threshold of energy
 700 availability. *American Journal of Physiology*, 266(3 Pt 2),
 701 817-823.
 702
- 703 Loucks, A. B., Kiens, B., and Wright, H. H. (2011). Energy availability
 704 in athletes. *Journal of Sports Sciences*,
 705 *29 Suppl 1*, S7-15.
 706
- 707 Manore, M. M. (2002). Dietary recommendations and athletic
 708 menstrual dysfunction. *Sports Medicine*, 32(14), 887-901.
 709
- 710 Mitchell, L., Hackett, D., Gifford, J., Estermann, F., and O'Connor, H.
 711 (2017). Do Bodybuilders Use Evidence-Based Nutrition
 712 Strategies to Manipulate Physique? *Sports*, 5(4), 76.
 713
- 714 Mitchell, L., Slater, G., Hackett, D., Johnson, N., and O'Connor, H.
 715 (2018). Physiological implications of preparing for a natural

- 716 male bodybuilding competition. *European Journal of Sport*
717 *Sciences*, 18(5), 619-629.
- 718
- 719 Mountjoy, M., Sundgot-Borgen, J., Burke, L., Carter, S., Constantini,
720 N., Lebrun, C., and Ljungqvist, A. (2014). The IOC consensus
721 statement: beyond the Female Athlete Triad—Relative
722 Energy Deficiency in Sport (RED-S). *British Journal of*
723 *Sports Medicine*, 48(7), 491-497.
- 724
- 725 Newton, L. E., Hunter, G., Bammon, M., and Roney, R. (1993).
726 Changes in Psychological State and Self-Reported Diet
727 During Various Phases of Training in Competitive
728 Bodybuilders. *The Journal of Strength and Conditioning*
729 *Research*, 7(3), 153-158.
- 730
- 731 Obel, C. (1996). Collapsing Gender in Competitive Bodybuilding:
732 Researching Contradictions and Ambiguity in Sport.
733 *International Review for the Sociology of Sport*, 31(2), 185-
734 202.
- 735
- 736 Parish, T., Bahgurst, T., and Turner, R. (2010). Becoming competitive
737 amateur bodybuilders: Identification of contributors.
738 *Psychology of Men and Masculinity*, 11(2), 152-159.
- 739
- 740 Petrizzo, J., DiMenna, F. J., Martins, K., Wygand, J., and Otto, R. M.
741 (2017). Case Study: The Effect of 32 Weeks of Figure-Contest
742 Preparation on a Self-Proclaimed Drug-free Female's Lean
743 Body and Bone Mass. *International Journal of Sport Nutrition*
744 *and Exercise Metabolism*, 27(6), 543-549.
- 745
- 746 Phillips, S. M. (2004). Protein requirements and supplementation in
747 strength sports. *Nutrition*, 20(7), 689-695.
- 748
- 749
- 750 Pope, H. G., Gruber, A. J., Choi, P., Olivardia, R., and Phillips, K. A.
751 (1997). Muscle Dysmorphia: An Underrecognized Form of
752 Body Dysmorphic Disorder. *Psychosomatics*, 38(6), 548-557.
- 753
- 754
- 755 Probert, A., Palmer, F., and Leberman, S. (2007). The Fine Line: An
756 insight into 'risky' practices of male and female competitive
757 bodybuilders. *Annals of Leisure Research*, 10(3-4), 272-290.
- 758
- 759 Reale, R., Slater, G., Cox, G. R., Dunican, I. C., and Burke, L. M.
760 (2018). The Effect of Water Loading on Acute Weight Loss
761 Following Fluid Restriction in Combat Sports Athletes.
762 *International Journal of Sport Nutrition and Exercise*
763 *Metabolism*, 28(6), 565-573.
- 764
- 765 Robinson, S. L., Lambeth-Mansell, A., Gillibrand, G., Smith-Ryan,
766 A., and Bannock, L. (2015). A nutrition and conditioning
767 intervention for natural bodybuilding contest preparation:
768 Case study. *Journal of the International Society of Sports*
769 *Nutrition*, 12(1), 20.
- 770

- 771 Rohrig, B. J., Pettitt, R.W., Pettitt, C.D., and Kanzenback, T.L. (2017).
 772 Psychophysiological tracking of a female physique competitor
 773 through competition preparation. *International Journal of*
 774 *Exercise Science*, 10(2), 301-311.
 775
- 776 Rossow, L. M., Fukuda, D. H., Fahs, C. A., Loenneke, J. P., and Stout,
 777 J. R. (2013). Natural bodybuilding competition preparation
 778 and recovery: a 12-month case study. *International Journal of*
 779 *Sports Physiology and Performance*, 8(5), 582-592.
 780
- 781 Rowbottom, M. (2017, December 1). IFBB competitors place their
 782 bets for World Fitness Championships. Retrieved from
 783 [https://www.insidethegames.biz/articles/1058593/ifbb-](https://www.insidethegames.biz/articles/1058593/ifbb-competitors-place-their-bets-for-world-fitness-championships)
 784 [competitors-place-their-bets-for-world-fitness-](https://www.insidethegames.biz/articles/1058593/ifbb-competitors-place-their-bets-for-world-fitness-championships)
 785 [championships](https://www.insidethegames.biz/articles/1058593/ifbb-competitors-place-their-bets-for-world-fitness-championships).
 786
- 787 Shephard, R. J. (1994). Electrolyte manipulation in female body-
 788 builders. *British Journal of Sports Medicine*, 28(1), 60-61.
 789
- 790 Slater, G., and Phillips, S. M. (2011). Nutrition guidelines for strength
 791 sports: sprinting, weightlifting, throwing events, and
 792 bodybuilding. *Journal of Sports Sciences*, 29 Suppl 1, S67-77.
 793
- 794 Steen, S. N. (1991). Precontest strategies of a male bodybuilder.
 795 *International Journal of Sport Nutrition*, 1(1), 69-78.
 796
- 797 Spendlove, J., Mitchell, L., Gifford, J., Hackett, D., Slater, G., Cobley,
 798 S., and O'Connor, H. (2015). Dietary Intake of Competitive
 799 Bodybuilders. *Sports Medicine*, 45(7), 1041-1063.
 800
- 801 Steele, I., Pope Jr, H., and Kanayama, G. Weightlifting. (2018). In I.D.
 802 Glick, S. Todd., and D. Kamis (Ed.), *The ISSP Manual of*
 803 *Sports Psychiatry* (pp. 91-116). New York: Routledge.
 804
- 805 Stiegler, P., and Cunliffe, A. (2006). The Role of Diet and Exercise for
 806 the Maintenance of Fat-Free Mass and Resting Metabolic Rate
 807 During Weight Loss. *Sports Medicine*, 36(3), 239-262.
 808
- 809 Sundgot-Borgen, J. (1994). Risk and trigger factors for the
 810 development of eating disorders in female elite athletes.
 811 *Medicine and Science in Sports and Exercise*, 26(4), 414-419.
 812
- 813 Sundgot-Borgen, J., and Garthe, I. (2011). Elite athletes in aesthetic
 814 and Olympic weight-class sports and the challenge of body
 815 weight and body compositions. *Journal of Sports Sciences*, 29
 816 *Suppl 1*, 101-114.
 817
- 818 Sundgot-Borgen, J., Meyer, N. L., Lohman, T. G., Ackland, T. R.,
 819 Maughan, R. J., Stewart, A. D., and Muller, W. (2013). How
 820 to minimise the health risks to athletes who compete in
 821 weight-sensitive sports review and position statement on
 822 behalf of the Ad Hoc Research Working Group on Body
 823 Composition, Health and Performance, under the auspices of
 824 the IOC Medical Commission. *British Journal of Sports*
 825 *Medicine*, 47(16), 1012-1022.

- 826
827 Sundgot-Borgen, J., and Torstveit, M. K. (2004). Prevalence of eating
828 disorders in elite athletes is higher than in the general
829 population. *Clinical Journal of Sport Medicine*, 14(1), 25-32
830
- 831 Tajrobehkar, B. (2016). Flirting With the Judges: Bikini Fitness
832 Competitors' Negotiations of Femininity in Bodybuilding
833 Competitions. *Sociology of Sport Journal*, 33(4), 294-304.
834
- 835 Tinsley, G. M., Trexler, E. T., Smith-Ryan, A. E., Paoli, A., Graybeal,
836 A. J., Campbell, B. I., and Schoenfeld, B. J. (2018). Changes
837 in Body Composition and Neuromuscular Performance
838 Through Preparation, 2 Competitions, and a Recovery Period
839 in an Experienced Female Physique Athlete. *The Journal of*
840 *Strength and Conditioning Research, Publish Ahead of Print.*
841
- 842 Trexler, E. T., Hirsch, K. R., Campbell, B. I., and Smith-Ryan, A. E.
843 (2017). Physiological Changes Following Competition in
844 Male and Female Physique Athletes: A Pilot Study.
845 *International Journal of Sport Nutrition and Exercise*
846 *Metabolism*, 27(5), 458-466.
847
- 848 Trexler, E. T., Smith-Ryan, A. E., and Norton, L. E. (2014). Metabolic
849 adaptation to weight loss: implications for the athlete. *Journal*
850 *of the International Society of Sports Nutrition*, 11(1), 7.
851
- 852 Van der Ploeg, G. E., Brooks, A. G., Withers, R. T., Dollman, J.,
853 Leaney, F., and Chatterton, B. E. (2001). Body composition
854 changes in female bodybuilders during preparation for
855 competition. *European Journal of Clinical Nutrition*, 55(4),
856 268-277.
857
- 858 Vallet, G. (2017). The gendered economics of bodybuilding.
859 *International Review of Sociology*, 27(3), 525-545.
860
- 861 Walberg-Rankin, J. E. C., and Gwazdauskas, F. C. (1993). Diet and
862 weight changes of female bodybuilders before and after
863 competition. *International Journal of Sport Nutrition.*, 3(1),
864 87-102.
865
- 866 Walberg, J. L., and Johnston, C. S. (1991). Menstrual function and
867 eating behavior in female recreational weight lifters and
868 competitive body builders. *Medicine and Science in Sports*
869 *and Exercise*, 23(1), 30-36.
870
- 871 West, R. V. (1998). The Female Athlete. *Sports Medicine*, 26(2), 63-
872 71.
873
- 874 Zanker, C. L., Osborne, C. Cooke, C. B., Oldroyd, B., and Trustcott,
875 J. G. (2004). Bone density, body composition and menstrual
876 history of sedentary female former gymnastics, aged 20-32
877 years *Osteoporosis International*, 15(2), 145-154.
878
879
880

881

882

883

884

885

886

887

888

889

890

Table 1: Overview of the recent studies of reproductive health of female physique athletes.

Study	N	Body weight change (Body Fat %)		Time period (weeks)	TEST		E ₂		T ₃		T ₄		CORT		Ghrelin		LP		TSH		IN		Method for menstrual status	Absence of menstruation	Bone mass density (DXA)
		C P	R C		C P	R C	C P	R C	C P	R C	C P	R C	C P	R C	C P	R C	C P	R C							
Haliday et al. 2016	1 ♀	-8.3kg; (15.1- 8.6%)	+5.2kg; (8.6- 14.8%)	20 CP; 20 RC	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Self-report	9 weeks pre- and up to 71 weeks post- competition	NA
Hulmi et al. 2016	27 ♀	-7.8kg (23.1- 12.7%)	+6.1kg (12.7- 20.1%)	20 CP; 17.5 RC	↓	(↑)	↓	↑	↓	(↑)	↓	↑	-	-	-	-	↓	↑	↓	↑	-	-	Serum and self-report	11.5% pre- competition and 28% post- competition	↓CP; ↑RC
Trexler al. 2017	8 ♀ 7 ♂	-	+3.9kg (12.5- 14.9%)	4-6 RC	-	↑	-	-	-	-	-	-	↑↓	↑	↓	-	-	-	-	-	↑	Saliva	-	-	
Petrizzo et al. 2017	1 ♀	-7.7kg (24.4- 11.3%)	-	24 CP	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Self-report	Oligomenorrhea	No change
Rohrig et al. 2017	1 ♀	-10.1kg (30.5- 15.9%)	-	24 CP	↑↓	-	↑↓	-	-	-	-	-	↑↓	-	-	-	↓	-	↓	-	-	-	Serum and self-report	8 weeks pre- competition	-
Tinsley et al., 2018	1 ♀	-6 kg (20.3- 11.6%)	+6.8kg (11.6- 18.8%)	18 CP (1) 7 CP (2) 9 RC	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Self-report	12 weeks pre- competition (1) and up to 12 weeks post- competition (2)	NA

♀ indicates female physique athletes, ♂ indicates male physique athletes, ↑↓ indicates fluctuation, CP indicates the pre-competition phase, RC indicates recovery phase, () indicates not recovered to initial baseline values, (1) indicates 1st competition and (2) indicates a 2nd competition. TEST = Testosterone, E2 = Estradiol, T₃ = Triiodothyronine, T₄ = Thyroxine, CORT = Cortisol; TSH= Thyroid stimulating hormone; LP= Leptin, IN = Insulin; DXA = Dual-energy X-ray absorptiometry. NA = Information not available.

Figure 1: An overview of the current female categories in women's physique competitions. The categories are progressive steps along a continuum between lean body mass and fat mass. 'Dry' refers to dehydration and the subsequent reduction in body water (Chappell et al., 2018). The number of height classes in each category is determined by the popularity of the single category. This figure was drawn using information retrieved from the International Federation of Bodybuilding and Fitness website (FBB Elite Pro Categories, 2017).