



LJMU Research Online

Stables, RG, Kasper, AM, Sparks, SA, Morton, JP and Close, GL

An Assessment of the Validity of the Remote Food Photography Method (Termed Snap-N-Send) in Experienced and Inexperienced Sport Nutritionists

<http://researchonline.ljmu.ac.uk/id/eprint/16728/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Stables, RG, Kasper, AM, Sparks, SA, Morton, JP and Close, GL (2021) An Assessment of the Validity of the Remote Food Photography Method (Termed Snap-N-Send) in Experienced and Inexperienced Sport Nutritionists. International Journal of Sport Nutrition and Exercise

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

**INTERNATIONAL JOURNAL OF
SPORT NUTRITION AND
EXERCISE METABOLISM**

**An assessment of the validity of the remote food
photography method (termed Snap-N-Send) in experienced
and inexperienced sport nutritionists**

Journal:	<i>International Journal of Sport Nutrition & Exercise Metabolism</i>
Manuscript ID	IJSNEM.2020-0216.R2
Manuscript Type:	Original Research
Keywords:	dietary intake, RED-S, LEA, exercise

SCHOLARONE™
Manuscripts

1 **ABSTRACT**

2 The remote food photography method (RFPM), often referred to
3 as ‘Snap-N-Send’ by sport nutritionists, has been reported as a
4 valid method to assess energy intake in athletic populations.
5 However, preliminary studies were not conducted in true free-
6 living conditions and dietary assessment was performed by one
7 researcher only. We therefore assessed the validity of ‘Snap-N-
8 Send’ to assess energy and macronutrient composition in
9 experienced (EXP, n=23) and inexperienced (INEXP, n=25)
10 sport nutritionists. Participants analysed two days of dietary
11 photographs, comprising eight meals. Day 1 consisted of
12 ‘simple’ meals based around easily distinguishable foods (i.e.
13 chicken breast and rice) and Day 2, ‘complex’ meals containing
14 ‘hidden’ ingredients (i.e. chicken curry). Estimates of dietary
15 intake were analysed for validity using one-sample t-tests and
16 typical error of estimates (TEE). INEXP and EXP nutritionists
17 underestimated energy intake for the simple day (Mean
18 difference, MD = -1.5 MJ, TEE = 10.1%; -1.2 MJ, TEE = 9.3%
19 respectively) and the complex day (MD = -1.2 MJ, TEE =
20 17.8%; MD = -0.6 MJ, 14.3% respectively). Carbohydrate intake
21 was underestimated by INEXP (MD = -65.5 g.day⁻¹, TEE =
22 10.8% and MD = -28.7 g.day⁻¹, TEE = 24.4%) and EXP (MD
23 = -53.4 g.day⁻¹, TEE = 10.1% and -19.9 g.day⁻¹, TEE = 17.5%)
24 for both simple and complex days, respectively. The inter-
25 practitioner reliability was generally ‘poor’ for energy and

Remote Food Photography Method in Sport Nutrition 2

26 macro-nutrients. Data demonstrate that the RFPM / ‘Snap-N-
27 Send’ under-estimates energy intake in simple and complex
28 meals and these errors are evident in experienced and
29 inexperienced sport nutritionists.

30

31 Key words: dietary intake, exercise, RED-S, LEA

Remote Food Photography Method in Sport Nutrition 3

32 **INTRODUCTION**

33 A fundamental activity for sport nutritionists is to estimate
34 energy and macronutrient intake from an athlete's self-reported
35 food intake (Braakhuis et al., 2003). Such dietary assessments
36 are important given the role of energy and macronutrient intake
37 in modulating training adaptation (Impey et al., 2018), body
38 composition (Kasper et al., 2018; Morton et al., 2010; Wilson et
39 al., 2015) and exercise performance (Burke & Hawley 2018).
40 Additionally, nutrient availability can also play a fundamental
41 role in growth and maturation (Hannon et al., 2020), mental
42 health (Wilson et al., 2014) and reducing the risk of illness and
43 injury (Kasper et al., 2018; Walsh, 2019; Wilson et al., 2014).
44 Despite the clear rationale to accurately assess an athlete's
45 energy intake, this remains a major methodological challenge
46 that is fraught with sources of error on both the athlete's and
47 sport nutritionist's part (Capling et al., 2017).

48

49 Broadly speaking, dietary assessment methods are classified as
50 'retrospective' (including 24-hour recall, food frequency
51 questionnaires, diet histories) or 'prospective' (including food
52 diaries with / without weighed inventory). Inaccuracies are
53 inherent with self-reported dietary assessments and include the
54 misreporting of food consumption alongside measurement error
55 (Gemming et al., 2014; Rollo et al., 2016; Westerterp et al.,
56 1986). Furthermore, most of the dietary assessment methods are

Remote Food Photography Method in Sport Nutrition 4

57 logistically complicated, especially when assessing multiple
58 athletes (e.g. sports teams) in free living conditions (Martin et
59 al., 2012). Validity and precision, in addition to practitioner and
60 participant burden, are cited as some of the main causes of
61 inaccuracies in dietary assessment (Livingstone & Black, 2003;
62 Thompson et al., 2010). In addition to the bias associated with
63 participant burden and self-reporting, the requirement of
64 accurate unbiased interpretation by a nutritionist or dietitian has
65 led to the criticism within the sports nutrition community that
66 systematic error in dietary analysis is neglected and somewhat
67 overlooked (Kirkpatrick & Collins, 2016).

68

69 In an attempt to improve participant reporting accuracy in
70 traditional pen and paper methods, Martin et al. (2009)
71 developed the remote food photograph method (RFPM)
72 whereby participants record dietary intake in real time via
73 ecological momentary assessment. In this approach, participants
74 take and transmit photographs (via camera enabled cell phones
75 with data transfer capability) of food selection and plate waste to
76 researchers for subsequent dietary analysis. In combining the
77 principles of the RFPM with elements of behavioural change
78 science to engage participants and all key stakeholders, Costello
79 et al. (2017) subsequently developed the 'Snap-N-Send'
80 methodology demonstrating that an athletic population was also
81 capable of adhering to self-reporting of dietary intake via smart

Remote Food Photography Method in Sport Nutrition 5

82 phone technology. However, whilst this preliminary study
83 concluded that ‘Snap-N-Send’ was valid and reliable as a
84 standalone dietary assessment method, there are several
85 limitations that should be noted. First, the experimental
86 conditions were not true free-living, given that participants were
87 restricted to consuming foodstuffs that were provided by the
88 researchers during the study period. In this way, the researcher
89 had prior knowledge of approximate portion sizes and
90 macronutrient profile of the foods consumed given that foods
91 were weighed by the research team before being distributed to
92 the participants. Second, the subsequent dietary analysis was
93 performed by one researcher only, an important methodological
94 factor considering the inherent variability that exists between
95 experienced sports dietitians when coding food records for
96 analysis (Braakhuis et al., 2003). Thus, the aim of the present
97 study was to assess the validity of utilising the RFPM / ‘Snap-
98 N-Send’ as a standalone method to assess energy and
99 macronutrient composition in experienced and inexperienced
100 sport nutritionists.

101

102

103 **METHODS**104 ***Participants***

105 Forty-eight participants were recruited to take part in this study.

106 Participants were non-randomly allocated to two independent

Remote Food Photography Method in Sport Nutrition 6

107 groups based upon the inclusion criteria: 1) Recent Sport and
108 Exercise Nutrition register (SENr) graduates with graduate
109 accreditation status (n=25) [termed INEXPERIENCED]; or 2)
110 Full SENr practitioner registrants with >3 years working within
111 elite sport (n=23) [termed EXPERIENCED]. All of the
112 'inexperienced' sport nutritionists had received recent training
113 in dietary assessment (including the RFPM) from experienced
114 sport nutritionists whilst all of the 'experienced' sport
115 nutritionists, as a criteria of their SENr registration, will have
116 demonstrated evidence of competency in dietary assessment.
117 This study was approved by the university ethics committee
118 (M20_SPS_767) and was conducted in accordance to the
119 Declaration of Helsinki.

120

121 ***Study Design***

122 Participants were provided with the same two days of dietary
123 images comprising of a total of eight meals (breakfast, morning
124 snack, lunch and evening meals). These foods, photographed
125 remotely, had been compiled by the research team with one day
126 being classed as 'simple' meals and the second day being
127 'complex' meals with the two days being similar in total energy
128 content. Dietary images and short descriptions were then sent to
129 each participant via email or over a free cellular picture
130 messaging smartphone application (WhatsApp Inc., California,
131 USA) for analysis. Participants were asked to analyse each meal

Remote Food Photography Method in Sport Nutrition 7

132 for its calorific and macronutrient content using Nutritics dietary
133 analysis software using the pre-set UK/Ireland database
134 (Nutritics version 5.5, Swords, Ireland) and return these data
135 files to the primary researcher to assess the ability of experienced
136 and inexperienced practitioners to estimate energy intake in
137 comparison to food labels.

138

139 ***Control***

140 To standardise perceived portion size, all meals were placed on
141 the same plate or bowl with cutlery on a 1 x 1 cm A3 reference
142 grid placemat as previously described (Costello et al., 2017). All
143 images were taken by the researcher at a height of sixty
144 centimetres at a ninety-degree angle. Images were later cropped
145 so that the reference grid filled the image (15.01 cm x 21.34 cm)
146 and added to a standard PowerPoint slide (19.05 cm x 25.4 cm)
147 with a brief description of the food in the image (e.g. Weetabix
148 cereal made with semi-skimmed milk).

149

150 ***Meal Design***

151 Day one of the diet diary was designed in a simplistic manner
152 whereby each individual food item could be easily identified and
153 distinguished by the participant, e.g. chicken breast and rice
154 [termed SIMPLE]. In this day, no extras were added to meals
155 such as butter on potatoes or condiments such as mayonnaise.
156 The second day was designed to contain a number of complex

Remote Food Photography Method in Sport Nutrition 8

157 meals whereby it was more difficult to ascertain a number of
158 individual ingredients and definite quantities of each food item,
159 e.g. chicken curry and rice [termed COMPLEX]. Again, no
160 hidden extras were added. For the purpose of this study, it was
161 presumed that all foods on the plate were consumed with no need
162 to attempt to calculate the left-over food items. An overview of
163 the meals and energy content can be found in Figure 1.

164

165 ***Statistical Analysis***

166 Data were assessed for normality using standard graphical
167 procedures and Shapiro-Wilk tests. Values of minimally
168 clinically important difference (MCID) have not been used in
169 this study because the use of hard anchors cannot be universally
170 applied for each variable in multiple scenarios (Cook et al.,
171 2014). For example, in an acute nutritional intervention,
172 differences in energy intake of $0.5 \text{ MJ}\cdot\text{day}^{-1}$ would have little
173 effect but would likely be clinically important in a chronic
174 setting. Likewise, a small change in nutrient content of diets that
175 have very low total energy may be important, but in an athlete
176 with much higher energy needs and intake, it will not be.
177 Therefore, the effect sizes of Cohen's d (for t-tests) and r -values
178 (for Wilcoxon signed rank tests) were used to help to determine
179 the magnitude of potential differences. These effect sizes were
180 interpreted as small, medium and large using the values of 0.2,
181 0.5, 0.8; and $0.1 < 0.4$, $0.4 < 0.6$, ≥ 0.6 for d and r respectively.

Remote Food Photography Method in Sport Nutrition 9

182

183 Consequently, differences between the actual nutrient data (as
184 obtained from food labels), the estimated energy intake, the
185 macronutrient content of the simple and complex days, and
186 individual meals and daily snacks, were assessed using one
187 sample t-tests or Wilcoxon signed rank tests where difference
188 data were non-parametric. Differences in the observed dietary
189 analysis data between the inexperienced and experienced groups
190 were assessed using independent t-tests for the energy and
191 macronutrient content of both the simple and complex days.

192

193 The validity of the observed data compared to the known
194 nutrient values was assessed using coefficient of variation (CV)
195 along with 95% limits of agreement (LoA), bias and 95%
196 confidence intervals (CI). Coefficient of variation was
197 interpreted using the following thresholds: <2% (excellent),
198 <5% (good), <10% (acceptable), >10% (poor), >20% (very
199 poor). Inter-rater reliability (termed inter-practitioner reliability
200 hereafter) was assessed using a two-way mixed effects model for
201 Cronbach's alpha, intra-class correlations (ICC) with 95% CI
202 and CV. All inferential statistical tests and validity calculations
203 were conducted using SPSS (v25 for Windows, Illinois, USA)
204 MS Excel (365 for Windows, Washington, USA) respectively.

205

206 **RESULTS**

Remote Food Photography Method in Sport Nutrition 1

207 *Estimated Dietary Intake*

208 The inexperienced, experienced, and whole sample
209 underestimated energy intake (Figure 2A and Table 2) for the
210 simple day (MD = -1.7 MJ, $w = 10.0$, $z = 4.1$, $p < 0.001$, $r = 0.58$;
211 MD = -1.2 MJ, $p < 0.001$, CI = -1.56, -0.81, $d = 1.36$ and MD =
212 -1.4 MJ, $p < 0.001$, CI = -64, -1.10, $d = 1.50$; respectively) and
213 the complex day (MD = -1.2 MJ, $p = 0.001$, CI = -1.80, -0.54, d
214 = 0.76; MD = -1.5 MJ, $w = 1140$, $z = 5.7$, $p < 0.001$, $r = 0.58$;
215 and, MD = -0.9 MJ, $p < 0.001$, CI = -1.32, -0.50, $d = 0.65$;
216 respectively). The estimated energy intake values were not
217 different between the groups for either the simple (MD = 0.35
218 MJ, $p = 0.186$, CI = -0.88, 0.18, $d = 0.59$) or complex days (MD
219 = $p = 0.185$, CI = -1.35, 0.27, $d = 0.39$).

220

221 Estimated carbohydrate (CHO) intake (Figure 2B) was
222 underestimated by the inexperienced (MD = -67.5 g, $w = 324.0$,
223 $z = 4.4$, $p < 0.001$, $r = 0.62$; and, MD = -26.9 g, $w = 217.0$, $z =$
224 2.4, $p = 0.016$, $r = 0.35$), the experienced (MD = -53.4 g, $p <$
225 0.001, CI = -62.7, -44.0, $d = 2.73$ and, MD = -64.2 g, $w = 1174$,
226 $z = 6.0$, $p < 0.001$, $r = 0.61$) and whole sample (MD = -62.3 g, p
227 < 0.001 , CI = -68.8, -55.8, $d = 2.79$; and, MD = -24.5 g, $p <$
228 0.001, CI = -37.3, -11.64, $d = 0.55$) for both the simple and
229 complex days respectively. There were again no differences in
230 the carbohydrate estimates between the groups for either the

Remote Food Photography Method in Sport Nutrition 1

231 simple (MD = 6.7 g, $p = 0.308$, CI = -19.6, 6.3, $d = 0.30$) or
232 complex (MD = 8.8 g, $p = 0.493$, CI = -34.7, 17.0, $d = 0.20$) days.

233

234 Estimates of fat intake (Figure 2D) made by the inexperienced
235 group were lower than the actual fat content of the simple day
236 (MD = -6.7 g, $w = 257.0$, $z = 2.5$, $p = 0.011$, $r = 0.36$), but this
237 was not the case for the experienced group (MD = -3.6g, $p =$
238 0.173 , CI = -8.8, 1.7, $d = 0.29$, respectively), and there were no
239 differences between the fat intake estimates of the two groups
240 combined (MD = -4.2 g, $p = 0.331$, CI = -12.9, 4.4, $d = 0.24$).
241 However, when two groups were combined for the whole
242 sample, fat intake was under-estimated by a small amount (MD
243 = -5.8 g, $p = 0.010$, CI = -10.1, -1.48, $d = 0.39$).

244

245 Fat intake estimates for the complex day were not different from
246 the actual value for either the inexperienced (MD = 5.38 g, $p =$
247 0.059 , CI = -10.9, 0.22, $d = 0.39$), experienced (MD = 3.95 g, $p =$
248 0.183 , CI = -2.0, 9.9, $d = 0.29$), or whole sample (MD = -1.0
249 g, $p = 0.630$, CI = -5.2, 3.2, $d = 0.08$). However, the
250 inexperienced group estimated fat intake to be lower than that of
251 the experienced group for the complex day (MD = -9.3 g, $p =$
252 0.023 , CI = -17.3, -1.4, $d = 0.69$).

253

254 The estimations of protein intake were not different between the
255 two groups (Figure 2C), for either the simple or complex days

Remote Food Photography Method in Sport Nutrition 1

256 (MD = 4.1 g, $p = 0.482$, CI = -15.8, 7.6, $d = 0.14$; and (MD = 2.4
257 g, $p = 0.791$, CI = -19.9, 15.2, $d = 0.13$, respectively).
258 Interestingly, the experienced group estimated protein intake to
259 be higher than the actual value for the simple day (MD = 10.1 g,
260 $p = 0.027$, CI = 2.1, 16.7, $d = 0.50$), but the inexperienced group
261 did not (MD, 5.4 g, $p = 0.070$, CI = -2.2, 14.1, $d = 0.38$). When
262 the whole sample was combined for the simple day, protein
263 intake was estimated to be higher than the actual value (MD =
264 7.9 g, $p = 0.009$, CI = 2.1, 13.7, $d = 0.44$). Conversely, for the
265 complex day protein intake estimates were lower than the actual
266 values for the inexperienced (MD = -18.0 g, $p = 0.011$, CI = -
267 31.5, -4.6, $d = 0.51$), experienced (MD = -15.7 g, $p = 0.012$, CI
268 = -27.7, -3.7, $d = 0.57$) and whole sample (MD = -16.9 g, $p <$
269 0.001 , CI = -25.7, -8.2, $d = 0.54$).

270

271 **Meal by Meal Estimates**

272 The complex day breakfast (figure 3A1-4) was underestimated
273 for energy (MD = -0.63 MJ, $p < 0.001$, CI = -0.82, -0.45, $d =$
274 1.40, and MD = -0.50 MJ, $p < 0.001$, CI = -0.67, -0.34, $d = 1.28$)
275 CHO (MD = -11.5 g, $w = 325.0$, $z = 4.4$, $p < 0.001$, $r = 0.62$, and
276 MD = -11.5 g, $w = 276.0$, $z = 4.2$, $p < 0.001$, $r = 0.62$), and protein
277 (MD = -22.1 g, $p < 0.001$, CI = -24.45, 1-.79, $d = 3.90$, and MD
278 = -18.5 g, $w = 276.0$, $z = 4.2$, $p < 0.001$, $r = 0.62$) by the
279 inexperienced and experienced groups. Notably the
280 inexperienced group also underestimated the energy (MD = -

Remote Food Photography Method in Sport Nutrition 1

281 0.18 MJ, $p = 0.005$, $w = 267.0$, $z = 2.8$, $r = 0.40$), protein (MD =
282 -3.5 g, $w = 240.0$, $z = 3.7$, $p < 0.001$, $r = 0.52$) and fat content
283 (MD = -1.5 g, $w = 236.0$, $z = 3.7$, $p < 0.001$, $r = 0.51$) of the
284 simple breakfast but this was not the case for the experienced
285 group.

286

287 Typically, the simple snack energy (MD = -0.80 MJ, $w = 324.0$,
288 $z = 4.4$, $p < 0.001$, $r = 0.62$, and 0.96 MJ, $p < 0.001$, CI = -1.11 ,
289 -0.81 , $d = 2.74$), CHO (MD = -12.6 g, $w = 324.0$, $z = 4.4$, $p <$
290 0.001 , $r = 0.62$, and MD = -12.9 g, $w = 254.0$, $z = 3.5$, $p < 0.001$,
291 $r = 0.52$) and fat (MD = 14.6 g, $w = 313.0$, $z = 4.1$, $p < 0.001$, r
292 = 0.57 , and MD = -15.8 g, $w = 276.0$, $z = 4.2$, $p < 0.001$, $r = 0.62$)
293 content was underestimated by the inexperienced and
294 experienced groups (figure 3B1-4). Conversely the
295 inexperienced and experienced groups overestimated energy
296 (MD = 0.29 MJ, $p = 0.001$, CI = $0.13-0.44$, $d = 0.76$, and MD =
297 0.34 MJ, $w = 234.0$, $z = 2.9$, $p = 0.004$, $r = 0.43$), protein (MD =
298 7.9 g, $w = 295$, $z = 3.6$, $p < 0.001$, $r = 0.50$, and MD = 8.0 g, w
299 = 228.0 , $z = 2.7$, $p = 0.006$, $r = 0.40$) and fat (MD = 4.3 g, $w =$
300 324.0 , $z = 4.3$, $p < 0.001$, $r = 0.62$, and MD = 4.4 g, $w = 272.0$, z
301 = 4.1 , $p < 0.001$, $r = 0.60$) for the complex snacks.

302

303 For the lunch meal, CHO content was underestimated by the
304 inexperienced (MD = 10.2 g, $w = 290.0$, $z = 3.4$, $p < 0.001$, $r =$
305 0.49 and MD = -20.1 g, $p < 0.001$, CI = -28.9 , -11.4 , $d = 0.95$)

Remote Food Photography Method in Sport Nutrition 1

306 and experienced (MD = 7.9 g, $p = 0.001$, CI = 12.4, -3.4, $d =$
307 0.76 and MD = 16.1 g, $p < 0.001$, CI = -23.6, -8.6, $d = 0.93$)
308 groups for both the simple and complex days respectively (figure
309 3 C1-4). The protein and fat content of the simple lunch were
310 overestimated by the inexperienced (MD = 5.2 g, $w = 253.0$, $z =$
311 2.4, $p = 0.015$, $r = 0.35$ and MD = 11.5 g, $w = 307.0$, $z = 3.9$, p
312 < 0.001 , $r = 0.55$) and experienced (MD = 6.2 g, $w = 222.0$, $z =$
313 2.6, $p = 0.011$, $r = 0.38$, and MD = 21.1 g, $w = 271.0$, $z = 4.0$, p
314 < 0.001 , $r = 0.60$) groups, whereas the fat (MD = 4.3 g $w = 324.0$,
315 $z = 4.3$, $p < 0.001$, $r = 0.62$ and MD = 7.1 g, $w = 248.0$, $z = 3.4$,
316 $p < 0.001$, $r = 0.49$) and energy content (MD = -0.8 MJ, $p <$
317 0.001, CI = -1.1, -0.5, $d = 1.21$ and MD = -0.6 MJ, $p < 0.001$, CI
318 = -0.8, -0.4, $d = 1.25$) of the complex lunch were underestimated
319 by the inexperienced and experienced groups, respectively.

320

321 The energy (MD = 0.15 MJ, $p = 0.024$, CI = 0.02, 0.28, $d = 0.48$,
322 and MD = 0.71 MJ, $w = 271.0$, $z = 4.1$, $p < 0.001$, $r = 0.60$), CHO
323 (MD = 46.9 g, $w = 325.0$, $z = 4.4$, $p < 0.001$, $r = 0.62$, and MD
324 = 45.9 g, $w = 276.0$, $z = 4.2$, $p < 0.001$, $r = 0.62$) and protein
325 content (MD = 5.0 g, $p = 0.004$, CI = 1.8, 8.1, $d = 0.64$, and MD
326 = 3.0 g, $w = 230.0$, $z = 2.8$, $p = 0.005$, $r = 0.41$) of the simple
327 evening meal (figure 3 D1-4) were overestimated, by the
328 inexperience and experienced groups respectively. Additionally,
329 the experienced group also overestimated the fat content for the
330 simple (MD = 4.5 g, $w = 256.0$, $z = 3.6$, $p < 0.001$, $r = 0.53$) and

Remote Food Photography Method in Sport Nutrition 1

331 the complex evening meal (MD = 18.6 g, w = 227.0, z = 2.7, p
332 = 0.006, r = 0.40).

333

334 *Assessment of Inter-Practitioner Reliability*

335 The inter-practitioner reliability (Table 2 and Figure 2) was
336 generally poor for the estimation of energy and nutrient intake.
337 Specifically, the only acceptable inter-practitioner reliability
338 was observed for the simple dietary intake day in both groups of
339 practitioners, and the sample as a whole. All of the complex
340 dietary intake day analysis resulted in poor or very poor inter-
341 practitioner reliability. The inexperienced group appeared to
342 have worse inter-practitioner reliability than their more
343 experienced counterparts, but even the experienced practitioners
344 displayed poor inter-practitioner reliability for energy intake and
345 carbohydrate, and very poor reliability for fat and protein
346 estimates. Furthermore, very poor inter-practitioner reliability
347 was observed in both groups, and the sample as a whole, for
348 estimates of fat and protein intake, with the exception of the
349 experienced group's estimate of fat in the simple day, which was
350 still poor.

351

352 **DISCUSSION**

353 The aim of the present study was to assess the validity of utilising
354 the RFPM / 'Snap-N-Send' as a standalone methodology to
355 assess energy and macronutrient composition. To this end, we

Remote Food Photography Method in Sport Nutrition 1

356 recruited 49 accredited sport nutritionists to analyse two days of
357 dietary images comprising four ‘simple’ meals or four ‘complex’
358 meals. We report that RFPM / ‘Snap-N-Send’ method has ‘poor’
359 validity compared with the known values for both total energy
360 intake and macronutrient composition. Additionally, the inter-
361 practitioner reliability was qualified as ‘poor’, even between the
362 experienced sport nutritionists. Taken together, our data provide
363 a reference point for practitioners when considering the typical
364 error associated with these methods of dietary assessment.

365

366 The design of the present study allowed for 24 different
367 assessments of validity (energy, carbohydrate, fat and protein; in
368 complex and simple days; by experienced, inexperienced,
369 combined nutritionists; 4x2x3). We report that only 8/24 of the
370 assessments were qualified as ‘adequate’ with the remaining
371 16/24 categorised as ‘poor’ or ‘very poor’. Moreover, no
372 assessments of validity classed as ‘good’ or ‘very good’. Overall,
373 the RFPM / ‘Snap-N-Send’ method significantly underreported
374 total energy content by 13% which is in line with previous
375 research who have reported 8.8%, 11.3% and 13.1% respectively
376 (Martin et al., 2012; Kikunga et al., 2007; Lassen et al., 2010).
377 More importantly, however, was the extreme variation observed
378 in the reporting of energy intake which ranged from -47% to
379 +18%. Indeed, ‘acceptable’ validity for energy intake was only
380 seen in the simple day when analysed by experienced

Remote Food Photography Method in Sport Nutrition 1

381 practitioners and this still resulted in a TEE of -9.3%. These data
382 are in contrast to the preliminary report assessing the validity of
383 the ‘Snap-N-Send’ methodology where variability was reported
384 as acceptable (<5%, Costello et al., 2017). It is noteworthy,
385 however, that these researchers combined digital photography
386 alongside a written food diary and all food items were weighed
387 by the researcher team pre- and post-consumption. This contrasts
388 with the present methodology where the individuals who
389 performed the dietary assessments had no prior knowledge of the
390 food being provided or portion sizes. As such, the data presented
391 herein likely represent a more ecologically valid assessment
392 scenario in which both practitioners and researchers are likely to
393 engage in dietary assessment activities. Indeed, in a further study
394 from Costello et al. (2019), the researchers compared ‘Snap-N-
395 Send’ derived estimates of energy intake obtained from free
396 living conditions (i.e. participants consumed their own food
397 choices with no prior researcher knowledge) with energy
398 expenditure (using doubly labelled water) and reported large
399 random error and reduced measurement accuracy at an
400 individual level. In this instance, the authors suggested that the
401 poor performance of ‘Snap-N-Send’ was a consequence of low
402 athlete adherence to submitting all of the food consumed.
403 However, when considered with the present data, we suggest that
404 it is likely due in part to the inability of practitioners to correctly
405 identify foods and quantities from dietary photographs. Indeed,

Remote Food Photography Method in Sport Nutrition 1

406 the limitation of using only one coder when performing dietary
407 assessments is an important methodological factor considering
408 the inherent variability that exists between experienced sports
409 dieticians when coding food records for analysis (Braakhius et
410 al., 2003). Our data could also suggest that the RFPM / Snap-N-
411 Send, requires a high level of specialist and specific training
412 prior to use in order to yield reliable data. We therefore suggest
413 that in free living conditions, practitioners should take into
414 consideration the limitations of this approach and interpret the
415 data accordingly.

416

417 In addition to total energy intake, we also provide the first report
418 of sport nutritionists using the RFPM / 'Snap-N-Send'
419 methodology to assess the validity of analysing macronutrient
420 composition. The validity of carbohydrate intake was 'poor' or
421 'very poor' in the experienced and inexperienced practitioners in
422 both the simple and complex days with the range being as much
423 as 75g-329g on one day. This 'poor' validity of carbohydrate
424 intake is of particular concern given the majority of the meals,
425 even on the complex day, used easily recognised carbohydrate
426 sources such as potatoes. Many sport nutritionists now look to
427 periodise carbohydrate intake based on the training of the athlete
428 utilising the 'fuel for the work required' concept (Impey et al.,
429 2018). The inability to accurately identify the amount of
430 carbohydrate from dietary photographs (even on simple days by

Remote Food Photography Method in Sport Nutrition 1

431 experienced practitioners) suggests that practitioners must be
432 cautious with regards to making carbohydrate alterations to their
433 athletes diets based purely upon pictures sent from their athletes.
434 Protein intake was 'acceptable' with both inexperienced and
435 experienced practitioners on the simple day however was 'poor'
436 on the complex day ranging from 68-203 g. On the simple day,
437 protein was easily identified with portion sizes easy to estimate
438 through using foods such as poached eggs. However, on the
439 more complex day, protein was in the form of scrambled eggs, a
440 food harder to quantify via images alone. It is therefore crucial
441 that in free living conditions practitioners are aware that
442 significant error may exist in protein intake estimated from
443 complex meals and advice should be tailored accordingly.
444 Interestingly the most valid macronutrient estimate was for fat
445 which was 'acceptable' in the experienced practitioners on both
446 the simple and complex days. This may be due to the food
447 choices being low fat meals, typically eaten by athletes, and
448 future studies may wish to assess this observation in meals with
449 a higher fat content.

450

451 In addition to quantifying total daily energy and macronutrient
452 composition, we also performed analysis on a meal-by-meal
453 basis. From a practical perspective, such analysis is highly
454 important given that nutritional periodisation is performed on a
455 meal-by-meal basis. In this regard, our data demonstrate

Remote Food Photography Method in Sport Nutrition 2

456 extreme variability on a meal-by-meal basis with no consistent
457 pattern of error in terms of the experience of practitioners,
458 complexity, or type of meals. It did appear that the snacks were
459 a particular problem with the complex snacks being over
460 estimated for both energy and protein intakes in experienced as
461 well as inexperienced practitioners. Given the high-reliance on
462 snacks by athletes to achieve total caloric intakes, as well as to
463 achieve suggested protein distribution (Areta et al., 2013) this
464 over estimation of energy and protein could be a particular
465 problem in athletic groups who often consume 3-4 snacks per
466 day.

467

468 The present study also assessed the inter-practitioner reliability
469 of RFPM / 'Snap-N-Send' in both the experienced and
470 inexperienced sport nutritionists on the complex and simple
471 days. With regards to the total energy intake, despite 'poor'
472 validity, there was 'acceptable' reliability in both the
473 inexperienced and experienced nutritionists on the simple food
474 day, however this became 'poor' on the complex food day.
475 Indeed, a CV of 20.2% and 15.4%, along with very low ICC's
476 was reported on the complex day for the inexperienced and
477 experienced nutritionists respectively. This pattern was also
478 observed for carbohydrate intakes. Taken together these data
479 suggest that when assessing anything apart from simple meals
480 that are atypical of many athletes in free living conditions, the

Remote Food Photography Method in Sport Nutrition 2

481 RFPM / ‘Snap-N-Send’ methodology lacks inter-practitioner
482 reliability even in experienced nutritionists. Given the lack of
483 differences reported between the experienced and inexperienced
484 sport nutritionists, our data suggests that experience in sport
485 nutrition *per se* does not improve the accuracy of the RFPM /
486 ‘Snap-N-Send’ methodology. Rather, sport nutritionists looking
487 to use this technique would benefit from enhanced specialist
488 training including targeted activities to address the components
489 underpinning the accuracy in quantifying meal and individual
490 food portions from pictures prior to use. It should be stressed,
491 however, that taking pictures alongside traditional dietary intake
492 methodologies could help to reduce participant burden, improve
493 the accuracy of food diaries and help with behaviour change
494 (Costello et al., 2019). It is therefore important not to dismiss the
495 benefit of pictures to help with dietary assessment, rather the
496 present data highlights the limitation of this technique as a
497 standalone methodology.

498

499 Despite presenting novel data, this study is not without
500 limitation, many of which are directly related to the controls
501 employed to improve internal validity. Only two days of meals
502 were analysed in an attempt to recruit high-performance
503 nutritionists working in the elite environment. Initial
504 conversations prior to testing suggested that this length of food
505 diary would be acceptable from a time perspective for applied

Remote Food Photography Method in Sport Nutrition 2

506 practitioners. Future studies may wish to assess more days with
507 a wider range of energy intakes. Given that underreporting is
508 further exacerbated in accordance with increases in total energy
509 expenditure (Barnard et al., 2002) it is possible that in sports with
510 higher energy intakes (e.g. rugby, Bradley et al., 2015), the
511 RFPM / 'Snap-N-Send' could have higher variability than
512 reported here. A second limitation is that the meals in the present
513 study (despite some being classed as complex) were relatively
514 plain with things such as sauces and deserts being left to a
515 minimum. Combined with the fact that it was not necessary to
516 account for uneaten food, there is a high possibility that when
517 used by athletes in the field as an assessment tool, the variability
518 could be more extreme than reported in the current data.
519 Likewise, the present study was based upon the diet histories
520 reporting 100% of the total food consumed. In the real-world it
521 is likely that athletes will forget to take pictures (or fail to
522 submit) all of the food and drinks consumed adding further error
523 to this method. The present study used only one dietary
524 assessment software (Nutritics) given that Nutritics is widely
525 used in sport nutrition in the UK and Ireland (where all
526 participants were based) and were familiar with the software
527 using it regularly in their daily jobs. To assess whether the error
528 reported was purely related to the software, the lead researcher
529 with specific knowledge of the foods and weights inputted all of
530 the data into Nutritics and gained values within 1% of the total

Remote Food Photography Method in Sport Nutrition 2

531 energy reported on the food labels, suggesting that the error was
532 not within the software but rather the interpretation of the food
533 from the pictures. Finally, the aim of the present study was to
534 assess the RFPM / ‘Snap-N-Send’ within sport nutrition and it
535 therefore cannot be excluded that specialist trained individuals
536 who are highly experienced in picture-based diet assessments
537 may achieve differing data to that reported in the present study.

538

539 In conclusion, we provide the first report to assess the validity of
540 the RFPM / ‘Snap-N-Send’ as a standalone methodology to
541 assess energy and macronutrient composition of dietary
542 photographs. Our data demonstrate ‘poor’ validity and inter-
543 practitioner reliability, even when dietary analysis was
544 performed by experienced sport nutritionists. The present data
545 therefore provide a reference point for practitioners when
546 considering the typical error associated with these methods of
547 dietary assessment. Such estimates of validity should therefore
548 be taken into account when utilising this method alongside the
549 requirement to use multiple coders when performing dietary
550 analysis of athletic populations.

551

552 **Acknowledgements**

553 The study was designed by RGS, AMK, JPM and GLC; data
554 were collected and analysed by RGS, GLC and SAS; data
555 interpretation and manuscript preparation were undertaken by

Remote Food Photography Method in Sport Nutrition 2

556 RGS, AMK, SAS, JPM and GLC. All authors approved the final
557 version of the paper.

Remote Food Photography Method in Sport Nutrition 2

558 **REFERENCES**

- 559 Areta, J.L., Burke, L.M., Ross, M.L., Camera, D.M., West,
560 D.W.D., Broad, E.M., Jeacocke, N.A., Moore, D.R.,
561 Stellingwerff, T., Phillips, S.M., Hawley, J.A., & Coffey, V.G.
562 (2013). Timing and distribution of protein ingestion during
563 prolonged recovery from resistance exercise alters myofibrillar
564 protein synthesis. *The Journal of Physiology*, *591*, 9, 2319-2331.
565
- 566 Barnard, J.A., Tapsell, L.C., Davies, P.S.W., Brenninger, V.L.,
567 & Storlien, L.H. (2002). Relationship of high energy expenditure
568 and variation in dietary intake with reporting accuracy on 7 day
569 food records and diet histories in a group of healthy adult
570 volunteers. *European Journal of Clinical Nutrition*, *56*, 358-367.
571
- 572 Braakhuis, A.J., Meredith, K., Cox, G.R., Hopkins, W.G., &
573 Burke, L.M. (2003). Variability in estimation of self-reported
574 dietary intake data from elite athletes resulting from coding by
575 different sports dieticians. *International Journal of Sport*
576 *Nutrition & Exercise Metabolism*, *13*, 152-165.
577
- 578 Bradley, W.J., Cavanagh, B., Douglas, W., Donovan, T.F.,
579 Twist, C., Morton J.P., & Close, G.L. (2015). Energy intake and
580 expenditure assessed 'in-season' in an elite European rugby
581 union squad. *European Journal of Sport Science*, *15*, 469-479.
582

Remote Food Photography Method in Sport Nutrition 2

- 583 Burke, L.M., & Hawley, J.A. (2018). Swifter, higher, stronger:
584 What's on the menu? *Science*, 362, 781-787.
585
- 586 Capling, L., Beck, K.L., Gifford, J.A., Slater, G., Flood, V.M.,
587 & O'Connor, H. (2017). Validity of dietary assessment in
588 athletes: A systematic review. *Nutrients*, 9, 1313.
589
- 590 Cook, J.A., Hislop, J., Adewuyi, T.E., Harrild, K., Altman, D.G.,
591 Ramsay, C.R., Fraser, C., Buckley, B., Fayers, P., Harvey, I.,
592 Briggs, A.H., Norrie, J.D., Fergusson, D., Ford, I., & Vale, L.D.
593 (2014). Assessing methods to specify the target difference for a
594 randomised controlled trial - DELTA (Difference ELicitation in
595 TriAls) review. *Health Technology Assessment*, 18(28).
596
- 597
- 598 Costello, N., Deighton, K., Dalton-Barron, N., Whitehead, S.,
599 Preston, T., & Jones, B. (2019). Can a contemporary dietary
600 assessment tool or wearable technology accurately assess the
601 energy intake of professional young rugby league players? A
602 doubly labelled water validation study. *European Journal of*
603 *Sport Science*, Online ahead of print.
604
- 605 Costello, N., Deighton, K., Dyson, J., McKenna, J., & Jones, B.
606 (2017). Snap-n-Send: A valid and reliable method for assessing

Remote Food Photography Method in Sport Nutrition 2

607 the energy intake of elite adolescent athletes. *European Journal*
608 *of Sport Science*, 1044-1055.

609

610 Gemming, L., Utter, J., & Ni Mhurchu, C. (2014). Image-
611 assisted dietary assessment: A systematic review of the
612 evidence. *Journal of the Academy of Nutrition & Dietetics*, 115,
613 64-77.

614

615 Hannon, M., Carney, D., Floyd, S., Parker, L.J.F., McKeown, J.,
616 Drust, B., Unnithan, V., Close, G.L., & Morton, J.P. (2020).
617 Cross-sectional comparison of body composition and resting
618 metabolic rate in Premier League academy soccer players:
619 Implications for growth and maturation. *Journal of Sports*
620 *Sciences, EPub*.

621

622 Impey, S.G., Hearris, M.A., Hammond, K.M., Bartlett, J.D.,
623 Louis, J., Close, G.L., & Morton, J.P. (2018). Fuel for the work
624 required: A theoretical framework for carbohydrate
625 periodisation and the glycogen threshold hypothesis. *Sports*
626 *Medicine*, 48, 1031-1048.

627

628 Kasper, A.M., Crighton, B., Langan-Evans, C., Riley, P.,
629 Sharma, A., Close, G.L., & Morton, J.P. (2018). Case study:
630 Extreme weight making causes relative energy deficiency,
631 dehydration and acute kidney injury in a male mixed martial arts

Remote Food Photography Method in Sport Nutrition 2

- 632 athlete. *International Journal of Sport & Exercise Metabolism*,
633 29, 1-20.
- 634
- 635 Kikunaga, S., Tin, T., Ishibashi, G., Wang, D.H., & Kira, S.
636 (2007). The application of a handheld personal digital assistant
637 with camera and mobile phone card (Wellnavi) to the general
638 population in a dietary survey. *Journal of Nutritional Science &*
639 *Vitaminology*, 53, 109-116.
- 640
- 641 Kirkpatrick, S.I., & Collins, C.E. (2016). Assessment of nutrient
642 intakes: Introduction to the special issue. *Nutrients*, 8, 184.
- 643
- 644 Lassen, A.D., Poulsen, S., Ernst, L., Andersen, K.K., Biloft-
645 Jensen, A., & Tetens, I. (2010). Evaluation of a digital method
646 to assess evening meal intake in a free-living adult population.
647 *Food & Nutrition Research*, 54, 1-9.
- 648
- 649 Livingstone, M.B.E., & Black, A.E. (2003). Markers of the
650 validity of reported energy intake. *The Journal of Nutrition*, 133,
651 895S-920S.
- 652
- 653 Martin, C.K., Correa, J.B., Han, H., Allen, H.R., Rood, J.,
654 Champagne, C.M., Gunturk, B.K., & Bray, G.A. (2012).
655 Validity of the remote food photography method (RFPM) for

Remote Food Photography Method in Sport Nutrition 2

656 estimating energy and nutrient intake in near real-time. *Obesity*
657 *(Silver Spring)*, 20, 891-899.

658

659 Martin, C.K., Kaya, S., & Gunturk, B.K. (2009). Quantification
660 of food intake using food image analysis. *Conference*
661 *Proceedings: Annual International Conference of the IEEE*
662 *Engineering in Medicine and Biology Society*, 6869-6872.

663

664 Morton, J.P., Robertson, C., Sutton, L., & MacLaren, D.P.M.
665 (2010). Making the weight: A case study from professional
666 boxing. *International Journal of Sport Nutrition & Exercise*
667 *Metabolism*, 20, 80-85.

668

669 Rollo, M.E., Williams, R.L., Burrows, T., Kirkpatrick, S.I.,
670 Bucher, T., & Collins, C.E. (2016). What are they really eating?
671 A review on new approaches to dietary intake assessment and
672 validation. *Current Nutrition Reports*, 5, 307-314.

673

674 Thompson, F.E., Subar, A.F., Loria, C.M., Reedy, J.L., &
675 Baranowski, T. (2010). Need for technological innovation in
676 dietary assessment. *Journal of the American Dietetic*
677 *Association*, 110, 48-51.

678

679 Walsh, N.P. (2019). Nutrition and athlete immune health: New
680 perspectives on an old paradigm. *Sports Medicine*, 49, 153-168.

Remote Food Photography Method in Sport Nutrition 3

681

682 Westerterp, K.R., Saris, W.H., Van Es, M., & Ten Hoor, F.
683 (1986). Use of the doubly labeled water technique in humans
684 during heavy sustained exercise. *Journal of Applied Physiology*,
685 *61*, 2162-2167.

686

687 Wilson, G., Drust, B., Morton, J.P., & Close, G.L. (2014).
688 Weight-making strategies in professional jockeys: Implications
689 for physical and mental health and well-being. *Sports Medicine*,
690 *44*, 785-796.

691

692 Wilson, G., Pritchard, P.P., Papageorgiou, C., Phillips, S.,
693 Kumar, P., Langan-Evans, C., Routledge, H., Owens, D.J.,
694 Morton, J.P., & Close, G.L. (2015). Fasted exercise and
695 increased dietary protein reduces body fat and improves strength
696 in jockeys. *International Journal of Sports Medicine*, *36*, 1008-
697 1014.

698 **FIGURE & TABLE LEGENDS**

699 **Figure 1.** Overview of diet diary provided for both simple and
700 complex days. This includes image and brief explanation
701 provided to participants (**non-italic**) alongside the calculated
702 energy and macronutrient breakdowns for each meal and overall
703 daily total may (**italics**). Mega joules, MJ; carbohydrate, CHO;
704 protein, PRO; and fat, FAT.

705

706 **Table 1.** Outcomes of the limits of agreement (LoA) and
707 coefficient of variation (CV) analysis. CI denotes 95%
708 Confidence interval.

709

710 **Table 2.** Outcomes of the inter-rater reliability analysis. (α):
711 Cronbach's alpha; (ICC): intra class correlation; (CI): 95%
712 confidence interval; (CV): coefficient of variation.

713

714 **Figure 2.** Total energy intake (**A**) estimated by inexperienced
715 (**black circles**) and experienced (**white circles**) accredited
716 practitioners on the simple and complex days. Macronutrient
717 intake estimated by practitioners for carbohydrate (**B**), protein
718 (**C**) and fat (**D**). Bars are representative of mean estimation with
719 the dashed line representing actual calculate energy intake for
720 energy. * represents a significant difference compared to actual
721 calculated intake. # indicates significant differences between
722 groups.

Remote Food Photography Method in Sport Nutrition 3

723

724

725 **Figure 3.** Meal by meal overview (**A**, Breakfast; **B**, Snack; **C**,
726 Lunch; **D**, Evening meal) of total energy, carbohydrate, protein
727 and fat content (**1-4** respectively) estimated by inexperienced
728 (**black circles**) and experienced (**white circles**) accredited
729 practitioners on the simple and complex days. * represents a
730 significant difference compared to actual calculated intake.

Table 1.

Dietary Variable	Inexperienced		Experienced		All	
	Simple	Complex	Simple	Complex	Simple	Complex
Daily Energy Intake (MJ)						
Bias	-1.5	-1.2	-1.2	-0.6	-1.4	-0.9
CI	-1.9, -1.2	-1.8, -0.5	-1.6, -0.8	-1.2, 0.1	-1.6, -1.1	-1.3, -0.5
LoA (upper)	0.3	1.8	5.0	1.8	0.4	1.8
LoA (lower)	-3.4	-4.3	-0.5	-3.0	-3.2	-3.7
CV (%)	10.1	17.8	9.3	14.3	9.8	16.4
Interpretation	Poor	Poor	Acceptable	Poor	Poor	Poor
Carbohydrate (g.day⁻¹)						
Bias	-65.5	-28.7	-53.4	-19.9	-62.6	-24.5
CI	-75.0, -56.0	-49.7, -7.8	-62.7, -44.0	-35.6, -4.2	-68.8, -55.8	-37.3, -11.6
LoA (upper)	-20.5	70.7	-7.5	51.7	-19.1	62.1
LoA (lower)	-110.5	-128.1	-110.2	-91.4	-106.1	-110.6
CV (%)	10.8	24.4	10.1	17.5	10.4	21.3
Interpretation	Poor	Very Poor	Poor	Poor	Poor	Very Poor
Fat (g.day⁻¹)						
Bias	-7.1	-5.8	-3.6	4.0	-5.8	-1.1
CI	-14.2, 0.0	-11.6, 0.0	-8.8, 1.7	-2.0, 9.9	-9.7, -1.1	-5.4, 3.1
LoA (upper)	26.5	21.7	20.2	31.0	23.7	27.5
LoA (lower)	-40.8	-33.2	-27.3	-23.0	-35.2	-29.7
CV (%)	19.3	20.4	5.7	6.6	7.1	7.0
Interpretation	Poor	Very Poor	Acceptable	Acceptable	Acceptable	Acceptable
Protein (g.day⁻¹)						
Bias	7.3	-17.2	10.1	-15.7	7.9	-16.5
CI	-0.6, 15.3	-31.2, -3.3	1.28, 18.9	-27.7, -3.7	2.9, 14.3	-25.4, -7.6
LoA (upper)	45.2	49.0	49.9	38.5	47.4	43.7
LoA (lower)	-30.5	-83.5	-29.7	-69.9	-31.6	-76.7
CV (%)	9.1	16.3	9.5	13.3	9.5	14.8
Interpretation	Acceptable	Poor	Acceptable	Poor	Acceptable	Poor

Table 2.

α : Cronbach's alpha; ICC: intra class correlation; CI: 95% confidence interval; CV: coefficient of variation.

Dietary Variable	Inexperienced		Experienced		All	
	Simple	Complex	Simple	Complex	Simple	Complex
Daily Energy Intake						
α	0.985	0.931	0.977	0.834	0.991	0.950
ICC	0.73	0.35	0.65	0.180	0.69	0.29
CI	0.32, 1.00	0.06, 1.00	0.23, 1.00	0.001, 0.99	0.29, 1.00	0.06, 1.00
CV (%)	12.1	20.6	10.7	15.4	11.5	18.3
Interpretation	Acceptable	Poor	Acceptable	Poor	Acceptable	Poor
Carbohydrate						
α	0.995	0.875	0.994	0.855	0.997	0.932
ICC	0.89	0.22	0.88	0.20	0.89	0.22
CI	0.60, 1.00	0.02, 0.99	0.57, 1.00	0.12, 0.99	0.60, 1.00	0.04, 1.00
CV (%)	15.6	28.6	14.0	19.3	14.8	24.1
Interpretation	Acceptable	Very Poor	Acceptable	Poor	Acceptable	Poor
Fat						
α	0.765	0.765	0.496	0.472	0.841	-2.562
ICC	0.12	0.12	0.04	0.04	0.10	-0.02
CI	-0.01, 0.99	-0.01, 0.99	-0.03, 0.99	-0.03, 0.99	0.04, 0.99	-0.02, 0.85
CV (%)	20.9	22.3	14.1	19.0	17.8	21.6
Interpretation	Very Poor	Very Poor	Poor	Very Poor	Very Poor	Very Poor
Protein						
α	0.722	0.846	0.823	0.865	0.892	0.928
ICC	0.09	0.18	0.17	0.218	0.15	0.21
CI	-0.01, 0.99	0.01, 1.00	0.002, 0.99	0.16, 0.99	0.02, 1.00	0.03, 1.00
CV (%)	14.3	27.4	14.7	22.2	14.4	24.8
Interpretation	Very Poor	Very Poor	Very Poor	Very Poor	Very Poor	Very Poor

	Simple Day	Complex Day
Breakfast	 <p>Weetabix cereal (made with semi-skimmed milk) <i>[MJ=1.08; CHO=39.5g; PRO=13.5g; FAT=5.1g]</i></p>	 <p>Scrambled eggs on toast (made with semi-skimmed milk) <i>[MJ=2.50; CHO 41.3g; PRO 45.3g; FAT 28.2g]</i></p>
Morning Snack	 <p>Avocado on toast with poached eggs <i>[MJ=3.19; CHO=44.0g; PRO=25.6g; FAT=53.7g]</i></p>	 <p>Overnight oats (made with chocolate milk and whey protein) <i>[MJ=1.56; CHO=55.0g; PRO=26.7g; FAT=4.8g]</i></p>
Lunch	 <p>Poached salmon with baby new potatoes and broccoli <i>[MJ= 2.49, CHO=34.7g; PRO=47.8g; FAT=26.0g]</i></p>	 <p>Chicken tikka masala with pilau rice <i>[MJ=2.48; CHO=62.1g; PRO=35.6g, FAT 20.4g]</i></p>
Evening Meal	 <p>Chicken breast fillet with basmati rice and mixed peppers <i>[MJ=2.57; CHO=94.9g; PRO=41.0g; FAT=4.4g]</i></p>	 <p>Chicken chow mein <i>[MJ= 1.99; CHO=49.2g; PRO=32.8g; FAT=15.2g]</i></p>
Total	<p><i>MJ=9.33; CHO=213.1g; PRO=127.9; FAT=89.2g</i></p>	<p><i>MJ=8.53; CHO=207.6g; PRO=140.4; FAT=68.6g</i></p>

Figure 1. Overview of diet diary provided for both simple and complex days. This includes image and brief explanation provided to participants (non-italic) alongside the calculated energy and macronutrient breakdowns for each meal and overall daily total may (italics). Mega joules, MJ; carbohydrate, CHO; protein, PRO; and fat, FAT.

178x238mm (400 x 400 DPI)




Figure 2. Total energy intake (A) estimated by inexperienced (black circles) and experienced (white circles) accredited practitioners on the simple and complex days. Macronutrient intake estimated by practitioners for carbohydrate (B), protein (C) and fat (D). Bars are representative of mean estimation with the dashed line representing actual calculate energy intake for energy. * represents a significant difference compared to actual calculated intake. # indicates significant differences between groups.

180x202mm (300 x 300 DPI)




Figure 3. Meal by meal overview (A, Breakfast; B, Snack; C, Lunch; D, Dinner) of total energy, carbohydrate, protein and fat content (1-4 respectively) estimated by inexperienced (black circles) and experienced (white circles) accredited practitioners on the simple and complex days.

209x247mm (300 x 300 DPI)