



LJMU Research Online

Buckley, BJR and Lip, GYH

Current Concepts: Comprehensive "Cardiovascular Health" Rehabilitation-An Integrated Approach to Improve Secondary Prevention and Rehabilitation of Cardiovascular Diseases.

<http://researchonline.ljmu.ac.uk/id/eprint/18000/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Buckley, BJR and Lip, GYH (2022) Current Concepts: Comprehensive "Cardiovascular Health" Rehabilitation-An Integrated Approach to Improve Secondary Prevention and Rehabilitation of Cardiovascular Diseases. Thrombosis and haemostasis. ISSN 0340-6245

LJMU has developed [LJMU Research Online](#) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

Current Concepts

'Cardiovascular Health' Rehabilitation: An integrated approach to improve secondary prevention and rehabilitation of cardiovascular diseases

Benjamin J. R. Buckley¹, Gregory Y. H. Lip^{1,2,3}

¹Liverpool Centre for Cardiovascular Science, University of Liverpool and Liverpool John Moores University, Liverpool, UK

²Liverpool Heart and Chest Hospital, Liverpool, UK

³Department of Clinical Medicine, Aalborg University, Aalborg, Denmark

Date June 17th, 2022

Manuscript wordcount 687 (excluding title page, table, figure, and references)

Corresponding author

Prof GYH Lip

gregory.lip@liverpool.ac.uk

Definitions

Physical activity - any bodily movement produced by skeletal muscles that increases energy expenditure.

Moderate-vigorous intensity exercise - activities purposely undertaken to improve a component of fitness with an energy expenditure of ≥ 3 metabolic equivalents (METs) eg brisk walking to running.

Exercise-based cardiac rehabilitation (ExCR) - exercise training alone (typically two sessions/wk) or in combination with psychological or educational interventions.

Physical activity and moderate-vigorous intensity exercise provides primary¹ and secondary² ‘cardio-protection’. Further, the benefits of regular exercise training go beyond improvement in traditional cardiovascular disease risk factors (*ie* blood pressure, cholesterol, glycaemia), and most notably, elicit an improvement in vascular endothelial function.³

As such, exercise-based cardiac rehabilitation (ExCR) is an essential component of routine care for patients with acute coronary syndrome, those undergoing revascularisation (coronary artery bypass graft or percutaneous coronary intervention), and those with heart failure.⁴ Indeed, a substantive evidence base supports ExCR as a clinically effective and cost-effective intervention for patients with acute coronary syndrome or heart failure with reduced ejection fraction and after coronary revascularization.² The evidence base for use of ExCR in other indications, including heart failure with preserved ejection fraction,⁵ atrial fibrillation,^{6,7} congenital heart disease,⁸ and stroke⁹, requires further research.

Although some evidence suggests comprehensive ExCR leads to improved outcomes for patients with AF,¹⁰ traditional ExCR infrastructure is already struggling to cope with existing demand. For example, our previous work suggested that only 1.6% of eligible patients with heart failure were referred or initiated an ExCR programme.⁵ Therefore, alternative models of rehabilitation support are needed for other cardiovascular conditions not yet eligible for ExCR referral (*ie* atrial fibrillation, stroke, and chronic coronary syndrome). This may provide an opportunity to go beyond exercise rehabilitation and focus on a more holistic cardiovascular health rehabilitation programme. Herein, we propose ‘cardiovascular health’ rehabilitation as a potential ‘concept’ approach to help cope with the growing demand for more holistic and integrated cardiovascular disease management. Such an integrated care approach is increasingly evident in various chronic long-term conditions.¹¹⁻¹³

‘Cardiovascular health’ rehabilitation could therefore go beyond exercise-based rehabilitation and incorporate three key pillars (**Central Figure**): 1. disease-specific management (medication optimisation and symptom burden), 2. lifestyle behaviour change support (exercise, physical activity, diet, smoking, alcohol, sleep, mental health, and social wellbeing), and 3. Cardiovascular comorbidity and risk management (individual patient comorbidities and cardiovascular event risk). This is highly relevant given the typical clustering

of cardiovascular disease and risk factors (for example, multimorbidity can be seen in 2/3 older adults),¹⁴ and the clustering of healthy/maladaptive lifestyle behaviours such as exercise, physical activity, smoking, alcohol consumption, and diet in people with cardiovascular disease.¹⁵


There are some existing examples of promising efforts towards a more comprehensive lifestyle/cardiovascular health lens such as the American Heart Associations' 'Life's simple 7'.¹⁶ This concept identified 7 key risk factors for cardiovascular disease (smoking, body mass index, physical activity, diet, total cholesterol, blood pressure, and fasting serum glucose) and demonstrated that maintaining ideal cardiovascular health in all 7 variables was related to lower lifetime risk of coronary heart disease.¹⁷ Further, the American Heart Association have recently highlighted a primary care agenda,¹⁸ highlighting modifiable risk factors for cognitive decline including depression, hypertension, physical inactivity, diabetes, obesity, hyperlipidaemia, poor diet, smoking, social isolation, excessive alcohol use, sleep disorders, and hearing loss. This focus on both heart and brain health is topical, since new-onset cardiovascular complications diagnosed following an ischaemic stroke (termed 'stroke-heart syndrome') are very common and associated with significantly worse 5-year prognosis, compared to stroke survivors without cardiovascular complications.¹⁹ This further reinforces the need for a holistic and personalised cardiovascular health rehabilitation pathway, especially for those with multimorbidity's such as brain-heart conditions.^{12 19}

Another example of integrated cardiovascular disease management includes the ABC (Atrial fibrillation Better Care) pathway, including three guiding concepts: "A"—avoid stroke (with anticoagulants); "B"—better symptom management (with rate or rhythm control); and "C"—cardiovascular and comorbidity risk optimisation.¹¹ This guideline-recommended approach²⁰ incorporates both disease-specific treatment (for atrial fibrillation and stroke prevention) and management of individual cardiovascular comorbidities associated with lower major adverse cardiovascular events.²¹

Conclusions

'Cardiovascular health rehabilitation' provides a simple integrated care pathway guide for secondary prevention and holistic 'vascular health' rehabilitation of patients with

cardiovascular disease. Promotion and evaluation of such an approach may provide an opportunity to improve patient-centred and integrated care pathways for patients with cardiovascular disease, especially those with multimorbidity.



Central Figure. Three key pillars of ‘Cardiovascular Health Rehabilitation’ to be incorporated within an integrated care pathway for patients with cardiovascular disease.

1 Disease-specific management: First, the focus should be on optimising the clinical management of the patient and their disease-specific symptoms. This includes optimising any medication and discussion of potential treatment options. It is important to also consider treatment burden on an individual patient basis.

2 Lifestyle behaviour change support: This should be patient-driven and allow choosing and tailoring of the cardiovascular health rehabilitation programme. For some, an exercise-based programme may be highly suitable whereas others may not yet be able to exercise and may want to focus on physical activity levels and diet, or in fact, others may first want to focus on their mental wellbeing and are not yet able to exercise. The type and intensity of rehabilitation should be malleable and able to progress with evolving patient goals. Some patients will need more support than others, whether that is face-to-face, virtual, or hybrid should be discussed with the patient and consideration given to available resources.

3 Cardiovascular comorbidity and risk management: Patients with cardiovascular disease often have comorbidities and are at an elevated risk of future cardiovascular events. Therefore, as part of an integrated rehabilitation approach, management of other risk factors and conditions such as obesity, hypertension, diabetes mellitus, sleep apnoea, and other secondary cardiovascular conditions should be managed to help reduce the risk of future adverse events and the underlying cardiovascular burden. This should also facilitate a patient with cardiovascular disease to better manage their multimorbidity and improve their overall quality of life. Education and counselling can be used to improve a patient’s understanding, adherence and compliance to rehabilitation.

References

1. Morris JN, Heady JA, Raffle PA, et al. Coronary heart-disease and physical activity of work. *Lancet* 1953;262(6796):1111-20; concl. doi: 10.1016/s0140-6736(53)91495-0 [published Online First: 1953/11/28]
2. Taylor RS, Dalal HM, McDonagh STJ. The role of cardiac rehabilitation in improving cardiovascular outcomes. *Nat Rev Cardiol* 2022;19(3):180-94. doi: 10.1038/s41569-021-00611-7 [published Online First: 2021/09/18]
3. Fiuza-Luces C, Santos-Lozano A, Joyner M, et al. Exercise benefits in cardiovascular disease: beyond attenuation of traditional risk factors. *Nat Rev Cardiol* 2018;15(12):731-43. doi: 10.1038/s41569-018-0065-1 [published Online First: 2018/08/18]
4. Ambrosetti M, Abreu A, Corrà U, et al. Secondary prevention through comprehensive cardiovascular rehabilitation: From knowledge to implementation. 2020 update. A position paper from the Secondary Prevention and Rehabilitation Section of the European Association of Preventive Cardiology. *Eur J Prev Cardiol* 2020;28(5):460-95. doi: 10.1177/2047487320913379
5. Buckley BJR, Harrison SL, Fazio-Eynullayeva E, et al. Cardiac rehabilitation and all-cause mortality in patients with heart failure: a retrospective cohort study. *Eur J Prev Cardiol* 2021;28(Supplement_1) doi: 10.1093/eurjpc/zwab035 [published Online First: 2021/08/02]
6. Buckley BJR, Harrison SL, Fazio-Eynullayeva E, et al. Association of Exercise-Based Cardiac Rehabilitation with Progression of Paroxysmal to Sustained Atrial Fibrillation. *J Clin Med* 2021;10(3) doi: 10.3390/jcm10030435 [published Online First: 2021/01/28]
7. Buckley BJR, Harrison SL, Fazio-Eynullayeva E, et al. Exercise-Based Cardiac Rehabilitation and All-Cause Mortality Among Patients With Atrial Fibrillation. *J Am Heart Assoc* 2021;10(12):e020804. doi: 10.1161/JAHA.121.020804 [published Online First: 2021/06/08]
8. Tikkanen AU, Oyaga AR, Riaño OA, et al. Paediatric cardiac rehabilitation in congenital heart disease: a systematic review. *Cardiol Young* 2012;22(3):241-50. doi: 10.1017/s1047951111002010 [published Online First: 2012/01/19]
9. Buckley BJR, Harrison SL, Fazio-Eynullayeva E, et al. Exercise-based cardiac rehabilitation and major adverse cardiovascular events in patients with cerebrovascular disease *Cerebrovasc Dis* In Press
10. Buckley BJR, Risom SS, Boidin M, et al. Atrial Fibrillation Specific Exercise Rehabilitation: Are We There Yet? *J Pers Med* 2022;12(4) doi: 10.3390/jpm12040610 [published Online First: 2022/04/24]
11. Lip GYH. The ABC pathway: an integrated approach to improve AF management. *Nat Rev Cardiol* 2017;14(11):627-28. doi: 10.1038/nrcardio.2017.153 [published Online First: 2017/09/30]
12. Lip GYH, Lane DA, Lenarczyk R, et al. Integrated care for optimizing the management of stroke and associated heart disease: a position paper of the European Society of Cardiology Council on Stroke. *European Heart Journal* 2022:ehac245. doi: 10.1093/eurheartj/ehac245

13. Lip GYH, Ntaios G. "Novel Clinical Concepts in Thrombosis": Integrated Care for Stroke Management-Easy as ABC. *Thromb Haemostasis* 2022;122(3):316-19. doi: 10.1055/a-1632-1777 [published Online First: 2021/09/03]
14. Jadad AR, To MJ, Emara M, et al. Consideration of multiple chronic diseases in randomized controlled trials. *Jama* 2011;306(24):2670-2. doi: 10.1001/jama.2011.1886 [published Online First: 2011/12/29]
15. Lee S-R, Choi E-K, Ahn H-J, et al. Association between clustering of unhealthy lifestyle factors and risk of new-onset atrial fibrillation: a nationwide population-based study. *Sci Rep-uk* 2020;10(1):19224. doi: 10.1038/s41598-020-75822-y
16. Folsom AR, Yatsuya H, Nettleton JA, et al. Community Prevalence of Ideal Cardiovascular Health, by the American Heart Association Definition, and Relationship With Cardiovascular Disease Incidence. *Journal of the American College of Cardiology* 2011;57(16):1690-96. doi: <https://doi.org/10.1016/j.jacc.2010.11.041>
17. Hasbani NR, Ligthart S, Brown MR, et al. American Heart Association's Life's Simple 7: Lifestyle Recommendations, Polygenic Risk, and Lifetime Risk of Coronary Heart Disease. *Circulation* 2022;145(11):808-18. doi: 10.1161/CIRCULATIONAHA.121.053730
18. Lazar RM, Howard VJ, Kernan WN, et al. A Primary Care Agenda for Brain Health: A Scientific Statement From the American Heart Association. *Stroke* 2021;52(6):e295-e308. doi: 10.1161/STR.0000000000000367
19. Buckley BJR, Harrison SL, Hill A, et al. Stroke-Heart Syndrome: Incidence and Clinical Outcomes of Cardiac Complications Following Stroke. *Stroke* 2022;53(5):1759-63. doi: 10.1161/strokeaha.121.037316 [published Online First: 2022/04/01]
20. Hindricks G, Potpara T, Dagres N, et al. 2020 ESC Guidelines for the diagnosis and management of atrial fibrillation developed in collaboration with the European Association of Cardio-Thoracic Surgery (EACTS): The Task Force for the diagnosis and management of atrial fibrillation of the European Society of Cardiology (ESC) Developed with the special contribution of the European Heart Rhythm Association (EHRA) of the ESC. *European Heart Journal* 2020 doi: 10.1093/eurheartj/ehaa612 [published Online First: 2020/08/30]
21. Romiti GF, Pastori D, Rivera-Caravaca JM, et al. Adherence to the 'Atrial Fibrillation Better Care' Pathway in Patients with Atrial Fibrillation: Impact on Clinical Outcomes- A Systematic Review and Meta-Analysis of 285,000 Patients. *Thromb Haemostasis* 2021 doi: 10.1055/a-1515-9630 [published Online First: 2021/05/22]