



LJMU Research Online

Ekele, JU

Phytochemical analysis of six anti-venom medicinal plants

<http://researchonline.ljmu.ac.uk/id/eprint/22543/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Ekele, JU (2023) Phytochemical analysis of six anti-venom medicinal plants. Journal of Medicinal Plants Studies, 11 (3). pp. 71-79. ISSN 2394-0530

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

Title: “Phytochemical analysis of six anti-venom medicinal plants”

Author: EKELE, Jiata Ugwah

KOGI STATE UNIVERSITY, ANYIGBA, KOGI STATE, P.M.B 1008 NIGERIA

Email: ekelejiata22@gmail.com

Phone number: +2348148800156

Orcid ID: 0009-0008-8349-5024

ABSTRACT

This present investigation was conducted to identify and screen the most common medicinal plants used to treat snake bites in Ankpa local government area of Kogi state, Nigeria. After an ethnobotanical survey, only the roots of six medicinal plants were analysed for their qualitative and quantitative phytochemical properties. Roots were pulverised and extracted using methanol solvent. Alkaloids were investigated using Mayer and Dragendorff test; Tannins - Ferric-Chloride test; Total Phenol - Dennis test; Glycosides - Fehling A and B test; Saponins - Frothing test; Flavonoids - NaOH-HCl test; and quantitative constituents were identified using spectrophotometric method. The phytochemical screening indicated the existence of different classes of secondary metabolites, specifically alkaloids, flavonoids, saponins, tannins, total phenols and glycosides, at different levels. Statistical analysis was investigated by Analysis of Variance (ANOVA), and the means were distinguished according to Duncan's New Multiple Range Test (DMRT) at 5% probability level. Alkaloid contents obtained were 0.5mg/g, 0.8mg/g, 0.8mg/g, 0.5mg/g, 0.6mg/g, 2.2mg/g of the extract; Flavonoids - 20.8mg/g, 42.2mg/g, 30.9mg/g, 20.7mg/g, 30.5mg/g, 23.6mg/g of the extract; Saponins - 16.5mg/g, 23.1mg/g, 29.6mg/g, 18.8mg/g, 28.8mg/g, 20.7mg/g of the extract; Tannins - 22.8mg/g, 32.7mg/g, 34.4mg/g, 10.6mg/g, 28.8mg/g, 30.2mg/g of the extract; Total phenols - 710.4mg/g, 704.2mg/g, 715.6mg/g, 719.6mg/g, 719.7mg/g, 718.1mg/g of the extract for the six medicinal plants *Annona senegalensis*, *Khaya senegalensis*, *Uvaria chamae*, *Lophira lanceolata*, *Phyllanthus muellerianus* and *Securidaca longipedunculata* respectively. The results of the study revealed the presence of valuable bioactive compounds with medicinal properties in the aqueous and organic solvent extracts of these plants. Thus, supporting their use in traditional medicine for treating various ailments, such as snakebites.

Keywords: phytochemicals, snakebites, saponins, alkaloids, medicinal plants, *Khaya senegalensis*

INTRODUCTION

All of human lives and survival would be challenging and even impossible without the symbiotic relationship with the enormous diversity of plants for sources of food, shelter, fragrance, clothing, flavours, fertilisers and even medicines (anti-inflammatory, anti-venom, anti-cancer, diuretic, anti-plasmodic, anti-hypertensive, anti-diabetic, laxative and anti-microbial function) [1,2,3,4,5]. From time immemorial, medicinal plants have actively contributed to the progress of new drug discovery, while remaining essential in healthcare, and representing the optimal and secure reservoir for generating forthcoming medicines that are both safe and effective [6,7,8].

Clearly defined by the WHO, the term "medicinal plant" refers to any kind of plant whose organ(s) contain substances that possess the potential to be utilised for the treatment and prevention of particular ailments or diseases, or valuable for synthesising pharmaceutical medicines [9,10,11]. And the therapeutic features of most plant parts, such as bark, root, seed, fruit and leaf utilised, have been associated with significant origins of bioactive compounds and secondary metabolite productions [12,13,14,15]. Plants produce secondary metabolites to protect themselves, but several researchers have demonstrated that these phytochemicals can protect against disease, even snakebites [16,17,18,19]. These bioactive substances include flavonoids, lignans, terpenoids, carbohydrates, tannins, cytogenic glycosides, saponins, lignins, alkaloids, phenolic compounds, steroids and flavonoids [20,21,22,23,24,19,25].

Meanwhile, snakebites represent an overlooked tropical injury that impacts numerous individuals globally, primarily concentrated in South Asia and sub-Saharan Africa, where the highest incidence is observed [26,27,28]. Annually, the occurrence of snakebites can reach up to an estimated five million cases, leading to approximately 2.5 million poisonings and fatalities between 20,000 to 125,000 [29,30]. According to the World Health Organisation's (2008) analysis of mortality data, snakebites contribute to 35% of all child deaths globally, and no single country in the world is free from the risk. In these settings, there is limited access to healthcare services, scarcity of anti-venom and high cost of treatment that often lead to poor outcomes, considerable unwholesomeness or mortality that exacerbates the challenges associated with snakebites [31,32]. However, the use of plants against the effects of snake bites seems to be the best alternate preferred solution to snakebites for people living in remote places; hence, more scientific attention has to be given to it [33]. Sadly, the potential use of these highly valued medicinal plants, which act as a better alternative to conventional synthetic drugs, is still poorly explored, especially as regards the phytochemicals associated with them [34,35,36].

In this study, comprehensive qualitative and quantitative phytochemical analyses were conducted on six medicinal plants originating from Ankpa local government area (LGA) of Kogi state, Nigeria, namely: *Lophira lanceolata*, *Uvaria chamae*, *Khaya senegalensis*, *Phyllanthus muellerianus*, *Annona senegalensis* and *Securidaca longepedunculata*.

MATERIALS AND METHODS

Study Area

First, an ethnobotanical survey was conducted in Ankpa LGA, Kogi State, Nigeria. Ankpa LGA is situated at Latitude 7°26'N and Longitude 7°38'E of the equator with a population of 359,300 (in 2016) and a total land area of 1 200km² (500sq mi) [37]. Ankpa experiences a hot and humid climate throughout the year, characterised by an average annual temperature of 27°C. The vegetation in Ankpa exhibits typical characteristics of a derived savannah, with a toposequence predominantly covered by grasses and scattered shrubs. The topography of the area is notably undulating, featuring significant variations in elevation. The soil is very fertile and suitable for cultivating several medicinal herbs and food crops such as beans, maize,

yam, cassava, vegetables and cash crops like oil palm and cashew. Ankpa is characterised by a distinct seasonal pattern, comprising two main seasons: the wet season, spanning from April to October, and the dry season, occurring from November to March. The region receives an average annual rainfall of approximately 1300mm [37].




Figure 1: Map showing the location and extent of the study area

Collection and Identification of Plant Materials

Fresh parts (particularly roots) of the six medicinal plants highlighted from the ethnobotanical survey were collected using sterile cutlass. The plants *Annona senegalensis* Pers. (Ukpokpo-Igala tribal name), *Khaya senegalensis* (Odopala), *Securidaca longepedunculata* (Iche-oko) and *Uvaria chamae* P. Beaux (Awuloko) were collected by Mr Joseph Yamusa, a traditional snakebite healer in Ogodo-Atteh, Ankpa LGA Kogi State. *Phyllanthus muellerianus* (Odegenehie) was collected by Mr Yakubu Akoja (Obo), a traditional snakebite healer in Akwu, Ankpa LGA, Kogi State. And *Lophira lanceolata* Van Tiegh (Okopi) was collected by Mr Samson Simon Ocholi, a traditional snakebite healer in Ogodo, Ankpa LGA, Kogi State.

The plant samples were taxonomically identified by Mr Theophilus Boniface Momoh at Botany Department, Kogi State University, Ayingba; six (6) plants were identified and used.

Sample preparation

The plant roots were cleaned with distilled water to remove soil dirt, dust and germs. The roots were cut into pieces using a sterile knife and were air-dried at room temperature before being pulverised using a mortar and pestle and then sieved. The powdered forms were kept in air-tight, properly labelled containers for further laboratory use.

Extraction Method

The crude extraction was done using ethanol as solvent. Two grams (2g) of each powdered plant sample were weighed accurately and macerated in 20 ml of 50% ethanol solvent in different conical flasks. Each sample was macerated in a separate conical flask with the solvent in such a way that the level of the solvent was above that of the plant powdered roots. The macerated mixtures were allowed to sit at room temperature for a duration of 48 hours.

The extracts were recovered from the macerated mixture through Whatman filter paper to remove the insoluble materials, and the following analyses were carried out on the filtrate.

Qualitative Phytochemical Screening of the Roots

The roots of *Annona senegalensis* Pers., *Khaya senegalensis*, *Phyllanthus muellerianus*, *Uvaria chamae* P. Beaux, *Lophira lanceolata* Van Tiegh and *Securidaca longepedunculata* were subjected to phytochemical screening using established protocols to identify the constituents previously described by Harris *et al.* and Adeoye *et al.* [38,39].

1. Alkaloids

For the Mayer test, 0.2g of the extract was heated with 2% H₂SO₄ for a duration of two minutes. After filtration, a few drops of Mayer's reagent were added to the filtrate. The formation of a cream-colored precipitate indicated the presence of alkaloids [38].

In the Dragendorff test, 0.2g of the extract was heated with 2% H₂SO₄ for two minutes. The resulting mixture was filtered, and a few drops of Dragendorff's reagent were added to the filtrate. The appearance of an orange-red precipitate indicated the presence of alkaloids [38].

2. Tannins

For the Ferric Chloride test, a small amount of each extract was combined with water, heated using a water bath, and then filtered. To the filtrate, a few drops of ferric chloride were added. The formation of a dark green solution indicated the presence of tannins [39].

3. Glycosides

In the Fehling A and B test, the extract was first hydrolysed with HCl and subsequently neutralised with NaOH solution. A few drops of equal amounts of Fehling A and B reagents were then added. The formation of red precipitates indicated the presence of glycosides [39].

4. Saponins

In the frothing test, 0.2g of the extract was vigorously shaken with 5 ml of distilled water and subsequently filtered. The resulting filtrate was then boiled. The presence of a frothy appearance with creamy white bubbles indicated the presence of saponins [39].

5. Flavonoids

In the NaOH-HCl test, 0.2g of the extract was dissolved in diluted NaOH solution, followed by the addition of HCl. The formation of a yellow solution indicated the presence of flavonoids [39].

6. Total phenol

For the Dennis test, a small amount of the extract was heated with 35% NaCO₃ for a duration of five minutes. Following this, a few drops of Folin-Ciocalteu reagent were added to the filtrate. The development of a blue-black color in the solution indicated the presence of total phenols [39].

Quantitative Phytochemical Screening of the Roots

1. Alkaloid determination using Harris *et al.* [38] method

A 5g portion of each sample was weighed and placed into a 250 ml beaker. To this, 200 ml of 20% acetic acid in ethanol was added, and the beaker was covered. The mixture was then left to stand for a period of 4 hours. Following this, the solution was filtered, and the volume of the extract was reduced to one-quarter of its original volume by concentrating it in a water bath. Concentrated ammonium hydroxide was slowly added drop by drop to the sample until precipitation was complete. The entire solution was allowed to settle, and the resulting precipitate was collected by filtration and weighed. The percentage of the total alkaloid was determined by calculating the weight of the precipitate relative to the original sample weight:

$$\text{Weight of Alkaloid} = \frac{\text{Residual Weight } (W_1 - W_2)}{\text{Weight of the Sample}} \times 100$$

$$W_1 = \text{Weight of Filter Paper} + \text{Alkaloid}$$

$$W_2 = \text{Weight of Filter Paper}$$

2. Total Phenol determination by spectrophotometric method

The determination of total phenolic content was conducted using the Folin-Ciocalteu method. A 20 μ l sample of the extract solution was mixed with 1.16ml of distilled water, and 100 μ l of Folin-Ciocalteu

reagent was added. Following this, 300µl of a 20% Na₂CO₃ solution was introduced to the mixture. The resulting solution was incubated in a shaking incubator at 40°C for 30 minutes, and its absorbance at 760nm was measured using a UV/visible spectrophotometer. Gallic acid served as the standard for the calibration curve. The total phenolic content was expressed as gallic acid equivalent (GAE) and was calculated using a linear equation based on the calibration curve [40].

$$A = 0.98C + 9.925 \times 10^{-3} \quad (R^2 = 0.9998)$$

A= Absorbance

C = Concentration mg GAEg⁻¹

3. Tannins determination using spectrophotometric method

Tannins were determined using the Folin-Ciocalteu method. A 0.1ml aliquot of the sample extract was added to a 10ml volumetric flask containing 7.5ml of distilled water and 0.5ml of Folin-Ciocalteu tannin reagent. To this mixture, 1ml of 35% Na₂CO₃ solution was added, and the flask was then filled to the 10ml mark with distilled water. The solution was thoroughly shaken and left at room temperature for 30 minutes. Reference standard solutions of gallic acid (20, 40, 60, 80, and 100µg/ml) were prepared using the same procedure as described earlier. The absorbance of the test samples at 725nm was measured using a UV/visible spectrophotometer, and the absorbance of the standard solutions against a blank was also measured at 725nm. The tannin content was expressed as milligrams of gallic acid equivalent (GAE) per gram of extract [40].

4. Saponins determination using the Spectrophotometric method

The total saponin content was determined following the method described by Makkar et al. with some modifications. To 50ml of the sample extract, 250ml of distilled water, 250ml of vanillin reagent (800mg vanillin in 10ml of 99.5% ethanol), and 2.5ml of 72% sulphuric acid were added. The mixture was thoroughly mixed and placed in a water bath at 60°C for 10 minutes. Subsequently, the solution was cooled in ice-cold water, and the absorbance was measured at 544nm. The values obtained were expressed as diosgenin equivalent (mg/DEg) extract, derived from a standard curve [40].

Extract: 1g of sample in 50ml of distilled water

absorbance × gradient factors × dilution factors

5. Flavonoids determination using the Spectrophotometric method

Flavonoid determination was performed using the aluminum chloride colorimetric method. A 0.5ml aliquot of each plant extract was mixed with 1.5ml of methanol, 0.1ml of 10% aluminum chloride, 0.1ml of 1M sodium acetate, and 2.8ml of distilled water. The mixture was left at room temperature for 30 minutes to allow the reaction to occur. The absorbance of the reaction mixture was measured at 510nm using a double-beam Perkin Elmer UV-Visible Spectrophotometer. To generate a calibration curve, quercetin solutions ranging from 12.5 to 100µg/ml were prepared in methanol [40].

Statistical Analysis

The mean levels of phytochemicals among different plant categories were analysed using One-way Analysis of Variance (ANOVA) with a confidence level of 95%. The upper and lower confidence limits

were also calculated. The statistical analysis was performed using IBM SPSS Statistics version 21.0 software. An interaction term was included in the model, and Post-hoc Tukey tests were conducted to assess the significance of the results using R Studio [41]. The data were presented as mean \pm standard deviation, based on three determinations. A p-value of ≤ 0.05 was considered statistically significant to determine differences between the groups.

RESULTS

Ethnobotanical survey results

Based on the conducted survey, the study area identified six plant species that were commonly used for the treatment of snake bites. Table 1 presents the common and scientific names of these plant species:

Table 1: Plants used for the treatment of Snakebite in Ankpa LGA

S/No	Common Names	Scientific Names	Igala Names	Family	Habit	Most frequent plant part used
1	Wild Sour Sop	<i>Annona senegalensis</i> Pers.	<u>Ukpokpo</u>	Annonaceae	Shrub	Roots
2	African Mahogany Red Oak Tree/ Red	<i>Khaya senegalensis</i>	<u>Odopala</u>	Meliaceae	Tree	Roots
3	Iron Wood	<i>Lophira lanceolata</i> Van. Tiegh	<u>Okopi</u>	Ochnaceae	Tree	Roots
4	Chewing Stick Plant	<i>Phyllanthus muellerianus</i>	<u>Odegenhie</u>	Euphorbiaceae	Climbing Shrub	Roots
5	Violet Tree/ Fibre Tree	<i>Securidaca longepedunculata</i> Fers.	<u>Iche-oko</u>	Polygalaceae	Shrub	Roots
6	Bush Banana	<i>Uvaria chamae</i> P. Beaux.	<u>Awuloko</u>	Annonaceae	Climbing Shrub	Roots

Phytochemical present (qualitative analysis)

Below is Table 2 summarising the phytochemical characteristics of the six medicinal plants. The results of the study demonstrated the presence of medically active compounds in the six plants examined. Table 2 indicates that total phenols, glycosides, flavonoids, and saponins were detected in all of the plants. However, tannins were not found in *Khaya senegalensis*, while alkaloids were exclusively present in

Uvaria chamae. These findings highlight the diverse phytochemical composition of the studied plants and suggest their potential therapeutic value.

Table 2: Qualitative phytochemicals constituents of six medicinal plant roots analysed

Plants	Alkaloids	Tannins	Flavonoids	Saponins	Total phenols	Glycosides
<i>Lophira lanceolata</i>	-	+	+	+	+	+
<i>Uvaria chamae</i>	+	+	+	+	+	+
<i>Khaya senegalensis</i>	-	-	+	+	+	+
<i>Phyllanthus muellerianus</i>	-	+	+	+	+	+
<i>Annona senegalensis</i>	-	+	+	+	+	+
<i>Securidaca longepedunculata</i>	-	+	+	+	+	+

- = Absent

+ = Present

Mean level of

phytochemicals in each plant

The mean levels of five (5) important phytochemicals were examined from the roots of the six (6) selected plants under study—namely, alkaloids, total phenols, saponins, flavonoids and tannins.

The level of alkaloids was highest in the root of *Uvaria chamae* with a mean of 2.233 mg/g, which was statistically significant from the levels observed in the roots of *Lophira lanceolata* (0.833 mg/g), *Securidaca longepedunculata* (0.807 mg/g), *Phyllanthus muellerianus* (0.620 mg/g), *Khaya senegalensis* (0.513 mg/g) and *Annona senegalensis* (0.511 mg/g) ($F_{5,30}$ 165.9; $p < 0.001$) (Figure 2). The level of alkaloids in *Khaya senegalensis* and *Phyllanthus muellerianus* were not significantly different from that of *Annona senegalensis* ($p = 0.574$).

Similarly, the level of flavonoids was highest in the root of *Lophira lanceolata* with a mean of 42.153 mg/g, which was significant from the levels observed in the roots of *Securidaca longepedunculata* (30.893 mg/g), *Phyllanthus muellerianus* (30.483mg/g), *Uvaria chamae* (23.550 mg/g), *Annona senegalensis* (20.813 mg/g) and *Khaya senegalensis* (20.670 mg/g) ($F_{5,30}$ 4047; $p < 0.001$) (Figure 3). The level of flavonoids in *Annona senegalensis* was not significantly different from that of *Khaya senegalensis* ($p = 0.826$).

The level of saponin was highest in the root of *Securidaca longepedunculata* with a mean of 29.537 mg/g, which was significant from the levels observed in the roots of *Phyllanthus muellerianus* (28.800 mg/g), *Lophira lanceolata* (23.033 mg/g), *Uvaria chamae* (20.693 mg/g), *Khaya senegalensis* (18.760 mg/g) and *Annona senegalensis* (16.457 mg/g) ($F_{5,30}$ 1.913; $p < 0.001$) (Figure 3).

The root of *Securidaca longepedunculata* exhibited the highest level of tannins among the tested plant samples with a mean of 34.370 mg/g, which was significant from the levels observed in the roots of *Uvaria chamae* (30.163mg/g), *Phyllanthus muellerianus* (28.733 mg/g), *Annona senegalensis* (22.770 mg/g) and *Khaya senegalensis* (10.567 mg/g) but was not significantly different from the level observed in the roots of *Lophira lanceolata* (32.723 mg/g) ($F_{5,30} 4095$; $p < 0.001$) (Figure 3). The level of tannins in *Lophira lanceolata* was not significantly different from that of *Uvaria chamae* ($p = 0.675$).

The root of *Phyllanthus muellerianus* with a mean of 719.72 mg/g, exhibited the highest level of phenols among the tested plant samples which was significant from the levels observed in the roots of *Uvaria chamae* (718.11 mg/g), *Securidaca longepedunculata* (715.61 mg/g), *Annona senegalensis* (710.43 mg/g) and *Lophira lanceolata* (704.23 mg/g) ($F_{5,30} 37.77$; $p < 0.001$) (Figure 4). But it was not significantly different from the level observed in the roots of *Khaya senegalensis* (719.61 mg/g) ($p = 0.991$).

Overall, phenols had the highest level among the five phytochemicals with a mean of 714.618 mg/g, followed by flavonoids (28.094 mg/g), tannins (26.554 mg/g), saponins (22.880 mg/g) while alkaloids had the least level (0.919 mg/g) ($F_{9,20} 7116$; $p < 0.001$).




Figure 2: It shows the alkaloid contents of the roots of *Annona senegalensis*, *Khaya senegalensis*, *Phyllanthus muellerianus*, *Uvaria chamae*, *Lophira lanceolata* and *Securidaca longepedunculata*, respectively




Figure 3: It shows the flavonoid, saponin and tannin contents of the roots of *Annona senegalensis*, *Khaya senegalensis*, *Phyllanthus muellerianus*, *Uvaria chamae*, *Lophira lanceolata* and *Securidaca longepedunculata* respectively




Figure 4: It shows the total phenolic contents of the roots of *Annona senegalensis*, *Khaya senegalensis*, *Phyllanthus muellerianus*, *Uvaria chamae*, *Lophira lanceolata* and *Securidaca longepedunculata*, respectively

DISCUSSION

Plant extracts are known to be highly abundant in pharmacologically active compounds and often possess multiple biochemical and pharmacological properties. These compounds have the ability to interact with toxins or enzymes, thereby neutralising or inhibiting their biochemical activities. These interactions play a crucial role in the pharmacological activity exhibited by plant extracts [42,43,44]. Several researchers have highlighted the potential of medicinal plants as anti-venoms, attributing their efficacy to the presence of various forms of flavonoids, steroids, and tannins. It is important to note that the full activity of an extract cannot be replicated by a single substance alone. Instead, the combined action of multiple active constituents present in the plants acts synergistically on different target structures such as receptors and enzymes. These chemical constituents are often referred to as 'multifunctional' due to their ability to exhibit more than one pharmacological or biochemical property [45,46,47,48].

Earlier studies on the phytochemical screening of *Annona senegalensis* have reported the presence of diverse secondary metabolites, including tannins [49], flavonoids [49], saponins [50], glycosides [51], and alkaloids [52]. However, it is worth noting that the alkaloids isolated from the roots of *Annona senegalensis* did not exhibit any biological activity [53]. These findings provide insights into the chemical composition of *Annona senegalensis* and shed light on the potential pharmacological properties associated with its secondary metabolites.

In line with the aforementioned studies, Audu *et al.* [54] conducted research on the phytochemical screening of *Lophira lanceolata*, revealing the presence of flavonoids, glycosides, phenols, and saponins. However, alkaloids were found to be absent in this species. Another study focused on the phytochemical screening of the leaves and bark of *Khaya senegalensis*, demonstrating the presence of tannins, steroids, flavonoids, saponins, and alkaloids in the ethanolic extracts. However, alkaloids and saponins were not detected in the methanolic extracts [55]. In contrast, the ethanolic extract of the root of *Uvaria chamae* exhibited significantly higher levels ($p \leq 0.05$) of total phenols, tannins, flavonoids, and saponin content compared to the findings reported by Komes *et al.* [56] and Iroabuchi [57]. These variations could be attributed to factors such as seasonal variations, maturity stage of the plants, and environmental conditions affecting the plant samples.

Phenols and flavonoids, which are essential constituents found in plants, have been identified as key contributors to the anti-venom activity of these plants, either directly or indirectly [58]. The high presence of phenols and flavonoids in these plants may account for their potent anti-venom properties. Saponins, known for their bitter taste, foaming properties, and ability to cause hemolysis in red blood cells, are utilised as emulsifying agents and expectorants [59]. Phenolic compounds possess a wide range of medicinal properties, including antioxidant, anti-venom, anti-inflammatory, and anti-cancer activities [60,61,62,63,64,65]. Flavonoids, on the other hand, represent a diverse and widely distributed group of natural compounds, exhibiting various chemical and biological activities, including radical scavenging, anti-allergic, anti-microbial, anti-venom, anti-inflammatory and vasodilating effects [66,67,68].

Quercetin-3-O- α -rhamnoside, a flavonoid, has been identified as an inhibitor of phospholipase A2 (PLA2) activity, as well as the haemolytic and haemorrhage-inducing activities of *Naja naja* venom. It has been found effective when used in a ratio of 1:20 (venom to quercetin-3-O- α -rhamnoside) [69]. Morelloflavone, a flavonone-(C-3 C-8'')-flavone biflavonoid, has also demonstrated inhibition of enzymatic, myotoxic, oedema-forming, and anticoagulant activities induced by *Crotalus durissus cumanensis* venom PLA2 [69]. Persimmon, which contains tannins, has been reported to exhibit activity against *Laticauda semifasciata* and *Trimeresurus flavoviridis* venoms. Ar-turmerone, a phenolic compound, has shown the ability to

neutralise haemorrhaging and lethality caused by *Bothrops jararaca* and *Crotalus durissus terrificus* snake venoms [70].

The methanol extract of the root bark of *Annona senegalensis* has been found to possess anti-venom properties against cobra (*Naja nigricollis* Wetch) venom in experimental rats. It showed a reduction in induced hyperthermia and detoxification of the snake venom by 16-33%. However, it did not restore liver functions [71]. The anti-venom properties of *Securidaca longepedunculata* root extract were evaluated by monitoring the levels of liver enzymes (ALT, AST, ALP), creatinine kinase (CPK), lactase dehydrogenase (LDH), and amylase in rats. The extract demonstrated a significant dose-dependent alteration in serum enzymes and urea levels [72]. A fraction of *Annona senegalensis* leaf methanol extract showed neutralising effects against lethal toxicity induced by *Echis ocellatus* venom. The key phytochemicals responsible for this activity were identified as flavonoids and tannins [73,74,75,76,77,78]. These findings suggest that the identified phytochemical compounds may be the bioactive constituents responsible for the observed effects. These plants hold promise as valuable sources of bioactive compounds with significant medicinal potential.

CONCLUSION

The outcomes of this investigation have demonstrated the presence of vital therapeutic components in the investigated plants. The found phytochemicals' bioactivity was supported by several findings from past investigations. Many studies have shown that the presence of these phytochemicals contributes to the pharmacological and physiological qualities of the plants investigated in the therapy of diverse ailments. Therefore, the extracts from these plants can be regarded as a trustworthy and dependable source for therapeutic medications. It is highly recommended to utilise these plants in native and modern medicine practices. It is also urged that additional research be done to identify, purify, and isolate the medicinal ingredients that give these plants their effects. Further research is also encouraged to clarify these extracts' potential mechanisms of action.

REFERENCES

- [1] Aadhan K, Kumar M and Anand SP. Traditional Herbal Medicine for the Treatment of Snakebite and Scorpion Sting by the Paliyar's tribe of Sathuragiri Hills. *Journal of Medicinal Plant Studies* 2017, 5(2): 1-5.
- [2] Ajali U and Chukwurah BKC. Antimicrobial Activity of *Securidaca longipedunculata*. *Journal of Phytomedicine* 2004, 11: 701-703.
- [3] Fabeku PO, Aibinu EI, Akinsulire RO and Odugbemi OT. Medicinal Plants useful for Malaria Therapy in Okeigbo, Ondo State, Southwest, Nigeria. *African Journal of Traditional, Complementaery and Alternative Medicine* 2007, 4(2): 191-198.
- [4] Saeed M, Arssad M, Ahmad E and Ishaque M. Ethnophytotherapies for the Treatment of Various Diseases by the local people of Selected Areas. *Pakistan Journal of Biological Science* 2004, 7: 1104-1108.
- [5] Okwu DE and Ekeke O. Phytochemical screening and mineral composition of chewing stick in South Eastern Nigeria. *Global Journal of Pure and Applied Sciences* 2003, 9: 235-238.
- [6] Ameen SA, Salihu T, Mbaoji CO, Anuoro-Dibia CA and Adedokun RAM. Medicinal Plants used to treat Snakebite by Fulani Herdsmen in Taraba State, Nigeria. *International Journal of Applied Agricultural and Apicultural Research* 2015, 11(1&2): 10-21.
- [7] Dar RA, Shahnawaz M and Qazi PH. Natural product medicines: A literature update. *Journal of Phytopharmacology* 2017, 6(6): 349-351.
- [8] Fakim AG. Medicinal plants: Traditions of yesterday and drugs of tomorrow. *Molecular Aspects of Medicine* 2006, 27: 1-93.
- [9] Alitonou GA, Koudoro AY, Dangou JS, Yehouenou B, Avlessi F, Adeoti S et al. Volatile Constituents and Biological Activities of Essential Oil from *Securidaca longepedunculata* Fresen. growing in Benin. *Scientific Study and Research* 2012, 13: 33-42.
- [10] Anselem A. Herbs for healing. Pax Herbals, Edo State, Nigeria 2004, 13-16.
- [11] World Health Organisation. Regulatory Situation of Herbal Medicines. A Worldwide Review, Geneva, Switzerland 1998, 1-5.
- [12] Cowley A. Alternative Medicine, New Integrative Carel News week 2002, 2.
- [13] Dapar LPM, Aguiyi CJ, Wannang NN, Gyang SS and Tanko MN. The Histopathologic effects of *Securidaca longepedunculata* on heart, liver, kidney and lungs of rats. *African Journal of Biotechnology* 2007, 6(5): 591-595.
- [14] Gomes A, Das R, Sarkhel S, Mishra R, Mukherjee S and Bhattacharya S. Herbs and Herbal constituent active against snakebite. *Indian journal of experimental biology* 2010, 48: 865-878.
- [15] Obeidat M, Shatnawi M, Al-Alawi M, Al-Zubi E and Al-Dmoor H. Anti-microbial activity of Crude extracts of Some plant leaves. *Research Journal of Microbiology* 2012, 59-67.

- [16] Gutiérrez JM, Rucavado A, Chaves F, Díaz C and Escalante T. Experimental pathology of local tissue damage induced by *Bothrops asper* snake venom. *Toxicon: Official Journal of the International Society on Toxinology* 2009, 54(7): 958–975 <https://doi.org/10.1016/j.toxicon.2009.01.038>.
- [17] Hassan SS, Al-Jabri AA, Sallam TA, Al-Balushi MS and Mothana RAA. Anti-snake Venom activity of *Hibiscus aethiopicus* L. against *Echis ocellatus* and *Naja nigricollis*. *Journal of Toxicology* 2010, 2(1): 1-8.
- [18] Okwu DE and Josiah C. Evaluation of the chemical composition of two Nigerian medicinal plants. *African Journal of Biotechnology* 2006, 5: 357–361.
- [19] Saidulu CH, Venkateshwar C and Gangadhar RS. Preliminary Phytochemical Studies of Medicinal Plant Drugs. *An International Quarterly Journal of Biology and Life Sciences* 2014, 2(1): 306.
- [20] Alaribe SI. A Survey of the Importance and Problems of Traditional Health care medicine: A case study of Ezinihitte Mbaise L.G.A. Imo State. Published B.Sc. project, Alvan Ikoku Federal College of Education, Owerri, Imo State 2008, 18-19.
- [21] Edoga HO, Okwu DE and Mbaebie BO. Phytochemicals constituents of some Nigerian medicinal plants. *African Journal Biotechnological* 2005, 4(7): 685-688.
- [22] Igwe SA and Nwobodo NN. Anticonvulsant Activity of Aqueous Root Extract of *Annona senegalensis* Pers. *International Journal of Advanced Biological and Biomedical Research* 2014, 2(8): 2441-2448.
- [23] Jada M, Usman W and Adamu Y. In vitro Anti-microbial Effect of Crude Tannins Isolated from the Leaf of *Annona senegalensis*. *International Journal of Biochemistry Research & Review* 2014, 4(6): 615–623.
- [24] Mann J. *Secondary Metabolism*. Oxford University press, London 1978, 154
- [25] Stamp N. Out of the quagmire of plant defense hypotheses. *The Quarterly Review of Biology* 2003, 78 (1): 23–55.
- [26] Abubakar SB, Habib AG and Mathew J. Amputation and Disability following Snakebite. *Tropical Doctor* 2010, 40(2): 114-116.
- [27] World Health Organization. *International Statistical Classification of Diseases and Related Health Problems* 2008. 10th revision version. <http://apps.who.int/classifications/icd10/browse/2010/en>.
- [28] World Health Organization. *A Multicomponent Strategy to Improve the Availibility of Anti-venom for Treating Snake bite Envenoming* 2016, 12-14.
- [29] Habib AG, Gebi UI and Onyemelukwe GC. Snake bite in Nigeria. *African Journal of Medicine and Medical Science* 2013, 30(3): 171-178.
- [30] Kasturiratne A, Wickremasinghe AR, De-Silva N, Gunawardena NK, Pathmeswaran A et al. A literature analysis and modelling based on Regional estimates of envenoming and deaths. *The Global Burden of Snakebite*. In Winkel Ken. *PLOS Medicine* 2008, 11(5): 218.
- [31] Lalloo G. Snake Envenoming: A Disease of Poverty. *PLoS Neglected Tropical Disease* 2009, 3(12): 569.

- [32] Meenatchisundaram S, Parameswari G, Subbaraj T and Michael A. Anti-venom activity of Medicinal Plants-A mini review. *Ethnobotanical Leaflets* 2008, 12: 1218-1220.
- [33] Omogbai EKI, Nworgu ZAM, Imhafidon MA, Ikpeme AA, Ojo DO and Nwako CN. Snake bite in Nigeria. A study of prevalence and treatment in Benin-city. *Tropical Journal of Pharmaceutical Research* 2002, 98: 312-315.
- [34] Atul-Kaushik D, Anghesom A, Kaushik JJ and Girmay B. Snake Venom Neutralisation Effects of African Medicinal Plants & Their Impact on Snakebites: A Review. *Asian Journal of Biomedical and Pharmaceutical Sciences* 2013, 3(24): 1-6.
- [35] Junaidu S, Kasim S, Abdulhadi BJ, Iliya M and Sanusi L. Anti Snake Venom Potential of *Securidaca longepedunculata* Leaf and Root Bark on Spitting Cobra (*Naja nigricollis* Hallowel) in Envenomed Wister Rats. *International Journal of Pharmacy and Biological Sciences* 2014, 9(6): 92-96.
- [36] Mustapha AA. *Annona senegalensis* Persoon: A multipurpose shrub, its Phytotherapeutic, Phytopharmacological and Phytomedicinal uses. *International Journal of Science and Technology* 2013, 2(12):862-865.
- [37] Kogi (State, Nigeria) - Population, Statistics, Charts, Map and Location. <http://www.citypopulation.de>. Retrieved 2019-09-22.
- [38] Harris KK, Meena S and Devshree V. Phytochemical Analysis of the Leaf, Stem and Seed Extracts of *Cajanus cajan*. *World Journal of Pharmacy and Pharmaceutical Science* 2014, 3(8): 694-733.
- [39] Adeoye AO, Boligon AA, Athayde ML and Bewaji CO. In Vitro Anti-oxidant Activity and High Performance Liquid Chromatography Fingerprint of *Adansonia digitata* Stem Bark Extracts. *International Journal of Pharmacognosy and Phytochemical Research* 2016, 8(12): 1951-1958.
- [40] Makkar HPP, Siddhuraju P and Becker K. *Methods in Molecular Biology. Plant Secondary Metabolites*. Humana Press Inc; Totowa New Jersey 2007, 393: 93-100.
- [41] R Core Team. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria 2021. URL <https://www.R-project.org/>
- [42] Makhija IK, Khamar D and Khan DE. *Der Pharmacia Lettre* 2010, 2, 399.
- [43] Ogbadoyi EO, Abdulganiy AO, Adama TZ and Okogun JI. In vivo trypanocidal activity of *Annona senegalensis* Pers. leaf extract against *Trypanosoma brucei brucei*. *Journal of Ethnopharmacology* 2007, 112(1): 85–89.
- [44] Okhale SE, Akpan EF, Fatokun OT, Esievo B and Kunle FO. *Annona senegalensis* Persoon (Annonaceae): A review of its ethnomedicinal uses, biological activities and phytochemicals. *Journal of Pharmacognosy and Phytochemistry* 2016, 5(2): 211-219.
- [45] Asad MHB, Murtaza G, Siraj S, Khan SA, Azhar S, Hussain MS, Ismail T, Hussain MS and Hussain I. Anti-venom Evaluation of Herbal Remedies. *African Journal of Pharmacy and Pharmacology* 2011, 5: 2292.

- [46] Okoli CO, Akah PA and Ezugworie U. Anti-inflammatory activity of extracts of root barks of *Securidaca longipedunculata* Fres (Polygalaceae). *African Journal of Traditional, Complementary and Alternative Medicine* 2005, 2(3): 54-63.
- [47] Okon JE, Esenowo GJ, Etim GJ and Umoh NS. Phytochemical Screening and Haemopoetic Study of the Ethanolic Root Extract of *Baphia nitida* (Lodd) on Albino Rats. *International Journal of Modern Biology Medicine* 2017, 3(1): 60-68.
- [48] Yeo D, Dinica R, Yapi HF, Furdui B, Praisler M, Djaman AJ and N'Guessan JD. Evaluation of the anti-inflammatory activity and phytochemical screening of *Annona senegalensis* leaves. *Thérapie* 2011, 66(1): 73–80.
- [49] Jada M, Usman W and Olabisi A. Crude Flavonoids Isolated from the Stem Bark of *Annona senegalensis* have Antimicrobial Activity. *Journal of Advances in Biology & Biotechnology* 2015, 2(1): 24–29.
- [50] Afolabi F and Afolabi O. Phytochemical Constituents of Some Medicinal Plants in South West, Nigeria. *International Research Journal of Applied Chemistry* 2013, 4(1): 76-78.
- [51] Ijaiya I, Arzika S and Abdulkadir M. Extraction and Phytochemical Screening of the Root and Leave of *Annona senegalensis* (Wild Custard Apple). *Academic Journal of Interdisciplinary Studies* 2014, 3(7): 9.
- [52] You M, Wickramaratne D, Silva G, Chai H, Chagwedera T and Farnsworth N. Roemerine, an Aporphine Alkaloid from *Annona senegalensis* that Reverses the Multidrug-Resistance Phenotype with Cultured Cells. *Journal of Natural Products* 1995, 58(4): 598–604.
- [53] Fall D, Sambou B, Seck M, Wélé A, Ndoye I and Gleye C. Enhancing the Anthelmintic Activity Roots of *Annona senegalensis*. *Dakar Medicinal Journal* 2008, 53(1): 61-67.
- [54] Audu SA, Mohammed I and Kaita HA. Phytochemical screening of the leaves of *Lophira lanceolata* (Ochanaceae). *Life Science Journal* 2007, 4(4):75-79.
- [55] Makut MD, Gyar SD, Pennap GRI and Anthony P. Phytochemical screening and anti-microbial activity of the ethanolic and methanolic extracts of the leaf and barks of *Khaya senegalensis*. *African Journal of Biotechnology* 2008, 7: 1216-1219.
- [56] Komes M, Toure A, Karamoko O and Adams C. Phytochemical Composition, Antioxidant and Antibacterial Activities of Root of *Uvaria chamae* P. Beauv. (Annonaceae) Used in Treatment of Dysentery in North of Côte d'Ivoire. *International Journal of Pharmacognosy and Phytochemistry Research* 2015, 7(6): 1047-1053.
- [57] Iroabuchi F. Phytochemical Constituents of *Clerodendron splendens* (A. Cheval) and *Uvaria chamae* (P. Beauv), M. Sc. Thesis, Michael Okpara University of Agriculture, Umudike, Nigeria 2004, 3-16.
- [58] Keles A, Koca I and Gencelep H. Anti-oxidant Properties of Wild Edible Mushrooms. *Journal of Food Processing and Technology* 2012, 2(1): 6.
- [59] Okwu DE. Phytochemicals, vitamins and mineral content of Nigerian medicinal plants. *International Journal of Molecular Medicine and Advanced Sciences* 2005, 1: 375-381.

- [60] Hamzah RU, Jigam AA, Makun HA and Egwim EC. Anti-oxidant Properties of Selected African Vegetables, Fruits and Mushrooms: A Review. *Mycotoxin and Food Safety in Developing Countries*, Dr. Hussaini Makun (Ed.), ISBN: 978-953-511096-5, InTech 2013, DOI: 10.5772/52771.
- [61] Onacha PA and Ali MS. Antileishmaniasis, Phytotoxicity and Cytotoxicity of Nigerian Euphorbiaceous Plants 2: *Phyllanthus amarus* and *Phyllanthus muellerianus* Extracts. *African Scientist* 2010, 11(2): 79-83.
- [62] Sarker SD and Nahar L. *Chemistry for Pharmacy Students. General, Organic and Natural Product Chemistry*. England: John Wiley and Sons 2007, 283-359.
- [63] Sarkhel S, Chakravarthy AK, Das R and Gomes A. A snake venom neutralising factor from the root extract of *Emblca officinalis* Linn. *Oriental Pharmacy Experimental Medicine* 2011, 11(1): 25-33.
- [64] Van-Wyk BE, Van-Oudtshoorn B and Gericke N. *Scientific Implications of Medicinal Plants. Medicinal plants of South Africa. Second Ed.* Briza Publications, Pretoria 2009, 268.
- [65] Vinod M, Verma H, Thakur R, Jain P and Shrivatava N. Medicinal Plants of Chhattisgarh with Anti-Snake Venom Property. *International Journal of Current Pharmaceutical Review and Research* 2012, 3(2): 1-10.
- [66] Omale J, Ebiloma UG and Idoko GO. In vivo neutralisation of *Naja nigricollis* venom by *Uvaria chamae*, *American Journal of Biochemistry and Biotechnology* 2013, 9 (3): 224-234.
- [67] Omodara NB, Amoko JS, Obijole OA and Ojo BM. Invaded and Ultraviolet Spectroscopic Analysis of Methanol Extract of *Phyllanthus muellerianus* Root. *Greener Journal of Physical Sciences* 2013, 3(4): 159-164.
- [68] Parajuli S, Tilja P, Parajuli N and Jamakattel-Pandit N. Antioxidant Activity, Total Phenol and Flavonoid Contents in Some Selected Medicinal Plants of Nepal. *Journal of Health and Applied Science* 2012, 2(1): 27-31.
- [69] Pereañez JA, Patiño AC, Núñez V and Osorio E. The Biflavonoid Morelloflavone inhibits the Enzymatic and Biological Activities of a Snake venom phospholipase A2. *Chemical and Biological Interaction* 2014, 220: 94-101.
- [70] Carvalho BMA, Santos JDL, Xavier BM, Almeida JR, Resende LM, Martins W et al. Snake Venom PLA2s Inhibitors Isolated from Brazilian Plants. *Synthetic and Natural Molecules. BioMedical Research International* 2013, 13(1): 4-6. <http://dx.doi.org/10.1155/2013/153045>.
- [71] Adzu B, Abubakar MS, Izebe KS, Akumka DD and Gamaniel KS. Effect of *Annona senegalensis* root bark extracts on *Naja nigricollis nigricollis* venom in rats. *Journal of Ethnopharmacology* 2004, 96: 507-513.
- [72] Wannang NN, Wudil AM, Dapar ML and Bichi LA. Evaluation of Antisnake venom activity of the aqueous root extract of *Securidaca longipedunculata* in rats. *Journal of Pharmacy and Bioresources* 2005, 3(1): 60-64. <https://doi.org/10.4314/jpb.v2i2.32067>.
- [73] Amlabu E, Ajagun E and Wandayi A. Detoxification of *Echis ocellatus* Venom Induced Toxicity by *Annona senegalensis* Pers. *Journal of Comparative and Integrated Medicine* 2014, 11(2): 93-97.

[74] Saswat GMG, Shah N, Nair S and Chowdhary A. In Vitro Neutralisation Effect of *Acorus calamus* and *Hibiscus rosa-sinensis* Extracts on Viper venom. *International Journal of Research and Review* 2016, 3(6): 47-55.

[75] Singh P, Yasir M, Tripathi MK and Shrivastava R. A Review on In-vitro Screening Assay for Inhibitory effect against Venom enzymes using Medicinal plants. *Toxicological International Journal* 2017, 23(3): 207-211.

[76] Soares AM, Tielia FK, Marcussi S, Lourenço MV, Januário AH and Sampaio SV. Medicinal Plants with Inhibitory Properties Against Snake Venoms. *Current Medicinal Chemistry* 2005, 12: 2625-2641.

[77] Traore MS, Baldé MA, Diallo MS, Baldé ES, Diané S and Camara A. Ethnobotanical survey on Medicinal Plants used by Guinean traditional healers in the treatment of Malaria. *Journal of Ethnopharmacology* 2013, 150(3): 1145–1153.

[78] Tshisikhawe MP, Van-Rooyen MW and Bhat RB. An Evaluation of the extent and threat of bark harvesting of Medicinal Plant species in the Venda region, Limpopo Province, South Africa. *Journal of Phyton* 2012, 81: 89-100.

ACKNOWLEDGEMENTS

Author's sincere appreciation goes to Mr Theophilus Momoh Boniface (project supervisor) and Mr Olusegun Olupinyo (project's chief lab technician) for their support towards this manuscript. I also thank the local communities, in particular the traditional healers of Ankpa LGA, for their warm welcome and for sharing their knowledge.