



LJMU Research Online

Woodward, J, Tooby, J, Tucker, R, Falvey, ÉC, Salmon, DM, Starling, L and Tierney, G

Instrumented mouthguards in elite-level men's and women's rugby union: characterising tackle-based head acceleration events

<http://researchonline.ljmu.ac.uk/id/eprint/23926/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Woodward, J, Tooby, J, Tucker, R, Falvey, ÉC, Salmon, DM, Starling, L and Tierney, G (2024) Instrumented mouthguards in elite-level men's and women's rugby union: characterising tackle-based head acceleration events. *BMJ Open Sport & Exercise Medicine*. 10 (3).





LJMU has developed [LJMU Research Online](#) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

Instrumented mouthguards in elite-level men's and women's rugby union: characterising tackle-based head acceleration events

James Woodward ¹, James Tooby,² Ross Tucker ³, Éanna C Falvey ^{4,5}, Danielle M Salmon,^{4,6} Lindsay Starling,⁴ Gregory Tierney ⁷

To cite: Woodward J, Tooby J, Tucker R, *et al*. Instrumented mouthguards in elite-level men's and women's rugby union: characterising tackle-based head acceleration events. *BMJ Open Sport & Exercise Medicine* 2024;**10**:e002013. doi:10.1136/bmjsem-2024-002013

► Additional supplemental material is published online only. To view, please visit the journal online (<https://doi.org/10.1136/bmjsem-2024-002013>).

Accepted 15 July 2024



© Author(s) or their employer(s) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹School of Sport, Ulster University, Belfast, UK

²Carnegie Applied Rugby Research (CARR) centre, Leeds Beckett University, Leeds, UK

³Institute of Sport and Exercise Medicine, University of Stellenbosch, Stellenbosch, South Africa

⁴World Rugby Limited, Dublin, Ireland

⁵Department of Medicine, University College Cork, Cork, Ireland

⁶Auckland University of Technology, Auckland, New Zealand

⁷Ulster University, Coleraine, UK

Correspondence to

James Woodward;
woodward-j2@ulster.ac.uk

ABSTRACT

Objectives To examine the propensity of tackle height and the number of tacklers that result in head acceleration events (HAEs) in elite-level male and female rugby tackles.

Methods Instrumented mouthguard data were collected from women (n=67) and men (n=72) elite-level rugby players from five elite and three international teams. Peak linear acceleration and peak angular acceleration were extracted from HAEs. Propensities for HAEs at a range of thresholds were calculated as the proportion of tackles/carries that resulted in an HAE exceeding a given magnitude for coded tackle height (low, medium, high) and number of tacklers. Propensity ratios with 95% CIs were calculated for tackle heights and number of tacklers.

Results High tackles had a 32.7 (95% CI=6.89 to 155.02) and 41.2 (95% CI=9.22 to 184.58) propensity ratio to cause ball carrier HAEs>30g compared with medium tackles for men and women, respectively. Low tackles had a 2.6 (95% CI=1.91 to 3.42) and 5.3 (95% CI=3.28 to 8.53) propensity ratio to cause tackler HAEs>30g compared with medium tackles for men and women, respectively. In men, multiple tacklers had a higher propensity ratio (6.1; 95% CI=3.71 to 9.93) than singular tacklers to cause ball carrier HAEs>30g but a lower propensity ratio (0.4; 95% CI=0.29 to 0.56) to cause tackler HAEs>30g. No significant differences were observed in female tacklers or carriers for singular or multiple tacklers.

Conclusion To limit HAE exposure, rule changes and coaching interventions that promote tacklers aiming for the torso (medium tackle) could be explored, along with changes to multiple tackler events in the male game.

INTRODUCTION

Rugby union is a contact sport that involves frequent high-intensity contact events (tackles, carries and rucks), resulting in a high injury risk.¹ There is growing concern around the relationship between head impacts and their potential short and long-term health risks.²⁻⁴ A head acceleration event (HAE) is defined as 'an event/incident that gives rise to an acceleration response of the head caused by an external short-duration collision force

WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ At the elite (professional) level, the tackle has the highest injury rate of all contact events in rugby union, with most concussions occurring during this contact event.
- ⇒ There is a growing concern surrounding exposure to high-magnitude and repeated head acceleration events (HAEs).
- ⇒ Law trials and modifications can improve player safety by mitigating risk factors for head injuries and exposure to HAEs.

WHAT THIS STUDY ADDS

- ⇒ An understanding of tackle characteristics, such as tackle height and the number of tacklers and how these factors influence HAE propensities in elite male and female rugby union players.
- ⇒ Tackle heights above the sternum of the ball-carrier had a greater propensity to result in HAEs>30g for both male and female ball-carriers. Low tackle heights had a higher propensity to result in HAEs>30g for male and female tacklers.
- ⇒ Tackles made by multiple players had a higher propensity to result in HAEs>30g for male ball-carriers. Tackles involving a singular tackler had a lower propensity to result in HAEs>30g for male tacklers. No significant differences were observed for tackle events that involved singular or multiple tacklers for female players.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- ⇒ These findings could contribute to developing improved coaching and officiating practice to help mitigate player exposure to HAEs.
- ⇒ To limit HAE exposure, rule changes and coaching interventions that promote tacklers aiming for the torso (medium tackle) could be explored, along with changes to multiple tackler events in the male game.

applied directly to the head or indirectly via the body in sport, recreational, military, or other activities of interest'.⁵

Repetitive HAEs have identified as a potential contributor to concussive injury

mechanisms and long-term brain health issues.⁶⁻⁸ Current research has yet to identify a specific magnitude or threshold of HAEs that will result in immediate or delayed clinical presentation consistent with concussion. Still, it has been proposed that exposure to repetitive HAEs should be reduced as far as possible to improve player welfare.^{9 10} Studies have used instrumented mouthguards (iMGs) to measure match HAE incidence and propensity in rugby players.¹¹⁻¹⁴ These studies highlighted tackle and carry as the match activities most likely to cause HAEs in male and female rugby players.^{11 12 14 15} The tackle in rugby union has the highest reported risk for general injury.^{1 16 17}

Analysing tackles resulting in a head injury has been common practice to identify law modifications and coaching interventions that may reduce their prevalence.^{3 18-20} Such research has identified factors such as the height of tackles and upright body positions of players at the time of contact as risk factors for head injuries.^{3 21 22} High tackles, where the tackler makes contact above the ball-carrier's sternum, often occur when the tackler is in an upright body position, resulting in a greater risk of head injury.^{3 22} These findings led to trials of law modifications to reduce the legal tackle height at the community level by World Rugby²³⁻²⁶ and their subsequent integration into community-level rugby in England by the Rugby Football Union.²⁷ Given the role of HAEs as a concussive injury mechanism, it is important to consider how law changes and trials may play a role in HAE exposure mitigation.⁶⁻⁸ Other research has identified an increased injury risk when multiple tacklers are involved in the tackle.^{17 28 29} Governing bodies such as the French Federation of Rugby (FFR) have trialled laws to reduce the number of tacklers involved in a tackle, in combination with a reduced legal tackle height, resulting in decreases in head-to-head contact.²⁷ However, little is known about how these law modifications can impact on HAEs experienced by male or female players.

It is important to consider law modifications and other interventions that may reduce exposure to the frequency and magnitude of HAEs.⁶ Identifying risk factors during the tackle, such as tackle height and a number of tacklers, presents an opportunity to gain insights into HAE aetiology. This study explored the propensity of HAEs sustained during the tackle event in elite male and female rugby matches by exploring tackle height and the number of tacklers involved in each tackle event.

METHODS

Study design and participants

A prospective observational cohort study was undertaken involving elite-level (professional) rugby union women (n=67) and men (n=72) players from European and international rugby competitions, including the Allianz Premier-15s (n=11), Gallagher Premiership (n=21), Top 14 (n=6) and United Rugby Championship (n=45), as well as Women's Autumn International Series in 2021 (n=13) and the Women's Rugby World Cup in

2022 (n=43). Video footage was obtained through Stats Perform (Chicago, Illinois, USA) and World Rugby.

All participants were equipped with custom-fit Prevent Biometrics (Prevent Biometrics, Minneapolis, Minnesota, USA) iMGs, following three-dimensional dental scans with a qualified dentist. These iMGs are equipped with triaxial gyroscopes and accelerometers, sampling at 3200 Hz with a measurement range of ± 35 rad/s and ± 200 g, respectively. Linear head kinematics collected at the iMG sensor location were transformed to the head centre of gravity.¹¹ Previous studies have validated the Prevent Biometrics iMGs, both in field and laboratory settings.^{15 30-32} HAE were identified when the linear acceleration exceeded an 8g trigger threshold on a single axis of the iMG accelerometer.¹² Time series data for HAEs were captured 10 ms before and 40 ms after the trigger event, with a recording threshold of 5g at the head centre of gravity. The level of noise/artefact in each kinematic signal was classified by an in-house Prevent Biometrics algorithm as minimal (class 0), moderate (class 1) or severe (class 2). A 4-pole (2x2), zero-phase, low-pass Butterworth filter was applied to each kinematic signal. A 200 Hz, 100 Hz and 50 Hz cut-off frequency (-6 dB) were applied to class 0, class 1 and class 2 signals, respectively.¹¹ Each iMG device was manually synchronised to video footage for each match to ensure to-the-frame synchronisation for analysis.

Tackles and carries were analysed from 304 player matches (161 men, 143 women) across 28 matches. Proximity sensor readings were used to verify that the iMG was worn during each contact event. Video analysis identified 2266 tackles, defined as tackler involvements and 1426 carries with associated HAEs. Tackle involvements and carries were then analysed in accordance with Tooby *et al*¹² and Woodward *et al*,¹¹ categorising the contact event and its HAE stage (initial collision, secondary contact, ground impact or breakdown) for each tackler in the tackle event.

Video analysis

All contact events in which the proximity sensors indicated the iMG was being worn were video verified and analysed via NacSport Scout Plus (V.7.0.0) by an experienced coder (3 years of rugby video analysis experience). The video had a minimum frame rate of 25 frames per second and 1080p resolution. All contact events were coded using an adapted framework from existing analysis frameworks by Tucker *et al*,²⁰ Davidow *et al*³³ and Hendricks *et al*,³⁴ see [table 1](#). The analysis framework was also adapted from law trial definitions by the FFR and World Rugby.²⁷ Intrarater reliability was conducted and Cohen's kappa calculated, see [table 1](#), on a randomly selected match, 1-month post-initial analysis in line with previous research, where a value of 0.8 or above indicates near-perfect agreement.³⁵⁻³⁷ All coded events were stored in an encrypted Microsoft Excel database and analysed using MATLAB (MATLAB_R2024a).

Table 1 Analysis framework variables and descriptors for tackle and carry head acceleration event and non-instrumented mouthguards triggered events, with intrarater reliability scores

Variable	Descriptor
Tackle height Intrarater reliability: 0.91	High – contact from the tackler is <u>above</u> the sternum, head/neck and shoulder/chest of the ball-carrier. Medium – contact is <u>below</u> the sternum and <u>above</u> the waistband of the ball-carrier's shorts (torso). Low – contact is <u>below</u> the waistband of the shorts (hip, upper leg and lower leg).
Number of tacklers Intrarater reliability: 0.95	Single – single tackler. Multiple (2+) – two or more tacklers.

Statistical analysis

Given that multiple HAEs may occur during each contact event, the HAEs used in the calculations in this study were the maximum from the initial collision HAE stage.^{11–13} Propensity was calculated across a range of peak linear acceleration (PLA) and peak angular acceleration (PAA) thresholds by dividing the number of tackle involvements that resulted in an HAE by the total number of tackle involvements that the individual was involved in while wearing the iMG, confirmed using the iMG infrared proximity sensor.¹² The total number of tackle involvements included the number of events resulting in an HAE, as well as the events that did not result in an HAE. Mean values across players and 95% CIs were calculated using a bootstrapping procedure, similar to Tooby *et al.*¹² HAEs were classified into lower (PLA<10g and PAA<1.0 krad/s²), medium (PLA>10g and <30g and PAA>1.0 krad/s²) and higher magnitude (PLA≥30g and PAA≥2.0 krad/s²) groups, to calculate a propensity ratio for tackle heights and number of tacklers for tackler and ball-carrier HAEs with statistical significance set at p<0.05 and non-overlapping 95% CIs.^{12 14}

s² and <2.0 krad/s²) and higher magnitude (PLA≥30g and PAA≥2.0 krad/s²) groups, to calculate a propensity ratio for tackle heights and number of tacklers for tackler and ball-carrier HAEs with statistical significance set at p<0.05 and non-overlapping 95% CIs.^{12 14}

RESULTS

Tackle height

Figure 1 shows the propensity of high, medium and low tackle heights to result in an HAE exceeding a given range of magnitude thresholds for male and female ball-carriers and tacklers. In males, high tackles had a propensity of 4.1% and 8.7%, resulting in an HAE>30g for the tackler and ball-carrier, respectively. In HAEs>30g, medium tackles had a propensity of 4.0% for the tackler and 0.3% for the ball-carrier, while low tackles had a propensity of 10.1% for the tackler and 1.1% for the




Figure 1 The propensity of tackles at different tackle heights (high, medium or low) to result in an HAE across peak linear acceleration thresholds for male and female tacklers and carriers. Shaded regions indicate the 95% CI. The number of tackles at each tackle height is shown as n. Online supplemental figure 1 presents the results with a peak angular acceleration recording threshold. HAE, head acceleration event.

		Tackle Height Propensity Ratios	PLA Magnitude Bands		
			Lower <10 g	Medium 10 - 30 g	Higher ≥30 g
Tackler	Male	High vs Medium	0.95	1.06	1.03
		High vs Low	0.89	0.72	0.40*
		Low vs Medium	1.06	1.47	2.55*
	Female	High vs Medium	0.99	1.09	1.86
		High vs Low	0.85	0.71	0.35
		Low vs Medium	1.16	1.54	5.29*
Carrier	Male	High vs Medium	2.16*	10.86*	32.69*
		High vs Low	1.61*	4.15*	7.76*
		Low vs Medium	1.34	2.62	4.22
	Female	High vs Medium	1.56*	3.57*	41.24*
		High vs Low	1.28	3.03*	3.54
		Low vs Medium	1.22	1.18	11.67

Figure 2 Propensity ratios of male and female tackler and carrier HAEs caused by tackle heights within lower, medium and higher HAE magnitude bands. Comparisons between tackle heights are included, where *denotes statistical significance. Online supplemental figure 2 presents the results with peak angular acceleration magnitude bands. HAE, head acceleration event.

ball-carrier. In females, the propensity of high tackles to result in an HAE>30 g was 2.8% for tacklers and 5.6% for ball-carriers, while medium tackles had a propensity of 1.5% and 0.1% and low tackles had a propensity of 8.0% and 1.6%, respectively.

High tackles had a 32.7 (95% CI=6.89 to 155.02) propensity ratio to cause male ball carrier HAEs>30 g compared with medium tackles (figure 2). Similarly, female ball-carriers had a 41.2 (95% CI=9.22 to 184.58) propensity ratio to experience an HAE>30 g from a high tackle compared with a medium tackle. High tackles had a 0.4 (95% CI=0.30 to 0.54) propensity ratio to cause male tackler HAEs>30 g compared with low tackles. Low tackles had a 2.6 (95% CI=1.91 to 3.42) propensity ratio to cause male tackler HAEs>30 g compared with medium tackles. In female tacklers, low tackles had a 5.3 (95% CI=3.28 to 8.53) propensity ratio to result in HAEs>30 g compared with medium tackles.

Number of tacklers in a tackle event

Figure 3 shows the propensity of singular and multiple tacklers involved in a tackle event to produce an HAE exceeding a given range of magnitude thresholds in male and female ball-carriers and tacklers. In males, 3.5% and 7.2% of tackles involving multiple tacklers caused HAEs above 30 g in tacklers and ball-carriers, respectively. Singular tacklers had an HAE>30 g propensity of 8.7% for the tackler and 1.2% for the ball-carrier.

In males, multiple tacklers had a significantly higher propensity ratio (6.1; 95% CI=3.71 to 9.93) (figure 4) to cause ball carrier HAEs>30 g than singular tacklers. Conversely, multiple tacklers had a significantly lower propensity ratio (0.4; 95% CI=0.29 to 0.56) to produce

HAEs>30 g compared with singular tacklers for male tacklers. In female players, there were no significant differences between single tackler events and multiple tackler events for either the ball-carrier or tackler.

DISCUSSION

This study is the first to explore how tackle height, and the number of tacklers influence the propensity of HAEs in the tackler and ball-carrier. Given the emphasis on HAE exposure reduction, it is important to explore these factors so that strategies can be targeted to reduce the incidence and magnitude of HAEs resulting from tackles in the sport.

Tackle height

Our first important finding is that tackle height significantly affects HAE propensity and does so differently in the ball-carrier and the tackler. For ball-carriers, tackles above the sternum (high) had a greater propensity to result in HAEs>30 g compared with tackles at the torso (medium tackle) in male and female carriers, respectively. These findings align with previous head injury assessment (HIA) and concussion research, where tackles above the sternum of the ball-carrier have shown an increased risk of HIA removal and concussion. However, this was true for both ball carriers and tacklers.^{3 20 28} This reinforces the potentially important role of tackle height laws in minimising the frequency of tackles above the sternum to mitigate ball carrier HAEs and head injuries to both players.

It is important to consider our finding that tacklers have a greater HAE propensity when tackling low compared with medium and high.¹² This contrasts with previous literature on HIA and concussion risk, which showed increased tackler injury risk when high tackles were made.^{3 20 28} While not as pronounced as the differences found for the ball-carrier, the greater propensity for >30 g HAEs when making low compared with medium tackles are of interest because they suggest a difference in HAE versus HIA removal risk characteristics. The mechanism for this difference is unclear, but further exploration is required since interventions to reduce HIA and concussion incidence may not successfully reduce HAE incidence, and vice versa, particularly if the shift to reduce tackle height results in an increased frequency of low tackles.

The trade-off between these two risk factors, where high and low tackles affect ball carrier and tackler HAE risk differently, is important for the sport since the objective is the protection of both players. When considering HAE risk to either player, the lowest risk would be achieved from medium (torso) tackles, which were least likely to cause HAEs exceeding 30 g in both male and female tacklers and ball carriers. The greatest overall propensity was found from high tackles by virtue of the very large increase in ball-carriers HAE propensity, despite the relatively lower but increased propensity for tackler HAEs from low tackles.




Figure 3 The propensity of tackles to result in an HAE across peak linear acceleration thresholds for male and female tacklers and carriers where there were multiple (2+) tacklers or a single tackler. Shaded regions indicate the 95% CI. The number of multiple (2+) or singular tackler tackles is shown as n. Online supplemental figure 3 presents the results with a peak angular acceleration recording threshold. HAE, head acceleration event.

The findings strongly suggest that the target of law interventions or coaching education should focus on medium tackles, which would most effectively reduce HAE prevalence in both male and female tacklers and ball-carriers. This would also align with the reduced

clinical outcome risk highlighted in previous concussion risk factor literature.^{3 20}

Number of tacklers

In a recent law trial, the FFR trialled a combination of lowering the legal tackle height and reducing the number of tacklers involved in the tackle event.²⁷ This trial focused on reducing the incidence of head-to-head contact between players while maintaining or improving enjoyment and participation. Previous literature has investigated the role of the number of tacklers in injurious events, finding that tackles involving multiple tacklers had a higher concussive injury risk than single-tackler events.^{28 29} This is, in part, logically explained by an increase in the number of physical heads involved during each tackle. Our findings support these previous results, with multiple tacklers having a greater propensity to produce an HAE>30 g in male ball-carriers. In contrast, male tacklers had a lower propensity to experience HAEs when involved in tackles with multiple tacklers. This

Player Role	Number of Tacklers	PLA Magnitude Bands		
		Lower <10 g	Medium 10-30 g	Higher ≥30 g
Tackler	Male Multiple vs Singular Tacklers	0.75*	0.68	0.41*
	Female Multiple vs Singular Tacklers	0.86	0.95	1.74
Carrier	Male Multiple vs Singular Tacklers	1.02	1.59	6.07*
	Female Multiple vs Singular Tacklers	1.28	1.61	1.58

Figure 4 Propensity ratios of male and female tackler and carrier HAEs caused by tackler numbers within lower, medium and higher HAE magnitude bands. Comparisons between tackler numbers are included, where *denotes statistical significance. Online supplemental figure 4 presents the results with peak linear acceleration magnitude bands. HAE, head acceleration event; PLA, peak linear acceleration.



may be due to the differing dynamics of multiple tackler events. This was not explored in the current study; however, future research should investigate the interaction between other technical elements of multiple versus singular tackler events and HAE propensity. Overall, the results suggest that limiting the number of tacklers may benefit male ball carriers.

No significant differences were observed between multiple or singular tackler events in the female cohort. These sex differences, which require further exploration, may result from technical elements in the tackle that differ between male and female rugby players.

Limitations

The data in this study was obtained from elite-level male and female rugby players and may not reflect the diversity in quality, quantity and level of rugby internationally, with all these factors potentially affecting HAE risk. Future research may investigate similar tackle characteristics and their propensities to result in HAEs at the community and age grade level of rugby. This study did not explore sociological or gendered factors affecting tackle technique and characteristics. Future research should explore tackle technique and factors that may modify tackle characteristics, including playing age and level, training history and conditioning.³⁸

The filtering of head kinematics was conducted through in-house Prevent Biometrics algorithms.¹² A common open-source signal processing approach would be beneficial for cross-study comparison.³⁹ This study also used an iMG trigger threshold of 8g. A prior simulation study suggested that false negatives may be influenced by a bias introduced through linear acceleration trigger mechanisms.^{12 40} Furthermore, while this study investigated peak resultant head kinematics (PLA, PAA), these do not consider directionality and temporal data, which may play a critical role in injury risk and mechanism.^{12 41} This study included data from a subset of matches in a playing season. Future research should consider HAEs throughout an entire playing season, including in a training environment, to account for potential variance and overall HAE exposure.

CONCLUSION

This study is the first to explore how tackle height, and the number of tacklers influence the propensity of HAEs in the tackler and ball-carrier. Tackle height affects HAE propensity and does so differently in the ball-carrier and the tackler. For ball-carriers, tackles above the sternum (high) had a significantly greater propensity ratio to result in HAEs>30g compared with tackles at the torso (medium tackle) in male and female carriers, respectively. Low tackles had a significantly greater propensity ratio to cause male and female tackler HAEs>30g compared with medium tackles. The current study also identified that tackles involving multiple tacklers had a greater propensity to result in HAEs>30g for male carriers. Conversely, for male tacklers, tackles involving a singular tackler had

a greater propensity to result in HAEs>30g. There were no significant differences in the number of tacklers and HAE exposure for female players. To limit HAE exposure, rule changes and coaching interventions that promote tacklers aiming for the torso (medium tackle) could be explored, along with changes to multiple tackler events in the male game.

X James Woodward @JamesWoodward97 and Ross Tucker @Scienceofsport

Acknowledgements The authors would like to thank all of the players and staff involved in this study for participating and contributing to the research. The authors would like to acknowledge Prevent Biometrics and World Rugby for their support and cooperation during the study.

Contributors All authors conceptualised the research project. GT and JW designed the study. JW, GT, DMS and JT collected data for the study. JW, GT and JT were responsible for the analysis and interpretation of the results. JW drafted the manuscript and is the guarantor. All authors critically reviewed and edited the manuscript prior to submission.

Funding World Rugby funded this research project.

Competing interests GT has received research funding from World Rugby and Prevent Biometrics and has previously conducted consultancy work for World Rugby. LS, RT, ÉCF and DMS are employed by or contracted as consultants for World Rugby.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and was approved by Ulster University's Research and Ethics Committee (UREC), University of Ulster (#REC-21-0061). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. Anonymous data are available at a reasonable request to the corresponding author.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

James Woodward <http://orcid.org/0000-0002-6402-2554>

Ross Tucker <http://orcid.org/0000-0002-4294-4822>

Éanna C Falvey <http://orcid.org/0000-0003-3961-1805>

Gregory Tierney <http://orcid.org/0000-0002-4666-4473>

REFERENCES

- 1 Burger N, Lambert M, Hendricks S. Lay of the land: narrative synthesis of tackle research in rugby union and rugby sevens. *BMJ Open Sport Exerc Med* 2020;6:e000645.
- 2 Tierney G. Concussion biomechanics, head acceleration exposure and brain injury criteria in sport: a review. *Sports Biomech* 2021;2021:1–29.
- 3 Tucker R, Raftery M, Kemp S, et al. Risk factors for head injury events in professional rugby union: a video analysis of 464 head injury events to inform proposed injury prevention strategies. *Br J Sports Med* 2017;51:1152–7.

- 4 Fuller CW, Taylor A, Douglas M, *et al*. Rugby World Cup 2019 injury surveillance study. *S Afr J Sports Med* 2020;32:v32i1a8062.
- 5 Kuo C, Patton D, Rooks T, *et al*. On-field deployment and validation for wearable devices. *Ann Biomed Eng* 2022;50:1372–88.
- 6 Stemper BD, Shah AS, Harezlak J, *et al*. Comparison of head impact exposure between concussed football athletes and matched controls: evidence for a possible second mechanism of sport-related concussion. *Ann Biomed Eng* 2019;47:2057–72.
- 7 Montenigro PH, Allosco ML, Martin BM, *et al*. Cumulative head impact exposure predicts later-life depression, apathy, executive dysfunction, and cognitive impairment in former high school and college football players. *J Neurotrauma* 2017;34:328–40.
- 8 Stern RA, Riley DO, Daneshvar DH, *et al*. Long-term consequences of repetitive brain trauma: chronic traumatic encephalopathy. *PM R* 2011;3:S460–7.
- 9 Ntikas M, Binkofski F, Shah NJ, *et al*. Repeated sub-concussive impacts and the negative effects of contact sports on cognition and brain integrity. *Int J Environ Res Public Health* 2022;19:7098.
- 10 Rowson S, Campoletano ET, Duma SM, *et al*. Accounting for variance in concussion tolerance between individuals: comparing head accelerations between concussed and physically matched control subjects. *Ann Biomed Eng* 2019;47:2048–56.
- 11 Woodward J, Tooby J, Tierney G. Comparing head acceleration events in elite men's and women's rugby union tackle events using instrumented mouthguards and video analysis. IRCOBI Conference Proceedings; 2022:2.
- 12 Tooby J, Woodward J, Tucker R, *et al*. Instrumented mouthguards in elite-level men's and women's rugby union: the incidence and propensity of head acceleration events in matches. *Sports Med* 2024;54:1327–38.
- 13 Tooby J, Weaving D, Al-Dawoud M, *et al*. Quantification of head acceleration events in rugby league: an instrumented mouthguard and video analysis pilot study. *Sensors (Basel)* 2022;22:584.
- 14 Allan D, Tooby J, Starling L, *et al*. The incidence and propensity of head acceleration events in a season of men's and women's English elite-level club rugby union matches. *Sports Med* 2024.
- 15 Tierney G, Weaving D, Tooby J, *et al*. Quantifying head acceleration exposure via instrumented mouthguards (IMG): a validity and feasibility study protocol to inform img suitability for the tackle project. *BMJ Open Sport Exerc Med* 2021;7:e001125.
- 16 Williams S, Trewartha G, Kemp S, *et al*. A meta-analysis of injuries in senior men's professional rugby union. *Sports Med* 2013;43:1043–55.
- 17 Cross M, Kemp S, Smith A, *et al*. Professional rugby union players have a 60% greater risk of time loss injury after concussion: a 2-season prospective study of clinical outcomes. *Br J Sports Med* 2016;50:926–31.
- 18 Raftery M, Tucker R, Falvey ÉC. Getting tough on concussion: how welfare-driven law change may improve player safety—a rugby union experience. *Br J Sports Med* 2021;55:527–9.
- 19 Stokes KA, Locke D, Roberts S, *et al*. Does reducing the height of the tackle through law change in elite men's rugby union (the championship, England) reduce the incidence of concussion? A controlled study in 126 games. *Br J Sports Med* 2021;55:220–5.
- 20 Tucker R, Raftery M, Fuller GW, *et al*. A video analysis of head injuries satisfying the criteria for a head injury assessment in professional rugby union: a prospective cohort study. *Br J Sports Med* 2017;51:1147–51.
- 21 Tierney GJ, Simms CK. Can tackle height influence head injury assessment risk in elite rugby union? *J Sci Med Sport* 2018;21:1210–4.
- 22 Quarrie KL, Hopkins WG. Tackle injuries in professional rugby union. *Am J Sports Med* 2008;36:1705–16.
- 23 Tonder R van, Starling L, Surmon S, *et al*. 11.35 effectiveness of a lowering in legal tackle height to reduce sport-related concussion in amateur adult male rugby. 6th International Conference on Concussion in Sport; 2024
- 24 van Tonder R, Brown JC, Surmon S, *et al*. Stakeholder perceptions of a tacklelaw variation to reduce concussion incidence in community rugby union: a qualitative study. *Int J Sports Sci Coach* 2024;17479541241227329.
- 25 van Tonder R, Starling L, Surmon S, *et al*. Tackling sport-related concussion: effectiveness of lowering the maximum legal height of the tackle in amateur male rugby - a cross-sectional analytical study. *Inj Prev* 2023;29:56–61.
- 26 van Tonder R, Hendricks S, Starling L, *et al*. Tackling the tackle 1: a descriptive analysis of 14,679 tackles and risk factors for high tackles in a community-level male amateur rugby union competition during a lowered tackle height law variation trial. *J Sci Med Sport* 2024;27:57–62.
- 27 World Rugby. Lower tackle height law trial introduction. Available: <https://passport.world.rugby/injury-prevention-and-risk-management/tackle-ready/lower-tackle-height-law-trial-introduction/> [Accessed 1 Dec 2023].
- 28 Cross MJ, Tucker R, Raftery M, *et al*. Tackling concussion in professional rugby union: a case-control study of tackle-based risk factors and recommendations for primary prevention. *Br J Sports Med* 2019;53:1021–5.
- 29 Hislop MD, Stokes KA, Williams S, *et al*. Reducing musculoskeletal injury and concussion risk in schoolboy rugby players with a pre-activity movement control exercise programme: a cluster randomised controlled trial. *Br J Sports Med* 2017;51:1140–6.
- 30 Kieffer EE, Begonia MT, Tyson AM, *et al*. A two-phased approach to quantifying head impact sensor accuracy: in-laboratory and on-field assessments. *Ann Biomed Eng* 2020;48:2613–25.
- 31 Liu Y, Domel AG, Yousefsani SA, *et al*. Validation and comparison of instrumented mouthguards for measuring head kinematics and assessing brain deformation in football impacts. *Ann Biomed Eng* 2020;48:2580–98.
- 32 Jones B, Tooby J, Weaving D, *et al*. Ready for impact? A validity and feasibility study of instrumented mouthguards (IMGs). *Br J Sports Med* 2022. 10.1136/bjsports-2022-105523. [Epub ahead of print 25 Jul 2022].
- 33 Davidow D, Quarrie K, Viljoen W, *et al*. Tackle technique of rugby union players during head impact tackles compared to injury free tackles. *J Sci Med Sport* 2018;21:1025–31.
- 34 Hendricks S, Till K, den Hollander S, *et al*. Consensus on a video analysis framework of descriptors and definitions by the rugby union video analysis consensus group. *Br J Sports Med* 2020;54:566–72.
- 35 Ross A, Gill N, Cronin J, *et al*. The relationship between physical characteristics and match performance in rugby sevens. *Eur J Sport Sci* 2015;15:565–71.
- 36 Menzies FA, Walsh J, Boyd AJ, *et al*. Assessment of contact involvements and scrums in international rugby union match-play using video analysis and microsensor technology methods. *J Sports Sci* 2023;41:2129–37.
- 37 Cohen J. A coefficient of agreement for nominal scales. *Educ Psychol Meas* 1960;20:37–46.
- 38 Dane K, Foley G, Wilson F. “Body on the line”: experiences of tackle injury in women's rugby union - a grounded theory study. *Br J Sports Med* 2023;57:1476–83.
- 39 Tierney G, Rowson S, Gellner R, *et al*. Head exposure to acceleration database in sport (HEADSport): a kinematic signal processing method to enable instrumented mouthguard (IMG) field-based inter-study comparisons. *BMJ Open Sport Exerc Med* 2024;10:e001758.
- 40 Wang T, Kenny R, Wu LC. Head impact sensor triggering bias introduced by linear acceleration thresholding. *Ann Biomed Eng* 2021;49:3189–99.
- 41 Gabler LF, Crandall JR, Panzer MB. Development of a second-order system for rapid estimation of maximum brain strain. *Ann Biomed Eng* 2019;47:1971–81.