



LJMU Research Online

Zeng, S, Jimenez-Serra, I, Cosentino, G, Viti, S, Barnes, AT, Henshaw, JD, Caselli, P, Fontani, F and Hily-Blant, P

15N Fractionation in Infrared-Dark Cloud Cores

<http://researchonline.ljmu.ac.uk/id/eprint/6443/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Zeng, S, Jimenez-Serra, I, Cosentino, G, Viti, S, Barnes, AT, Henshaw, JD, Caselli, P, Fontani, F and Hily-Blant, P (2017) 15N Fractionation in Infrared-Dark Cloud Cores. *Astronomy and Astrophysics* (A22). ISSN 0004-6361

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

^{15}N Fractionation in Infrared-Dark Cloud Cores

S. Zeng¹, I. Jiménez-Serra¹, G. Cosentino², S. Viti², A. T. Barnes^{3,4}, J. D. Henshaw³, P. Caselli⁴, F. Fontani⁵, P. Hily-Blant⁶

¹ School of Physics and Astronomy, Queen Mary University of London, Mile End Road, E1 4NS London, United Kingdom

² University College London, 132 Hampstead Road, London, NW1 2PS, UK

³ Astrophysics Research Institute, Liverpool John Moores University, Liverpool, L3 5RF, UK

⁴ Max-Planck Institute for Extraterrestrial Physics, Giessenbachstrasse 1, 85748 Garching, Germany

⁵ INAF-Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, 50125, Firenze, Italy

⁶ Institut de Planétologie et d'Astrophysique de Grenoble, 414 rue de la Piscine, F-38041 Grenoble, France

Received date / Accepted date

ABSTRACT

Context. Nitrogen is one of the most abundant elements in the Universe and its $^{14}\text{N}/^{15}\text{N}$ isotopic ratio has the potential to provide information about the initial environment in which our Sun formed. Recent findings suggest that the Solar System may have formed in a massive cluster since the presence of short-lived radioisotopes in meteorites can only be explained by the influence of a supernova.

Aims. To determine the $^{14}\text{N}/^{15}\text{N}$ ratio towards a sample of cold, massive dense cores at the initial stages in their evolution.

Methods. We have observed the $J=1\rightarrow 0$ transitions of HCN, H^{13}CN , HC^{15}N , HN^{13}C and H^{15}NC toward a sample of 22 cores in 4 Infrared-Dark Clouds (IRDCs). IRDCs are believed to be the precursors of high-mass stars and star clusters. Assuming LTE and a temperature of 15 K, the column densities of HCN, H^{13}CN , HC^{15}N , HN^{13}C and H^{15}NC are calculated and their $^{14}\text{N}/^{15}\text{N}$ ratio is determined for each core.

Results. The $^{14}\text{N}/^{15}\text{N}$ ratio measured in our sample of IRDC cores range between ~ 70 and ≥ 763 in HCN and between ~ 161 and ~ 541 in HNC. They are consistent with the terrestrial atmosphere (TA) and protosolar nebula (PSN) values, and with the ratios measured in low-mass pre-stellar cores. However, the $^{14}\text{N}/^{15}\text{N}$ ratios measured in cores C1, C3, F1, F2 and G2 do not agree with the results from similar studies toward the same massive cores using nitrogen bearing molecules with nitrile functional group (-CN) and nitrogen hydrides (-NH) although the ratio spread covers a similar range.

Conclusions. Amongst the 4 IRDCs we measured relatively low $^{14}\text{N}/^{15}\text{N}$ ratios towards IRDC G which are comparable to those measured in small cosmomaterials and protoplanetary disks. The low average gas density of this cloud suggests that the gas density, rather than the gas temperature, may be the dominant parameter influencing the initial nitrogen isotopic composition in young PSN.

Key words. ISM: molecules – Astrochemistry – Stars: formation

1. Introduction

For decades, our Solar System was believed to have resided in a relatively isolated, low-mass molecular cloud core during its formation. However the detection of short-lived radioactive species in meteorites have suggested a different scenario in which the birthplace of the Sun may have been a massive cluster affected by a supernova event (Adams 2010; Dukes & Krumholz 2012; Pfalzner 2013; Nicholson & Parker 2017). If so, the initial chemical composition of our Solar System, and thus of planets, meteorites and comets, may have been affected by the same physical process.

Measurements of the abundance isotopic ratios of the elements can be used to unveil the initial chemical composition of the proto-solar nebulae (PSN) from which the Solar System formed. The isotopic ratios of carbon ($^{12}\text{C}/^{13}\text{C}$) and oxygen ($^{16}\text{O}/^{18}\text{O}$) show a remarkable agreement among cometary materials, the local interstellar medium (ISM) and the Solar value (Manfroid et al. 2009; Milam et al. 2005; Wilson & Rood 1994). Nitrogen, by contrast, has a peculiar behaviour since its $^{14}\text{N}/^{15}\text{N}$ isotopic ratio exhibits discrepancies across various environments within the Solar System. The $^{14}\text{N}/^{15}\text{N}$ ratio measured in Jupiter's atmosphere (450 ± 100 , Fouchet et al. 2004) is considered as the most representative value of the PSN and it matches the present

day solar wind value (441 ± 6 , Marty et al. 2010). However, the ratios measured in the terrestrial atmosphere (TA) (~ 272 in Earth, Junk & Svec 1958; 272 ± 54 in Venus, Hoffman et al. 1979; 173 ± 11 in Mars, Wong et al. 2013), comets (147.8 ± 5.7 , Manfroid et al. 2009, 139 ± 26 from HCN and 165 ± 40 from CN, Bockelée-Morvan et al. 2008), Interplanetary Dust Particles (or IDPs; values of 180-305; Floss et al. 2006) and meteorites (192-291, Alexander et al. 2007), are lower than that measured in Jupiter's atmosphere.

In molecular clouds, the discrepancies in the $^{14}\text{N}/^{15}\text{N}$ isotopic ratios spread over a larger range. In contrast to many molecular species (e.g. CO), N-bearing molecules do not suffer significant freeze-out onto grains in the coldest, densest regions of IRDCs, and are therefore reliable tracers of the gas chemistry and kinematics in cores. The nitrogen fractionation mechanisms are either due to chemical fractionation (Terzieva & Herbst 2000; Rodgers & Charnley 2008; Wirström et al. 2012; Hily-Blant et al. 2013) or selective photodissociation effect (Lyons et al. 2009; Heays et al. 2014). IRDCs are dense and highly extinguished (with visual extinctions > 10 mag; Kainulainen & Tan 2013), and therefore selective photodissociation is not expected to play an important role since this process becomes inefficient at $A_v \geq 3$ mag (Heays et al. 2014). As for chemical fractionation, these tracers can be cat-

egorized into: 1) Hydride-bearing molecules with an amine (-NH) functional group believed to have originated from reactions with N_2 ; and 2) Nitrile-bearing molecules with a nitrile (-CN) functional group that form via reactions with atomic N (Rodgers & Charnley 2008; Hily-Blant et al. 2013). Numerous measurements of the $^{14}N/^{15}N$ ratio exist towards low-mass pre-stellar cores (334 ± 50 , 1000 ± 200 and 230 ± 90 from NH_3 , N_2H^+ and HCN, respectively; Lis et al. 2010; Bizzocchi et al. 2013; Hily-Blant et al. 2013) and protostars (~ 160 -290 from HCN and HNC; Wampfler et al. 2014), but observations of this ratio towards their massive counterparts are lacking.

A re-investigation of the fractionation processes of nitrogen in the ISM by Roueff et al. (2015) showed that nitrogen chemistry depends on the temperature and density of the primordial gas in the parental cloud. Since the Sun may have formed in a massive cluster, and since low-mass and high-mass star-forming regions present gas temperatures and densities that differ respectively by ~ 5 -10 K and by factors of 10 (Pillai et al. 2006; Crapsi et al. 2007; Henshaw et al. 2013), measurements of the $^{14}N/^{15}N$ ratio in high-mass star-forming regions could provide insight into the initial bulk composition of the PSN.

Recently, Adande & Ziurys (2012) and Fontani et al. (2015) have measured the $^{14}N/^{15}N$ ratio towards a sample of high-mass star-forming regions. While in the Adande & Ziurys (2012) sample the $^{14}N/^{15}N$ ratios measured from CN and HNC lie between ~ 120 -400, in Fontani et al. (2015)'s work these measurements range from ~ 180 to ~ 1300 in N_2H^+ and ~ 190 to ~ 450 in CN. In both studies, the $^{14}N/^{15}N$ ratios obtained from CN are comparable and fall between the TA and PSN values. However, only a few of these objects were pre-stellar in nature and larger samples of high-mass starless/pre-stellar cores are needed to measure the $^{14}N/^{15}N$ isotopic ratio in regions with physical conditions resembling those of the early stages of the Solar System formation.

We present measurements of the $^{14}N/^{15}N$ isotopic ratio in HCN and HNC obtained toward a sample of 22 high-mass cold cores embedded in 4 IRDCs. These cores are believed to represent the nurseries of high-mass stars and star clusters and have physical properties (densities 10^4 - 10^6 cm^{-3} and temperatures ≤ 20 K; Pillai et al. 2006; Butler & Tan 2012) similar to those expected for the initial conditions of the Solar System. In Section 2, we describe the observations and data analysis. The results are presented in Section 3.1 whilst the uncertainties involved in our calculations are discussed in Section 3.2. In Sections 4.1, 4.2 and 4.3 we investigate the correlation of IRDC cores with star formation activity and compare our results with previous measurements of the $^{14}N/^{15}N$ isotopic ratio in Solar System objects, low-mass and high-mass star-forming regions. In Sections 4.4 and 4.5, we discuss the effects of ^{13}C depletion on the chemistry of nitrogen fractionation and the systematic trend observed in young and quiescent IRDCs with respect to more evolved, star-forming IRDCs. Our conclusions are presented in Section 5.

2. Observations

Observations of the $J=1\rightarrow 0$ rotational transition of HCN, $H^{13}CN$, $HC^{15}N$, $HN^{13}C$ and $H^{15}NC$ were obtained with the IRAM-30m telescope¹ towards 22 massive cores embedded in IRDCs G028.37+00.07, G034.43+00.24, G034.77-00.55 and G035.39-00.33 (hereafter Clouds C, F, G and H respectively, as in Butler & Tan 2012). The frequencies of the transitions and

¹ Based on observations carried out under projects number 134-12 and 027-13 with the IRAM 30m Telescope. IRAM is supported by INSU/CNRS (France), MPG (Germany) and IGN (Spain).

Table 1. Observed HCN and HNC isotopologue transitions. Molecular data extracted from the JPL and CDMS molecular catalogues (Pickett et al. 1998; Müller et al. 2005).

Molecules	Transition	Frequency [GHz]	A_{ul} [s^{-1}]	E_u [K]	g_u
$H^{13}CN$	$J = 1\rightarrow 0, F = 1\rightarrow 1$	86.33877	2.4×10^{-5}	4.14	3
$H^{13}CN$	$J = 1\rightarrow 0, F = 2\rightarrow 1$	86.34018	2.4×10^{-5}	4.14	5
$H^{13}CN$	$J = 1\rightarrow 0, F = 0\rightarrow 1$	86.34227	2.4×10^{-5}	4.14	1
$HC^{15}N$	$J = 1\rightarrow 0$	86.05496	2.4×10^{-5}	4.13	3
$HN^{13}C$	$J = 1\rightarrow 0$	87.09085	1.9×10^{-5}	4.18	3
$H^{15}NC$	$J = 1\rightarrow 0$	88.86571	2.0×10^{-5}	4.26	3
HCN	$J = 1\rightarrow 0, F = 1\rightarrow 1$	88.63041	8.1×10^{-6}	4.25	3
HCN	$J = 1\rightarrow 0, F = 2\rightarrow 1$	88.63160	8.1×10^{-6}	4.25	5
HCN	$J = 1\rightarrow 0, F = 0\rightarrow 1$	88.63393	8.1×10^{-6}	4.25	1




Fig. 1. From left to right, top to bottom panels: Emission lines of $H^{15}NC$, $HN^{13}C$, $HC^{15}N$, and hyperfine transitions of $H^{13}CN$ observed with IRAM-30m toward IRDCs C, F, G and H. Red lines indicate the best Gaussian fit to the lines. Note that the hyperfine components of $H^{13}CN$ were initially fitted but given the bad results of the fit, in a second step only the main $F=2\rightarrow 1$ component of $H^{13}CN$ was fitted using a single-Gaussian component profile (see red line in bottom panels and Section 3.1 for details). See also Figures A.1, A.2, A.3 and A.4 in Appendix for all spectra taken.

the molecular data are included in Table 1 whereas the properties of each IRDC are listed in Table 2. The EMIR receivers were tuned at 87 GHz and the FTS spectrometer provided a spectral resolution of 200 kHz (or ~ 0.68 kms^{-1}). We note that the HNC($J=1\rightarrow 0$) transition was not covered within our frequency range. Typical system temperatures ranged from 106 K to 199 K. The half-power beam width (HPBW) of the telescope was $28''$ at 87 GHz. The spectra were measured in units of antenna temperature, T_A^* , and converted into main beam temperature, T_{mb} , by using a beam efficiency of 0.81. Data reduction was carried out using the GILDAS/CLASS software package². The spectra of HCN ($J=1\rightarrow 0$), $H^{13}CN$ ($J=1\rightarrow 0$), $HC^{15}N$ ($J=1\rightarrow 0$), $HN^{13}C$ ($J=1\rightarrow 0$) and $H^{15}NC$ ($J=1\rightarrow 0$) were obtained towards all cores reported by Butler & Tan (2012) within Clouds F, G and H. For Cloud C, however, all cores were observed except C7 that laid outside our map. C7 will thus not be considered in our analysis.

3. Results

3.1. $^{14}N/^{15}N$ ratios derived from the ^{13}C isotopologues of HCN and HNC

In Fig. 1, we show a sample of spectra of $H^{13}CN$, $HC^{15}N$, $HN^{13}C$ and $H^{15}NC$ obtained towards one massive core in each IRDC

² See <http://www.iram.fr/IRAMFR/GILDAS>.

Table 2. Properties of IRDCs: galactic coordinates l and b , average peak radial velocity V_{LSR} , mass surface density $\Sigma(\text{sat})$, mass M in Solar mass (Butler & Tan 2012) and Galactocentric distance R_{gc} .

IRDCs	l [°]	b [°]	V_{LSR} [kms $^{-1}$]	$\Sigma(\text{sat})$ [gcm $^{-2}$]	M [M_{\odot}]	R_{gc} [kpc]	$^{12}\text{C}/^{13}\text{C}$
C(G028.37+00.07)	28.373	0.076	78.6	0.520	45000	4.65	40.2
F(G034.43+00.24)	34.437	0.245	57.1	0.370	4460	5.74	46.8
G(G034.77-00.55)	34.771	-0.557	43.5	0.347	2010	6.24	49.8
H(G035.39-00.33)	35.395	-0.336	44.7	0.416	13340	6.27	50.0

(the rest of spectra are shown in the Appendix in Figures A.1, A.2, A.3 and A.4). We assume that the emission from these isotopologues is optically thin and consider LTE conditions when calculating their column densities. For the excitation temperature T_{ex} of the molecular gas in these clouds, we assume a lower limit of 10K and an upper limit of 20K based on NH_3 measurements obtained toward other IRDCs (Pillai et al. 2006). Because the hyperfine structure of HN^{13}C cannot be resolved, we considered only one velocity component for this species, in the same way as for HC^{15}N and H^{15}NC . These lines were therefore fitted by using a single-component Gaussian profile. For H^{13}CN , the hyperfine structure of the $J=1\rightarrow 0$ transition could be resolved in the spectra and the HFS line fitting method implemented in CLASS was initially used to obtain the optical depth of the H^{13}CN emission. We note however that the HFS fits presented large uncertainties (see the optical depth values in Table A.5) and therefore, in a second step, we fitted the main $F=2\rightarrow 1$ component of H^{13}CN with a single-Gaussian component profile, calculated the column densities of H^{13}CN assuming optically thin emission, and corrected them by the statistical weight of the $F=2\rightarrow 1$ transition. The measured integrated intensities, radial velocities, linewidths and peak intensities of the lines are listed in Appendix Tables A.1, A.2, A.3, and A.4. We consider that a line is detected when its peak intensity is $\geq 3\sigma$, with σ the rms noise level measured in the spectra (see column 2 in Tables A.1, A.2, A.3 and A.4). For the cores with no molecular detections, we used the 3σ noise level as upper limits to their peak intensities. Note that if we take into account the optical depths derived from the HFS method (of ~ 0.1 -5.8), the resulting H^{13}CN column densities and hence the $^{14}\text{N}/^{15}\text{N}$ ratio are increased by factors of 3–6. Nevertheless, they are consistent with the previously determined results within the uncertainties.

The corresponding $^{14}\text{N}/^{15}\text{N}$ ratios were computed from the molecular column densities of the ^{14}N and ^{15}N HCN and HNC isotopologues after correcting by the $^{12}\text{C}/^{13}\text{C}$ ratio for each IRDC. This ratio is calculated by using the Galactic $^{12}\text{C}/^{13}\text{C}$ gradient as a function of Galactocentric distance derived by Milam et al. (2005) from CN measurements. The $^{12}\text{C}/^{13}\text{C}$ ratios for IRDCs C, F, G and H are, respectively, 40.2, 46.8, 49.8 and 50.0 (see Table 2). The molecular column densities together with the derived $^{14}\text{N}/^{15}\text{N}$ ratios at $T_{ex}=15$ K are listed in Table 3.

In our measurements, the uncertainties in the $^{14}\text{N}/^{15}\text{N}$ ratios were derived by propagating errors and by using the 1σ uncertainties in the line integrated intensities calculated as $\text{rms} \times \sqrt{\Delta v \times \delta v}$, with Δv the average linewidth of the line for all cores with emission and with δv the velocity resolution of the spectrum (~ 0.68 km s $^{-1}$). The derived uncertainties for the $^{14}\text{N}/^{15}\text{N}$ ratios are approximately $\sim 35\%$. This may due to the weak detection of molecules in some of the cores. For the cores with no detections, the $^{14}\text{N}/^{15}\text{N}$ ratios have been estimated using the 3σ upper limits to the integrated intensities of H^{15}NC and HC^{15}N (see column 3 in Tables A.1, A.2, A.3 and A.4), and they should be considered as lower limits.

From Table 3, we find that the $^{14}\text{N}/^{15}\text{N}$ ratios obtained towards Clouds C, F, G and H vary over a large range of val-

Table 3. Column densities and nitrogen ratios obtained from the ^{13}C isotopologues of HCN and HNC

Core	H^{13}CN	HC^{15}N	HN^{12}C	H^{15}NC	HCN	HNC
	$N_{\text{tot}}(T_{ex}=15\text{K})$ [$\times 10^{12}\text{cm}^{-2}$]	$N_{\text{tot}}(T_{ex}=15\text{K})$ [$\times 10^{11}\text{cm}^{-2}$]	$N_{\text{tot}}(T_{ex}=15\text{K})$ [$\times 10^{11}\text{cm}^{-2}$]	$N_{\text{tot}}(T_{ex}=15\text{K})$ [$\times 10^{11}\text{cm}^{-2}$]	$^{14}\text{N}/^{15}\text{N}$ ($T_{ex}=15\text{K}$)	$^{14}\text{N}/^{15}\text{N}$ ($T_{ex}=15\text{K}$)
Cloud C						
C1	1.94	≤ 1.37	3.19	3.07	$\geq 571\pm 65$	418 ± 75
C2	3.03	4.09	4.25	3.57	298 ± 52	478 ± 63
C3	0.89	≤ 2.1	0.98	≤ 2.45	$\geq 170\pm 64$	$\geq 161\pm 14$
C4	2.02	4.20	3.42	5.07	193 ± 30	271 ± 31
C5	2.34	2.80	3.02	3.54	337 ± 89	342 ± 40
C6	1.92	2.26	2.44	2.34	343 ± 83	420 ± 82
C7	-	-	-	-	-	-
C8	1.23	≤ 1.33	1.94	2.41	$\geq 371\pm 60$	325 ± 73
C9	2.74	9.01	3.94	7.59	122 ± 20	209 ± 31
Cloud F						
F1	1.82	≤ 1.12	2.05	3.99	$\geq 763\pm 82$	240 ± 26
F2	1.15	≤ 1.25	2.05	2.53	$\geq 431\pm 63$	378 ± 77
F3	1.11	1.51	2.25	1.95	346 ± 80	541 ± 79
F4	1.83	3.03	2.78	3.47	282 ± 60	374 ± 57
Cloud G						
G1	0.35	2.47	0.92	≤ 1.94	70 ± 28	$\geq 237\pm 21$
G2	≤ 0.31	≤ 1.54	1.71	4.14	-	206 ± 31
G3	0.63	≤ 1.72	1.74	3.65	$\geq 181\pm 54$	237 ± 36
Cloud H						
H1	0.98	≤ 1.39	1.91	4.09	$\geq 353\pm 51$	234 ± 20
H2	0.97	≤ 1.32	1.90	3.11	$\geq 366\pm 132$	306 ± 40
H3	1.84	2.00	2.19	2.24	458 ± 98	488 ± 68
H4	0.86	3.03	1.86	2.86	142 ± 34	326 ± 44
H5	1.13	1.44	1.96	2.96	395 ± 97	331 ± 36
H6	0.94	2.18	2.71	4.56	216 ± 77	297 ± 62

ues. In particular, for IRDCs C, F and H, the nitrogen ratios in HCN range between $122\text{--}\geq 571$, $282\text{--}\geq 763$, $142\text{--}458$, respectively, while in HNC they range between $\geq 161\text{--}478$ for Cloud C, $240\text{--}541$ for Cloud F, and $234\text{--}488$ for Cloud H. On the other hand, the $^{14}\text{N}/^{15}\text{N}$ ratios measured toward the cores in Cloud G are systematically lower ranging between $70\text{--}\geq 181$ in HCN and $206\text{--}\geq 237$ in HNC. We have tested the effects of T_{ex} on our results, and have found that if we use $T_{ex}=10$ K or $T_{ex}=20$ K instead of $T_{ex}=15$ K, the derived $^{14}\text{N}/^{15}\text{N}$ isotopic ratios for both HCN and HNC do not vary significantly, lying within the $\sim 30\%$ uncertainties. Higher T_{ex} (e.g. 50 K and 100 K) also confirm this behaviour, with the $^{14}\text{N}/^{15}\text{N}$ ratios changing within a factor of 1.2. Nevertheless, Roueff et al. (2015) have recently pointed out that species such as HN^{13}C and H^{13}CN may suffer significant depletion in molecular clouds, challenging the interpretation of $^{14}\text{N}/^{15}\text{N}$ isotopic ratios derived from ^{13}C containing isotopologues. In Section 3.2, we explore this possibility by directly measuring the $^{14}\text{N}/^{15}\text{N}$ ratios using HCN and its ^{15}N isotopologue toward the IRDCs cores in our sample with optically thin HCN emission.

3.2. $^{14}\text{N}/^{15}\text{N}$ ratios derived from HCN and its ^{15}N isotopologue

In this section, we test whether the $^{14}\text{N}/^{15}\text{N}$ ratios derived in Section 3.1 are significantly affected by ^{13}C depletion as proposed by the modeling of Roueff et al. (2015). We have thus carried out direct measurements of the $^{14}\text{N}/^{15}\text{N}$ ratios by using the $J=1\rightarrow 0$ rotational transitions of HCN and HC^{15}N , which were observed simultaneously within our frequency setup. HCN ($J=1\rightarrow 0$) is optically thick in IRDC star-forming cores such as the cores in Clouds C and F, or core H1 in Cloud H. Therefore for this

Table 4. Column densities and nitrogen ratios obtained in HCN and its ^{12}C isotopologue

Core	HCN		HC ^{15}N	HCN
	$N_{\text{tot}}(T_{\text{ex}}=15\text{K})$ [$\times 10^{13}\text{cm}^{-2}$]	$N_{\text{tot}}(T_{\text{ex}}=15\text{K})$ [$\times 10^{11}\text{cm}^{-2}$]	$N_{\text{tot}}(T_{\text{ex}}=15\text{K})$ [$\times 10^{11}\text{cm}^{-2}$]	$^{14}\text{N}/^{15}\text{N}$ ($T_{\text{ex}}=15\text{K}$)
Cloud G				
G1	1.06	2.47		43 ± 9
G3	1.15	≤ 1.72		$\geq 67\pm 3$
Cloud H				
H2	3.72	≤ 1.32		$\geq 282\pm 5$
H3	5.26	2.00		263 ± 49
H4	3.65	3.03		121 ± 24
H5	3.71	1.44		259 ± 57

test we only use the IRDC cores within our sample that show optically-thin or moderately optically-thick emission (i.e. with $\tau \leq 1-2$). These cores are G1 and G3 in Cloud G, and H2, H3, H4, and H5 in Cloud H. The rms noise level, integrated intensity, central radial velocity, linewidth, peak intensity and derived optical depth of the HCN ($J=1\rightarrow 0$) lines, are shown in Table A.5 in the Appendix.

Following the same analysis procedures as for H^{13}CN in Section 3.1, the $^{14}\text{N}/^{15}\text{N}$ ratios were calculated from the column densities of HCN and HC^{15}N assuming optically thin emission, $T_{\text{ex}}=15\text{K}$, and LTE conditions (see Table 4). For the HC^{15}N non-detections, the upper limits to the column density of this molecule were estimated from the 3σ rms noise level in the HC^{15}N spectra. The derived $^{14}\text{N}/^{15}\text{N}$ ratios range from ≥ 67 to ≥ 282 . If we compare these values with those from column 6 in Table 3, we find that the $^{14}\text{N}/^{15}\text{N}$ ratios inferred from HCN are systematically lower (by factors 1.2-2.7) with respect to those obtained from H^{13}CN . This is in contrast with the results from Roueff et al. (2015) since, from their models, the $^{12}\text{C}/^{13}\text{C}$ isotopic ratio measured from HCN should be a factor of ~ 2 higher than that derived from CN (i.e. at time-scales $\geq 1\text{Myr}$ for the typical densities of IRDC cores of $\sim 10^5\text{cm}^{-3}$; see Figure 4 in their paper). We note that this also holds if the $^{14}\text{N}/^{15}\text{N}$ ratios of the moderately-optically thick cores are corrected by their HCN optical depths (with $\tau(\text{HCN})=0.71-1.84$, which corresponds to correction factors of $\sim 1.4-2.2$). Except for core H5, the corrected values of the $^{14}\text{N}/^{15}\text{N}$ ratios for cores H2, H3 and H4 are, respectively, ≥ 503 , 445 and 168 , which are consistent with those inferred from H^{13}CN and lie within the uncertainties. Although our sub-sample of optically-thin/moderately-optically thick IRDC cores is small, this test suggests that the $^{14}\text{N}/^{15}\text{N}$ ratios obtained using the ^{13}C containing isotopologues are not strongly affected by ^{13}C depletion as proposed by the models of Roueff et al. (2015). In Section 4.5, we discuss the possible reasons for this.

4. Discussion

4.1. Correlation with star formation activity

The chemistry of HCN and HNC is known to be temperature dependent (Pineau des Forets et al. 1990) and any star formation activity in the core could locally heat the molecular gas enhancing the abundance of HCN (and its isotopologues) over HNC. Therefore, it is important to investigate whether the measured $^{14}\text{N}/^{15}\text{N}$ isotopic ratios in HCN and HNC show any correlation with the level of star formation activity in the observed IRDC cores. For this purpose, we adopted the classification of the embedded cores in IRDCs C, F, G and H proposed by Chambers et al. (2009) and Rathborne et al. (2010). Each of the cores is classified as quiescent, intermediate or active based upon their colour in Spitzer/IRAC 3-8 μm images as well as the pres-




Fig. 2. Column densities of HCN (top panel) and HNC (bottom panel) ^{15}N isotopologues plotted against those of the ^{14}N species for all the cores in the sample. The cores are classified as active (red), intermediate (green), and quiescent (blue). Black indicate cores with no known classification. Different symbols are used to denote different clouds: IRDC C (circle), IRDC F (star), IRDC G (square) and IRDC H (triangle). The straight lines indicate the lowest and highest value of the corresponding $^{14}\text{N}/^{15}\text{N}$ ratio for each molecule.

ence or absence of $24\mu\text{m}$ point source emission. The summary of each core's classification is listed in Table 5.

In Fig.2, we report the column densities of the ^{15}N isotopologues against those of the ^{14}N species in relation to their star-formation classification. This Figure shows that there is no correlation between the column densities of the HCN or HNC isotopologues with the level of star formation activity in the IRDC cores. Such conclusion is also confirmed by plotting the column densities of HC^{15}N against that of H^{15}NC . In other words, the measurements of $^{14}\text{N}/^{15}\text{N}$ ratio towards IRDCs C, F, G and H from the $J=1\rightarrow 0$ transitions of HCN and HNC indeed probe the chemical composition of the envelope of these IRDCs cores. As such, it is in general not affected by local star formation feedback, although the highest $^{15}\text{N}/^{14}\text{N}$ ratio is found towards one of the active cores. Therefore, we cannot rule out that higher- J transitions and higher-angular resolution observations give $^{15}\text{N}/^{14}\text{N}$ ratios that are correlated with star formation activity.

4.2. Comparison with Solar System objects and low-mass star-forming regions

To understand whether IRDC cores have a nitrogen chemical composition consistent with that of the Solar System's birth-place, it is essential to compare the results obtained toward our sample of IRDC cores with those measured in Solar System objects (see Fig 3). For completeness, Fig. 3 also reports the $^{14}\text{N}/^{15}\text{N}$ ratios obtained towards low-mass pre-stellar/starless and star-forming cores. Overall, the $^{14}\text{N}/^{15}\text{N}$ ratios from Clouds C, F and H show values similar to those observed in the Sun, planets and pre-stellar/star-forming regions, and consistent with the TA and the PSN values. On the other hand, the $^{14}\text{N}/^{15}\text{N}$ ratios measured towards Cloud G lie mostly at the level of the TA value and are significantly lower than the PSN value in Fig 3. The results in Cloud G are also in agreement with the measurements

Table 5. Summary of IRDC cores classification

IRAC 3-8 μm	24 μm emission	Core category	Cores in IRDCs C, F, G and H
8.0 μm	Yes/No	Red	-
Green Fuzzy	Yes	Active	C2, C3, C6, C9, H2, H5, H6
Green Fuzzy	No	Intermediate	C4, F1
None	Yes	-	-
None	No	Quiescent	C1, C5, C8, F2, F3, G2, H1, H3, H4
3.6 μm emission	Yes/No	Blue	-

Note: Cores F4, G1 and G3 have not yet been classified (Rathborne et al. 2010; Chambers et al. 2009).

obtained in Comets, IDPs, Meteorites and especially with protoplanetary disks (80–160) as recently measured by Guzmán et al. (2017). Furthermore, they marginally agree with the lower end of the ratios derived in starless/pre-stellar and star-forming cores. We also caution that half of the $^{14}\text{N}/^{15}\text{N}$ ratios in Cloud G are lower limits (HC^{15}N has not been detected in cores G2 and G3; and H^{15}NC has not been detected in core G1; see Tables A.2 and A.4), and therefore we may be lacking enough statistics to draw a firm conclusion.

4.3. Comparison with high-mass star-forming regions

The comparison with the measurements from Adande & Ziurys (2012) and Fontani et al. (2015) towards high-mass star-forming regions, shows that our measurements are consistent with their results as a whole. Especially, the $^{14}\text{N}/^{15}\text{N}$ ratios from HNC in IRDCs almost lie in the same range as those measured by Adande & Ziurys (2012) in CN and HNC and by Fontani et al. (2015) in CN. The $^{14}\text{N}/^{15}\text{N}$ ratios from HCN are also compatible with the results measured by Fontani et al. (2015) in N_2H^+ emission. Since Fontani et al. (2015) also measured the $^{14}\text{N}/^{15}\text{N}$ ratio towards cores C1, F1, F2 and G2 included in our sample, we have compared the results between each individual core and they show some discrepancies as a result. Indeed, the $^{14}\text{N}/^{15}\text{N}$ ratios obtained by Fontani et al. (2015) in N^{15}NH^+ and $^{15}\text{NNH}^+$ (CN was not detected towards these cores) are, respectively, ~ 1445 and ~ 1217 for C1, ~ 672 and ~ 566 for F1 and ~ 872 and ~ 856 for G2, i.e. overall significantly higher than those measured in this work. In contrast, F2 shows a lower value of ~ 232 and ≥ 195 in $^{15}\text{NNH}^+$ and N^{15}NH^+ respectively. We note that these large discrepancies have also been found in low-mass pre-stellar cores and could be associated with the different chemistries involved in the formation of N_2H^+ and HNC/HCN (Wirström et al. 2012; Hily-Blant et al. 2013; Bizzocchi et al. 2013). More recently, cores C1, C3, F1, F2 and G2 have been studied independently by Colzi et al. (2017, submitted) using isotopologues of HCN and HNC. A similar range for the $^{14}\text{N}/^{15}\text{N}$ ratios has been found in HCN (≥ 150 -748) whilst results in HNC lie in a slightly higher range (263-813). In both samples, core G2 shows one of the smallest ratios in HCN and HNC.

4.4. ^{13}C depletion and its effects on nitrogen fractionation

In Section 3.2, we evaluated whether the depletion of ^{13}C for species such as HCN and HNC (as predicted by the models of Roueff et al. 2015) could affect our derived values of the $^{14}\text{N}/^{15}\text{N}$ ratios. Our test revealed that the $^{14}\text{N}/^{15}\text{N}$ values inferred from HCN are either consistent, or lower, than those measured from the ^{13}C isotopologue, in contrast to the modelling predictions. This could be due to two reasons: i) the time-scales (age) of IRDC cores, and ii) the kinetic temperature of the gas within them.

Regarding the time-scales, Kong et al. (2017) have modelled the chemistry of deuterated species such as N_2D^+ in IRDC cores to provide constraints to the dynamical age of these cores. Their modelling shows that the enhanced D/H ratio in these objects can be reproduced for time-scales of $\sim 10^5$ yrs (note that these authors model the N_2D^+ emission arising from the C1 core in Cloud C). On the other hand, Roueff et al. (2015) predict similar $^{12}\text{C}/^{13}\text{C}$ ratios associated with CN and with HCN/HNC at these time-scales (of $\sim 10^5$ yrs) for the typical H_2 gas densities of IRDC cores (of $\sim 10^5, \text{cm}^{-3}$; see Butler & Tan 2012). The large differences in $^{12}\text{C}/^{13}\text{C}$ ratios associated with HCN/HNC, such as those discussed in Section 3.2, would therefore not be expected.

Nevertheless, the definition of a "core" formation timescale is somewhat ambiguous, for example Barnes et al. (2016) found that the D/H fraction within Cloud F would have taken several 10^6 yr to form. In light of this, a more self-consistent comparison between timescales inferred by chemical models is required.

Concerning the gas temperature of IRDC cores, measurements of the emission of NH_3 toward these cores give kinetic temperatures of the gas of 15-20 K (Pillai et al. 2006), which are higher than those assumed in the models of Roueff et al. (2015, of 10 K). Since carbon depletion is strongly dependent on gas/dust temperature, it is unclear whether these results can be compared directly to IRDC cores (note that no models are provided for temperatures higher than 10 K). Therefore, additional modeling is needed to test the effects of ^{13}C depletion in the chemistry of nitrogen fractionation at slightly higher temperatures similar to those found in IRDCs.

4.5. Systematic trend of $^{14}\text{N}/^{15}\text{N}$ ratio between IRDCs

Table 3 and Figure 3 show that the $^{14}\text{N}/^{15}\text{N}$ ratios observed in Cloud G are systematically lower than those measured in Clouds C, F and H. This may be due to the properties of Cloud G itself. As discussed in Section 4.1, cores G1, G2 and G3 do not show any trace of star-formation activity, whilst the other three IRDCs show several cores that are actively forming stars (see e.g. cores C2 or H2). In addition, Cloud G is the least massive, the most diffuse (it has the weakest emission in high-density tracers; Cosentino et al., in prep.), and it has the lowest peak H_2 mass surface density amongst the four targeted IRDCs (see Table 2 and Butler & Tan 2012). Given that the kinetic temperature of the gas is similar across IRDCs (~ 15 -20 K), we propose that density could be one of the important parameters that is responsible for the discrepancies found between Cloud G and the other IRDCs, although the models do not agree with this scenario. Therefore, we speculate that the PSN may have formed in an IRDC with properties similar to those of cloud G. However, we note that the properties of this sample of IRDC cores need to be further investigated along with relevant chemical models in order to confirm the proposed idea.




Fig. 3. Nitrogen isotope ratio variations measured in different sources starting from small Solar System bodies (in red; Busemann et al. 2006; Floss et al. 2006; Alexander et al. 2007; Bockelée-Morvan et al. 2008; Manfroid et al. 2009; Mumma & Charnley 2011; Guzmán et al. 2017) to planets (in black; Junk & Svec 1958; Hoffman et al. 1979; Fouchet et al. 2004; Niemann et al. 2005; Marty et al. 2010; Wong et al. 2013; Fletcher et al. 2014), low-mass pre-stellar cores (in yellow; Gerin et al. 2009; Bizzocchi et al. 2010; Lis et al. 2010; Bizzocchi et al. 2013; Hily-Blant et al. 2013), low-mass star-forming cores (in light grey; Wampfler et al. 2014) and high-mass star-forming cores (in blue; Adande & Ziurys 2012; Fontani et al. 2015). Our measurements in IRDCs are shown on the right. On the upper right corner, we show a representative error bar for our measurements of the $^{14}\text{N}/^{15}\text{N}$ ratio. Black arrows indicate the lower limits in our measurements. Note that the $^{14}\text{N}/^{15}\text{N}$ ratio measured in HCN from core G2 is shown as zero with no lower limit indication due to both of the column densities of H^{13}CN and HC^{15}N are only given as upper limits.

5. Conclusions

We have measured the nitrogen isotopic ratio $^{14}\text{N}/^{15}\text{N}$ toward a sample of cold IRDC cores. This ratio ranges between ~ 70 and ≥ 763 in HCN and between ~ 161 and ~ 541 in HNC. In particular, Cloud G systematically shows lower nitrogen isotopic ratios than the other three clouds, with values being consistent with the ratio measured toward small Solar System bodies such as Comets, IDPs, Meteorites and also proto-planetary disks. Since Cloud G shows lower overall gas densities, and since it likely is at an earliest stage of evolution, we propose that gas density is the key parameter in nitrogen fractionation in IRDCs. Higher angular resolution observations, as well as chemical modeling of the nitrogen fractionation of HCN and HNC at temperatures similar to those found in IRDCs, are needed to establish the origin of the discrepancies in the measured $^{14}\text{N}/^{15}\text{N}$ ratios found in Cloud G with respect to Clouds C, F and H. The comparison between the modeling predictions and our IRDC measurements may allow to constrain the main chemical reactions involved in the fractionation process of Nitrogen in the proto-solar nebula.

Acknowledgements. We would like to thank an anonymous referee for the valuable comments to a previous version of the manuscript. I.J.-S. acknowledges the financial support received from the STFC through an Ernest Rutherford Fellowship (proposal number ST/L004801/2). P.C. acknowledges financial support of the European Research Council (ERC; project PALs 320620). The research leading to these results has also received funding from the European Commission (FP/2007-2013) under grant agreement No. 283393 (RadioNet3)."

References

Adams, F. C. 2010, ARA&A, 48, 47
 Adande, G. R. & Ziurys, L. M. 2012, ApJ, 744, 194
 Alexander, C. M. O. D., Fogel, M., Yabuta, H., & Cody, G. D. 2007, Geochim. Cosmochim. Acta, 71, 4380
 Barnes, A. T., Kong, S., Tan, J. C., et al. 2016, MNRAS, 458, 1990
 Bizzocchi, L., Caselli, P., & Dore, L. 2010, A&A, 510, L5
 Bizzocchi, L., Caselli, P., Leonardo, E., & Dore, L. 2013, A&A, 555, A109
 Bockelée-Morvan, D., Biver, N., Jehin, E., et al. 2008, ApJ, 679, L49
 Busemann, H., Young, A. F., O'D. Alexander, C. M., et al. 2006, Science, 312, 727

Butler, M. J. & Tan, J. C. 2012, ApJ, 754, 5
 Chambers, E. T., Jackson, J. M., Rathborne, J. M., & Simon, R. 2009, ApJS, 181, 360
 Crapsi, A., Caselli, P., Walmsley, M. C., & Tafalla, M. 2007, A&A, 470, 221
 Dukes, D. & Krumholz, M. R. 2012, ApJ, 754, 56
 Fletcher, L. N., Greathouse, T. K., Orton, G. S., et al. 2014, Icarus, 238, 170
 Floss, C., Stadermann, F. J., Bradley, J. P., et al. 2006, Geochim. Cosmochim. Acta, 70, 2371
 Fontani, F., Caselli, P., Palau, A., Bizzocchi, L., & Ceccarelli, C. 2015, ApJ, 808, L46
 Fouchet, T., Irwin, P. G. J., Parrish, P., et al. 2004, Icarus, 172, 50
 Gerin, M., Marcelino, N., Biver, N., et al. 2009, A&A, 498, L9
 Guzmán, V. V., Öberg, K. L., Huang, J., Loomis, R., & Qi, C. 2017, ArXiv e-prints [arXiv:1701.07510]
 Heays, A., van Dishoeck, E., Visser, R., et al. 2014, in 69th International Symposium on Molecular Spectroscopy
 Henshaw, J. D., Caselli, P., Fontani, F., et al. 2013, MNRAS, 428, 3425
 Hily-Blant, P., Bonal, L., Faure, A., & Quirico, E. 2013, Icarus, 223, 582
 Hoffman, J. H., Hodges, R. R., McElroy, M. B., Donahue, T. M., & Kolpin, M. 1979, Science, 205, 49
 Junk, G. & Svec, H. J. 1958, Geochim. Cosmochim. Acta, 14, 234
 Kainulainen, J. & Tan, J. C. 2013, A&A, 549, A53
 Kong, S., Tan, J. C., Caselli, P., et al. 2017, ArXiv e-prints [arXiv:1701.05953]
 Lis, D. C., Wootten, A., Gerin, M., & Roueff, E. 2010, ApJ, 710, L49
 Lyons, J. R., Bergin, E. A., Ciesla, F. J., et al. 2009, Geochim. Cosmochim. Acta, 73, 4998
 Manfroid, J., Jehin, E., Hutsemékers, D., et al. 2009, A&A, 503, 613
 Marty, B., Zimmermann, L., Burnard, P. G., et al. 2010, Geochim. Cosmochim. Acta, 74, 340
 Milam, S. N., Savage, C., Brewster, M. A., Ziurys, L. M., & Wyckoff, S. 2005, ApJ, 634, 1126
 Müller, H. S. P., Schlöder, F., Stutzki, J., & Winnewisser, G. 2005, Journal of Molecular Structure, 742, 215
 Mumma, M. J. & Charnley, S. B. 2011, ARA&A, 49, 471
 Nicholson, R. B. & Parker, R. J. 2017, MNRAS, 464, 4318
 Niemann, H. B., Atreya, S. K., Bauer, S. J., et al. 2005, Nature, 438, 779
 Pflanzner, S. 2013, A&A, 549, A82
 Pickett, H. M., Poynter, R. L., Cohen, E. A., et al. 1998, J. Quant. Spectr. Rad. Transf., 60, 883
 Pillai, T., Wyrowski, F., Carey, S. J., & Menten, K. M. 2006, A&A, 450, 569
 Pineau des Forets, G., Roueff, E., & Flower, D. R. 1990, MNRAS, 244, 668
 Rathborne, J. M., Jackson, J. M., Chambers, E. T., et al. 2010, ApJ, 715, 310
 Rodgers, S. D. & Charnley, S. B. 2008, ApJ, 689, 1448
 Roueff, E., Loison, J. C., & Hickson, K. M. 2015, A&A, 576, A99
 Terzieva, R. & Herbst, E. 2000, MNRAS, 317, 563
 Wampfler, S. F., Jørgensen, J. K., Bizzarro, M., & Bisschop, S. E. 2014, A&A, 572, A24
 Wilson, T. L. & Rood, R. 1994, ARA&A, 32, 191
 Wirstrom, E. S., Charnley, S. B., Cordiner, M. A., & Milam, S. N. 2012, ApJ, 757, L11
 Wong, M. H., Atreya, S. K., Mahaffy, P. N., et al. 2013, Geophys. Res. Lett., 40, 6033

Appendix A: Spectra and fitting parameters of isotopologues of HCN and HNC




Fig. A.1. Spectra of HC^{15}N , HN^{13}C , H^{15}NC and H^{13}CN observed with IRAM-30m toward IRDC C. The red line indicates the best Gaussian fit. Note that the hyperfine components of H^{13}CN were initially fitted but given the bad results of the fit, in a second step only the main $F=2\rightarrow 1$ component of H^{13}CN was fitted using a single-Gaussian component profile (see red line in bottom panels and Section 3.1 for details).




Fig. A.2. Spectra of HC^{15}N , HN^{13}C , H^{15}NC and H^{13}CN observed with IRAM-30m toward IRDC F. The red line indicates the best Gaussian fit. Note that the hyperfine components of H^{13}CN were initially fitted but given the bad results of the fit, in a second step only the main $F=2\rightarrow 1$ component of H^{13}CN was fitted using a single-Gaussian component profile (see red line in bottom panels and Section 3.1 for details).




Fig. A.3. Spectra of HC^{15}N , HN^{13}C , H^{15}NC and H^{13}CN observed with IRAM-30m toward IRDC G. Note that only the main component of H^{13}CN was fitted with a single-Gaussian component profile, which was then used to calculate the column densities. The red line indicates the best Gaussian fit and these spectra have been scaled to fit into each individual panel.




Fig. A.4. Spectra of HC^{15}N , HN^{13}C , H^{15}NC and H^{13}CN observed with IRAM-30m toward IRDC H. The red line indicates the best Gaussian fit. Note that the hyperfine components of H^{13}CN were initially fitted but given the bad results of the fit, in a second step only the main $F=2\rightarrow 1$ component of H^{13}CN was fitted using a single-Gaussian component profile (see red line in bottom panels and Section 3.1 for details).

Table A.1. The root mean square noise in the spectra rms, integrated intensities $\int T_{\text{mb}}d\nu$, radial velocity ν , line widths $\Delta\nu$ and peak temperature T_{peak} of $\text{HC}^{15}\text{N}(1\rightarrow 0)$ in cores within Clouds C, F, G and H.

Core	RMS [10^{-2} K]	$\int T_{\text{mb}}\Delta\nu$ [K kms $^{-1}$]	ν [kms $^{-1}$]	$\Delta\nu$ [kms $^{-1}$]	T_{peak} [K]
Cloud C					
C1*	1.8	≤ 0.07	-	-	≤ 0.05
C2	1.8	0.2 ± 0.05	79.36 ± 0.62	5.02 ± 1.07	0.04
C3*	2.7	≤ 0.10	-	-	≤ 0.08
C4	1.6	0.21 ± 0.05	80.31 ± 0.59	5.12 ± 1.71	0.04
C5	1.9	0.14 ± 0.05	77.89 ± 0.89	4.90 ± 2.16	0.03
C6	1.8	0.11 ± 0.04	79.19 ± 0.50	2.81 ± 1.01	0.04
C7	-	-	-	-	-
C8*	1.7	≤ 0.06	-	-	≤ 0.05
C9	3.3	0.44 ± 0.07	78.30 ± 0.17	2.26 ± 0.37	0.18
Cloud F					
F1*	1.5	≤ 0.05	-	-	≤ 0.04
F2*	1.6	≤ 0.06	-	-	≤ 0.05
F3	1.5	0.07 ± 0.02	57.9 ± 0.22	1.49 ± 0.49	0.05
F4	1.8	0.15 ± 0.04	57.74 ± 0.54	3.93 ± 1.05	0.04
Cloud G					
G1	2.1	0.12 ± 0.04	42.85 ± 0.35	2.20 ± 0.67	0.05
G2*	2.0	≤ 0.08	-	-	≤ 0.06
G3*	2.2	≤ 0.08	-	-	≤ 0.07
Cloud H					
H1*	1.8	≤ 0.07	-	-	≤ 0.05
H2*	1.7	≤ 0.06	-	-	≤ 0.05
H3	1.5	0.10 ± 0.03	44.97 ± 0.33	2.28 ± 0.86	0.04
H4	1.8	0.15 ± 0.04	45.37 ± 0.56	3.99 ± 1.17	0.03
H5	1.8	0.07 ± 0.02	45.74 ± 0.19	1.13 ± 0.48	0.06
H6	3.3	0.11 ± 0.05	46.03 ± 0.43	1.68 ± 0.63	0.06

(a) * indicates 3σ upper-limit values for T_{peak} .

(b) The integrated intensity upper limits are calculated as $3\sigma \times \sqrt{\delta\nu} \times \Delta\nu$, with $\delta\nu$ the spectral resolution (0.68 km s^{-1}) and $\Delta\nu$ the average linewidth measured considering all cores.

Table A.2. The root mean square noise in the spectra rms, integrated intensities $\int T_{\text{mb}}d\nu$, radial velocity ν , line widths $\Delta\nu$ and peak temperature T_{peak} of $\text{HN}^{13}\text{C}(1\rightarrow 0)$ in cores within Clouds C, F, G and H.

Core	RMS [10^{-2} K]	$\int T_{\text{mb}}\Delta\nu$ [K kms $^{-1}$]	ν [kms $^{-1}$]	$\Delta\nu$ [kms $^{-1}$]	T_{peak} [K]
Cloud C					
C1	1.8	1.22 ± 0.03	80.12 ± 0.04	2.84 ± 0.09	0.41
C2	1.7	1.62 ± 0.04	79.32 ± 0.04	3.32 ± 0.08	0.46
C3	2.3	0.37 ± 0.05	81.09 ± 0.11	1.93 ± 0.34	0.18
C4	1.6	1.30 ± 0.03	80.26 ± 0.03	2.35 ± 0.07	0.52
C5	1.6	1.15 ± 0.04	79.31 ± 0.04	2.57 ± 0.10	0.42
C6	1.7	0.93 ± 0.03	79.47 ± 0.04	2.15 ± 0.09	0.41
C7	-	-	-	-	-
C8	1.7	0.74 ± 0.04	78.94 ± 0.06	2.44 ± 0.14	0.29
C9	3.2	1.50 ± 0.07	78.59 ± 0.07	3.16 ± 0.17	0.45
Cloud F					
F1	1.5	0.78 ± 0.03	57.61 ± 0.07	3.39 ± 0.15	0.22
F2	1.6	0.78 ± 0.03	58.09 ± 0.04	2.22 ± 0.10	0.33
F3	1.5	0.86 ± 0.03	58.25 ± 0.04	2.12 ± 0.09	0.38
F4	1.7	1.06 ± 0.03	58.85 ± 0.04	2.35 ± 0.08	0.42
Cloud G					
G1	1.9	0.35 ± 0.04	43.28 ± 0.15	2.47 ± 0.34	0.13
G2	2.1	0.65 ± 0.04	42.15 ± 0.08	2.67 ± 0.19	0.23
G3	2.2	0.66 ± 0.03	41.68 ± 0.05	1.79 ± 0.11	0.35
Cloud H					
H1	1.7	0.73 ± 0.03	45.13 ± 0.02	1.43 ± 0.06	0.48
H2	1.6	0.73 ± 0.04	45.57 ± 0.03	1.50 ± 0.09	0.46
H3	1.6	0.84 ± 0.05	45.48 ± 0.06	2.00 ± 0.16	0.39
H4	1.6	0.71 ± 0.03	45.58 ± 0.03	1.53 ± 0.07	0.44
H5	1.7	0.75 ± 0.03	45.64 ± 0.03	1.46 ± 0.06	0.48
H6	2.8	1.03 ± 0.05	45.73 ± 0.04	1.77 ± 0.09	0.55

(a) The integrated intensity upper limits are calculated as $3\sigma \times \sqrt{\delta\nu} \times \Delta\nu$, with $\delta\nu$ the spectral resolution (0.68 km s^{-1}) and $\Delta\nu$ the average linewidth measured considering all cores.

Table A.3. The root mean square noise in the spectra rms, integrated intensities $\int T_{\text{mb}}d\nu$, radial velocity ν , line widths $\Delta\nu$ and peak temperature T_{peak} of $\text{H}^{15}\text{NC}(1\rightarrow 0)$ in cores within Clouds C, F, G and H.

Core	RMS [10^{-2} K]	$\int T_{\text{mb}}\Delta\nu$ [K kms $^{-1}$]	ν [kms $^{-1}$]	$\Delta\nu$ [kms $^{-1}$]	T_{peak} [K]
Cloud C					
C1	1.4	0.12 \pm 0.03	79.58 \pm 0.47	3.24 \pm 0.72	0.04
C2	1.6	0.14 \pm 0.03	79.06 \pm 0.20	1.88 \pm 0.57	0.07
C3*	2.6	\leq 0.10	-	-	\leq 0.07
C4	1.5	0.20 \pm 0.04	80.16 \pm 0.30	3.17 \pm 0.85	0.06
C5	1.5	0.14 \pm 0.02	79.30 \pm 0.14	1.57 \pm 0.32	0.08
C6	1.7	0.09 \pm 0.03	79.37 \pm 0.27	1.66 \pm 0.79	0.05
C7	-	-	-	-	-
C8	1.5	0.10 \pm 0.03	78.17 \pm 0.47	2.76 \pm 0.83	0.03
C9	3.3	0.30 \pm 0.07	78.92 \pm 0.27	2.58 \pm 0.07	0.11
Cloud F					
F1	0.9	0.09 \pm 0.02	56.14 \pm 0.13	1.21 \pm 0.30	0.03
F2	1.6	0.10 \pm 0.03	57.92 \pm 0.42	2.50 \pm 1.12	0.04
F3	1.3	0.08 \pm 0.02	57.90 \pm 0.12	1.00 \pm 0.21	0.07
F4	1.5	0.14 \pm 0.03	58.95 \pm 0.32	2.69 \pm 0.61	0.05
Cloud G					
G1*	2.0	\leq 0.08	-	-	\leq 0.06
G2	1.9	0.16 \pm 0.03	42.26 \pm 0.26	2.22 \pm 0.46	0.07
G3	2.3	0.14 \pm 0.03	41.42 \pm 0.13	1.24 \pm 0.26	0.11
Cloud H					
H1	1.5	0.16 \pm 0.02	45.05 \pm 0.07	1.20 \pm 0.22	0.13
H2	1.5	0.12 \pm 0.03	45.44 \pm 0.15	1.60 \pm 0.51	0.07
H3	1.4	0.09 \pm 0.02	45.56 \pm 0.12	1.04 \pm 0.37	0.08
H4	1.5	0.11 \pm 0.02	45.49 \pm 0.15	1.50 \pm 0.21	0.07
H5	1.4	0.12 \pm 0.02	45.57 \pm 0.11	1.23 \pm 0.26	0.09
H6	2.7	0.18 \pm 0.04	45.75 \pm 0.24	1.90 \pm 0.38	0.09

(a) * indicates 3σ upper-limit values for T_{peak} .(b) The integrated intensity upper limits are calculated as $3\sigma \times \sqrt{\delta\nu} \times \Delta\nu$, with $\delta\nu$ the spectral resolution (0.68 km s $^{-1}$) and $\Delta\nu$ the average linewidth measured considering all cores.**Table A.4.** The root mean square noise in the spectra rms, integrated intensities $\int T_{\text{mb}}d\nu$, radial velocity ν , line widths $\Delta\nu$, peak temperature T_{peak} of the $\text{H}^{13}\text{CN}(1\rightarrow 0, F=2\rightarrow 1)$ hyperfine component and optical depth τ of the $\text{H}^{13}\text{CN}(1\rightarrow 0)$ emission in cores within Clouds C, F, G and H.

Core	RMS [10^{-2} K]	$\int T_{\text{mb}}\Delta\nu$ [K kms $^{-1}$]	ν [kms $^{-1}$]	$\Delta\nu$ [kms $^{-1}$]	T_{peak} [K]	τ
Cloud C						
C1	1.8	0.53 \pm 0.05	80.08 \pm 0.14	3.46 \pm 0.39	0.14	1.47 \pm 1.18
C2	1.8	0.83 \pm 0.13	78.63 \pm 0.10	2.97 \pm 0.31	0.26	1.79 \pm 0.57
C3	3.4	0.24 \pm 0.06	81.08 \pm 0.27	2.13 \pm 0.46	0.11	2.86 \pm 4.00
C4	1.6	0.55 \pm 0.07	79.98 \pm 0.07	2.11 \pm 0.22	0.25	1.75 \pm 0.63
C5	1.9	0.63 \pm 0.07	79.24 \pm 0.13	2.95 \pm 0.35	0.20	1.33 \pm 0.81
C6	1.7	0.52 \pm 0.07	79.36 \pm 0.12	2.69 \pm 0.30	0.18	1.87 \pm 1.00
C7	-	-	-	-	-	-
C8	1.8	0.33 \pm 0.04	78.60 \pm 0.15	2.61 \pm 0.39	0.12	2.11 \pm 2.09
C9	4.8	0.75 \pm 0.15	77.62 \pm 0.13	2.05 \pm 0.29	0.34	5.88 \pm 2.22
Cloud F						
F1	1.5	0.50 \pm 0.04	57.77 \pm 0.14	3.57 \pm 0.40	0.13	2.29 \pm 1.25
F2	1.6	0.31 \pm 0.03	58.04 \pm 0.11	2.36 \pm 0.26	0.13	1.17 \pm 1.43
F3	1.5	0.30 \pm 0.03	58.21 \pm 0.08	1.95 \pm 0.18	0.15	1.32 \pm 1.36
F4	1.5	0.50 \pm 0.05	58.81 \pm 0.07	2.36 \pm 0.18	0.20	0.81 \pm 0.66
Cloud G						
G1	2.1	0.09 \pm 0.03	41.79 \pm 0.16	1.11 \pm 0.36	0.08	5.35 \pm 5.78
G2*	2.2	\leq 0.08	-	-	\leq 0.07	0.1 \pm 2.34
G3	2.0	0.17 \pm 0.03	41.27 \pm 0.20	1.84 \pm 0.39	0.09	1.25 \pm 3.04
Cloud H						
H1	1.5	0.27 \pm 0.03	45.09 \pm 0.10	1.89 \pm 0.28	0.13	4.87 \pm 2.44
H2	1.5	0.26 \pm 0.03	45.47 \pm 0.08	1.62 \pm 0.32	0.15	1.38 \pm 1.44
H3	1.5	0.50 \pm 0.05	45.41 \pm 0.23	5.13 \pm 0.63	0.09	0.10 \pm 3.98
H4	1.6	0.23 \pm 0.02	45.51 \pm 0.08	1.57 \pm 0.18	0.14	0.81 \pm 1.43
H5	1.5	0.31 \pm 0.03	45.43 \pm 0.10	2.06 \pm 0.24	0.14	0.56 \pm 1.69
H6	2.8	0.26 \pm 0.03	45.37 \pm 0.06	0.74 \pm 0.22	0.33	0.10 \pm 1.58

(a) * indicates 3σ upper-limit values for T_{peak} .(b) The integrated intensity upper limits are calculated as $3\sigma \times \sqrt{\delta\nu} \times \Delta\nu$, with $\delta\nu$ the spectral resolution (0.68 km s $^{-1}$) and $\Delta\nu$ the average linewidth measured considering all cores.

Table A.5. The root mean square noise in the spectra rms, integrated intensities $\int T_{\text{mb}} d\nu$, radial velocity ν , line widths $\Delta\nu$, peak temperature T_{peak} of the HCN ($1\rightarrow 0$, $F=2\rightarrow 1$) hyperfine component and optical depth τ of the HCN($1\rightarrow 0$) emission in cores within Clouds G and H.

Core	rms [10^{-2} K]	$\int T_{\text{mb}} d\nu$ [K kms $^{-1}$]	ν [kms $^{-1}$]	$\Delta\nu$ [kms $^{-1}$]	T_{peak} [K]	τ
Cloud G						
G1	1.4	0.95 \pm 0.07	42.66 \pm 0.16	4.05 \pm 0.31	0.22	0.10 \pm 0.16
G3	2.3	1.03 \pm 0.05	40.43 \pm 0.05	1.17 \pm 0.09	0.82	0.10 \pm 0.02
Cloud H						
H2	1.6	3.32 \pm 0.05	44.06 \pm 0.02	3.19 \pm 0.05	0.98	1.29 \pm 0.18
H3	1.4	4.71 \pm 0.06	43.80 \pm 0.02	3.48 \pm 0.05	1.27	1.16 \pm 0.13
H4	1.5	3.26 \pm 0.06	44.47 \pm 0.03	3.34 \pm 0.07	0.92	0.71 \pm 0.04
H5	2.0	3.32 \pm 0.05	44.43 \pm 0.04	3.36 \pm 0.06	0.93	1.84 \pm 0.29