

**Post-exercise cold-water immersion modulates skeletal muscle PGC-1 $\alpha$  mRNA expression in immersed and non-immersed limbs: evidence of systemic regulation**

Allan R <sup>1,3</sup>, Sharples AP<sup>1</sup>, Close GL<sup>1</sup>, Drust B<sup>1</sup>, Shepherd SO<sup>1</sup>, Dutton J<sup>2</sup>, Morton JP<sup>1</sup> and Gregson W<sup>1</sup>

<sup>1</sup>Research Institute for Sport and Exercise Sciences  
Liverpool John Moores University  
Byrom St Campus  
Liverpool  
UK

<sup>2</sup>Norwich Medical School  
University of East Anglia  
Norwich Research Park  
Norwich  
NR4 7TJ

<sup>3</sup>Centre for Applied Sport and Exercise Sciences  
University of Central Lancashire  
Preston  
UK

**Running Head:** Cold-water immersion and PGC-1 $\alpha$

Address for correspondence:  
Robert Allan  
Football Exchange, Research Institute for Sport and Exercise Sciences  
Liverpool John Moores University  
Tom Reilly Building  
Byrom St Campus  
Liverpool  
L3 3AF  
United Kingdom  
Email: [r.j.allan@2013.ljmu.ac.uk](mailto:r.j.allan@2013.ljmu.ac.uk) / [rallan1@uclan.ac.uk](mailto:rallan1@uclan.ac.uk)

## **New & Noteworthy**

We report for the first time that post-exercise cold-water immersion of one limb also enhances PGC-1 $\alpha$  expression in a contra-lateral non-immersed limb. We suggest increased systemic  $\beta$ -adrenergic stimulation, and not localised cooling per se, exerts regulatory effects on local signalling cascades thereby modulating PGC-1 $\alpha$  expression. These data therefore have important implications for research designs that adopt contralateral non-immersed limbs as a control condition, whilst also increasing our understanding of potential mechanisms underpinning cold-mediated PGC-1 $\alpha$  responses.

## **Author Contributions**

Conception and design of the experiments: RA, WG, JM, APS, BD; Collection, analysis and interpretation of data: RA, APS, JD, JM, WG, SS, GC; Drafting the article and Critical Revision of the article for important intellectual content: RA, APS, GC, SS, BD, JD, JM, WG. All authors approved the final version for publication and agree to be accountable for all aspects of the work.

## Abstract

Mechanisms mediating post-exercise cold-induced increases in PGC-1 $\alpha$  gene expression in human skeletal muscle are yet to be fully elucidated, but may involve local cooling effects on AMPK and p38 MAPK related signalling and/or increased systemic  $\beta$ -adrenergic stimulation. We aimed to therefore examine whether post-exercise cold-water immersion enhancement of PGC-1 $\alpha$  mRNA is mediated through local or systemic mechanisms. Ten subjects completed acute cycling (8x5 min at ~80% peak power output) followed by seated-rest (CON) or single-leg cold-water immersion (CWI; 10 min, 8°C). Muscle biopsies were obtained pre-, post- and 3 h post-exercise from a single limb in the CON condition but from both limbs in CWI (thereby providing tissue from a CWI and non-immersed limb, NOT). Muscle temperature decreased up to 2 h post-exercise following CWI (-5°C) in the immersed limb, with lesser changes observed in CON and NOT (-3°C;  $P<0.05$ ). No differences between limbs were observed in p38MAPK phosphorylation at any time point ( $P<0.05$ ), whilst a significant interaction effect was present for AMPK phosphorylation ( $P=0.031$ ). Exercise (CON) increased gene expression of PGC-1 $\alpha$  3 h post-exercise (~5-fold;  $P<0.001$ ). CWI augmented PGC-1 $\alpha$  expression above CON in both the immersed (CWI; ~9-fold;  $P=0.003$ ) and NOT limbs (~12-fold;  $P=0.001$ ). Plasma Normetanephrine concentration was higher in CWI vs. CON immediately post-immersion (860 vs. 665 pmol/L;  $P=0.034$ ). We report for the first time that local cooling of the immersed limb evokes transcriptional control of PGC1- $\alpha$  in the non-immersed limb, suggesting increased systemic  $\beta$ -adrenergic activation of AMPK may mediate, in part, post-exercise cold-induction of PGC-1 $\alpha$  mRNA.

**Key Words:** PGC-1 $\alpha$ , CWI, Normetanephrine, AMPK

## 87    **Introduction**

88    It is well established that the transcriptional co-activator peroxisome proliferator-activated  
89    receptor  $\gamma$  coactivator-1 $\alpha$  (PGC-1 $\alpha$ ), the proposed “master regulator” of skeletal muscle  
90    mitochondrial biogenesis (37), is sensitively controlled by acute and chronic exercise (3, 4, 10,  
91    36). Upstream control of PGC-1 $\alpha$  includes phosphorylation by energy and stress sensing  
92    kinases AMP-activated protein kinase (AMPK) (23) and p38 mitogen-activated protein kinase  
93    (p38 MAPK) (2). Consistent with its initial discovery as cold-inducible (34, 38), recent studies  
94    have examined the potential of acute post-exercise cold exposure to also modulate PGC-1 $\alpha$   
95    expression. For example, in human tissue, both cold-ambient temperatures (42, 43) and post  
96    exercise cold-water immersion (18, 20) enhances the skeletal muscle PGC-1 $\alpha$  gene  
97    transcription and protein translational response versus exercise alone. The precise  
98    mechanism(s) mediating cold-induced regulation of the PGC-1 $\alpha$  transcriptional pathway have  
99    yet to be fully determined, though the cooling-induced alterations in muscle blood flow (14,  
100    30) are unlikely to mediate these effects (33, 45). It is noteworthy, however, that chronic cold-  
101    induced changes in PGC-1 $\alpha$  protein content arise in conjunction with increased activity of  
102    signalling kinases AMPK and p38 MAPK (18). These data suggest that cooling of the skeletal  
103    muscle tissue may play a role in mediating the post-exercise cold-induction of PGC-1 $\alpha$  mRNA  
104    through activation of local signalling kinases.

105    Alternatively, systemic control via increased  $\beta$ -adrenergic activity is suggested to play a potent  
106    role in mediating the effects of cold exposure on PGC-1 $\alpha$  expression via AMPK (29) and  $\beta$ 2-  
107    adrenergic receptor (31) mechanisms. Indeed, plasma norepinephrine concentrations remain  
108    higher following high-intensity exercise after cold-water immersion versus control conditions  
109    (13). Studies to date have utilised the non-immersed limb as the control condition without the  
110    use of a true control (no cooling) condition (11, 18, 20). By assuming the response to cold-  
111    water immersion is mediated locally, such experimental designs do not permit examination of

the role of systemic versus localised mechanisms in mediating cold-induced changes in PGC-1 $\alpha$ . Indeed, Ihsan et al. (20) observed that PGC-1 $\alpha$  gene transcription was not induced in a non-immersed control limb, despite the limb having previously been exercised, and exercise being a potent stimulus to induce PGC-1 $\alpha$  expression. This would therefore suggest that an induction of PGC-1 $\alpha$  expression in an immersed limb occurred by way of cold-induced mechanisms. In this regard, we suggest that the increased systemic  $\beta$ -adrenergic activity associated with post-exercise cooling of the immersed limb could also modulate PGC-1 $\alpha$  expression in the non-immersed contralateral limb.

Therefore, the aim of the present study was to examine whether the cold-water induced increase in PGC-1 $\alpha$  mRNA observed post exercise is mediated through local or systemic mechanisms. To this end, we employed a novel experimental design (in a repeated measures crossover design) where ten recreationally active males completed an acute cycling protocol (8x5 min at ~80% peak power output) followed by a seated-rest condition (CON) or single-leg cold-water immersion (CWI; 10 min, 8°C). Muscle biopsies were obtained pre-, post- and 3 h post-exercise from a single limb in the CON condition but from both limbs post-exercise in the CWI trial (thereby providing tissue from a CWI and non-immersed limb, NOT). In this way, our design allowed us to obtain tissue from true control conditions but yet, also sample tissue from an immersed and non-immersed limb that was subject to the same hormonal milieu.

## Methods

*Subjects.* Ten recreationally active healthy males (age  $26 \pm 4$  y; body mass  $79.29 \pm 6.73$  kg; height  $180 \pm 5$  cm;  $\dot{V}O_{2\text{peak}}$   $51.46 \pm 9.07$  mL.kg<sup>-1</sup>.min<sup>-1</sup>; peak power output (PPO)  $265.2 \pm 38.33$  W; mean  $\pm$  SD) participated in this study. Subjects were instructed to refrain from exercise, alcohol and caffeine 48 hours prior to the test day. All subjects gave written informed consent

to participate after details and procedures of the study had been fully explained. Subjects had no history of neurological disease or musculoskeletal abnormality and none were under any pharmacological treatment during the course of the study. Each subject was medically screened by a practising GP prior to participation for their individual risk associated with high-intensity exercise, muscle biopsy procedures and immersion in cold water. All procedures performed in the study were approved by the Ethics Committee of Liverpool John Moores University and in accordance with the 1964 Helsinki declaration and its later amendments.

*Experimental Design.* In a repeated-measures crossover design, subjects completed two high-intensity intermittent cycling protocols followed either by 10 min seated rest (CON) or single-leg CWI (10 min at 8°C). Muscle biopsies were obtained from the vastus lateralis immediately before, post- and 3 h post-exercise. During the CWI trial muscle biopsies in the post-exercise period were taken from both the immersed and non-immersed leg. This design allowed us to assess the impact of the cold stimulus locally (immersed leg) and systemically (non-immersed leg) against a relevant resting control. Both experimental trials were conducted in a counterbalanced, randomized order with at least 10 days between conditions.

*Experimental Protocol.* Subjects attended the laboratory on 3 separate occasions. On the first occasion, subjects completed an incremental exercise test to fatigue for the determination of  $\dot{V}O_{2max}$  and PPO (15). Results from this test were used to determine the power output necessary for cycling at a proportion of PPO on subsequent test days (detailed below). Prior to the first test day, subjects completed a 24-hour food diary to be replicated before the second trial. Upon arrival at the laboratory (0900h) subjects were fitted with a heart-rate monitor (Polar RS400, Kempele, Finland), skin and rectal temperature probes (MHF-18050-A and MRV-55044-A, Ellab, Rodovre, Denmark). Legs were marked for subsequent insertion of muscle temperature needles; area of insertion was calculated as half the length of the femur, over the ‘belly’ of the vastus lateralis. The needle thermistor was then placed at a depth of 3cm, plus one-half of the

161 skinfold measurement, for the determination of deep muscle temperature (3cm). Following 10-  
162 min resting in a supine position, baseline measures of heart rate (HR), temperature and oxygen  
163 uptake ( $\dot{V}O_2$ ; Oxycon Pro, Jaeger, Wuerzburg, Germany) were assessed. Resting venous blood  
164 samples were drawn from a superficial vein in the anti-cubital crease of the forearm using  
165 venepuncture cannulation (BD Nexiva Closed IV Catheter 22G Blue, Becton Dickinson,  
166 Oxford, UK). Resting muscle temperature was assessed using a needle thermistor (13050;  
167 Ellab, Rodovre, Denmark) inserted into the vastus lateralis at 3cm depth as previously  
168 described (30). Finally, resting muscle biopsy samples from the vastus lateralis (~30–50 mg  
169 wet wt) were obtained under local anaesthesia (0.5% marcaine) using a Pro-Mag 2.2 biopsy  
170 gun (MD-TECH, Manan Medical Products, Northbrook, IL). At rest only a single leg was  
171 biopsied in the cooling trial (CWI) in attempt to reduce the stress experienced by subjects. This  
172 leg was randomised between the immersed and non-immersed limbs to exclude potential  
173 variation caused by leg dominance; all subsequent biopsies were completed in both legs, at  
174 each time point, 2cm proximal to the previous incision.

175 Following the resting biopsy, subjects completed a high-intensity intermittent cycling protocol  
176 consisting of  $8 \times 5$  min bouts at 82.5% PPO separated by 1 min rest (adapted from 44) followed  
177 by either single-legged CWI (CWI: 10 min at 8°C; NOT: non-immersed leg i.e. 10 min at room  
178 temperature) or a control condition (CON; seated rest). CWI was conducted using a specialised  
179 seated mechanical hoist to lower the subject so that one leg was placed inside the cold-bath,  
180 allowing the other (NOT) to remain outside, and dry. Subjects then recovered in a semi-reclined  
181 position under normal laboratory temperatures (~21°C) until 3-hours post-exercise. Measures  
182 of heart rate, skin temperature (thigh and calf) and rectal temperature were recorded throughout  
183 the exercise and recovery periods. Oxygen uptake was measured during the final minute of  
184 each high-intensity bout of exercise, during immersion, immediately post-immersion and again  
185 at 1, 2 and 3 h post-exercise. Ratings of perceived exertion (RPE) were assessed during the

final minute of each exercise bout (5), whilst subjective measures of perceived shivering were assessed throughout water immersion and the 3h recovery period using a visual analogue scale from 1 (No shivering) to 5 (Intense Shivering) (24). Laboratory temperatures remained stable throughout ( $21.18 \pm 0.76^{\circ}\text{C}$ ) and at no point were subjects allowed to rub themselves dry or shower (changing into dry shorts after immersion was allowed). Subjects were advised to wear the same clothes between trials. Muscle temperature was assessed post-exercise, immediately post-immersion and at 1, 2 and 3h post-exercise. Venous blood samples were also drawn at these times. Bi-lateral muscle biopsies occurred immediately after exercise and 3h post-exercise.

Venous blood samples were drawn from a superficial vein in the anti-cubital crease of the forearm using standard venepuncture techniques (BD Nexiva Closed IV Catheter 22G Blue, Becton Dickinson, Oxford, UK). Blood samples (~10ml) were collected into vacutainer tubes (Becton Dickinson, Oxford, UK) containing EDTA and stored on ice until centrifugation at  $1500 \text{ rev.min}^{-1}$  for 15-min at  $4^{\circ}\text{C}$ . Following centrifugation, aliquots of plasma were stored at  $-80^{\circ}\text{C}$  for later analysis. Plasma metanephrine and Normetanephrine concentrations were measured using liquid chromatography tandem mass spectrometry as previously described (35). All samples were analysed in duplicate, with the mean value employed.

*rt-qRT-PCR*. Skeletal muscle samples (~30 mg) were transferred to 2 ml lysing tubes containing 1.4 mm ceramic beads (Lysing Matrix D, MP Biomedicals, UK) containing 1 ml ice-cooled TRI- reagent (Life Technologies Ltd, UK) and homogenised at 6 m/s for 3 x 40 seconds, separated by 5 minutes cooling on ice (MP Fastprep-24, MP Biomedicals, UK). RNA was extracted according to the TRI-reagent manufacturer's instructions. RNA concentration and purity were assessed by UV spectroscopy at ODs of 260 and 280 nm using a Nanodrop 3000 (Fisher, Roskilde, Denmark). A target of  $A_{260} / A_{280}$  ratio was set at 1.8 to 2.2. Seventy ng RNA was used for each PCR reaction. Primer sequences (Table 1) were identified using



211 Gene (NCBI, <http://www.ncbi.nlm.nih.gov/gene>) and designed using Primer-BLAST  
212 (NCBI, <http://www.ncbi.nlm.nih.gov/tools/primer-blast>). Sequence homology searches  
213 ensured specificity and that all primers had no potential unintended targets. The primers were  
214 ideally designed to yield products spanning exon-exon boundaries to prevent any amplification  
215 of gDNA. Three or more GC bases in the last five bases at the 3' end of the primer were  
216 avoided. Secondary structure interactions (hairpins, self-dimer and cross dimer) within the  
217 primer were avoided. All primers were between 16 and 25 bp, and amplified a product of  
218 between 67-212 bp. Primers were purchased from Sigma (Suffolk, UK).

219 rt-qRT-PCR amplifications were performed using QuantiFast™ SYBR® Green RT-PCR one  
220 step kit on a Rotor-gene 3000Q (Qiagen, Crawley, UK) supported by rotor-gene software  
221 (Hercules, CA, USA). rt-qTR-PCR was performed as follows: hold 50°C for 10 min (reverse  
222 transcription/cDNA synthesis), 95°C for 5 min (transcriptase inactivation and initial  
223 denaturation step) and PCR steps of 40 cycles; 95°C for 10s (denaturation), 60°C for 30s  
224 (annealing and extension). Upon completion, dissociation/melting curve analysis were  
225 performed to reveal and exclude non-specific amplification or primer-dimer issues (all melt  
226 analysis in this study presented single reproducible peaks for the reference gene and each target  
227 gene suggesting amplification of a single product). Following initial screening of suitable  
228 reference genes, GAPDH showed the most stable  $C_t$  values across all RT-PCR runs, subjects  
229 and regardless of experimental condition ( $23.54 \pm 1.69 C_t$ ; 7% Co-efficient of variation) and  
230 was selected as the reference gene in all RT-PCR assays. The average PCR efficiency was 90%  
231 and variation for all genes was less than 4.3%. The relative gene expression levels were  
232 calculated using the comparative  $C_t$  ( $\Delta\Delta C_t$ ) equation (41) where the relative expression was  
233 calculated as  $2^{-\Delta\Delta C_t}$  and where  $C_t$  represents the threshold cycle. mRNA expression for all target  
234 genes was calculated relative to the reference gene (GAPDH; subject's own samples reference)  
235 within same subject and condition and to a calibrator of Pre-exercise.

236 *SDS-PAGE and Western Blotting.* Approximately 30 mg of frozen muscle was homogenized  
237 using 2.4 mm ceramic beaded tubes (6 m/s for 3 x 40 seconds, separated by 5 minutes cooling  
238 on ice; MP Fastprep-24, MP Biomedicals, UK), in 500 µl of ice-cold lysis buffer [25 mM  
239 Tris·HCl (pH 7.4), 50 mM NaF, 100 mM NaCl, 5 mM EGTA, 1 mM EDTA, 10 mM Na-  
240 pyrophosphatase, 1% Triton X-100] and supplemented with a protease phosphatase inhibitor  
241 cocktail (Halt Protease and Phosphatase Inhibitor 186 Cocktail, Thermo Scientific, # 78442).  
242 The resulting homogenates were centrifuged at 14,000 g for 10 min at 4°C, and the supernatant  
243 was collected. The protein content of the supernatant was determined using a bicinchoninic  
244 acid assay (Sigma, UK). Each sample was diluted with an equal volume of 2X Laemmli buffer  
245 (National Diagnostics) and boiled for 10 min at 100°C. Forty µg of total protein from each  
246 sample was loaded (65 µg for phosphorylated analytes) and then separated in Tris-glycine  
247 running buffer (10x Tris/glycine, Geneflow, Staffordshire, UK) using self-cast 10% separating  
248 [33% Protogel; (30% w/v acrylamide: 0.8% (w/v) Bis-Acrylamide stock solution (37.5:1)),  
249 25% Protogel resolving buffer (1.5M Tris-HCL, 0.4% SDS, pH 8.8), 41% ddH<sub>2</sub>O, 100µl 10%  
250 APS, 20µl TEMED] and 4% stacking [13% Protogel, 25% Protogel Stacking buffer (0.5M Tris  
251 HCL, 0.4% SDS, pH 6.8), 61% ddH<sub>2</sub>O, 100µl 10% APS, 20µl TEMED] gels (National  
252 Diagnostics, Geneflow, UK). Gels were transferred semidry onto nitrocellulose membrane  
253 (Transblot Turbo, BioRad) for 30 min at 25V and 1.0 mA in transfer buffer [10% TRIS/glycine  
254 (Sigma), 20% methanol, 70% ddH<sub>2</sub>O). After transfer, membranes were briefly washed in TBST  
255 (0.19 M Tris pH 7.6, 1.3 M NaCl, 0.1% Tween-20] before being blocked for 1 h at room  
256 temperature in TBST with 1% BSA. The membranes were then washed for 3 x 5 min in TBST  
257 before being incubated overnight at 4°C with antibodies for anti-phospho-AMPK Thr172 (cat  
258 no: 2532), p38 MAPKThr180/Tyr182 (cat no: 9211) (Cell Signalling) as well as total protein  
259 content of AMPK (cat no: 2531), p38 MAPK (cat no: 9212) (Cell Signalling, UK), GAPDH  
260 (25778; Santa Cruz), and PGC-1α (Calbiochem, Merck Chemicals, UK) all at concentrations

of 1:1000 in 1 x TBS. The following morning, membranes were washed for a further 3 x 5 min in TBST and subsequently incubated with goat anti-rabbit horseradish peroxidase-conjugated secondary antibody (Bio-Rad, UK) for 1h at room temperature. After a further 3 x 5 min washes in TBST, membranes were exposed in a chemiluminescence liquid (SuperSignal, Thermo Fisher Scientific, Rockford, IL) for 5 min (30 sec for GAPDH). Membranes were visualized using a Bio-Rad Chemi-doc system, and band densities were determined using Image Lab image-analysis software (Bio Rad, UK). Samples from each subject for all exercise conditions were run on the same gel and statistical analysis conducted on raw densitometry data. Phosphorylated AMPK<sup>Thr172</sup> and p38 MAPK<sup>Thr180/Tyr182</sup> were normalised to their total protein, as these did not change significantly across blots or samples ( $P > 0.05$ ). PGC-1 $\alpha$  was normalised to GAPDH.

*Statistical analysis.* All data are presented as mean  $\pm$  SD, unless otherwise stated. Baseline data, distance cycled, exercise HR and RPE were compared between conditions using a Paired Samples T-test. A two-factor (two condition  $\times$  time) within-participants general linear model was used to evaluate the effect of time (baseline v post exercise) with shared baseline data used for NOT and CWI (Statistical Package for the Social Sciences version 21.0; SPSS Inc., Chicago, IL). A two-factor (three condition  $\times$  time) within-participants general linear model was subsequently used to evaluate the influence of the cooling intervention following exercise and the 3h post exercise period. The main effects for condition and time was followed up using planned LSD multiple contrasts. Where a significant condition by time interaction was observed, the post exercise to 3h post exercise change scores were calculated and compared across the 3-conditions using LSD multiple contrasts. The ES magnitude was classified as trivial ( $<0.2$ ), small ( $>0.2-0.6$ ), moderate ( $>0.6-1.2$ ), large ( $>1.2-2.0$ ) and very large ( $>2.0-4.0$ ) (17). The  $\alpha$  level for evaluation of statistical significance was set at  $P < .05$ .

286

## 287 **Results**

### 288 *Exercise Response.*

289 Distance cycled (CON  $32.52 \pm 4.21$  km, CWI  $32.33 \pm 4.33$  km;  $P = 0.629$ , ES 0.04 Trivial),  
290 heart rate ( $P = 0.309$ , ES 0.13 Trivial),  $\dot{V}O_2$  (ml/kg.min<sup>-1</sup>;  $P = 0.855$ , ES 0.02 Trivial) and RPE  
291 ( $P = 0.637$ , ES 0.08 Trivial) were similar between CON and CWI trials (data not shown). Mean  
292 HR during the final minute of exercise was  $182 \pm 8$  beats.min<sup>-1</sup> in CON and  $183 \pm 8$  beats.min<sup>-1</sup>  
293 <sup>1</sup> in CWI, equating to ~94% HR max. The RPE in the final exercise bout was  $19 \pm 1$  AU and  
294  $19 \pm 1$  AU in the CON and CWI trials respectively.

295

### 296 *Recovery Response.*

297 *Metabolic Responses.* Heart rate (ES 0.90 Moderate) and  $\dot{V}O_2$ , (ES 1.20 Moderate) was higher  
298 in CWI vs CON during the post-exercise recovery period (Table 2;  $P < 0.001$ ). The change in  
299 HR and  $\dot{V}O_2$ , over time was also different between conditions ( $P < 0.001$ ) with increases in HR  
300 and  $\dot{V}O_2$ , occurring during the initial 2 minutes of immersion. Following immersion, HR and  
301  $\dot{V}O_2$  dropped below pre-immersion values and remained lower throughout the 3h recovery  
302 period (HR, ES  $>0.92$  Moderate;  $\dot{V}O_2$ , ES  $>1.25$  Large) ( $P < 0.05$ ).

303 *Thermoregulatory Responses.* Rectal temperature was similar between conditions throughout  
304 the post-exercise period (CON  $37.52 \pm 0.24^\circ\text{C}$ , CWI  $37.48 \pm 0.05^\circ\text{C}$ , ES 0.49 Small,  $P = 0.217$ ).  
305 The change in rectal temperature over time was different between conditions, with a small  
306 decline in rectal temperature occurring after 3 minutes of immersion until 3h post-exercise  
307 ( $P=0.034$ , ES  $>0.22$  Small). Thigh skin temperature was generally lower throughout the post-  
308 exercise recovery period in CWI versus CON (ES 6.26 Very Large) and NOT (ES 6.46 Very

Large) (Figure 1a,  $P < 0.001$ ). The change over time was also different between conditions, with thigh skin temperature continually decreasing in CWI and remaining lower than pre-immersion values until 1h post exercise (ES 3.0 Very Large,  $P < 0.001$ ). Values in CON and NOT limbs remained similar to pre-immersion throughout the 3h recovery period ( $P = 0.10$ , ES 0.57 Small).

Post-exercise muscle temperature (3 cm depth) was similar between CON (38.75°C), CWI (38.86°C) and NOT (38.54°C) (Figure 1b;  $P > 0.05$ ). During the 3 h recovery period muscle temperature (3cm) was lower in CWI versus CON (ES 1.60 Large) and NOT (ES 1.77 Large) ( $P < 0.001$ ). The change in muscle temperature over time was also different between conditions ( $P < 0.001$ ). Muscle temperature declined to a large extent immediately after immersion in the CWI limb, followed by a further gradual reduction during the remaining 3h post-exercise period ( $P < 0.001$ ). In CON and NOT conditions, muscle temperature was reduced to a lesser extent immediately following immersion followed by a further gradual reduction during the 3h post-exercise period ( $P = 0.246$ , ES 0.34 Small) (See Figure 1b).

Subjective shivering ratings were greater in CWI vs CON during the post-exercise recovery period (ES 1.20 Large,  $P = 0.067$ ). The change in shivering over time also tended to be different between conditions ( $P = 0.062$ ), with 'slight' shivering observed in the CWI condition during the first 2 minutes following immersion (ES  $>0.60$  Moderate). Slight shivering was also observed 2h post exercise in the CWI condition (ES 0.95 Moderate).

#### *AMPK and P38 MAPK activity and total abundance*

Phosphorylation of AMPK<sup>Thr172</sup> was not increased post-exercise ( $P = 0.242$ , ES 0.20 Small). At post-exercise and 3h post-exercise phosphorylation of AMPK<sup>Thr172</sup> was similar between conditions ( $P = 0.846$ , ES 0.03 Trivial). However, the change in AMPK<sup>Thr172</sup> between these time points was different between conditions ( $P = 0.031$ ; Figure 2). AMPK<sup>Thr172</sup>

phosphorylation increased in CWI vs. CON ( $P = 0.027$ , ES 1.22 Large) with a moderate increase in AMPK<sup>Thr172</sup> phosphorylation also observed in NOT vs. CON ( $P = 0.145$ , ES 0.70 Moderate). Representative Western blots are shown in Figure 5.

Exercise induced a small increase in phosphorylation of p38MAPK<sup>Thr180/Tyr182</sup> ( $P = 0.056$ , ES 0.44 Small, Figure 3). At post exercise and 3h post-exercise phosphorylation of p38MAPK<sup>Thr180/Tyr182</sup> was similar between conditions ( $P = 0.672$ ; ES 0.03, Trivial). No differences in the change in phosphorylation between these time points was observed between conditions ( $P = 0.268$ , Figure 3). Representative Western blots are shown in Figure 5.

#### *PGC-1alpha mRNA and protein abundance.*

PGC-1 $\alpha$  mRNA expression was moderately increased with exercise ( $P = 0.066$ , ES 0.92 Moderate, Figure 4a). At 3h post-exercise, expression was greater in CWI (ES 1.2 Moderate,  $P = 0.003$ ) and NOT (ES 1.6 Large,  $P = 0.001$ ) versus CON, but was similar between CWI and NOT (ES 0.6 Small,  $P = 0.141$ ) (Figure 4a). This reflected the greater change in expression in CWI and NOT conditions between post exercise and 3h post exercise time points ( $P = 0.001$ , Figure 4a). PGC-1 $\alpha$  protein content was not influenced by exercise ( $P = 0.092$ ) or CWI ( $P = 0.471$ , Figure 4b). Representative Western blots are shown in Figure 5.

#### *Additional gene expression*

Exercise induced increases in SIRT1 ( $P = 0.057$ , ES 0.8 Moderate) and NRF2 ( $P = 0.028$ , ES 0.6 Moderate) mRNA (data not shown). No changes were seen between conditions, or between conditions over time ( $P > 0.05$ ). Gene expression analysis for p53, COXIV, CS, TFAM, SIRT1, NRF2 and ERR $\alpha$  mRNA was not influenced by exercise or CWI ( $P > 0.05$ ; data not shown).

#### *Plasma Metanephrine and NorMetanephrine.*

Metanephrine concentrations were similar between conditions ( $P = 0.159$ , ES 0.15 Trivial). The change in metanephrine over time was also similar between conditions ( $P = 0.299$ ). Metanephrine concentration was increased post-exercise (ES 2.46 Very Large) and post-immersion (ES 0.77 Moderate) vs. baseline ( $P \leq 0.02$ ). Normetanephrine values were greater in CWI vs. CON ( $P = 0.034$ , ES 0.43 Small) with the largest difference seen post-immersion (860 vs. 665 pmol/L, CWI vs. CON, respectively). The change in Normetanephrine over time was similar between conditions ( $P = 0.821$ ). Normetanephrine concentrations increased with exercise (ES >4.70;  $P < 0.001$ ) and remained above baseline post-immersion (ES 1.52 Large) and 1hr post-exercise (ES 1.06 Moderate) ( $P < 0.001$ ). Concentrations returned to baseline at 2hr post-exercise (See Table 3).

## Discussion

The aim of the present study was to examine whether the post-exercise cold-water induced increase of PGC-1 $\alpha$  mRNA is mediated through local or systemic mechanisms. Using a novel experimental design, we report for the first time the appearance of systemic “cross-talk” between immersed and non-immersed limbs, as evidenced by the similar increase in PGC-1 $\alpha$  mRNA in these limbs after single-legged CWI. Additionally, we suggest that this effect could be mediated by  $\beta$ -adrenergic induced stimulation of AMPK. In addition to providing novel data on the potential mechanisms mediating post-exercise cold-induced enhancement of PGC-1 $\alpha$  expression, our data also have potential implications for research designs that utilise as non-immersed limbs as control conditions.

Since its initial discovery (38) the importance of the post-exercise PGC-1 $\alpha$  response to the oxidative adaptive process has been examined extensively, with mRNA increases ranging 5 to 10-fold commonly observed at 3-4 hours following exercise (3, 4, 21, 36). More recently, cold-

380 ambient temperatures (42, 43) and post exercise cold-water immersion (18-20, 24), have also  
381 been shown to enhance (~2- to 4- fold greater) the skeletal muscle PGC-1 $\alpha$  gene transcription  
382 and protein translational response versus exercise alone. In line with such observations, the  
383 high-intensity intermittent cycling protocol used in the present study elicited a ~5-fold increase  
384 in PGC-1 $\alpha$  mRNA at 3 hours post-exercise. Furthermore, CWI enhanced this response to  
385 a greater extent than exercise alone (~9-fold increase vs. Pre in the CWI immersed limb).

386 There have been suggestions that reduced tissue temperature may be responsible for the  
387 differences observed between cold-treated and control limbs. This stems from the initial  
388 discovery that PGC-1 $\alpha$  was cold inducible in animals (6, 9, 38). In humans, recent data  
389 from Ihsan and colleagues (18, 20) implicates a reduction in tissue temperature in the cold-  
390 induced increases of PGC-1 $\alpha$ , as increases in mRNA (3h post-exercise) and protein content  
391 (after 4 weeks of training) were seen only in a cooled limb, and not in the contralateral non-  
392 immersed limb. Potential mechanisms underpinning such responses include activation of non-  
393 noxious thermoreceptors via reduced skin temperature (Hensel & Boman, 1960, 22). Within  
394 the present study, both the NOT and CWI limbs displayed similar acute PGC-1 $\alpha$  mRNA  
395 expression in the 3 h recovery period, whilst skin and muscle temperature were significantly  
396 reduced in the CWI limb only. Indeed, the non-immersed limb (NOT) showed a similar  
397 temperature profile (skin and muscle temperature) to that of CON, where the magnitude of  
398 PGC-1 $\alpha$  mRNA response was almost half when compared with CWI and NOT. When taken  
399 together, these data suggest that alterations to local muscle temperature do not play a significant  
400 role in cold-induced regulation of PGC-1 $\alpha$  expression.

401 As a result of increased local PGC-1 $\alpha$  gene expression in the non-immersed exercised limb to  
402 a similar magnitude as the immersed limb, we sought to consolidate the role of the upstream  
403 kinases AMPK and p38-MAPK (7, 23, 47) in their ability to regulate PGC-1 $\alpha$  transcription.  
404 p38MAPK is a stress activated kinase that has been shown extensively to be phosphorylated



after acute exercise, independent of intensity (10). Moreover, p38MAPK can exert its effect upon PGC-1 $\alpha$  transcription via upregulated ATF2 activity at the PGC-1 $\alpha$  promoter (2). In the current study, exercise induced a small (ES 0.44, 1.5-fold) increase in phosphorylation with no further response to cooling. Moreover, our data supports previous data showing acute post-exercise phosphorylation of p38MAPK locally in skeletal muscle tissue occurs systemically (46). Exercise-induced intensity dependant AMPK phosphorylation is a well reported phenomenon (8, 12, 28) in rodent and human studies (34, 10). Moreover, AMPK is implicated in PGC-1 $\alpha$  activity via direct phosphorylation, initiating many of the important gene regulatory functions of PGC-1 $\alpha$  in skeletal muscle (23). The post-exercise increase in phosphorylation of AMPK in the present study was similar in magnitude to previous work (3) from our laboratory, albeit failing to achieve statistical significance. Notwithstanding this, large and moderate effect sizes were observed at 3 h post-exercise in the CWI and NOT limbs vs. CON, respectively. These greater increases from post-exercise to 3 h post-exercise in the immersion trial (CWI and NOT limbs), compared to a slight decline in CON suggest the increases in phosphorylation of AMPK during the post-cooling period are controlled by a systemic mechanism, possibly adrenergic control via cold-augmented plasma Normetanephrine.

Epinephrine and norepinephrine are both dual  $\alpha$ - and  $\beta$ - adrenergic agonists. Previous in vivo and in vitro incubation techniques utilising  $\alpha$ - and  $\beta$ - adrenergic agonists have reported increased AMPK activation in rodent skeletal muscle (32), adipose tissue (26) and cell cultures (48) implicating catecholamines as a potential AMPK activator. Despite this, support for the above hypothesis is currently conflicting, and is limited by distinct differences in methodological design and species studied. In rodent muscle, acute infusion of adrenergic agonists/antagonists have previously shown to be ineffective at altering PGC-1 $\alpha$  transcriptional activity (39, 40) and its upstream effector p38-MAPK (25). In contrast, treatment with  $\beta$ - adrenergic agonists/antagonists has been shown to induce and inhibit PGC-1 $\alpha$  respectively

(31), whilst incubation of rodent skeletal muscle with the adrenergic agonist phenylephrine increased the activity of the upstream regulator of PGC-1 $\alpha$ , AMPK (32). When results from these studies are taken together, it could be suggested that adrenergic activation of AMPK is a potential mechanism to explain the systemic increases in PGC-1 $\alpha$  gene expression described herein.

In humans, one study has investigated the impact of higher catecholamine levels on AMPK phosphorylation in human skeletal muscle (27). These authors assessed muscle biopsies from an exercised and non-exercised limb in conditions of heightened catecholamine release. Results showed AMPK activity was restricted to contracting muscle only, with no systemic effects notable in the non-exercised limb despite the increased catecholamine levels. Importantly, in our study all limbs were exercised before undergoing cold exposure. It therefore may be that the cold induction of  $\beta$ -adrenergic pathways (via increased catecholamines) presented in this manuscript allows an additive response to exercise stimulated AMPK phosphorylation. Further studies are now required to verify this signalling response in related experimental conditions in human skeletal muscle.

Another pathway by which increased catecholamine's may enhance PGC-1 $\alpha$  transcription is via increased activation of  $\beta$ -adrenergic receptors. Activation of these receptors increases intracellular cAMP, which could ultimately activate CREB function on the PGC-1 $\alpha$  promoter (1). However, evidence exists to show  $\beta$ -adrenergic stimulation does not activate a p38MAPK - ATF2 - CREB - PGC-1 $\alpha$  signalling axis in skeletal muscle (25). These findings resonate with results from the present study as no changes in phosphorylation of p38MAPK were observed alongside increased plasma Normetanephrine concentrations. Further work is required to investigate the influence of the dual stress of exercise and cold temperature upon this signalling axis.

Downstream of PGC-1 $\alpha$ , Slivka and colleagues (42) noted that recovery in cold ambient temperatures (4h in 7°C ambient air) reduced the expression of the transcription factors Nuclear-respiratory factor 2 (NRF2) and estrogen-related receptor  $\alpha$  (ERR $\alpha$ ), whilst having no effect on mitochondrial transcription factor a (TFAM). Their importance to the adaptive response is highlighted by their roles in oxidative metabolism. Our data suggest the immersion protocol used herein was not sufficient to induce such changes. Indeed, 10 minutes of single-legged immersion offers a much smaller cooling stimulus than the 4 hours in cool ambient temperatures, as used by Slivka et al. (42). In addition, an acute increase in PGC-1 $\alpha$  gene expression was not followed by changes in PGC-1 $\alpha$  total protein content, perhaps due to the acute time-frame of sampling applied in the present study. With this lack of change in total protein content it is however, unlikely that changes in gene expression of downstream genes such as NRF2, TFAM, COXIV, CS, ERR $\alpha$  presented here, would be affected by its upstream protein function as a transcription factor (PGC-1 $\alpha$ ). More research is required to understand the effect and dose response of a cold stimulus on downstream targets of PGC-1 $\alpha$ .

It is difficult to explain the differences in PGC-1 $\alpha$  mRNA results between the present study and Ihsan et al. (20), particularly the difference in non-immersed limbs. Moreover, reasons as to why Ihsan and colleagues failed to see the expected exercise-induced response in their non-immersed control limb, with minor changes from baseline at 3h post-exercise (1.5-fold increase vs. 3-5-fold increase usually seen), remain unclear. One possible explanation might include the differing muscle recruitment patterns occurring between the exercise protocols utilised (cycling vs. running). Ultimately, results from the present study have future implications on scientific study designs. Those wishing to investigate cold induced post-exercise responses in skeletal muscle must be aware of the evidence that implicates a systemic response of Normetanephrine, and a local response of p-AMPK in both immersed and non-immersed limbs. Further evidence is required to support the impact of systemic transcriptional responses in unilateral research

designs, as such designs may be liable to error; potentially underestimating the actual response occurring in the immersed limb if relativized to a non-immersed limb instead of a resting control. Ultimately, the choice of scientific design lies with the question posed, as contralateral designs remain useful for understanding both local and systemic responses.

In summary, the present study characterises for the first time the mechanistic control of cold induced PGC-1 $\alpha$  mRNA expression. Data herein indicate a reduction in tissue temperature (2-3°C) plays a limited role as similar levels of PGC-1 $\alpha$  mRNA expression are observed in an immersed and non-immersed limb despite a reduction in tissue temperature in the CWI limb only. Moreover, a cold-induced systemic increase in plasma Normetanephrine may impact localised phosphorylation events of the signalling kinase AMPK<sup>Thr172</sup>, with potential downstream effects upon rates of PGC-1 $\alpha$  mRNA expression. Future studies should investigate the role of  $\beta$ -adrenergic receptors in Normetanephrine induced AMPK phosphorylation and the signalling role of MEF2 and CRE/ATF2 sites to confirm a link between catecholamines and PGC-1 $\alpha$ . Moreover, due to the acute nature of the present study more work is required to investigate whether the response seen herein is maintained over a more chronic term.

#### **Acknowledgements**

All work was conducted at Liverpool John Moores University. Analysis of catecholamines was conducted by John Dutton at UEA. The authors would like to thank Dean Morrey and Gemma Miller for their assistance in laboratory co-ordination and Kelly Hammond and Chris Mawhinney for their assistance in data collection.

#### **Disclosures**

The authors report no conflicts of interest.

## 502    **References**

- 503    **1. Akimoto T, Li P, Yan Z.** Functional interaction of regulatory factors with the Pgc-1alpha  
504    promoter in response to exercise by in vivo imaging. *Am J Physiol Cell Physiol* 295: C288-  
505    292, 2008.
- 506    **2. Akimoto T, Pohnert SC, Li P, Zhang M, Gubs C, Rosenberg PB, Williams RS, Yan Z.**  
507    Exercise stimulates PGC-1 alpha transcription in skeletal muscle through activation of the  
508    p38 MAPK pathway. *J Biol Chem* 280: 19587-19593, 2005.
- 509    **3. Bartlett JD, Joo CH, Jeong T-S, Louhelainen J, Cochran AJ, Gibala MJ, Gregson W,**  
510    **Close GL, Drust B, Morton JP.** Matched work high-intensity interval and continuous  
511    running induce similar increases in PGC-1alpha mRNA, AMPK, p38, and p53  
512    phosphorylation in human skeletal muscle. *J Appl Physiol* 112: 1135–1143, 2012.  
513
- 514    **4. Bartlett JD, Louhelainen J, Iqbal Z, Cochran AJ, Gibala MJ, Gregson W, Close GL,**  
515    **Drust B, Morton JP.** Reduced carbohydrate availability enhances exercise-induced p53  
516    signaling in human skeletal muscle: implications for mitochondrial biogenesis. *Am J Physiol*  
517    *Regul Integr Comp Physiol* 304: R450–8, 2013.  
518
- 519    **5. Borg G.** Psychophysical bases of perceived exertion. *Med Sci Sports Exerc* 14(5): 377-  
520    381, 1982.  
521
- 522    **6. Bruton JD, Aydin J, Yamada T, Shabalina IG, Ivarsson N, Zhang S-J, Wada M, Tavi**  
523    **P, Nedergaard J, Katz A, others.** Increased fatigue resistance linked to Ca<sup>2+</sup>-stimulated  
524    mitochondrial biogenesis in muscle fibres of cold-acclimated mice. *J Physiol* 588: 4275–  
525    4288, 2010.  
526
- 527    **7. Cantó C, Auwerx J.** AMP-activated protein kinase and its downstream transcriptional  
528    pathways. *Cell Mol Life Sci* 67: 3407–23, 2010.  
529
- 530    **8. Cochran AJR, Little JP, Tarnopolsky MA, Gibala MJ.** Carbohydrate feeding during  
531    recovery alters the skeletal muscle metabolic response to repeated sessions of high-intensity  
532    interval exercise in humans. *J Appl Physiol* 108: 628–36, 2010.  
533
- 534    **9. Eddy SF & Storey KB.** Differential expression of Akt, PPARgamma and PGC-1 during  
535    hibernation in bats. *Biochem Cell Biol* 81(4): 269-274, 2003.  
536
- 537    **10. Egan B, Carson BP, Garcia-Roves PM, Chibalin AV, Sarsfield FM, Barron N,**  
538    **McCaffrey N, Moyna NM, Zierath JR, O’Gorman DJ.** Exercise intensity-dependent  
539    regulation of peroxisome proliferator-activated receptor gamma coactivator-1alpha mRNA  
540    abundance is associated with differential activation of upstream signalling kinases in human  
541    skeletal muscle. *J Physiol* 588: 1779–1790, 2010.  
542
- 543    **11. Frohlich M, Faude O, Klein M, Pieter A, Emrich E, Meyer T.** Strength training  
544    adaptations after cold-water immersion. *J Strength Cond Res* 28(9): 2628-2633, 2014.  
545
- 546    **12. Gibala MJ, McGee SL, Garnham AP, Howlett KF, Snow RJ, Hargreaves M.** Brief  
547    intense interval exercise activates AMPK and p38 MAPK signaling and increases the  
548

expression of PGC-1alpha in human skeletal muscle. *J Appl Physiol* 106: 929–34, 2009.

**13. Gregson W, Allan R, Holden S, Phibbs P, Doran D, Campbell I, Waldron S, Joo CH, Morton JP.** Postexercise cold-water immersion does not attenuate muscle glycogen resynthesis. *Med Sci Sports Exerc* 45: 1174–81, 2013.

**14. Gregson W, Black MA, Jones H, Milson J, Morton J, Dawson B, Atkinson G, Green DJ.** Influence of cold water immersion on limb and cutaneous blood flow at rest. *The American J Sports Med* 39: 1316–1323, 2011.

**15. Hawley JA & Noakes TD.** Peak power output predicts maximal oxygen uptake and performance time in trained cyclists. *Eur J Appl Physiol* 65(1): 79-83, 1992.

**16. Hensel H & Boman KK.** Afferent impulses in cutaneous sensory nerves in human subjects. *J Neurophysiol* 23: 564-578, 1960.

**17. Hopkins WG, Marshall SW, Batterham AM, Hanin J.** Progressive statistics for studies in sports medicine and exercise science. *Med Sci Sports Exerc* 41: 3-12, 2009.

**18. Ihsan M, Markworth JF, Watson G, Choo HC, Govus A, Pham T, Hickey AJ, Cameron-Smith D, Abbiss CR.** Regular Post-Exercise Cooling Enhances Mitochondrial Biogenesis through AMPK and p38 MAPK in Human Skeletal Muscle. *Am J Physiol Regul Integr Comp Physiol* 309(3):R286-294, 2015.

**19. Ihsan M, Watson G, Abbiss C.** PGC-1alpha Mediated Muscle Aerobic Adaptations to Exercise, Heat and Cold Exposure. *Cellular and Molecular Exercise Physiology* 3: e7, 2014a.

**20. Ihsan M, Watson G, Choo HC, Lewandowski P, Papazzo A, Cameron-Smith D, Abbiss CR.** Postexercise Muscle Cooling Enhances Gene Expression of PGC-1α. *Med Sci Sports Exerc* 46(10): 1900-1907, 2014b.

**21. Impey SG, Hammond KM, Shepherd SO, Sharples AP, Stewart C, Limb M, Smith K, Philp A, Jeromson S, Hamilton DL, Close GL, Morton JP.** Fuel for the work required: a practical approach to amalgamating train-low paradigms for endurance athletes. *Physiol Rep* 4(10): e12803, 2016.

**22. Ishida K, Nakamura T, Kimura K, Kanno N, Takahashi N, Kamijo YI, Tajima F.** Suppression of activation of muscle sympathetic nerve during non-noxious local cooling after the end of local cooling in normal adults. *Eur J Appl Physiol* 116(4): 851-858, 2016.

**23. Jäger S, Handschin C, St-Pierre J, Spiegelman BM.** AMP-activated protein kinase (AMPK) action in skeletal muscle via direct phosphorylation of PGC-1alpha. *Proc Natl Acad Sci U.S.A.* 104: 12017–22, 2007.

**24. Joo CH, Allan R, Drust B, Close GL, Jeong TS, Bartlett JD, Mawhinney C, Louhelainen J, Morton JP, Gregson W.** Passive and post-exercise cold-water immersion augments PGC-1α and VEGF expression in human skeletal muscle. *Eur J Appl Physiol* 116: 2315-2326, 2016.

- 599 **25. Kim SH, Asaka M, Higashida K, Takahashi Y, Holloszy JO, Han D-H.** beta-  
600 Adrenergic stimulation does not activate p38 MAP kinase or induce PGC-1alpha in skeletal  
601 muscle. *Am J P Endocrinol Metab* 304: E844–E852, 2013.
- 602
- 603 **26. Koh HJ, Hirshman MF, He H, Manabe Y, Balschi JA, Goodyear LJ.** Adrenaline is a  
604 critical mediator of acute exercise-induced AMP-activated protein kinase activation in  
605 adipocytes. *Biochem J* 403(3): 473–481, 2007.
- 606
- 607 **27. Kristensen JM, Johnsen AB, Birk JB, Nielsen JN, Jensen BR, Hellsten Y, Richter**  
608 **EA, Wojtaszewski JF.** Absence of humoral mediated 5'AMP-activated protein kinase  
609 activation in human skeletal muscle and adipose tissue during exercise. *J Physiol* 585: 897-  
610 909, 2007.
- 611
- 612 **28. Little JP, Safdar A, Bishop D, Tarnopolsky MA, Gibala MJ.** An acute bout of high-  
613 intensity interval training increases the nuclear abundance of PGC-1alpha and activates  
614 mitochondrial biogenesis in human skeletal muscle. *Am J Physiol Regul Integr Comp Physiol*  
615 300: R1303–R1310, 2011.
- 616
- 617 **29. Manfredi LH, Zanon NM, Garófalo M, Navegantes LC, Kettelhut I.** Effect of short-  
618 term cold exposure on skeletal muscle protein breakdown in rats. *J Appl Physiol* 115: 1496–  
619 1505, 2013.
- 620
- 621 **30. Mawhinney C, Jones H, Joo CH, Low DA, Green DJ, Gregson W.** Influence of cold-  
622 water immersion on limb and cutaneous blood flow after exercise. *Med Sci Sports Exerc* 45:  
623 2277–85, 2014.
- 624
- 625 **31. Miura S, Kawanaka K, Kai Y, Tamura M, Goto M, Shiuchi T, Minokoshi Y, Ezaki**  
626 **O.** An increase in murine skeletal muscle peroxisome proliferator-activated receptor-gamma  
627 coactivator-1alpha (PGC-1alpha) mRNA in response to exercise is mediated by beta-  
628 adrenergic receptor activation. *Endocrinology* 148: 3441–8, 2007.
- 629
- 630 **32. Minokoshi Y, Kim YB, Peroni OD, Fryer LGD, Müller C, Carling D, Kahn B.** Leptin  
631 stimulates fatty-acid oxidation by activating AMP-activated protein kinase. *Nature* 415: 39-  
632 343, 2002.
- 633
- 634 **33. Norrbom J, Sundberg CJ, Ameln H, Kraus WE, Jansson E, Gustafsson T.** PGC-  
635 1alpha mRNA expression is influenced by metabolic perturbation in exercising human  
636 skeletal muscle. *J Appl Physiol* 96: 189–94, 2004.
- 637
- 638 **34. Oliveira RLGS, Ueno M, de Souza CT, Pereira-da-Silva M, Gasparetti AL, Bezzera**  
639 **RMN, Alberici LC, Vercesi AE, Saad MJA, Velloso LA.** Cold-induced PGC-1alpha  
640 expression modulates muscle glucose uptake through an insulin receptor/Akt-independent,  
641 AMPK-dependent pathway. *Am J Physiol Endocrinol Metab* 287: E686–95, 2004.
- 642
- 643 **35. Peaston RT, Graham KS, Chambers E, van der molen JC, Ball S.** Performance of  
644 plasma free metanephrines measured by liquid chromatography-tandem mass spectrometry in  
645 the diagnosis of pheochromocytoma. *Clin Chim Acta* 411:546–52, 2010.
- 646
- 647 **36. Perry CG, Lally J, Holloway GP, Heigenhauser GJ, Bonen A, Spriet LL.** Repeated  
648 transient mRNA bursts precede increases in transcriptional and mitochondrial proteins during

training in human skeletal muscle. *J Physiol* 588: 4795–4810, 2010.

**37. Puigserver P, Spiegelman BM.** Peroxisome proliferator-activated receptor- $\gamma$  coactivator 1 $\alpha$  (PGC-1 $\alpha$ ): transcriptional coactivator and metabolic regulator. *Endocr Rev* 24: 78–90, 2003.

**38. Puigserver P, Wu Z, Park CW, Graves R, Wright M, Spiegelman BM.** A cold-inducible coactivator of nuclear receptors linked to adaptive thermogenesis. *Cell* 92: 829–839, 1998.

**39. Robinson MM, Richards JC, Hickey MS, Moore DR, Phillips SM, Bell C, Miller BF.** Acute beta-adrenergic stimulation does not alter mitochondrial protein synthesis or markers of mitochondrial biogenesis in adult men. *Am J Physiol Regul Integr Comp Physiol* 298(1): R25–R33, 2010.

**40. Robinson MM, Bell C, Peelor FF 3<sup>rd</sup>, Miller BF.**  $\beta$ -Adrenergic receptor blockade blunts postexercise skeletal muscle mitochondrial protein synthesis rates in humans. *American J Physiol Regul Integr Comp Physiol* 301(2): R327–334, 2011.

**41. Schmittgen TD, Livak KJ.** Analyzing real-time PCR data by the comparative C(T) method. *Nat Protoc* 3(6): 1101–1108, 2008.

**42. Slivka D, Heesch M, Dumke C, Cuddy J, Hailes W, Ruby B.** Effects of post-exercise recovery in a cold environment on muscle glycogen, PGC-1 $\alpha$ , and downstream transcription factors. *Cryobiology* 66: 250–5, 2013.

**43. Slivka DR, Dumke CL, Tucker TJ, Cuddy JS, Ruby B.** Human mRNA response to exercise and temperature. *Int J Sports Med* 33: 94–100, 2012.

**44. Stepto NK, Martin DT, Fallon KE, Hawley JA.** Metabolic demands of intense aerobic interval training in competitive cyclists. *Med Sci Sports Exerc* 33: 303–10, 2001.

**45. Taylor CW, Ingham SA, Ferguson RA.** Acute and chronic effect of sprint interval training combined with postexercise blood-flow restriction in trained individuals. *Exp Physiol* 101: 143–154, 2016.

**46. Widegren U, Jiang XJ, Krook A, Chibalin AV, Bjornholm M, Tally M, Roth RA, Henriksson J, Wallberg-henriksson H, Zierath JR.** Divergent effects of exercise on metabolic and mitogenic signalling pathways in human skeletal muscle. *FASEB J* 12(13): 1379–1389.

**47. Wright DC, Geiger PC, Han D-H, Jones TE, Holloszy JO.** Calcium induces increases in peroxisome proliferator-activated receptor gamma coactivator-1 $\alpha$  and mitochondrial biogenesis by a pathway leading to p38 mitogen-activated protein kinase activation. *J Biol Chem* 282: 18793–18799, 2007.

**48. Yin W, Mu J, Birnbaum MJ.** Role of AMP-activated protein kinase in cyclic AMP-dependent lipolysis in 3T3-L1 adipocytes. *J Biol Chem* 278(44): 43074–43080, 2003.