



LJMU Research Online

Prieto, JL, Chen, P, Dong, S, Shappee, BJ, Seibert, M, Bersier, D, Holoien, TW-S, Kochanek, CS, Stanek, KZ and Thompson, TA

The Rise and Peak of the Luminous Type II In SN 2017hcc/ATLAS17Isn from ASAS-SN and Swift UVOT Data

<http://researchonline.ljmu.ac.uk/id/eprint/7678/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Prieto, JL, Chen, P, Dong, S, Shappee, BJ, Seibert, M, Bersier, D, Holoien, TW-S, Kochanek, CS, Stanek, KZ and Thompson, TA (2017) The Rise and Peak of the Luminous Type II In SN 2017hcc/ATLAS17Isn from ASAS-SN and Swift UVOT Data. Research Notes of the American Astronomical

LJMU has developed [LJMU Research Online](#) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

THE RISE AND PEAK OF THE LUMINOUS TYPE IIn SN 2017hcc/ATLAS17LSN FROM ASAS-SN AND SWIFT UVOT DATA

J. L. Prieto,¹ Ping Chen,² Subo Dong,² B. J. Shappee,³ M. Seibert,⁴ D. Bersier,⁵ T. W.-S. Holoién,⁴ C. S. Kochanek,⁶ K. Z. Stanek,⁶ and T. A. Thompson⁶

¹*Núcleo de Astronomía, Facultad de Ingeniería y Ciencias, Universidad Diego Portales, Ejército 441, Santiago, Chile*

²*Kavli Institute for Astronomy and Astrophysics, Peking University, Yi He Yuan Road 5, Hai Dian District, Beijing 100871, China*

³*Institute for Astronomy, University of Hawai'i, 2680 Woodlawn Drive, Honolulu, HI 96822, USA*

⁴*The Observatories of the Carnegie Institution for Science, 813 Santa Barbara St., Pasadena, CA 91101, USA*

⁵*Astrophysics Research Institute, Liverpool John Moores University, 146 Brownlow Hill, Liverpool L3 5RF, UK*

⁶*Department of Astronomy, The Ohio State University, 140 W. 18th Avenue, Columbus, OH 43210, USA.*

Keywords: stars: mass-loss, supernovae: general, supernovae: individual: SN 2017hcc

INTRODUCTION

Type IIn supernovae (SNe) are a heterogeneous class of stellar explosions associated with the deaths of massive stars surrounded by dense circumstellar environments created by strong mass-loss episodes (e.g., [Smith 2014](#)). In these objects, the interaction between the SN ejecta and a dense, H-rich circumstellar envelope dominates the evolution of their light curves and spectra.

SN 2017hcc/ATLAS17lsn was discovered as an optical transient on UT 2017 Oct. 2.38 by the Asteroid Terrestrial-impact Last Alert System (ATLAS; [Tonry 2011](#)) at an orange filter magnitude of $o = 17.44$ mag (TNS 13920¹). Its non-detection by ATLAS on 2017 Sep. 30.43 ($o \lesssim 19.04$ mag) showed that the transient was caught early. We obtained a low resolution optical spectrum on Oct. 7.4 with FLOYDS mounted on the FTS 2m telescope at the Siding Spring Observatory that showed the characteristics of a young Type IIn SN at $z = 0.0173$ (TNS 1284²).

¹ <https://wis-tns.weizmann.ac.il/object/2017hcc/discovery-cert>

² <https://wis-tns.weizmann.ac.il/object/2017hcc/classification-cert>

OBSERVATIONS AND ANALYSIS

We have been obtaining observations of SN 2017hcc as part of the All-Sky Automated Survey for Supernovae (ASAS-SN; Shappee et al. 2014) using the quadruple 14-cm “Brutus” telescopes in Haleakala, Hawaii, and the recently installed quadruple 14-cm “Bohdan Paczynski” telescopes in CTIO, Chile. The ASAS-SN images are reduced by an automated difference imaging pipeline and the magnitudes of sources are calibrated using standard V (“Brutus”) or SDSS g -band (“Paczynski”) photometry of local standard stars from the APASS DR9 (Henden et al. 2012). The SN was undetected by ASAS-SN down to $V/g \lesssim 17 - 18$ mag on 2017 Sep. 29-30 and first detected rising at $V = 15.98 \pm 0.15$ on Oct. 7.4. Now it has reached peak magnitude at $V \simeq 13.7$ mag. We show the current ASAS-SN light curve in Fig. 1.

We triggered *Swift* ToO observations that started on Oct. 28.4. The near-UV and optical colors of SN 2017hcc are consistent with a hot (but cooling) blackbody temperature with $W1 - U \simeq -0.4$ mag on Oct. 28.4 and $W1 - U \simeq 0.0$ mag on Nov. 19.6. The SN is undetected in X-rays with the *Swift* XRT (Chandra et al. 2017) and shows very strong continuum polarization at optical wavelengths (Mauerhan et al. 2017). We obtained new spectra on 2017 Oct. 20-21 with WFCCD on the du Pont 2.5m telescope at Las Campanas Observatory. The spectra are characterized by a strong blue continuum with Balmer and He I emission lines (see top right panel of Fig. 1), with line profiles characteristic of Type IIn. We measure a better constrained redshift of $z = 0.0168$ ($D \simeq 73$ Mpc).

We used a blackbody function to fit the spectral energy distribution (SED) of the SN using the *Swift* data and correcting for Galactic extinction (bottom left panel of Fig. 1). The fits show a hot, decreasing blackbody temperature and an increasing blackbody radius, consistent with early observations of SNe. Assuming a fixed temperature for the early rise of the light curve where there is only ASAS-SN data, we obtain the bolometric light curve of SN 2017hcc shown in the bottom right panel of Fig. 1. The bolometric luminosity at peak $L_{\text{bol,peak}} = (1.34 \pm 0.14) \times 10^{44}$ erg s $^{-1}$ ($M_{V,\text{peak}} \simeq -20.7$ mag), makes SN 2017hcc one of the most luminous Type IIn SNe known (e.g., Smith et al. 2007; Ofek et al. 2014; Jencson et al. 2016). From the bolometric light curve we constrain the risetime to be ~ 27 days and the total radiated energy of the event to date is $\sim 4 \times 10^{50}$ erg.

We thank Dr. Brad Cenko for approving the *Swift* ToO observations. We thank Las Cumbres Observatory and its staff for their continued support of ASAS-SN. ASAS-SN is funded in part by the Gordon and Betty Moore Foundation through grant GBMF5490 to the Ohio State University, NSF grant AST-1515927, the Mt. Cuba Astronomical Foundation, the Center for Cosmology and AstroParticle Physics (CCAPP) at OSU, and the Chinese Academy of Sciences South America Center for Astronomy (CASSACA).

REFERENCES

- | | |
|---|---|
| Chandra, P., et al. 2017, ATel, 10936 | Shappee, B. J., et al. 2014, ApJ, 788, 48 |
| Henden, A. A., et al. 2012, JAAVSO, 40, 430 | Smith, N., et al. 2007, ApJ, 666, 1116 |
| Jencson, J. E., et al. 2016, MNRAS, 456, 2622 | Smith, N. 2014, ARA&A, 52, 487 |
| Mauerhan, J. C., et al. 2017, ATel, 10911 | Tonry, J. L. 2011, PASP, 123, 58 |
| Ofek, E. O., et al. 2014, ApJ, 788, 154 | |




Figure 1. *Top Left:* ASAS-SN V (red) and *g*-band (blue) light curves of SN 2017hcc. *Top Right:* du Pont spectrum obtained on UT 2017 Oct. 20.08 showing a blue continuum and the characteristic features of a Type IIIn SN. *Bottom Left:* SED evolution obtained from observations at near-UV and optical wavelengths with the *Swift* UVOT. The lines show blackbody model fits with $T_{\text{bb}} = 16539 \pm 343 \text{ K}$ and $R_{\text{bb}} = (1.59 \pm 0.07) \times 10^{15} \text{ cm}$ on UT 2017 Oct. 28.4 (red) and $T_{\text{bb}} = 11657 \pm 238 \text{ K}$ and $R_{\text{bb}} = (2.83 \pm 0.08) \times 10^{15} \text{ cm}$ on UT 2017 Nov. 19.6 (blue). *Bottom Right:* Evolution of the bolometric luminosity as a function of time. For the early rise (blue and red points) of the light curve we assume a blackbody SED with the temperature fixed to $T_{\text{bb}} = 16539 \text{ K}$ obtained from the earliest *Swift* UVOT photometry to estimate a bolometric correction to the ASAS-SN V and *g*-band photometry. For the later part of the light curve (black points), we derive the bolometric luminosity directly from blackbody fits to the *Swift* UVOT photometry.