

Meloro, C, Hudson, A and Rook, L

Feeding habits of extant and fossil canids as determined by their skull geometry

<https://researchonline.ljmu.ac.uk/id/eprint/875/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Meloro, C, Hudson, A and Rook, L (2015) Feeding habits of extant and fossil canids as determined by their skull geometry. Journal of Zoology, 295 (3). pp. 178-188. ISSN 1469-7998

LJMU has developed [LJMU Research Online](https://researchonline.ljmu.ac.uk/) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

Feeding habits of extant and fossil canids as determined by their skull geometry

C. Meloro¹, A. Hudson¹ & L. Rook²

¹ Research Centre in Evolutionary Anthropology and Palaeoecology, School of Natural Sciences and Psychology, Liverpool John Moores University, Liverpool, UK

² Dipartimento di Scienze della Terra, Università di Firenze, Firenze, Italy

Keywords

Canidae; geometric morphometrics; skull shape; diet; hypercarnivore.

Correspondence

Carlo Meloro, Research Centre in Evolutionary Anthropology and Palaeoecology, School of Natural Sciences and Psychology, Liverpool John Moores University, James Parsons Building, Byrom Street, Liverpool L3 3AF, UK.
Email: C.Meloro@ljmu.ac.uk

Editor: Andrew Kitchener

Received 29 May 2014; revised 30 October 2014; accepted 12 November 2014

doi:10.1111/jzo.12196

Abstract

The canids belong to one of the most prominent families of mammalian carnivores. Feeding adaptations of extant species is well documented by field observations; however, we are still missing palaeoecological insights for many enigmatic fossil specimens. We employ geometric morphometrics to quantify skull size and shape in extant and fossil members of the Canini tribe, inclusive of jackals and wolf-like taxa. Skull data are tested to identify correlates of dietary adaptations in extant species for predicting adaptations in fossils. Main vectors of shape variation correlate with the relative skull-palatal length, the position of the upper carnassial tooth and the anterior tip of the secondary palate. Allometry occurs in the palatal shape but size explains only a small fraction (about 4%) of shape variance. Although we quantified only palatal and tooth shape for the inclusion of fragmentary fossils, discriminant function analysis successfully classify extant Canini in dietary groups (small, medium and large prey specialist) with 89% of accuracy. The discriminant functions provide insights into many enigmatic specimens such as *Eucyon adoxus* (=small prey), fossil jackal-like from Koobi Fora formation (=small prey) and the Plio-Pleistocene Old World canid guild (*Canis etruscus*, *C. arnensis* and *Lycaon falconeri*). Clearly, both skull size and shape are excellent predictors of feeding habits in Canini thus also provide information about fossil taxonomic affinities.

Introduction

Members of the family Canidae have successfully invaded every continent, except Antarctica, occupying a multitude of ecological niches, which is a testament to their adaptability in the present and in the past (Sillero-Zubiri, Hoffmann & Macdonald, 2004). The most updated molecular phylogeny (Lindblad-Toh *et al.*, 2005) identified distinct clades within the Canidae: the redfox-like clade, the South American clade, the wolf-like clade and the grey and island fox clade. This study will focus on the wolf-like clade (tribe Canini), which exhibit one of the most complete fossil record in the Old World (Tedford, Taylor & Wang, 1995; Tedford, Wang & Taylor, 2009). Tedford *et al.* (2009) recently provided a morphological phylogeny merging both extant and fossil species, although functional morphology of many enigmatic fossil specimens is still obscure and difficult to characterize (e.g. the genus *Eucyon*, or the wolf-like *Canis etruscus*; Cherin *et al.*, 2014).

The wolf-like clade had an explosion of forms during the Plio-Pleistocene so that biochronology considers such a

proliferation of species in the Old World into a separate faunal event (the wolf event, *c.* 2.0 Ma; Azzaroli, 1983; Azzaroli *et al.*, 1988; Torre *et al.*, 1992, 2001; Rook & Torre, 1996; Sardella & Palombo, 2007; Rook & Martínez-Navarro, 2010; Sotnikova & Rook, 2010). Palaeoecology of many of these canids represented by a coyote-like (*C. arnensis*), a small wolf-like (*C. etruscus*) and an African hunting dog-like (*Lycaon falconeri* group; Rook, 1994; Martínez-Navarro & Rook, 2003) was pioneered by Kurtén (1974) and Palmqvist, Arribas & Martínez-Navarro (1999) and later reconsidered by Meloro (2011) in a study on mandible shape. Here we aim to investigate skull shape that is expected to provide better insights into feeding ecology of extant, hence fossil Canini.

There have been numerous studies on the relationship between diet and craniodental form in Carnivora and canids in particular (Biknevicius & Ruff, 1992; Van Valkenburgh, Sacco & Wang, 2003; Sacco & Van Valkenburgh, 2004; Christiansen & Adolfssen, 2005; Christiansen & Wroe, 2007). Within canids, a shorter snout indicates larger moment arms for the temporalis and masseter muscles (Damasceno,

Hignst-Zaher & Astúa, 2013) and the canines are closer to the fulcrum, both creating a more powerful bite force (Christiansen & Adolfssen, 2005; Christiansen & Wroe, 2007). This is interpreted as an adaptation to kill large prey and can be detected in living and extinct canid tribes (Van Valkenburgh & Koepfli, 1993; Van Valkenburgh *et al.*, 2003; Andersson, 2005; Slater, Dumont & Van Valkenburgh, 2009).

Early morphometric attempts on Canidae general morphology already elucidated cophenetic similarities in relation to their taxonomy and ecology (Clutton-Brock, Corbet & Hills, 1976). By focusing on palatal and upper teeth morphology with geometric morphometric techniques, we intend to capture both size and shape aspects relevant to interpret fossil species. Geometric morphometrics has the advantage of allowing clear data visualization in multivariate shape space (Adams, Rohlf & Slice, 2004, 2013; Lawing & Polly, 2009). In addition, shape distances can be employed to infer morphological similarities: this is a straightforward way to compare data between living and fossil specimens (Caumul & Polly, 2005; Meloro *et al.*, 2008; Meloro, 2011). Due to the tendency in canids of increasing body mass towards their evolution in relation to ecological feeding specialization (Van Valkenburgh, Wang & Damuth, 2004), we will also explore skull size as possible proxy for predicting diet in extant and fossil species.

Materials and methods

Sample size

Skulls belonging to 102 specimens (85 extant and 17 fossils) were included in this study (Supporting Information Appendix S1). Our sample is representative of the broad diversity within the *Canis* clade including jackals and wolf-like ecomorphs (nine extant and 10 fossil species, Table 1). All extant specimens belong to wild captured individuals. Both male and female skulls were used indistinctively because sexual dimorphism is considered a negligible source of variance to infer dietary adaptations from the skulls. Indeed, sexual dimorphism within canids is generally small (Gittleman & Van Valkenburgh, 1997) and the gender is unknown for many fossil specimens.

For fossil species, we used the nomenclature finalized by Tedford *et al.* (2009). The small genera *Eucyon* and *Cynotherium* (with the species *Eucyon adoxus* and *Cynotherium sardous*) were also considered for their unequivocal affinities with extant *Canis*-like species (Lyras *et al.*, 2006; Rook, 1992, 2009).

Data capture

Digital photographs were collected on skulls positioned in ventral view by Meloro using a Nikon 995 at a 1-metre distance. A spirit level was positioned on the palate of the skull to ensure parallelism between camera optical plan and the flat-test region of the skull. On each skull, 15 landmarks were recorded by Hudson in the palate region to capture details of

Table 1 Skull sample sizes of extant and fossil canid species together with assigned dietary grouping

Species	Status	No. of specimens	Diet
<i>Canis lupus</i> ^a	Extant	14	Large
<i>Canis dingo</i>	Extant	3	Medium
<i>Canis latrans</i>	Extant	12	Medium
<i>Canis aureus</i>	Extant	10	Small
<i>Canis adustus</i>	Extant	10	Small
<i>Canis mesomelas</i>	Extant	9	Small
<i>Canis simensis</i>	Extant	8	Small
<i>Cuon alpinus</i> ^b	Extant	9	Large
<i>Lycan pictus</i>	Extant	10	Large
<i>Eucyon adoxus</i>	Fossil	1	
<i>Cynotherium sardous</i>	Fossil	1	
<i>Canis africanus</i>	Fossil	1	
<i>Canis antonii</i>	Fossil	1	
<i>Canis arnensis</i>	Fossil	2	
<i>Canis chiliensis</i>	Fossil	1	
<i>Canis dirus</i>	Fossil	2	
<i>Canis etruscus</i>	Fossil	3	
<i>Canis cf. mesomelas</i>	Fossil	1	
<i>Canis mosbachensis</i>	Fossil	1	
<i>Canis lupus</i> (Grotta Romanelli)	Fossil	1	
<i>Canis lupus</i> (Spain)	Fossil	1	
<i>Lycan falconeri</i>	Fossil	1	

^aIncludes subspecies (*Canis lupus gigas* and *Canis lupus pambasileus*).

^bIncludes subspecies (*Cuon alpinus dukhnensis* and *Cuon alpinus javanicus*).

Small, mesocarnivore feeding on small prey; Medium, mesocarnivore feeding on medium prey; Large, hypercarnivore feeding on large prey.

tooth and cusp positioning using the software tpsDig2 ver. 2.17 (Rohlf, 2013a; Fig. 1). Landmarks 1–2 record the width of the incisor arch, 3–4 the relative size of canine, landmark 5 is at the anterior tip of P3, 6 to 10 relative size of the upper carnassial (P4) together with the positioning of the main cusps, 10–14 covers the M1 morphology and landmark 15 is the most posterior point delimiting the end of the palate.

Cusp positions were recorded on P4 and M1 as good proxy for dietary adaptations but also to understand possible phylogenetic affinities between extant and fossil taxa (cf. Rook & Torre, 1996; Brugal & Boudadi-Maligne, 2010). The posterior part of the skull and the zygomatic arch were not covered by landmarks because they were not present in many of the analysed fossils.

Intra-individual error in landmarking was assessed using three landmarked replicas for three specimens. There were no differences in the variance of coordinates' values between replicas [analysis of variance (ANOVA) and multivariate analysis of variance (MANOVA) $P > 0.9$].

Geometric morphometrics

Landmark coordinates were aligned using generalized Procrustes superimposition (Rohlf & Slice, 1990) with the software tpsRelw ver. 1.53 (Rohlf, 2013b). The software




Figure 1 Skull of *Canis adustus* showing the landmark locations placed on each specimen. (1) tip of the snout defined by middle point between the first two frontal incisors, (2) posterior tip of the third incisor, (3) anterior tip of canine, (4) posterior tip of canine, (5) anterior tip of the third premolar, (6, 7, 8, 9, 10) outline of carnassial tooth, (11, 12) cusps of molar, (13) anterior tip of molar, (14) posterior tip of molar, (15) junction of the stiff and hard palate. The distance between 3 and 4 describe canine length. The distance between 6 and 10 describe carnassial tooth length. The distance between 1 and 15 describes snout length. Deviation of the specimens analysed from the consensus configuration of landmarks are shown below the skull. Scale bar equals 1 cm.

performed three operations: translation, rotation and scaling to transform the original 2D coordinates of landmarks into shape coordinates. A principal component analysis of the covariance matrix of the shape coordinates was then computed. Shape variation along each principal component axis was visualized using a thin-plate spline (Bookstein, 1991). Thin-plate splines visualize shape variation assuming that the average consensus configuration has no deformation and line on an infinite metal plane whose bending describe shape changes (Zelditch, Swiderski & Sheets, 2004).

The size of landmark configuration was extrapolated from the raw coordinates via centroid size (=the square root of the mean squared distance from each landmark to centroid of the landmark configuration Bookstein, 1989). In order to scale centroid size to the mean, natural log transformation was used (cf. Meloro *et al.*, 2008).

Feeding categories

For each extant species, a feeding category was assigned following multiple references. Van Valkenburgh (1989) grouped extant carnivores into three dietary categories: hypercarnivores, mesocarnivores and hypocarnivores. However, because there are no hypocarnivores in the sample for this study, Palmqvist *et al.*'s (1999) grouping of canids was also considered. Using both categorizations as a template, diet categories were assigned as small prey (mesocarnivore, mostly feeding on rodents and lagomorphs), medium prey (mesocarnivore that can include a wider range of prey sizes) and large prey (hypercarnivore, mostly preying on large ungulates). Extant jackals and the Ethiopian wolf belong to the category 'small prey', while the grey wolf, the African wild dog and the dhole are categorized as 'large prey' (cf. Slater *et al.*, 2009). The coyote and the dingo were categorized as 'medium prey' because of their broad adaptability in also hunting large prey

in group (Gese, Rongstad & Mytton, 1988; Lingle, 2002; Sillero-Zubiri *et al.*, 2004; Christiansen & Wroe, 2007; Letnic, Ritchie & Dickman, 2012).

Data analyses

Differences in skull size and shape due to diet were preliminary tested using ANOVA and parametric and non-parametric MANOVA. Due to the large number of independent shape variables, a selection of principal components (the one explaining at least 95% of variance) was employed to validate MANOVA models based on the full set of shapes (cf. Meloro & O'Higgins, 2011).

Additionally, allometry was tested in order to identify the possible influence of size on shape data (Mitteroecker *et al.*, 2013). A multivariate regression was employed to identify and visualize allometric signal in the whole sample of 102 skulls using thin-plate spline.

Discriminant function analysis was employed to provide prediction for fossil species using diet categories as factor and shape coordinates and natural log centroid size as independent variables. To considerably reduce the number of independent dietary predictors, a stepwise procedure was applied: a variable was entered into the model if the probability of its *F*-value was bigger than 0.05 and was removed if the probability was less than 0.10. Meloro (2011) consistently demonstrated the importance of including mandibular size as a predictor of feeding adaptation in Carnivora. We expect this to also hold for skull size in canids.

An UPGMA cluster analysis was employed to identify cophenetic similarities between fossil and extant specimens. Averaged shape coordinates were first computed for each extant and fossil species, then Procrustes distances calculated to construct the clustering UPGMA tree (cf. Meloro, 2011).

Results

Skull shape

Variability in skull shape is significantly reduced by using principal component analysis, with the first 12 PC axes explaining 95.26% of total shape variance. PC1 and PC2 explain 45.76% and 15.60% of total variance, respectively, and their combination show substantial differences between small jackal-like and large wolf-like species (Fig. 2). At the extreme negative of PC1, *C. simensis* is represented by a thin and slender palate with relatively short incisor row and canine but long snout, on the opposite of PC1 *L. pictus* together with *Cuon* share a much larger palate with relatively larger upper carnassial and M1. PC2 is highly influenced by position of landmark 15 and separates jackals and hypercarnivore *Lycaon-Cuon* from grey wolf and coyote. Fossil canids are evenly spaced in different areas of the morphospace and tend generally to occupy less extreme scores with the exception of *L. falconeri* (at the extreme positive PC1 and negative PC2).

MANOVA shows significant differences between diet in skull shape (represented by the first 12 PCs; Wilk's lambda = 0.164, $F = 8.677$, d.f. = 24, 142, $P < 0.0001$). Same applies when non-parametric MANOVA is computed after permuting Euclidean distances between dietary groups 9999 times ($F = 16.74$, $P < 0.0001$).

Skull shape differs significantly also between dietary categories (Wilk's lambda = 0.050, $F = 3.88$, d.f. = 52, 58, $P < 0.0001$).

Skull size and allometry

Skull size (here represented by ln centroid size of the landmark configuration) was normally distributed across dietary categories (P -values after Kolmogorov–Smirnov always > 0.06). This allowed us to perform an ANOVA test that shows significant differences between small, medium and large prey consumers ($F = 22.963$, d.f. = 2, 82, $P < 0.0001$; Fig. 3a). Due to significant differences in homogeneity of variance test (Levene statistic 5.702, d.f. = 2, 82, $P = 0.005$), Dunnett's T3




Figure 2 Plot of the first and second principal components. Thin-plate spline diagrams illustrate patterns of landmark displacements along each warp. Triangles indicate canids in the large dietary category, ellipsoids indicate canids in the medium dietary category and circles indicate canids in the small dietary category. Crosses and stars indicate fossil specimens with an unknown diet category. Below deformation grids from positive to negative RW scores.




Figure 3 (a) Box plot showing differences in natural log transformed centroid size (LnCS) between diet categories of extant specimens of canid skull (the outlier in the 'medium prey' category is a specimen of *C. latrans*); (b) skull shape deformation related to size from the smallest (*C. mesomelas*) to the largest (*C. dirus*) canid species. Values in parentheses are Ln centroid size.

was employed. This test shows significant differences in size between all the diet categories ($P < 0.025$ in all pairwise comparisons).

A significant allometric component was also detected even if Ln centroid size explains only a very small fraction of total shape variance (Wilks' lambda = 0.343, $F = 5.531$, d.f. = 26, 75, $P < 0.0001$; 4.11% of variance). Indeed, deformation grids depicted only a small deformation occurring mostly in the canine and upper carnassial (P4) areas (Fig. 3b). A closer inspection of allometry shows significant negative correlation only between Ln CS and PC3 (10.12% of variance, Spearman $r = -0.541$), PC8 (1.85% of variance, Spearman $r = -0.281$) and PC10 (1.20% of variance, Spearman $r = -0.119$).

Dietary discrimination

After stepwise, only five out of 30 shape coordinates and Ln centroid size were selected by the discriminant function analysis. Two significant discriminant functions (DFs) were extracted to differentiate dietary groups (DF1: 93.8% variance, Wilk's lambda = 0.113, $\chi^2 = 173.66$, d.f. = 12, $P < 0.0001$; DF2: 6.2% variance, Wilk's lambda = 0.733, $\chi^2 = 24.691$, d.f. = 5, $P < 0.0001$).

Percentage of correctly classified cases after cross-validation is high (small = 86.5%; medium = 86.7% and large = 93.9%).

DF1 was positively and significantly loaded on Ln CS ($r = 0.314$), Procrustes coordinate X of the landmark 6 (the anterior tip of P4, $r = 0.251$), and negatively on coordinate Y

for landmark 1 (tip of the snout, $r = -0.586$). DF2 correlated positively with coordinate Y of landmark 3 (anterior tip of the canine, $r = 0.841$) and negatively on coordinate X of landmark 11 (M1 paracone, $r = 0.478$), Y for landmark 13 (anterior tip of M1, $r = 0.398$).

The deformation grids were obtained after regressing DF scores versus shape coordinates. They show how species adapted to kill large prey at the positive DF1 are characterized by a shorter and thicker muzzle opposite to species adapted to kill small prey (Fig. 4). Medium prey specialists exhibit intermediate DF1 scores and negative DF2 scores. They are discriminated by 'small prey' due to a thin and long muzzle with relatively bigger carnassial (P4) and M1 (Fig. 4).

Fossil specimens are predicted to cover the whole range of dietary adaptations of extant Canini (Table 2). Species represented by multiple specimens are sometimes predicted into more than one category with the exception of the dire wolf for which both specimens are consistently categorized as predators of large prey. *Eucyon adoxus*, *Cynotherium sardous*, *C. cf. mesomelas* and one specimen of *C. arnensis* and *C. etruscus* follow within the 'small prey' category, while *C. lupus* from Romanelli, one specimen of *C. arnensis* and one of *C. chihliensis* follow within category 'medium prey'. All large fossil hypercarnivores are classified as 'large' (Table 2).

Clustering

The UPGMA based on Procrustes distances yields a cophenetic cluster with a high cophenetic correlation




Figure 4 Plot of the first two discriminant functions (DF) extracted from a combination of shape and size variables. Extant specimens are labelled according to their diet categorization. Fossil specimens are labelled individually. Below deformation grids from positive to negative DF scores.

($r = 0.882$). There is a mix of ecological and taxonomic signal with some fossil taxa clustering together due to their unique affinities (e.g. *E. adoxus* with *C. cf. mesomelas* from Olduvai Gorge). The fossil hunting dog *L. falconeri* is clearly an out-group that allows identifying three main groups: (1) a cluster showing the affinity of the extant Ethiopian wolf (*C. simensis*) with the prehistoric *C. arnensis*; (2) a cluster that separates extant jackal-like forms (inclusive of the fossil hypercarnivore *C. antonii* and wolf-like *C. etruscus* and *C. mosbachensis*) from grey wolf cluster inclusive of the dingo and the dire wolf; (3) hypercarnivore cluster inclusive of fossil *C. africanus*, extant *Lycaon* and *Cuon* and a fossil grey wolf from Spain.

Discussion

With no doubt, skull size and shape of extant Canini can strongly be linked to their feeding habits (Van Valkenburgh &

Koepfli, 1993; Van Valkenburgh *et al.*, 2003; Andersson, 2005; Slater *et al.*, 2009; Damasceno *et al.*, 2013). By investigating only the palate, we critically limited the amount of size and shape information, but demonstrate that this area is ecologically and taxonomically informative. Indeed, MANOVA and ANOVA show significant differences between feeding categories redefined to fit the broad dietary variation observed in the Canini tribe (Sillero-Zubiri *et al.*, 2004).

The palate of species adapted to hunt small prey is thin, longer and characterized by relatively shorter P4 and M1. All these adaptations can be observed in extant jackals and especially in the Ethiopian wolf (*C. simensis*) that occupy the extreme morphological variation on the first PC (Fig. 2). This confirms early morphometric observation by Rook & Azzaroli Puccetti (1996) and functional morphology by Slater *et al.* (2009). In contrast, the grey wolf, African hunting dog and the dhole cluster together in the morphospace (Fig. 2) for

Table 2 Dietary classification provided for fossil specimens using discriminant function analysis

	Most likely group	P(D G)	P (G D)	Second most likely group
<i>Eucyon adoxus</i>	Small	0.726	0.796	Medium
<i>Canis africanus</i>	Large	0.280	1.000	Medium
<i>Canis antonii</i>	Large	0.852	0.998	Medium
<i>Canis arnensis</i> IGF 601V	Small	0.015	0.991	Medium
<i>Canis arnensis</i> IGF 867	Medium	0.006	0.935	Small
<i>Canis chileensis</i>	Medium	0.192	0.503	Large
<i>Canis dirus</i> cast M11960	Large	0.003	1.000	Medium
<i>Canis dirus</i> cast unknown	Large	0.078	1.000	Medium
<i>Canis etruscus</i> cast MNCN an5006	Small	0.522	0.867	Medium
<i>Canis etruscus</i> SBAU337628	Large	0.839	0.995	Medium
<i>Canis etruscus</i> SBAU398989	Large	0.922	0.996	Medium
<i>Canis. cf. mesomelas</i>	Small	0.101	0.941	Medium
<i>Canis mosbachensis</i>	Large	0.126	0.677	Medium
<i>Canis lupus</i> (Romanelli)	Medium	0.208	0.599	Large
<i>Canis lupus</i> (Spain)	Large	0.000	0.975	Small
<i>Cynotherium sardous</i>	Small	0.073	0.932	Medium
<i>Lycaon falconeri</i>	Large	0.276	1.000	Medium

P (D | G) is the probability of membership in a group given the discriminant function score. P (G | D) is the posterior probability based on the sample employed to generate the discriminant functions.

their typical hypercarnivorous traits (Van Valkenburgh, 1991): a short and broad muzzle with larger incisors and canine (cf. Andersson, 2005) and relatively larger upper carnassial. All these features correlate with higher bite forces (Christiansen & Wroe, 2007; Damasceno *et al.*, 2013), hence the ability to kill prey much larger than themselves. Not surprisingly, these morphologies are well separated from the other feeding groups, supporting the highest classification rate in the discriminant function analysis.

In agreement with previous findings on the mandible, it is not only palatal shape that is a good discriminator of diet in extant Canini but also its size (cf. Meloro, 2011). The ecological continuum observed in Canini diet is reflected into skull morphology so that intermediate sized dogs (the coyote and the dingo) show intermediate skull shapes allowing them to expand feeding niches under different circumstances. Indeed, the medium size canid hunters possess relatively larger upper carnassial and M1 but retain a longer and thin snout (in the case of the coyote) or have a broad palate but not so extreme as in *Cuon* or *Lycaon* (the dingo in Fig. 2).

It is important to note that although an allometric component was detected in our data, it accounts only for a small percentage of shape variance. When size generally explains large portion of shape variance, it is common practice to use 'size-free' shape residuals, although this correction generally does not provide additional insights (cf. Meloro *et al.*, 2014). Mitteroecker *et al.* (2013) recently argued the necessity to take size into account by actually adding, and not removing this variable from subsequent analyses. Our results confirm such assertion, thus supporting the combined interpretation of palatal size and shape to infer palaeoecology of fossil species.

Fossil genera *Eucyon* and *Cynotherium* cluster well within the morphological variation of extant Canini confirming pre-

vious taxonomic observations on their affinities (Rook, 2009; Lyras, van Der Geer & Rook, 2010). The principal component plot shows similar scores between these taxa and the extant jackals, both clustering within the range of the side-striped jackal (Fig. 2). Consequently, the dietary reconstruction as specialist hunter of small prey fits well with previous attempts for the *Cynotherium* (cf. Abbazzi *et al.*, 2005; Lyras *et al.*, 2006) and underlines the strong affinity of *Eucyon* (at least for the species *E. adoxus*) with jackals.

Dietary reconstruction for Plio-Pleistocene dogs confirms the puzzling evolution of the Etruscan wolf (*C. etruscus*) and the coyote-like *C. arnensis* while supporting the hypercarnivorous traits of *L. falconeri*, *C. antonii* and *C. africanus* (cf. Rook, 1994; Tedford *et al.*, 2009). Both *C. etruscus* and *C. arnensis* specimens occupy more than one dietary classification in agreement with previous studies (Meloro, 2011; Cherin *et al.*, 2014; Flower & Shreve, 2014). However, there is a clear size partitioning with the Arno dog being classified as 'small-medium', while only one *C. etruscus* is predicted as 'small prey' with the others grouped into 'large prey' category. Due to ecological character displacement, it is possible that morphological variation in these taxa was broad and influenced by presence or absence of larger competitors (García & Virgós, 2007).

Diet of the large American dire wolf fits consistently with previous palaeoecological reconstructions (Anyonge & Baker, 2006; Meloro, 2011, 2012), while new insights emerge for *C. chihliensis* from the lower Pleistocene of China. Tong, Hu & Wang (2012) identified a mosaic of features combining hypercarnivorous dentition with a relatively small size compared with the grey wolf. Consequently, the size constraint on hunting behaviour supports our prediction of *C. chihliensis* as an adaptable hunter within the 'medium' category (cf. dingo,

see also Fig. 2). For the middle Pleistocene *C. mosbachensis*, a large-size categorization also seems likely based on its morphofunctional similarity to the grey wolf (cf. Flower & Shreve, 2014). Diet prediction for the wolf of Romanelli cave also fits within the category 'medium'. Although Sardella *et al.* (2014) confirmed its taxonomic affinity to the grey wolf, they also pointed out how its smaller size confounded previous taxonomic attempts of this species into golden jackal or *C. mosbachensis*. The grey wolf is highly flexible in size and ecology (Sillero-Zubiri *et al.*, 2004). Such flexibility has been observed in prehistoric specimens (Flower & Shreve, 2014) as well as ancestral forms supporting possible ecogeographical differentiation in the past. Comfortably the fossil grey wolf from Spain is predicted as large prey specialist.

The enigmatic *Canis* cf. *mesomelas* from Koobi Fora deserves a separate note. Werdelin & Lewis (2005) and Werdelin & Peigné (2010) reviewed the rich Plio-Pleistocene East African carnivore fauna. Taxonomy of jackals is not clear yet and there seems to be evidence for different ecomorphotypes in hominin fossil sites. Our analysis suggests the Koobi Fora specimen being adapted for hunting small-sized prey. Interestingly, the UPGMA analysis (Fig. 5) supports shape similarity not with extant jackals, but with the Mio-Pliocene genus *Eucyon* suggesting that it was a distinct (but ecologically equivalent to the extant jackal) morphotype.

For the other taxa, the UPGMA cluster analysis shows a mixed signal based on shape data. P4 and M1 morphology are phylogenetic characters in Canini (Tedford *et al.*, 2009) although the presented UPGMA (Fig. 5) cannot disentangle the ecological from the phylogenetic signal (cf. Meloro, 2011).




Figure 5 UPGMA Cluster analysis obtained on Procrustes distances of averaged sample for 23 canid species.

The clustering of *C. africanus* within *Lycaon-Cuon* confirms the grouping proposed by Rook (1994). However, the palate of *L. falconeri* from Valdarno and that of *C. antonii* are highly distinct from *C. africanus*. Ecogeographical and temporal variation could explain such a pattern even if larger and more complete sample is needed to prove this assertion. The grouping of *E. adoxus* with the jackal from Koobi Fora suggests how distinct the morphology is from these Plio-Pleistocene forms with no extant relatives, even if their smaller size supports ecological similarities with jackals and coyotes. *Cynotherium* is also enigmatically positioned (although outside of the wolf cluster) while the cluster of *C. etruscus* with *C. adustus* also does not support the wolf phylogenetic hypothesis (cf. Tedford *et al.*, 2009). Interestingly, recent research on African jackals supports the identification of a North African wolf subspecies (*C. lupus lupaster*) that was morphologically ascribed to the golden jackal (Gaubert *et al.*, 2012), suggesting how puzzling morphological characters can be, not only in fossil but also in extant species. The Romanelli grey wolf is an out-group within the wolf cluster while the dire wolf is grouped with the dingo and grey wolf. Extant *Lycaon* and *Cuon* cluster together consistently with their hyper-carnivorous feeding habits.

Members of Canini clearly occupied a broad range of ecological niches since the Pliocene then differentiating during Early Pleistocene with the evolution of modern taxa (Sotnikova & Rook, 2010). Such a rapid differentiation resulted in a high flexibility of ecomorphological skull traits whose combination provides robust palaeoecological insights.

Acknowledgements

We are grateful to the curators of different museum institutions for kindly providing access to osteological collections: P. Jenkins, L. Tomsett, R. Portela-Miguez, A. Salvador, D. Hills, J.J. Hooker, P. Brewer and A. Currant (British Museum of Natural History, London); E. Cioppi (Museo di Storia Naturale Università di Firenze, Florence); M. Reilly and J. Liston (Hunterian Museum and Art Gallery, University of Glasgow, Glasgow); B. Sanchez, J. Morales, J. Cabarga and J. B. Rodríguez (Museo Nacional de Ciencias Naturales, Madrid); A. Kitchener (Royal Museum of Scotland, Edinburgh); E. Gilissen and W. Wendelen (Royal Museum for Central Africa, Tervuren); E. Mbua, M. Mungu, F. Nderitu and O. Mwebi (Kenya National Museum, Nairobi). Visits to the Museo Nacional de Ciencias Naturales and Royal Museum of Central Africa were supported by the Synthesys grants 'Feeding habits in extinct European carnivores' (ES-TAF 858) and 'Ecomorphology of extant African carnivores' (BE-TAF 4901) to C. Meloro while part of the comparative analyses were supported by Synthesys grants 'Craniodental morphometric analysis of living and fossil jackals' FR-TAF 3311 and BE-TAF 3607 to L. Rook. Access to the collections of National Museum of Kenya was kindly granted by the Governments of Kenya and Tanzania and the Leverhulme Trust project 'Taxon-Free Palaeontological Methods for Reconstructing

Environmental Change' (F/00 754/C). Finally, we would like to thank two anonymous reviewers and the journal editor for their insights and care that considerably improved the quality of our manuscript.

References

- Abbazzi, L., Arca, M., Tiveri, C. & Rook, L. (2005). The endemic canid *Cynotherium* (Mammalia, Carnivora) from the Pleistocene deposits of Monte Tuttavista (Nuoro, Eastern Sardinia). *Riv. Ital. Paleontol. S.* **111**, 493–507.
- Adams, D.C., Rohlf, F.J. & Slice, D.E. (2004). Geometric morphometrics: ten years of progress following the 'revolution'. *Ital. J. Zool.* **71**, 5–16.
- Adams, D.C., Rohlf, F.J. & Slice, D.E. (2013). A field comes of age: geometric morphometrics in the 21st century. *Hystrix* **24**, 7–14.
- Andersson, K. (2005). Were there pack-hunting canids in the tertiary, and how can we know? *Paleobiology* **31**, 56–72.
- Anyonge, W. & Baker, A. (2006). Craniofacial morphology and feeding behavior in *Canis dirus*, the extinct Pleistocene dire wolf. *J. Zool. (Lond.)* **269**, 309–316.
- Azzaroli, A. (1983). Quaternary mammals and the end-Villafranchian dispersal event – a turning point in the history of Eurasia. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **44**, 117–139.
- Azzaroli, A., De Giuli, C., Ficarelli, G. & Torre, D. (1988). Late Pliocene to early mid-Pleistocene mammals in Eurasia: faunal succession and dispersal events. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **66**, 77–100.
- Biknevicius, A.R. & Ruff, C.B. (1992). The structure of the mandibular corpus and its relationship to feeding behaviors in extant carnivores. *J. Zool. (Lond.)* **228**, 479–507.
- Bookstein, F.L. (1989). 'Size and Shape': a comment on semantics. *Syst. Zool.* **38**, 173–180.
- Bookstein, F.L. (1991). *Morphometric tools for landmark data. Geometry and Biology*. New York: Cambridge University Press.
- Brugal, J.P. & Boudadi-Maligne, M. (2010). Quaternary small to large canids in Europe: taxonomic status and biochronological contribution. *Quat. Int.* **243**, 171–182.
- Caumul, R. & Polly, P.D. (2005). Phylogenetic and environmental components of morphological variation: skull, mandible and molar shape in marmots (*Marmota*, Rodentia). *Evolution* **59**, 2460–2472.
- Cherin, M., Berté, D.F., Rook, L. & Sardella, R. (2014). Re-defining *Canis etruscus* (Canidae, Mammalia): a new look into the evolutionary history of Early Pleistocene dogs supported by the outstanding fossil record from Pantalla (Perugia, central Italy). *J. Mamm. Evol.* **21**, 95–110.
- Christiansen, P. & Adolfsson, J.S. (2005). Bite forces, canine strength and skull allometry in carnivores (Mammalia, Carnivora). *J. Zool. (Lond.)* **266**, 133–151.
- Christiansen, P. & Wroe, S. (2007). Bite forces and evolutionary adaptations to feeding ecology in carnivores. *Ecology* **88**, 347–358.
- Clutton-Brock, J., Corbet, G.B. & Hills, M. (1976). A review of the family Canidae, with a classification by numerical methods. *Br. Mus. (Nat. Hist.) J.* **29**, 119–199.
- Damasceno, E.M., Hignst-Zaher, E. & Astúa, D. (2013). Bite force and encephalization in the Canidae (Mammalia: Carnivora). *J. Zool. (Lond.)* **290**, 246–254.
- Flower, L.O.H. & Shreve, D.C. (2014). An investigation of palaeodietary variability in European Pleistocene canids. *Quat. Sci. Rev.* **96**, 188–203.
- García, N. & Virgós, E. (2007). Evolution of community composition in several carnivore paleoguilds from the European Pleistocene: the role of interspecific competition. *Lethaia* **40**, 33–44.
- Gaubert, P., Bloch, C., Benyacoub, S., Abdelhamid, A., Pagani, P., Djagoun, C.A.M.S., Coloux, A. & Dufour, S. (2012). Reviving the African wolf *Canis lupus lupaster* in North and West Africa: a mitochondrial lineage ranging more than 6,000 km wide. *PLoS ONE* **7**, e42740.
- Gese, E.M., Rongstad, O.J. & Mytton, W.R. (1988). Relationship between coyote group size and diet in South-eastern Colorado. *J. Wildl. Manage.* **52**, 647–653.
- Gittleman, J.L. & Van Valkenburgh, B. (1997). Sexual size dimorphism in the canines and skulls of carnivores: effects of size, phylogeny, and behavioural ecology. *J. Zool. (Lond.)* **242**, 97–117.
- Kurtén, B. (1974). A history of coyote-like dogs (Canidae, Mammalia). *Acta Zool. Fenn.* **140**, 1–38.
- Lawing, A.M. & Polly, P.D. (2009). Geometric morphometrics: recent applications to the study of evolution and development. *J. Zool. (Lond.)* **280**, 1–7.
- Letnic, M., Ritchie, E.C. & Dickman, C.R. (2012). Top predators as biodiversity regulators: the dingo *Canis lupus dingo* as a case study. *Biol. Rev.* **87**, 390–413.
- Lindblad-Toh, K. *et al.* (2005). Genome sequence, comparative analysis and haplotype structure of the domestic dog. *Nature* **438**, 803–819.
- Lingle, S. (2002). Coyote predation and habitat segregation of white-tailed deer and mule deer. *Ecology* **83**, 2037–2048.
- Lyras, G., van Der Geer, A.A.E. & Rook, L. (2010). Body size of insular carnivores: evidence from the fossil record. *J. Biogeogr.* **37**, 1007–1021.
- Lyras, G.A., van der Geer, A.A.E., Dermitzakis, M. & De Vos, J. (2006). *Cynotherium sardous*, an insular canid (Mammalia: Carnivora) from the Pleistocene of Sardinia (Italy), and its origin. *J. Vert. Paleontol.* **26**, 735–745.
- Martínez-Navarro, B. & Rook, L. (2003). Gradual evolution in the African hunting dog lineage systematic implications. *C. R. Palevol* **2**, 695–702.
- Meloro, C. (2011). Feeding habits of Plio-Pleistocene large carnivores as revealed by the mandibular geometry. *J. Vert. Paleontol.* **31**, 428–446.

- Meloro, C. (2012). Mandibular shape correlates of tooth fracture in extant Carnivora: implications to inferring feeding behaviour of Pleistocene predators. *Biol. J. Linn. Soc. Lond.* **106**, 70–80.
- Meloro, C. & O'Higgins, P. (2011). Ecological Adaptations of mandibular form in fissiped carnivore. *J. Mammal. Evol.* **18**, 185–200.
- Meloro, C., Raia, P., Piras, P., Barbera, C. & O'Higgins, P. (2008). The shape of the mandibular corpus in large fissiped carnivores: allometry, function and phylogeny. *Zool. J. Linn. Soc.* **154**, 832–845.
- Meloro, C., Cáceres, N., Carotenuto, F., Sponchiado, J., Melo, G.L., Passaro, F. & Raia, P. (2014). In and out the Amazonia: evolutionary ecomorphology in howler and capuchin monkeys. *Evol. Biol.* **41**, 38–51.
- Mitteroecker, P., Gunz, P., Windhager, S. & Schaefer, K. (2013). A brief review of shape, form, and allometry in geometric morphometrics, with applications to human facial morphology. *Hystrix* **24**, 59–66.
- Palmqvist, P., Arribas, A. & Martínez-Navarro, B. (1999). Ecomorphological study of large canids from the lower Pleistocene of southeastern Spain. *Lethaia* **32**, 75–88.
- Rohlf, F.J. (2013a). tpsDig2 ver.2.16. Ecology & Evolution, SUNY at Stony Brook.
- Rohlf, F.J. (2013b). tpsRelw ver.1.53. Ecology & Evolution, SUNY at Stony Brook.
- Rohlf, F.J. & Slice, D.E. (1990). Extensions of the Procrustes method for the optimal superimposition of landmarks. *Syst. Zool.* **39**, 40–59.
- Rook, L. (1992). '*Canis*' *monticinensis* sp. nov., a new Canidae (Carnivora, Mammalia) from the late Messinian of Italy. *Boll. Soc. Paleontol. I.* **31**, 151–156.
- Rook, L. (1994). The Plio-Pleistocene Old World *Canis* (*Xenocyon*) ex. gr. *falconeri*. *Boll. Soc. Paleontol. I.* **33**, 71–82.
- Rook, L. (2009). The wide ranging genus *Eucyon* Tedford & Qiu, 1996 (Mammalia, Carnivora, Canidae) in Mio-Pliocene of the Old World. *Geodiversitas* **31**, 723–743.
- Rook, L. & Azzaroli Puccetti, M.L. (1996). Remarks on the skull morphology of the endangered Ethiopian jackal, *Canis simensis* Rüppel, 1838. *Memorie Fisiche della Accademia Nazionale dei Lincei, ser.* **9**, 277–302.
- Rook, L. & Martínez-Navarro, B. (2010). Villafranchian: The long story of a Plio-Pleistocene European large mammal biochronologic unit. *Quat. Int.* **219**, 134–144.
- Rook, L. & Torre, D. (1996). The wolf event in western Europe and the beginning of the Late Villafranchian. *Neues Jahrb. Geol. P-A.* **8**, 495–501.
- Sacco, T. & Van Valkenburgh, B. (2004). Ecomorphological indicators of feeding behaviour in the bears (Carnivora: Ursidae). *J. Zool. (Lond.)* **263**, 41–54.
- Sardella, R. & Palombo, M.R. (2007). The Plio-Pleistocene boundary: which significant for the so-called 'wolf-event'? Evidence from Western Europe. *Quaternaire* **18**, 65–71.
- Sardella, R., Bertè, D., Lurino, D.A., Cherin, M. & Tagliacozzo, M. (2014). The wolf from Grotta Romanelli (Apulia, Italy) and its implications in the evolutionary history of *Canis lupus* in the Late Pleistocene of Southern Italy. *Quat. Int.* **328–329**, 179–195.
- Sillero-Zubiri, C., Hoffmann, M. & Macdonald, D.W. (2004). *Canids: foxes, wolves, jackals and dogs. Status survey and conservation action plan.* IUCN/SSC Canid Specialist Group.
- Slater, G., Dumont, E.R. & Van Valkenburgh, B. (2009). Implications of predatory specialization for cranial form and function in canids. *J. Zool. (Lond.)* **278**, 181–188.
- Sotnikova, M.V. & Rook, L. (2010). Dispersal of the Canini (Mammalia, Canidae: Caninae) across Eurasia during the Late Miocene to Early Pleistocene. *Quat. Int.* **212**, 86–97.
- Tedford, R.H., Taylor, B.E. & Wang, X. (1995). Phylogeny of the Canidae (Carnivora: Canidae): the living taxa. *Am. Mus. Novit.* **3146**, 1–37.
- Tedford, R.H., Wang, X. & Taylor, B.E. (2009). Phylogenetic systematics of the North American fossil Caninae (Carnivora: Canidae). *Bull. Am. Mus. Nat. Hist.* **325**, 1–218.
- Tong, H., Hu, N. & Wang, X. (2012). New remains of *Canis chihliensis* from Shianshenmiazou, a lower Pleistocene site in Yanguai, Hebei. *Vertebrat. Palasiatic.* **50**, 335–360.
- Torre, D., Ficarelli, G., Masini, F., Rook, L. & Sala, B. (1992). Mammal dispersal events in the Early Pleistocene of Western Europe. *Cour. Forsch.-Inst. Senckenberg* **153**, 51–58.
- Torre, D., Abbazzi, L., Bertini, A., Fanfani, F., Ficarelli, G., Masini, F., Mazza, P. & Rook, L. (2001). Structural changes in Italian Late Pliocene – Pleistocene large mammal assemblages. *Boll. Soc. Paleontol. I.* **40**, 303–306.
- Van Valkenburgh, B. (1989). Carnivore dental adaptations and diet: a study of trophic diversity within guilds. In *Carnivore behavior, ecology, and evolution*, Vol. 1: 410–436. Gittleman, J.L. (Ed.). Ithaca: Cornell University Press.
- Van Valkenburgh, B. (1991). Iterative evolution of hypercarnivory in canids (Mammalia: Carnivora): evolutionary interactions among sympatric predators. *Paleobiology* **17**, 340–362.
- Van Valkenburgh, B. & Koepfli, K. (1993). Cranial and dental adaptations for predation in canids. *Symp. Zool. Soc. Lond.* **65**, 15–37.
- Van Valkenburgh, B., Wang, X. & Damuth, J. (2004). Cope's rule, hypercarnivory, and extinction in North American canids. *Science* **306**, 101–104.
- Van Valkenburgh, B.V., Sacco, T. & Wang, X. (2003). Pack hunting in Miocene borophagine dogs; evidence from craniodental morphology and body size. *Bull. Am. Mus. Nat. Hist.* **279**, 147–162.
- Werdelin, L. & Lewis, M.E. (2005). Plio-pleistocene carnivora of eastern Africa: species richness and turnover patterns. *Zool. J. Linn. Soc.* **144**, 121–144.

- Werdelin, L. & Peigné, S. (2010). Carnivora, chapter: 32. In *Cenozoic mammals of Africa*: 603–657. Werdelin, L. & Sanders, W.J. (Eds). Berkeley and London: University of California Press.
- Zelditch, M.L., Swiderski, D.L. & Sheets, H.D. (2004). *Geometric morphometrics for biologists: a primer*. 2nd edn. Amsterdam: Elsevier.

Supporting Information

Additional Supporting Information may be found in the online version of this article at the publisher's web-site:

Appendix S1. List of extant and fossil skull specimens of Canidae.