



LJMU Research Online

Hartman, YAW, Karssemeijer, EGA, van Diepen, LAM, Olde Rikkert, MGM and Thijssen, DHJ

Dementia Patients Are More Sedentary and Less Physically Active than Age- and Sex-Matched Cognitively Healthy Older Adults.

<http://researchonline.ljmu.ac.uk/9165/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Hartman, YAW, Karssemeijer, EGA, van Diepen, LAM, Olde Rikkert, MGM and Thijssen, DHJ (2018) Dementia Patients Are More Sedentary and Less Physically Active than Age- and Sex-Matched Cognitively Healthy Older Adults. *Dementia and Geriatric Cognitive Disorders*. 46 (1-2). pp. 81-89.

LJMU has developed [LJMU Research Online](#) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

Dementia patients are more sedentary and less physically active than age- and sex-matched cognitively healthy older adults

Y.A.W. (Yvonne) Hartman*, MSc ^a

E.G.A. (Esther) Karssemeijer*, MD ^{b,c}

E.A.M. (Lisanne) van Diepen ^a

M.G.M. (Marcel) Olde Rikkert, MD PhD ^{b,c}

D.H.J. (Dick) Thijssen, PhD ^{a,d}

* Shared first authorship: these authors contributed equally to this manuscript

^a Radboud university medical center, Radboud Institute for Health Sciences, Department of Physiology, Nijmegen, The Netherlands.

^b Radboud university medical center, Donders Institute for Brain Cognitive function and Behavior, Department of Geriatric Medicine, Behavior, Nijmegen, The Netherlands.

^c Radboudumc Alzheimer Centre, Radboud university medical center, Nijmegen, The Netherlands

^d Research institute for Sport and Exercise Sciences, Liverpool John Moores University, Liverpool, UK

Corresponding author

Prof. Dr. Dick H.J. Thijssen

Department of Physiology

Radboud University Medical Center

Philips van Leijdenlaan 15

dick.thijssen@radboudumc.nl

Tel: (+31)24 36 14 209, Fax: (+31)24 366 8340

Key words: sedentary behavior, dementia, physical activity, actigraphy

Declarations

Conflict-of-interest statement: None declared.

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

2 ABSTRACT

3 **Aims:** To examine physical activity and sedentary behavior characteristics of ambulatory and
4 community dwelling patients with dementia compared to cognitively healthy age-, sex- and weight-
5 matched controls.

6 **Methods:** In this cross-sectional study we included community dwelling dementia patients (N=45,
7 age=79.6±5.9 years, MMSE=22.8±3.2) and matched controls (N=49, age=80.0±7.7 years,
8 MMSE=29.0±1.2). Participants wore a wrist accelerometer for seven days to assess sedentary time,
9 sedentary bout duration and time spent in very light, light-to-moderate and moderate-to-vigorous
10 physical activities.

11 **Results:** Relative sedentary time and sedentary bout duration was significantly higher in dementia
12 patients compared to controls (median and interquartile range: 57% (49 - 68) vs 55% (47 - 59) and
13 18.3 (16.4–21.1) minutes vs 16.6 (15.3–18.4) minutes, P=0.042 and P=0.008 respectively). In addition,
14 dementia patients spent a lower percentage of their waking time in light-to-moderate and moderate-to-
15 vigorous intensity physical activities (20% (15–23) vs 22% (18–25) and 5% (2–10) vs 10% (5–13),
16 P=0.017 and P=0.001 respectively).

17 **Conclusion:** We revealed that dementia patients are more sedentary and perform less physical activity
18 compared to cognitively healthy controls. This may have clinically important consequences, given the
19 observation that sedentary behavior and little physical activity independently predicts all-cause
20 mortality and morbidity.

21

22

23

24 INTRODUCTION

25 The incidence and prevalence of dementia is rising [1]. Pharmacological treatment to slow disease
26 progression shows limited benefits on cognitive functioning [2]. Therefore, non-pharmaceutical
27 therapies are needed to attenuate or slow cognitive decline. Engagement in moderate to vigorous
28 physical activity (i.e. exercise) is one of the most important modifiable risk factors for dementia [3].
29 Moreover, exercise interventions have beneficial effects on cognitive function in older adults with
30 dementia [4]. Interestingly, recent research showed that sedentary behavior (activities requiring low
31 levels of energy expenditure, e.g. sitting and lying), independent of performance of physical activity,
32 is strongly related to negative health outcomes and mortality [5,6]. Moreover, sedentary behavior is
33 associated with lower cognitive performance [7], which stresses the relevance to understand the
34 prevalence and characteristics of sedentary behavior in the context of dementia. Therefore, we aim to
35 objectively determine physical activity and sedentary behavior characteristics of community-dwelling
36 dementia patients compared to cognitive healthy age-, sex- and weight-matched controls. Secondary,
37 we will explore whether increasing age attenuates physical activity and sedentary behavior in
38 dementia.

39

40 MATERIALS AND METHODS

41 Participants and design. In this cross-sectional study persons with a dementia diagnosis, aged >60
42 years, that were ambulatory and community dwelling were included. Dementia diagnosis was based
43 on comprehensive clinical assessment by a physician, typically including neuropsychological
44 assessment and imaging. We used the mini-mental state examination (MMSE) to indicate severity of
45 cognitive impairment [8]. Baseline measurements of a longitudinal trial examining the effects of
46 exercise on cognitive functioning in dementia were used for the current study [9]. Cognitively healthy
47 controls were age-, sex- and weight-matched to dementia patients and had no history of cognitive
48 impairment (MMSE >24 [8]). All participants were ambulatory and community-dwelling. The study
49 protocol was approved by the local Medical Ethics Committee in accordance with the latest revision of
50 the declaration of Helsinki. Written informed consent was obtained from all participants.

51 Physical activity monitoring. Directly after screening, physical activity and sedentary behavior were
52 assessed by the Philips Actiwatch 2, a wrist-worn accelerometer validated in middle-aged females
53 [10]. The Philips Actiwatch 2 contains an acceleration-responsive piezoelectric sensor which measures
54 wrist accelerations in three directions every 30 seconds. These wrist accelerations were translated into
55 a number of counts that were used to estimate physical activity and sedentary behavior. The
56 accelerometer was worn for seven days on the non-dominant wrist, to provide a reliable estimate of
57 physical activity and sedentary behavior. the [11,12]. All participants wore the actiwatch 24 hours per
58 day. The accelerometer was waterproof and participants did not take it off during swimming or taking
59 a shower. Therefore, non-wear time was not expected. Sleep intervals, including daytime naps, were
60 filled in by the participants or their caregivers in a sleep diary.

61
62 Data analysis. Data was uploaded using the Philips Actiware 6 software. Data from the first day of
63 testing were excluded from analysis to give participants the opportunity to familiarize with the device.
64 Sleep intervals were manually set by the researcher using the Philips Actiware 6 software [13] and
65 excluded by custom software written in MATLAB R2014b (MathWorks, USA). Hereby only the data
66 of the waking hours remained. Participants were included if they provided at least six valid days (>10
67 h of waking data). Data was converted from counts per epoch to counts per minute (CPM). Cut-off
68 points of 145 counts per minute (CPM), 145-274 CPM, 274-597 CPM and >597 CPM were used for
69 sedentary behavior, very light, light-to-moderate and moderate-to-vigorous physical activity
70 respectively [10]. To account for individual differences in waking time, our primary analysis
71 expressed activity levels as a percentage of total (awake) measuring time. Interruptions in sedentary
72 behavior were defined as spending one minute ≥ 145 CPM after 5 minutes < 145 CPM. Prolonged
73 sedentary behavior was defined as spending 30 minutes < 145 CPM without one minute above 145
74 CPM. Duration of average sedentary bout was defined by total time spent sedentary divided by
75 number of interruptions in sedentary behavior.

76
77 Statistical analysis. Statistical analyses were performed in IBM SPSS Statistics 20.0 (IBM SPSS; IBM
78 Corp., Armonk, New York, USA). We performed a complete case analysis, including only those

79 participants that wore the actiwatch for a minimum of 7 days. Chi²-tests for categorical variables and
80 independent samples t-tests for continuous variables were used to compare dementia patients and
81 controls. Non-parametric tests were used for not normally distributed data (including physical activity
82 and sedentary behavior). To evaluate the impact of age, the same analyses were performed between
83 participants aged <80 years and ≥80 years. All data are presented as median (interquartile range)
84 unless stated otherwise. Level of significance was set at P<0.05.

85

86 RESULTS

87 45 dementia patients (mean±SD age=79.6±5.9 years, MMSE=22.8±3.2) and 49 controls (mean±SD
88 age=80.0±7.7 years, MMSE=29.0±1.2) were included (Figure 1). Sex, BMI, walking aid use and
89 number of comorbidities did not differ between groups (Table 1). MMSE was significantly lower in
90 dementia patients (P<0.001), and dementia patients received significantly more home care (P<0.001).
91 The majority of dementia patients and controls lived independently (93% and 98% respectively).

92

93 Total waking time tended to be lower in dementia patients compared to controls (mean±SD 14.9±1.3
94 h/day vs 15.4±1.0 h/day, P=0.053, Table 2). Dementia patients had significantly lower activity counts
95 and spent more hours in categories reflecting lower-intensity activity (Table 2). Relative sedentary
96 time was significantly higher in dementia patients compared to controls (57% (49 - 68) vs 55% (47 -
97 59), P=0.042, Table 2). In addition, dementia patients spent a lower percentage of their waking time in
98 light-to-moderate and moderate-to-vigorous intensity physical activity (20% (15-23) vs 22% (18-25)
99 and 5% (2-10) vs 10% (5-13), P=0.017 and P=0.001, respectively). Number of interruptions in
100 sedentary behavior and prolonged sedentary bouts did not differ. Duration of sedentary bouts was
101 significantly longer in dementia patients compared to controls (18.3 (16.4-21.1) minutes vs 16.6 (15.3-
102 18.4) minutes, P=0.008, Table 2).

103

104 When comparing younger versus older subgroups, older participants showed more walking aid users.
105 Whilst older controls showed significantly less physical activity and more sedentary behavior
106 compared to their younger peers, no such changes were present between dementia patients aged ≥80

107 years and aged <80 years (Figure 2). More specifically, younger controls had a significantly higher
108 activity counts, lower relative sedentary time, and spent more hours in categories reflecting light-to-
109 moderate and moderate-to-vigorous intensity activity compared to older controls. Since no such
110 differences were present between young and older dementia patients, differences in physical activity
111 and sedentary behavior characteristics between populations was most prominent in the younger groups
112 (see Figure 2).

113

114 **DISCUSSION**

115 The aim of our study was to objectively investigate physical activity and sedentary behavior
116 characteristics of dementia patients compared to controls, and assess whether age affects this
117 comparison. First, in our relatively large sample we found that dementia patients spent significantly
118 more of their waking hours in a sedentary state and significantly less time in light-to-moderate and
119 moderate-to-vigorous intensity activities. This may have clinically important consequences, given the
120 observation of previous prospective studies that sedentary behavior independently predicts all-cause
121 mortality and morbidity [5,14]. Secondly, we found that older age was associated with a decline in
122 physical activity and increase in sedentary behavior in controls, whilst no such age-related changes
123 were found in dementia patients. Consequently, negligible differences in physical activity and
124 sedentary behavior characteristics were present when comparing older dementia patients and controls.

125

126 In line with observations from previous work [15,16], our data confirm that community-dwelling
127 dementia patients spend a large amount of time in sedentary behavior and have low levels of physical
128 activity. We add the novel finding that differences between dementia patients and controls remain
129 when corrected for sleep time. Even though we used an accelerometer validated to measure physical
130 activity and sedentary behavior [10], time spent in moderate-to-vigorous intensity activity seems
131 unusually high in both groups [17]. We have compared our findings to other studies that assessed
132 physical activity and sedentary behavior in older adults with and without dementia, and we note that
133 sedentary behavior ranged from 6.7 to 10.7 hours between studies [15,18,19]. Moreover, percentage of
134 elderly meeting physical activity guidelines (150min/week of moderate-to-vigorous physical activity)

135 ranged from 27-69% [17,20]. Differences in sedentary behavior and physical activity duration between
136 studies might relate to the use of different types of accelerometers, given that the reported studies
137 used hip-worn accelerometers, wrist-worn accelerometers or questionnaires to estimate physical
138 activity and sedentary behavior [21]. However, this does not invalidate our primary comparison
139 between subjects with dementia and healthy peers. Moreover, this highlights the importance of
140 including a control group to provide valid interpretation of the results.

141
142 Our observation raises the question if differences in physical activity and sedentary behavior are
143 simply a consequence of dementia. A decline in executive functioning (i.e. necessary for goal directed
144 behavior such as physical activity) could lead to apathy [22], which is known to lower activity levels
145 in Alzheimer's' patients [23]. However, it is important to realize that lower physical activity and
146 higher sedentary behavior have already been reported in the preclinical stages of dementia [24] and in
147 subjects with mild cognitive impairment) [25]. This might suggest that differences in physical activity
148 and sedentary behavior are causally linked to progression from mild cognitive impairment to later
149 stages of dementia. Future research is necessary to answer this question on cause of effect.
150 Nevertheless, since higher physical activity and lower sedentary behavior are associated with better
151 cognitive performance [7,25], benefits of interventions promoting physical activity and reducing
152 sedentary behavior should be investigated.

153
154 In response to our second research question, we found that older age was associated with a decline in
155 physical activity and increase in sedentary behavior in controls. This observation can partly be
156 explained by deterioration of walking and mobility and increased disability with older age [26].
157 Interestingly, no further decline in physical activity and increase in sedentary behavior was found in
158 dementia patients, despite the age-related increased number of walking aid users. This striking result
159 suggests that cognitive impairment in dementia has a great impact on physical activity and sedentary
160 behavior, and may be more important than the impact of other factors such as deterioration of walking
161 and mobility. An alternative explanation is that a (near) minimum level is achieved in the decline in
162 levels of physical activity and sedentary behavior in community-dwelling subjects [17].

163

164 In addition to the duration of sedentary behavior, previous work revealed that the frequency of
165 breaking up sitting (and therefore duration of each sedentary bout) may have clinical relevance.
166 Breaks in sedentary behavior can prevent cardiovascular impairments [5] and plays a role in
167 maintaining glycemic control, which may positively influence brain health [27]. Our study found that
168 the average duration of a sedentary bout was higher in dementia patients compared to controls. This
169 observation suggests that not only reducing sedentary time, but also preventing prolonged sedentary
170 bouts by regularly breaking up sedentary behavior, can be targeted as a lifestyle intervention [28].

171

172 Limitations Since we only included community-dwelling patients, our results cannot be generalized to
173 institutionalized dementia patients. Furthermore, dementia patients in our study were enrolled in an
174 exercise trial [9]. Therefore, this group may be more motivated to be physically active. Nonetheless,
175 significant differences in physical activity characteristics were observed. Furthermore, physical
176 activity characteristics were measured by wrist-worn accelerometry, which is associated with limited
177 discriminative capacity between sedentary and very light intensity activities [29]. Consequently,
178 differences between dementia patients and controls might even be more pronounced. Another
179 limitation relates to our accelerometer, which has only been validated in a group of middle-aged
180 females [10] and was unable to correct for potential presence of short (~1-min) periods of non-wear
181 time. Nonetheless, we do not expect this will invalidate our findings of between-group differences in
182 physical activity and sedentary behavior. In addition, using wrist-worn accelerometry may explain the
183 two outliers in the younger dementia patients (Figure 2), which could relate to restless arm
184 movements. Whilst this may affect exploring individual differences in physical activity versus
185 sedentary behavior characteristics, significant differences remained present at group level.
186 Furthermore, we did not discriminate between types of sedentary activities. Cognitively challenging
187 sedentary activities, such as reading, might have a protective effect on cognition and are therefore less
188 harmful than passive sedentary activities (e.g. television-viewing) [30]. The final limitation is use of
189 the MMSE as a cognitive screening instrument since this measure, especially in healthy highly
190 educated older adults, has limited discriminative power to detect mild cognitive deficits [31,32]. It is

191 important to indicate that diagnosis of dementia was not made using the MMSE, but included standard
192 clinical procedures (including imaging if required).

193

194 Clinical relevance. Knowledge of physical activity characteristics across the entire activity spectrum
195 in dementia is highly relevant given that physical activity is an important factor accelerating
196 development and progression of dementia [3,7]. In addition, a high amount of sedentary behavior and
197 low amount of physical activity are associated with higher mortality and morbidity [5,6]. Even though
198 we found relatively small differences in physical activity characteristics between dementia patients
199 and controls, these may be highly relevant. For example, even very short breaks of light intensity
200 activity (i.e. 2 minutes walking) can already prevent acute metabolic [33] and cardiovascular [5]
201 impairments. This underlines the importance to develop interventions suitable for this vulnerable
202 patient group to safely engage in light intensity activities. However, future research should first
203 explore the role of physical activity and sedentary behavior in the progression and prevention of
204 dementia.


205

206 Conclusion. In the current study we objectively demonstrated that dementia patients spend
207 significantly more of their waking hours in sedentary behavior and spend less time in light-to-
208 moderate and moderate-to-vigorous intensity physical activity. Moreover, we found that older age
209 attenuated sedentary behavior and physical activity in controls, whilst this age-related decline is absent
210 in dementia patients. This means that patients with dementia (independent of age) lead a physically
211 inactive lifestyle characterized with significant time spent sedentary.

212 Taken together, these data improve our understanding of physical activity and sedentary behavior
213 characteristics in this highly relevant patient group and implies that targeting sedentary behavior and
214 physical activity may be relevant in dementia patients, especially at a younger age.

215 **FIGURE CAPTIONS**


216 **Fig. 1: Flowchart of participants**



217

218

219 Fig 2: Activity levels of dementia patients and cognitively healthy controls



220 Activity levels of younger and older age groups. Mean activity score (counts per minute) grouped by age.
 221 Controls Age<80 (n=23), Controls Age ≥80 (n=26), Dementia patients Age <80 (n=22), Dementia patients Age
 222 ≥80 (n=23). Values represent median ± interquartile range. * Significant difference between: Controls Age <80
 223 and Controls Age ≥80 P=0.044, Controls Age <80 and Dementia patients Age <80 P=0.004, Controls Age <80
 224 and Dementia patients Age ≥ 80 P=0.002.
 225

226
 227
 228
 229
 230

Sedentary behavior in dementia

Table 1. Baseline characteristics

	Control (n=49)	Dementia (n=45)	P-value
Age (years), mean \pm SD	80.0 \pm 7.7	79.6 \pm 5.9	0.744*
Females, n (%)	25 (51.0%)	22 (48.9%)	0.836†
Body Mass Index (kg/m ²), mean \pm SD	25.5 (4.0)	26.3 (5.0)	0.411*
Mini Mental State Examination‡, mean \pm SD	29.0 \pm 1.2	22.8 \pm 3.2	<0.001*
Number of walking aid users, n (%)	10 (20.4%)	16 (35.6%)	0.101†
Number of home-care receivers, n (%)	6 (12.2%)	24 (53.3%)	<0.001†
Number of comorbidities§, mean \pm SD	2.7 \pm 1.8	3.3 \pm 1.9	0.150*
Residence, n (%)			0.629†
- Independent, alone	18 (36.7%)	14 (31.1%)	
- Independent, together	30 (61.2%)	28 (62.2%)	
- Care home	1 (2.0%)	3 (6.7%)	
- Nursing home	0	0	
Dementia type, n (%)			
- Alzheimer's disease	n/a	25 (55.6%)	
- Vascular dementia	n/a	2 (4.4%)	
- Alzheimer's Disease/Vascular Dementia (%)	n/a	12 (26.7%)	
- Dementia type not specified (%)	n/a	6 (13.3%)	

*Differences between groups were tested with independent samples t-test; †Differences between groups were tested with Chi-Square Test; ‡ Scores on the Mini-Mental State Examination (MMSE) range from 0 (severe impairment) to 30 (no impairment); §Comorbidities are scored using the Older Persons and Informal Caregivers Survey-Minimum Dataset (TOPICS-MDS) with a theoretical range of 0-17 and a higher score indicates more comorbidities [34]

Sedentary behavior in dementia

Table 2: Physical activity and sedentary behavior characteristics

	Control (n=49)	Dementia (n=45)	P-value
Total waking time (h/day)	15.4 ± 1.0*	14.9 ± 1.3*	0.053†
Counts per minute (day ⁻¹)	226 ± 61*	186 ± 76*	0.005 †
Absolute values (h/day)			
Sedentary time	8.1 (7.2 – 9.2)	8.5 (7.2 – 10.0)	0.216
Very light intensity activity	2.2 (1.9 – 2.6)	2.3 (1.7 – 2.9)	0.748
Light-to-moderate intensity activity	3.5 (2.7 – 4.0)	2.7 (2.0 – 3.7)	0.006
Moderate-to-vigorous intensity activity	1.5 (0.8 – 2.0)	0.8 (0.4 – 1.5)	0.001
Relative values (% of total measuring time)			
% sedentary time	55 (47 – 59)	57 (49 – 68)	0.042
% very light intensity activity	15 (12 – 16)	16 (12 – 19)	0.284
% light-to-moderate intensity activity	22 (18 – 25)	20 (15 – 23)	0.017
% moderate-to-vigorous intensity activity	10 (5 – 13)	5 (2 – 10)	0.001
Sedentary behavior characteristics			
Number of interruptions in sedentary behavior (day ⁻¹)	28.2 (26.2 – 32.5)	27.2 (24.5 – 31.0)	0.195

Sedentary behavior in dementia

Number of 30 minutes prolonged sedentary bouts (day ⁻¹)	2.0 (0.9 – 3.3)	2.3 (1.0 – 4.1)	0.227
Duration of average sedentary bout (minutes)	16.6 (15.3 – 18.4)	18.3 (16.4 –21.1)	0.008

Median and Interquartile range are presented unless reported otherwise. P-values represent Mann-Whitney U test; *Mean ± SD; †Differences between groups were tested with independent samples t-test

REFERENCES

- 1 Perera G, Pedersen L, Ansel D, Alexander M, Arrighi HM, Avillach P, et al: Dementia prevalence and incidence in a federation of European Electronic Health Record databases-The European Medical Informatics Framework resource. *Alzheimers Dement* 2018;14:130-139.
- 2 Kavirajan H, Schneider LS: Efficacy and adverse effects of cholinesterase inhibitors and memantine in vascular dementia: a meta-analysis of randomised controlled trials. *Lancet Neurol* 2007;6:782-792.
- 3 Ashby-Mitchell K, Burns R, Shaw J, Anstey KJ: Proportion of dementia in Australia explained by common modifiable risk factors. *Alzheimers Res Ther* 2017;9:11.
- 4 Groot C, Hooghiemstra A, Raijmakers P, Van Berckel B, Scheltens P, Scherder E, et al: The effect of physical activity on cognitive function in patients with dementia: a meta-analysis of randomized control trials. *Ageing Res Rev* 2016;25:13-23.
- 5 Carter S, Hartman Y, Holder S, Thijssen DH, Hopkins ND: Sedentary Behavior and Cardiovascular Disease Risk: Mediating Mechanisms. *Exerc Sport Sci Rev* 2017;45:80-86.
- 6 Ekelund U, Steene-Johannessen J, Brown WJ, Fagerland MW, Owen N, Powell KE, et al: Does physical activity attenuate, or even eliminate, the detrimental association of sitting time with mortality? A harmonised meta-analysis of data from more than 1 million men and women. *Lancet* 2016;388:1302-1310.
- 7 Falck RS, Davis JC, Liu-Ambrose T: What is the association between sedentary behaviour and cognitive function? A systematic review. *Br J Sports Med* 2017;51:800-811.
- 8 Folstein MF, Folstein SE, McHugh PR: "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975;12:189-198.
- 9 Karssemeijer EGA, Bossers WJR, Aaronson JA, Kessels RPC, Olde Rikkert MGM: The effect of an interactive cycling training on cognitive functioning in older adults with mild dementia: study protocol for a randomized controlled trial. *BMC Geriatr* 2017;17:73.
- 10 Neil-Sztramko SE, Rafn BS, Gotay CC, Campbell KL: Determining activity count cut-points for measurement of physical activity using the Actiwatch2 accelerometer. *Physiol Behav* 2017;173:95-100.
- 11 Hart TL, Swartz AM, Cashin SE, Strath SJ: How many days of monitoring predict physical activity and sedentary behaviour in older adults? *Int J Behav Nutr Phys Act* 2011;8:62.
- 12 Sasaki JE, Junior JH, Meneguci J, Tribess S, Marocolo Junior M, Stabelini Neto A, et al: Number of days required for reliably estimating physical activity and sedentary behaviour from accelerometer data in older adults. *J Sports Sci* 2018;36:1572-1577.
- 13 Ancoli-Israel S, Cole R, Alessi C, Chambers M, Moorcroft W, Pollak CP: The role of actigraphy in the study of sleep and circadian rhythms. *Sleep* 2003;26:342-392.
- 14 van der Ploeg HP, Chey T, Korda RJ, Banks E, Bauman A: Sitting time and all-cause mortality risk in 222 497 Australian adults. *Arch Intern Med* 2012;172:494-500.
- 15 van Alphen HJ, Volkers KM, Blankevoort CG, Scherder EJ, Hortobagyi T, van Heuvelen MJ: Older Adults with Dementia Are Sedentary for Most of the Day. *PLoS One* 2016;11:e0152457.
- 16 James BD, Boyle PA, Bennett DA, Buchman AS: Total daily activity measured with actigraphy and motor function in community-dwelling older persons with and without dementia. *Alzheimer Dis Assoc Disord* 2012;26:238-245.
- 17 Keadle SK, McKinnon R, Graubard BI, Troiano RP: Prevalence and trends in physical activity among older adults in the United States: A comparison across three national surveys. *Prev Med* 2016;89:37-43.
- 18 Harvey JA, Chastin SF, Skelton DA: How Sedentary are Older People? A Systematic Review of the Amount of Sedentary Behavior. *J Aging Phys Act* 2015;23:471-487.
- 19 Manns P, Ezeugwu V, Armijo-Olivo S, Vallance J, Healy GN: Accelerometer-Derived Pattern of Sedentary and Physical Activity Time in Persons with Mobility Disability: National Health and Nutrition Examination Survey 2003 to 2006. *J Am Geriatr Soc* 2015;63:1314-1323.

- 20 Visser M, Wijnhoven HAH, Comijs HC, Thomese F, Twisk JWR, Deeg DJH: A Healthy Lifestyle in Old Age and Prospective Change in Four Domains of Functioning. *J Aging Health*: 2018;898264318774430.
- 21 Falck RS, McDonald SM, Beets MW, Brazendale K, Liu-Ambrose T: Measurement of physical activity in older adult interventions: a systematic review. *Br J Sports Med* 2016;50:464-470.
- 22 McPherson S, Fairbanks L, Tiken S, Cummings JL, Back-Madruga C: Apathy and executive function in Alzheimer's disease. *J Int Neuropsychol Soc* 2002;8:373-381.
- 23 David R, Mulin E, Friedman L, Le Duff F, Cygankiewicz E, Deschaux O, et al: Decreased daytime motor activity associated with apathy in Alzheimer disease: an actigraphic study. *Am J Geriatr Psychiatry* 2012;20:806-814.
- 24 Sabia S, Dugravot A, Dartigues JF, Abell J, Elbaz A, Kivimaki M, et al: Physical activity, cognitive decline, and risk of dementia: 28 year follow-up of Whitehall II cohort study. *BMJ* 2017;357:j2709.
- 25 Falck RS, Landry GJ, Best JR, Davis JC, Chiu BK, Liu-Ambrose T: Cross-Sectional Relationships of Physical Activity and Sedentary Behavior With Cognitive Function in Older Adults With Probable Mild Cognitive Impairment. *Phys Ther* 2017;97:975-984.
- 26 Koolhaas CM, van Rooij FJA, Schoufour JD, Cepeda M, Tiemeier H, Brage S, et al: Objective Measures of Activity in the Elderly: Distribution and Associations With Demographic and Health Factors. *J Am Med Dir Assoc* 2017;18:838-847.
- 27 Wheeler MJ, Dempsey PC, Grace MS, Ellis KA, Gardiner PA, Green DJ, et al: Sedentary behavior as a risk factor for cognitive decline? A focus on the influence of glycemic control in brain health. *Alzheimer's & dementia (New York, N Y)* 2017;3:291-300.
- 28 Dempsey PC, Owen N, Biddle SJH, Dunstan DW: Managing Sedentary Behavior to Reduce the Risk of Diabetes and Cardiovascular Disease. *Current Diabetes Reports* 2014;14:522.
- 29 Tudor-Locke C, Barreira TV, Schuna JM: Comparison of step outputs for waist and wrist accelerometer attachment sites. *Med Sci Sports Exerc* 2015;47:839-842.
- 30 Kikuchi H, Inoue S, Sugiyama T, Owen N, Oka K, Nakaya T, et al: Distinct associations of different sedentary behaviors with health-related attributes among older adults. *Prev Med* 2014;67:335-339.
- 31 Tangalos EG, Smith GE, Ivnik RJ, Petersen RC, Kokmen E, Kurland LT, et al: The Mini-Mental State Examination in general medical practice: clinical utility and acceptance: *Mayo Clin Proc, Elsevier*, 1996, pp 829-837.
- 32 O'Bryant SE, Humphreys JD, Smith GE, Ivnik RJ, Graff-Radford NR, Petersen RC, et al: Detecting dementia with the mini-mental state examination in highly educated individuals. *Arch Neurol* 2008;65:963-967.
- 33 Dunstan DW, Kingwell BA, Larsen R, Healy GN, Cerin E, Hamilton MT, et al: Breaking up prolonged sitting reduces postprandial glucose and insulin responses. *Diabetes Care* 2012;35:976-983.
- 34 Lutomski JE, Baars MA, Schalk BW, Boter H, Buurman BM, den Elzen WP, et al: The development of the Older Persons and Informal Caregivers Survey Minimum DataSet (TOPICS-MDS): a large-scale data sharing initiative. *PLoS One* 2013;8:e81673.