



LJMU Research Online

Bachelet, E, Bozza, V, Han, C, Udalski, A, Bond, IA, Beaulieu, J-P, Street, RA, Kim, H-I, Bramich, DM, Cassan, A, Dominik, M, Jaimes, RF, Horne, K, Hundertmark, M, Mao, S, Menzies, J, Ranc, C, Schmidt, R, Snodgrass, C, Steele, IA, Tsapras, Y, Wambsganss, J, Mroz, P, Soszynski, I, Szymanski, MK, Skowron, J, Pietrukowicz, P, Kozlowski, S, Poleski, R, Ulaczyk, K, Pawlak, M, Abe, F, Barry, R, Bennett, DP, Bhattacharya, A, Donachie, M, Fukui, A, Hirao, Y, Itow, Y, Kawasaki, K, Kondo, I, Koshimoto, N, Li, MCA, Matsubara, Y, Muraki, Y, Miyazaki, S, Nagakane, M, Rattenbury, NJ, Suematsu, H, Sullivan, DJ, Sumi, T, Suzuki, D, Tristram, PJ and Yonehara, A

First Assessment of the Binary Lens OGLE-2015-BLG-0232

<http://researchonline.ljmu.ac.uk/id/eprint/10211/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Bachelet, E, Bozza, V, Han, C, Udalski, A, Bond, IA, Beaulieu, J-P, Street, RA, Kim, H-I, Bramich, DM, Cassan, A, Dominik, M, Jaimes, RF, Horne, K, Hundertmark, M, Mao, S, Menzies, J, Ranc, C, Schmidt, R, Snodgrass, C, Steele, IA, Tsapras, Y, Wambsganss, J, Mroz, P, Soszynski, I, Szymanski, MK, Skowron, J, Pietrukowicz, P, Kozlowski, S, Poleski, R, Ulaczyk, K, Pawlak, M, Abe, F, Barry, R, Bennett, DP, Bhattacharya, A, Donachie, M, Fukui, A, Hirao, Y, Itow, Y, Kawasaki, K, Kondo, I, Koshimoto, N, Li, MCA, Matsubara, Y, Muraki, Y, Miyazaki, S, Nagakane, M, Rattenbury, NJ, Suematsu, H, Sullivan, DJ, Sumi, T, Suzuki, D, Tristram, PJ and Yonehara, A

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

<http://researchonline.ljmu.ac.uk/>

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

FIRST ASSESSMENT OF THE BINARY LENS OGLE-2015-BLG-0232.

E. BACHELET^{R0}, V. BOZZA^{S0,S1}, C. HAN^{C0}, A. UDALSKI^{O0}, I.A. BOND^{M0}, J.-P. BEAULIEU^{P0,P1}, R.A. STREET^{R0}, J.-I. KIM^{C1}

AND,

D. M. BRAMICH^{R1}, A. CASSAN^{R2}, M. DOMINIK^{R3,†}, R. FIGUERA JAIMES^{R7,R3,R4}, K. HORNE^{R3}, M. HUNDERTMARK^{R5,R6}, S. MAO^{R7,R8,R9}, J. MENZIES^{R10}, C. RANC^{R2}, R. SCHMIDT^{R6}, C. SNODGRASS^{R11}, I. A. STEELE^{R12}, Y. TSAPRAS^{R6}, J. WAMBSGANSS^{R6},

(THE ROBONET COLLABORATION)

P. MRÓZ^{O0}, I. SOSZYŃSKI^{O0}, M.K. SZYMAŃSKI^{O0}, J. SKOWRON^{O0}, P. PIETRUKOWICZ^{O0}, S. KOZŁOWSKI^{O0}, R. POLESKI^{O1}, K. ULACZYK^{O0}, M. PAWLAK^{O0},

(THE OGLE COLLABORATION)

F. ABE^{M2}, R. BARRY^{M3}, D. P. BENNETT^{M3,M10}, A. BHATTACHARYA^{M3,M10}, M. DONACHIE^{M4}, A. FUKUI^{M5,M13}, Y. HIRAO^{M1}, Y. ITOW^{M2}, K. KAWASAKI^{M1}, I. KONDO^{M1}, N. KOSHIMOTO^{M11,M12}, M. CHEUNG ALEX LI^{M4}, Y. MATSUBARA^{M2}, Y. MURAKI^{M2}, S. MIYAZAKI^{M1}, M. NAGAKANE^{M1}, N. J. RATTENBURY^{M4}, H. SUEMATSU^{M1}, D. J. SULLIVAN^{M6}, T. SUMI^{M1}, D. SUZUKI^{M9}, P. J. TRISTRAM^{M7}, A. YONEHARA^{M8}

(THE MOA COLLABORATION)

^{R0}Las Cumbres Observatory, 6740 Cortona Drive, Suite 102, Goleta, CA 93117 USA

^{R1}New York University Abu Dhabi, PO Box 129188, Saadiyat Island, Abu Dhabi, UAE

^{R2}Sorbonne Universités, UPMC Univ Paris 6 et CNRS, UMR 7095, Institut d'Astrophysique de Paris, 98 bis bd Arago, 75014 Paris, France

^{R3}SUPA, School of Physics & Astronomy, University of St Andrews, North Haugh, St Andrews KY16 9SS, UK

^{R4}European Southern Observatory, Karl-Schwarzschild-Str. 2, 85748 Garching bei München, Germany

^{R5}Niels Bohr Institute & Centre for Star and Planet Formation, University of Copenhagen, Øster Voldgade 5, 1350 - Copenhagen K, Denmark

^{R6}Zentrum für Astronomie der Universität Heidelberg, Astronomisches Rechen-Institut, Mönchhofstr. 12-14, 69120 Heidelberg, Germany

^{R7}Physics Department and Tsinghua Centre for Astrophysics, Tsinghua University, Beijing 100084, China

^{R8}National Astronomical Observatories, Chinese Academy of Sciences, 20A Datun Road, Chaoyang District, Beijing 100012, China

^{R8}Jodrell Bank Centre for Astrophysics, School of Physics and Astronomy, The University of Manchester, Oxford Road, Manchester M13 9PL, UK

^{R9}South African Astronomical Observatory, PO Box 9, Observatory 7935, South Africa

^{R10}Planetary and Space Sciences, Department of Physical Sciences, The Open University, Milton Keynes, MK7 6AA, UK

^{R11}Astrophysics Research Institute, Liverpool John Moores University, Liverpool CH41 1LD, UK

^{C0}Department of Physics, Chungbuk National University, Cheongju 28644, Korea

^{C1}Korea Astronomy and Space Science Institute, Daejeon 34055, Korea

^{O0}Warsaw University Observatory, Al. Ujazdowskie 4, 00-478 Warszawa, Poland

^{O1}Department of Astronomy, Ohio State University, 140 W. 18th Ave., Columbus, OH 43210, USA

^{M0}Institute of Natural and Mathematical Sciences, Massey University, Auckland 0745, New Zealand

^{M1}Department of Earth and Space Science, Graduate School of Science, Osaka University, Toyonaka, Osaka 560-0043, Japan

^{M2}Institute for Space-Earth Environmental Research, Nagoya University, Nagoya 464-8601, Japan

^{M3}Code 667, NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA

^{M4}Department of Physics, University of Auckland, Private Bag 92019, Auckland, New Zealand

^{M5}Subaru Telescope Okayama Branch Office, National Astronomical Observatory of Japan, NINS, 3037-5 Honjo, Kamogata, Asakuchi, Okayama 719-0232, Japan

^{M6}School of Chemical and Physical Sciences, Victoria University, Wellington, New Zealand

^{M7}University of Canterbury Mt. John Observatory, P.O. Box 56, Lake Tekapo 8770, New Zealand

^{M8}Department of Physics, Faculty of Science, Kyoto Sangyo University, 603-8555 Kyoto, Japan

^{M9}Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, 3-1-1 Yoshinodai, Chuo, Sagamihara, Kanagawa, 252-5210, Japan

^{M10}Department of Astronomy, University of Maryland, College Park, MD 20742, USA

^{M11}Department of Astronomy, Graduate School of Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

^{M12}National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan

^{M13}Instituto de Astrofísica de Canarias, Vía Láctea s/n, E-38205 La Laguna, Tenerife, Spain

^{S0}Dipartimento di Fisica "E.R. Caianiello", Università di Salerno, Via Giovanni Paolo II 132, 84084, Fisciano, Italy

^{S1}Istituto Internazionale per gli Alti Studi Scientifici (IIASS), Via G. Pellegrino 19, 84019 Vietri sul Mare (SA), Italy

^{P0}School of Physical Sciences, University of Tasmania, Private Bag 37 Hobart, Tasmania 7001 Australia; jeanphilippe.beaulieu@utas.edu.au

^{P1}Sorbonne Universités, UPMC Université Paris 6 et CNRS, UMR 7095, Institut d'Astrophysique de Paris, 98http://www.plt.axvline/ bis bd Arago, 75014 Paris, France; beaulieu@iap.fr and

† Royal Society University Research Fellow

(Received receipt date; revised revision date; Accepted acceptance date)

Draft version November 8, 2018

ABSTRACT

We present an analysis of the microlensing event OGLE-2015-BLG-0232. This event is challenging to characterize for two reasons. First, the light curve is not well sampled during the caustic crossing due to the proximity of the full Moon impacting the photometry quality. Moreover, the source brightness is difficult to estimate because this event is blended with a nearby K dwarf star. We found that the light curve deviations are likely due to a close brown dwarf companion (i.e., $s = 0.55$ and $q = 0.06$), but the exact nature of the lens is still unknown. We finally discuss the potential of follow-up observations to estimate the lens mass and distance in the future.

Keywords: gravitational microlensing

1. INTRODUCTION

Twenty years after the first exoplanet detection, it is clear that planets are abundant in the Milky Way (Cassan et al.

2012; Fressin et al. 2013; Bonfils et al. 2013; Clanton & Gaudi 2016; Suzuki et al. 2016). But the dividing line between super-Jupiter and brown dwarfs is still uncertain. Burrows et al. (2001) define brown dwarfs as objects within mass limits $[13, 73] M_J$. As underlined by Schlaufman (2018), this definition is problematic because the critical mass for deuterium burning depends on the object composition (Spiegel et al. 2011). More recently, an alternative definition has been proposed based on the formation mechanisms (Schneider et al. 2011): planets are formed by core accretion while brown dwarfs are a result of gas collapse. The former is motivated by exoplanet formation models and by the observational evidence that giant planets tend to form more frequently around metal-rich stars (Mordasini et al. 2012; Buchhave et al. 2012; Mortier et al. 2012). In contrast, Latham et al. (2002) found no significant correlation between metallicity and stellar binary occurrence. But this definition is also problematic because it is nearly impossible to distinguish the two scenarios observationally (Wright et al. 2011; Bryan et al. 2018). Recently, Schlaufman (2018) revisited the mass definition by combining and clustering samples of low-mass stars, brown dwarfs and planets orbiting Solar-type stars and ultimately derived a surprisingly low upper planetary mass limit of $\sim 6 M_J$.

Brown dwarf detections are therefore important to understand the planetary regime boundaries but these objects are intrinsically difficult to detect directly, due to their low-luminosity. Moreover, the radii of brown dwarfs and Jupiter-like planets are very similar due to the degeneracy pressure (Zapolsky & Salpeter 1969; Burrows & Liebert 1993). It is therefore difficult to distinguish them with the transit method alone. Microlensing on the other hand can detect brown dwarfs several kpc away, either in binary systems or as single objects (Zhu et al. 2016; Chung et al. 2017; Shvartzvald et al. 2018), because the method does not need flux measurements from the lens. Several brown dwarfs and brown dwarfs candidates have been discovered through this method (Bachelet et al. 2012a; Bozza et al. 2012; Ranc et al. 2015; Han et al. 2016; Poleski et al. 2017; Mróz et al. 2017).

In this work, we present the analysis of OGLE-2015-BLG-0232/MOA-2015-BLG-046. The data presented in Section 2 show clear signatures of a binary lens event. In Section 3, we present the modeling procedure and find that the mass ratio of the lens system favors a brown dwarf companion (close model) or a low-mass M dwarf companion (wide model). We present a detailed study of both the microlensing source and the bright blend in Section 4. Because no parallax was measured, we discuss in Section 6 the possible follow-up observations to unlock the final solution of this microlensing puzzle.

2. OBSERVATIONS

The microlensing event OGLE-2015-BLG-0232 ($\alpha = 18^{\text{h}}06^{\text{m}}43.84^{\text{s}}$, $\delta = -32^{\circ}54^{\text{m}}27.3^{\text{s}}$; $l = -1^{\circ}.172199$, $b = -5^{\circ}.9060$) was an early event of the 2015 microlensing season first discovered by the Optical Gravitational Lens Experiment (OGLE) (Udalski 2003) on 2015 March 2 UT 17:50 and also detected later by the Microlensing Observations in Astrophysics (MOA) collaboration (Bond et al. 2001) as MOA-2015-BLG-046 on 2015 March 10 at UT 16:42. C. Han first delivered an email alert indicating an ongoing anomaly on 2015 March 15 at UT 02:16. Independently, the RoboNet team, based on the SIGNALMEN anomaly detector (Dominik et al. 2007) and the RoboTAP algorithm (Hundertmark et al. 2017), automatically triggered observations on the Las Cumbres Observatory network of robotic telescopes (Tsapras et al.

2009). Unfortunately, the Moon was nearly full during this period, preventing surveys from acquiring more data during the anomaly. This event was also observed in the near infrared by the VISTA Variables in the Via Lactea (VVV) survey (Minniti et al. 2010). Real-time modeling conducted independently by C.Han and V.Bozza indicated that this event was probably due to a low-mass binary lens ($q \sim 0.01$). All teams reprocessed their photometry at the end of the season using the difference image analysis (DIA) technique : RoboNet used DanDIA (Bramich 2008; Bramich et al. 2013), OGLE and MOA used their own implementation of DIA (Udalski et al. 2015; Bond et al. 2001). The K band of VVV was re-reduced using pySIS (Albrow et al. 2009). The VVV pySIS photometry were roughly calibrated to an independent VVV catalog (Beaulieu et al. 2016) by adding an offset of 0.6 mag. Note that the VVV_K light curve is nearly flat, so we did not use this dataset in the first round of modeling. In total, 7659 data points are available for the analysis, as summarized in Table 1.

3. MODELING

3.1. Description

This event is clearly anomalous and real-time models found that a binary lens with a small mass ratio accurately reproduces the observations. A static binary model is described with seven parameters : t_0 the time of the minimum impact parameter u_0 , $t_E = \theta_E/\mu$ the angular Einstein radius crossing time, $\rho = \theta_*/\theta_E$ the normalized angular source radius, s the normalized projected separation, q the mass ratio between the two lens components and finally α the lens/source trajectory angle relative to the binary axis. Here, μ is the relative proper motion between the source and the lens and θ_E is the angular Einstein ring, see for example Gould (2000). Note that we restrict the modeling of the data points to the time window $t \in [2456850, 2457200]$ to speed-up the modeling. For events like OGLE-2015-BLG-0232 that exhibit caustic crossings, the limb-darkening of the source star has to be considered. Unfortunately, in this case, the observations taken around HJD ~ 2457087 were in SDSS- i' band only in order to reduce the impact of the moonlight. Moreover, the caustic crossings are not intensively covered by the data. For these reasons, we investigated a simpler model, the Uniform Source Binary Lens (USBL)(Bozza 2010; Bozza et al. 2012) and use pyLIMA (Bachelet et al. 2017) to perform the modeling. A detailed description of this binary fitting code is given in Bachelet (2018, in prep). We did not use the standard grid approach to locate the global minimum, but instead ran a global search on all parameters using the differential evolution method (Storn & Price 1997; Bachelet et al. 2017). Briefly, this method uses a set of starting points in parameter space and maintains an ordered population of candidate solutions while exploring potential new solutions by combining existing ones. This algorithm was successfully tested by applying it to previously published events. In practice, we split the parameter space in two regions: $s < 1$ and $s > 1$. This is motivated by the dramatic change of the caustics topology between these two regimes but also the presence of the close/wide degeneracy, see for example Erdl & Schneider (1993); Dominik (1999); Bozza (2000); Cassan (2008). We ran the algorithm several times and found that it converged to similar solutions. This event was also modeled in real time by V.Bozza using

Name	Collaboration	Location	Aperture(m)	Filter	Code	N_{data}	Longitude(deg)	Latitude(deg)
OGLE _I	OGLE	Chile	1.3	I	Woźniak	525	289.307	-29.015
MOA _{Red}	MOA	New Zealand	1.8	Red	Bond	6569	170.465	43.987
MOA _V	Boller&Chivens	New Zealand	0.6	V	Bond	184	170.465	43.987
VVV _K	VISTA	Chile	4.1	K	pySIS	198	289.6081	-24.616
LSCA _i	RoboNet	Chile	1.0	SDSS-i	DanDIA	30	289.195	-30.167
LSCB _i	RoboNet	Chile	1.0	SDSS-i	DanDIA	23	289.195	-30.167
LSCC _i	RoboNet	Chile	1.0	SDSS-i	DanDIA	21	289.195	-30.167
CPTA _i	RoboNet	South Africa	1.0	SDSS-i	DanDIA	21	220.810	-32.347
CPTB _i	RoboNet	South Africa	1.0	SDSS-i	DanDIA	21	220.810	-32.347
CPTC _i	RoboNet	South Africa	1.0	SDSS-i	DanDIA	12	220.810	-32.347
COJA _i	RoboNet	Australia	1.0	SDSS-i	DanDIA	29	149.065	-31.273
COJB _i	RoboNet	Australia	1.0	SDSS-i	DanDIA	18	149.065	-31.273

Table 1
Summary of observations.

RTModel¹. This system uses a different method to explore the parameter space: a template matching approach (Mao & Di Stefano 1995; Liebig et al. 2015). It also found similar solutions, raising confidence in our results. Results relative to this first exploration can be seen Table 3.

3.2. Error bar rescaling

It is common practice to rescale the uncertainties in microlensing using (in mag unit in the present work):

$$\sigma' = k \sqrt{\sigma^2 + e_{\min}^2} \quad (1)$$

where σ' is the rescaled uncertainty, k and e_{\min} are parameters that need to be tuned to reach a certain metric to optimize. The usual metric used is to force the χ^2/dof for each dataset to converge to 1 (Bachelet et al. 2012b; Miyake et al. 2012; Yee et al. 2013). However, Andrae et al. (2010) show that the use of the reduced χ^2 , for model diagnostic, is relevant only for linear models, which is not the case in the present work. Instead, they recommend the use of normality tests of residuals, like Bachelet et al. (2015).

The physical reasons that motivate the rescaling are to account for photometric low-level systematics and potential underestimation of the uncertainties. There are multiple causes coming from both instrumentation and software reductions. The impact is expected to be different for each dataset and therefore, instead of automatically rescaling the errorbars of each dataset blindly, we assessed whether this was necessary. To do so, we use the approach describe below.

First, we rescaled OGLE-IV uncertainties using the custom method of Skowron et al. (2016)². We then analyzed the residuals around the best model using three test of normality : a Kolmogorov-Smirnov test, an Anderson-Darling test and a Shapiro-Wilk test. We considered to rescale a dataset if any of these test were not successful (i.e, the p-value associated to the test was less than 1%). All datasets, except MOA_{Red}, passed the three normality tests. The majority of datasets present a relative small number of observations (≤ 100), any deviations to normality would be then hard to detect. On the other hand, it might indicate that uncertainties reproduce the data scatter accurately. Note that the OGLE-IV dataset also passed the three tests after the rescaling process.

Name	N_{data}	k	e_{\min}
OGLE _I ^a	68	0.97 ± 0.06	0.0
MOA _{Red}	467	3.51 ± 0.08	0.0
MOA _V	43	2.1 ± 0.2	0.0
VVV _K	14	3.5 ± 0.5	0.0
LSCA _i	30	1.1 ± 0.1	0.0
LSCB _i	23	1.9 ± 0.2	0.0
LSCC _i	21	1.7 ± 0.2	0.0
CPTA _i	21	1.1 ± 0.1	0.0
CPTB _i	21	1.2 ± 0.1	0.0
CPTC _i	12	1.2 ± 0.2	0.0
COJA _i	29	1.7 ± 0.1	0.0
COJB _i	18	2.1 ± 0.3	0.0

Table 2

Error bar rescaling coefficients used in this paper. ^a Note that the OGLE_I uncertainties receive a special treatment before this rescaling step, see text.

As a secondary check, we follow the same approach as Dominik et al. (2018) and fit the parameters of Equation 1 around the best model from the previous section, using the modified $\chi^{2'}$:

$$\chi^{2'} = \sum_i \frac{(f_i - m_i)^2}{\sigma_i'^2} + 2 \ln(\sigma_i') \quad (2)$$

with f_i the observed flux, m_i the microlensing model in flux and σ_i' is the modified error in flux relative to Equation 1. It appeared rapidly that the term e_{\min} was not constrained, due to the relative small range of magnification in the light curves. We therefore delete this term from Equation 1 and fit only the first term k . The results presented in the Table 2 are consistent with the previous analysis and indicate a soft rescaling, with the exception of the MOA_{Red} dataset.

3.3. Results

Both algorithms converged to models with similar geometries: the strong anomalies seen in Figure 1 are due to a central caustic crossing. However the data do not constrain strongly the models, leading to significant difference in the model parameters given in the Table 3. To obtain a more comprehensive picture, we run two sets of Monte-Carlo Markov Chain (MCMC) explorations around these best models, using the **emcee** algorithm (Foreman-Mackey et al. 2013) implemented

¹ <http://www.fisica.unisa.it/GravitationAstrophysics/RTModel.htm>

² <http://ogle.astrouw.edu.pl/ogle4/errorbars/blg/errcorr-OIV-BLG-I.dat>

in pyLIMA. Note that during this optimization process, we modified the model parameters so that we model t_c and u_c , the time and closest approach to the central caustic, instead of t_0 and u_0 . The idea is to use parameters more directly related to the main features of the light curve. This is a standard practice that significantly improves the model convergence (Cas-san 2008; Han 2009; Penny 2014).

The geometry of the best fitting model is sensitive to the close/wide degeneracy (Griest & Safizadeh 1998; Bozza 2000; Dominik 2009). However, close models are slightly favored. The mass ratio of this event is not well constrained. This is due to a lack of observations during the anomaly, especially during the central caustic entrance and exit.

We tried to model second-order effects, such as annual parallax and the orbital motion of the lens (Gould & Loeb 1992; Dominik 1998; Albrow et al. 2000; Gould 2004; Bachelet et al. 2012b). Due to the relatively short timescale of the event and the relatively low coverage of the anomaly features, these second order effects were not constrained well enough to be considered as a solid detection.

4. PROPERTIES OF THE SOURCE

4.1. Optical observations

Following Bond et al. (2017), we calibrated the MOA_R and MOA_V magnitudes to the OGLE-III system using the relation in the Appendix. The resulting color-magnitude diagram (CMD thereafter) is presented in Figure 2, and we summarize information from the various catalogs used in Table 4. We found that the color of the red giant clump (RGC) centroid is $(V - I)_{RGC} = 1.75 \pm 0.05$ mag and its brightness is $I_{RGC} = 15.3 \pm 0.1$ mag. Knowing the intrinsic color of the RGC $(V - I)_{0,RGC} = 1.06$ mag and its intrinsic brightness $I_{0,RGC} = 14.45$ mag (Nataf et al. 2013), we estimate the absorption $A_I = 0.9 \pm 0.1$ mag and the extinction $E(V - I) = 0.69 \pm 0.05$ mag toward the microlensing event. We found a good agreement with an independent determination using the Interstellar Extinction Calculator on the OGLE website³, based on Nataf et al. (2013) and Gonzalez et al. (2012), with $A_I = 0.79 \pm 0.1$ mag and $E(V - I) = 0.68 \pm 0.05$ mag. From the best model and the color transformations in the Appendix, the source magnitudes are $V_{s,OGLE_{III}} = 21.2 \pm 0.1$ mag and $I_{s,OGLE_{III}} = 19.15 \pm 0.09$ mag (and a color of $(V - I)_{s,OGLE_{III}} = 2.0 \pm 0.1$ mag). In principle, it is possible to obtain a model-independent color using linear regression between two bands λ_1 and λ_2 since the microlensing magnification is achromatic (Dong et al. 2006; Bond et al. 2017):

$$f_{\lambda_1} = \frac{f_{s,\lambda_1}}{f_{s,\lambda_2}}(f_{\lambda_2} - f_{b,\lambda_2}) + f_{b,\lambda_1} \quad (3)$$

where f_s and f_b are the source and blending flux respectively. However, this requires simultaneous observations which are difficult in practice. Here, we consider MOA_R and MOA_V as simultaneous if the acquisition time was within 15 minutes. We found a model independent source color of $(V - I)_{s,OGLE_{III}} = 2.0 \pm 0.1$ mag, in agreement with the previous estimation. Finally, we obtained the intrinsic color $(V - I)_{0,s,OGLE_{III}} = 1.4 \pm 0.1$ mag and brightness $I_{0,s,OGLE_{III}} = 18.4 \pm 0.1$ of the source in the OGLE-III system (i.e., in the Johnson-Cousins system).

Because this event was also observed by OGLE-IV, we conducted a similar analysis using the OGLE-IV CMD. The cor-

responding CMD is presented in Figure 2. In this CMD, we found that the color of the red giant clump (RGC) centroid is $(V - I)_{RGC} = 1.67 \pm 0.05$ mag and its brightness is $I_{RGC} = 15.3 \pm 0.1$ mag. The best model and the V -band transformation in Equation A4 lead to $V_{s,OGLE_{IV}} = 21.2 \pm 0.1$ mag and $I_{s,OGLE_{IV}} = 19.47 \pm 0.01$ mag (and a color of $(V - I)_{s,OGLE_{IV}} = 1.7 \pm 0.1$ mag). Assuming the source suffers the same extinction as the RGC, we measured an offset between the source and the RGC $\Delta((V - I)_{s,OGLE_{IV}}, I_{s,OGLE_{IV}}) = (0.03 \pm 0.1, 4.2 \pm 0.1)$. However, the OGLE-IV system is not perfectly calibrated, and the difference in the colors need to be multiplied by a factor 0.93 (for the CCD 24 of the OGLE camera mosaic) (Udalski et al. 2015). Based on the OGLE-IV CMD, the source color is $(V - I)_{0,s,OGLE_{IV}} = 1.09 \pm 0.1$ mag and the brightness is $I_{0,s,OGLE_{IV}} = 18.7 \pm 0.1$ mag.

While the two studies converge to a similar conclusion, we use for the source properties $(V - I)_{0,s,OGLE_{IV}} = 1.09 \pm 0.1$ mag and $I_{0,s,OGLE_{IV}} = 18.7 \pm 0.1$ mag, because they rely on a single color transformation and also because the color term in Equation A4 is smaller than the one in Equation A2. From optical observations, the source is probably a K-dwarf (Bessell & Brett 1988) or, potentially, a K subgiant that lies behind the Galactic Bulge.

Using Kervella & Fouqué (2008) and the optical color, we can obtain the angular source radius θ_* . We obtain 13% precision on $\theta_* = 0.8 \pm 0.1 \mu\text{as}$. Finally, we can then estimate the angular Einstein ring radius $\theta_E = \theta_*/\rho = 0.8 \pm 0.2$ mas (using the best model) and $\mu = 7.0 \pm 3$ mas/yr. This provides one mass and distance constraint to the lens system since (Gould 2000):

$$M_{tot} = \frac{\theta_E^2}{\kappa \pi_{rel}} \quad (4)$$

with $\pi_{rel} = \frac{1-x}{D_s x}$ au, $x = D_l/D_s$ (the distance to the lens and the source respectively) and the constant $\kappa = 8.144 \text{ mas.M}_\odot^{-1}$.

4.2. Near infrared

Thanks to VVV observations, we can perform a similar study using K -band data and construct a near-infrared CMD, as shown in Figure 3. Gonzalez et al. (2012) provide extinction maps toward the Galactic Bulge. Using their online tool⁴, we found $A_K = 0.10 \pm 0.06$ mag and $E(J - K) = 0.19 \pm 0.11$ mag. This agrees relatively well with the 3D Maps toward the Galactic Bulge of Schultheis et al. (2014) (i.e., $E(J - K) = 0.30 \pm 0.06$ mag and $A_K = 0.16 \pm 0.04$ mag assuming Nishiyama et al. (2009) extinction law). From the best model, the source brightness is $K_{s,s} = 17 \pm 1$ mag and the blend brightness is $K_{s,b} = 13.61 \pm 0.03$ mag. The relatively low precision on the source magnitude in K_s is again due to the lack of observations during the event high-magnification phase of the event. Unfortunately, the maximum observed magnification was only $A \sim 1.6$, while the secondary maximum observed magnification was $A \sim 1.05$. The color of the source is $(I_{OGLE_{IV}} - K_{VVV}) = 2 \pm 1$ mag, leading to an extinction corrected color of $(I_{OGLE_{IV}} - K_{VVV})_o = 2.0 - A_I + A_K = 1 \pm 1$ mag, and a magnitude of $K_{0,VVV} = 17 \pm 1$ mag. Using Bessell & Brett (1988), we found that this color is consistent with the optical colour and corresponds to a K-type source star.

4.3. Does the source belong to the Sagittarius Dwarf Galaxy?

³ <http://ogle.astrouw.edu.pl/>

⁴ <http://mill.astro.puc.cl/BEAM/calculator.php>




Figure 1. *Left:* Light curves and best-fit model for OGLE-2015-BLG-0232. *Right:* Central caustic (red curve), source trajectory (blue line), and source positions at the epoch of observations. The insets show zooms around the caustic crossings. There was one OGLE₁ measurement during the caustic entry and four CPTA₁ points (only two are visible in the inset) during the cusp exit. This allows a reasonable constraint on the normalised source radius ρ .

Parameters	pyLIMA ($s < 1$)	RT Model [†] ($s < 1$)	MCMC ($s < 1$)	pyLIMA ($s > 1$)	RT Model [†] ($s > 1$)	MCMC ($s > 1$)
$t_c - 2450000$	7087.20(1)	7087.49(4)	7087.2(2)	7086.68(4)	7086.93(4)	7086.76(8)
u_c	-0.00048(4)	0.00135(7)	-0.0005(6)	0.00320(8)	0.0020(2)	0.0026(5)
t_E	41.7(3)	46.1(3)	42(6)	34.7(1)	35(3)	39(3)
ρ (10^{-4})	9.9(3)	19.9(8)	10(1)	7.0(5)	7.0(2)	7.3(9)
s	0.545(2)	0.699(2)	0.55(7)	3.05(1)	2.58(2)	2.9(2)
q	0.0597(8)	0.0180(1)	0.06(2)	0.338(3)	0.17(1)	0.24(6)
α	-3.031(3)	-3.061(2)	-3.03(2)	3.017(4)	3.045(5)	3.008 (9)
χ^2	766	799	764	822	843	812

[†] The parameters are obtained from the online RTModel website (<http://www.fisica.unisa.it/gravitationAstrophysics/RTModel/2015/RTModel.htm>).

Table 3

Close/Wide best models of pyLIMA, RTModel and MCMC explorations. Models from RTModel were used as starting point for a Levenberg-Marquardt (LM) optimization with pyLIMA. Numbers in bracket in the table represents 1σ errors from LM and 68% range for the MCMC explorations.

Due to the relatively large galactic latitude of the event (i.e, $b = -5^\circ.9060$), the line of sight does not go through much of the Galactic Disk. This raises the possibility that the source is located in the stream of the Sagittarius Dwarf galaxy (Ibata et al. 1994). If this were the case, the source would be located very far away, $D_s \sim 25$ kpc. Cseresnjejs & Alard (2001) predicted that events due to the Sagittarius dwarf galaxy should represent roughly 1% of the total events detected toward the Galactic Bulge fields each year. They also predicted that these events should mainly occur for main-sequence source stars with $V \geq 21$ mag and that the median Einstein ring radius crossing time would be 1.3 times larger than the one observed for Milky Way sources. To test this, we constructed a map of the Sagittarius Dwarf galaxy in Figure 4. We followed the method of Majewski et al. (2003) and

selected stars with $E(B-V) < 0.555$, $0.95 < (J-K_s)_o < 1.10$ and $10.5 < K_{s,o} < 12$ combined with the extinction maps from Schlegel et al. (1998) (with a low resolution of 0.5 deg)⁵. However, the line of sight ($\ell = -1^\circ.17$, $b = -5^\circ.90$) is quite distant from the highest density of the Sagittarius Dwarf galaxy: M54. The Sagittarius dwarf star population has been studied in great detail, see for example Marconi et al. (1998); Monaco et al. (2002, 2004); Giuffrida et al. (2010). Several features can be used to distinguish stars from the Milky Way and the dwarf galaxy. In particular, the CMD of the dwarf galaxy presents several horizontal branches and red-giant branches, signatures of different star populations. The

⁵ We use the python implementation available at <https://github.com/gregreen/dustmaps>




Figure 2. Optical color-magnitude diagrams of stars within $2'$ of the line of sight of this event. OGLE and the transformed MOA are in blue (filled and empty respectively), the source is in red, the blend is in orange, and the position of the RGC is in magenta. The star symbol represents the star presented in the Table 4. The grey squares represent the region used to estimate the position of the RGC. Left: OGLE_{III} photometric system (i.e., Johnson-Cousins (Szymański et al. 2011)). Right: instrumental OGLE_{IV} photometric system.




Figure 3. Color magnitude diagram of stars within $2'$ of the line of sight to this event, using $I_{\text{OGLE}_{\text{IV}}}$ and K_{VVV} .

optical CMD of OGLE-2015-BLG-0232 does not show these signatures, indicating that there is no significant contamination from the dwarf galaxy.

Due to the large distance to the center of the Sagittarius Dwarf galaxy ($\geq 10^\circ$) and the absence of particular features in the CMD, we discount this hypothesis and assume that the source star belongs to the Milky Way.

5. INFORMATION ON THE BLEND

Results from our modeling indicate that this event was highly blended. It is clear from Figure 2 and Figure 3 that the blend belongs to the foreground stars branch of the CMD, indicating a close blend. In the following, we consider the blend as a single star and neglect the potential contamination from the source because the blend ratio is substantial with $g \sim 50$.




Figure 4. Map of the Sagittarius Dwarf galaxy from the 2MASS catalog (Cutri et al. 2003; Skrutskie et al. 2006).

5.1. Gaia measurements

The Gaia mission (Gaia Collaboration et al. 2016, 2018; Luri et al. 2018) recently released a vast catalog of parallax and proper motions measurements for more than a billion of stars. In addition to this goldmine, effective temperatures, radii and luminosities are also estimated. We summarized the Gaia measurements for OGLE-2015-BLG-0232 in Table 4. Recent studies indicate biases in Gaia parallax measurements of several μas (Lindgren et al. 2018; Zinn et al. 2018; Riess et al. 2018). We therefore use the estimation of the blend distance $D_b = 1023^{+86}_{-75}$ pc by Bailer-Jones et al. (2018), and so the blend is a late-type G or an early-type K dwarf. For this target, we also found $T_{\text{eff}} = 4707^{+269}_{-228}$ K, $R = 1.0^{+0.1}_{-0.1} R_\odot$, $\mathcal{L} = 0.42^{+0.07}_{-0.07} \mathcal{L}_\odot$ and ultimately estimated the mass of this blend $M \sim \mathcal{L}^{1/4} \sim 0.8 M_\odot$, typical of a K-dwarf. However, Andrae et al. (2018) note that these parameters are estimated by neglecting the extinction toward the target. While this ap-

proximation is reasonable for this target because the blend is relatively close and the extinction along the line of sight is relatively small, these fundamental parameters are probably biased.

The brightnesses of the blend in the Gaia bands are $G = 15.918 \pm 0.001$ mag, $G_{BP} = 16.48 \pm 0.01$ mag and $G_{RP} = 15.15 \pm 0.01$ mag. Using the system transformation in the Appendix, we convert these magnitudes to the Johnson-Cousins system to find $V = 16.25 \pm 0.05$ mag and $I = 15.08 \pm 0.05$ mag. Given the blend distance, we assumed half extinction and found an intrinsic color $(V-I)_{o,b,G} = 1.33 - 0.69/2 = 1.0 \pm 0.1$ and brightness $I_{o,b,G} = 15.08 - 0.79/2 = 14.7 \pm 0.1$ mag, typical of a K2 dwarf star (Bessell & Brett 1988). Using the color-effective temperature relation of Casagrande et al. (2010), the blend effective temperature is $T_{\text{eff}} = 4900 \pm 400$ K. We estimated the blend physical radius $R_b = 0.8 \pm 0.1 R_{\odot}$, the luminosity $\mathcal{L} = 0.3 \pm 0.1 \mathcal{L}_{\odot}$ and finally derived the blend mass $M_b \sim 0.7 M_{\odot}$ (Boyajian et al. 2012). Knowing that the angular radius of the blend is $\theta_b = 4.8 \pm 0.5 \mu\text{as}$ (Kervella & Fouqué 2008), one can derive an independent estimate of the blend distance $D_b = 800 \pm 200$ pc, in good agreement with the Gaia parallax measurement.

If the blend were the lens and assuming that the source is at 8 kpc, a blend mass of $M_b \sim 0.7 M_{\odot}$ at a distance $D_b \sim 1000$ pc, the angular Einstein ring would be $\theta_{E,b} \sim 2.2$ mas. This is in strong disagreement with the value of $\theta_E = 0.8 \pm 0.2$ mas derived in Section 4.1. This is a first clue that the bright blend is likely not the lens.

From Table 4, the proper motion of the bright blend is $\mu_G(N, E) = (2.2 \pm 0.1, 9.9 \pm 0.1)$ mas/yr. The speed of the Sun in the Galactic frame is $\mathbf{V}_{\odot}(U, V, W) \sim (11, 12, 7) + (0, 220, 0)$ km/s (Fich et al. 1989; Schönrich et al. 2010): the first term is the intrinsic Sun velocity and the second term is the speed of the Galactic disk in the Galactic coordinates system. Assuming the source is at 8 kpc, the expected proper motion of the source is about $\mu_s(l, b) \sim (-6 \pm 3, 0 \pm 3)$ mas/yr, see Kuijken & Rich (2002) and Kozłowski et al. (2006) for the estimation of the uncertainties. The Galactic proper motion transform to $\mu_s(\alpha, \delta) \sim (-3 \pm 3, -5 \pm 3)$ mas/yr (Binney & Merrifield 1998; Poleski 2013; Bachelet et al. 2018). Therefore, if the bright blend were the lens, one would expect a relative proper motion of $\mu_{\text{rel}}(N, E) = \mu_s - \mu_G \sim (-5 \pm 3, -15 \pm 3)$ mas/yr. The relative proper motion would be $\mu_{\text{rel}} = 16 \pm 4$ mas/yr, in disagreement with the estimation $\mu_{\text{rel}} = 7 \pm 3$ mas/yr of the Section 4.1. This is the second clue that the blend is not the lens.

5.2. Blend brightness from models

Using our best-fit model and the color relationships given in the Appendix, we derived the brightnesses of the blend: $I_{b,\text{OGLEIV}} = 15.1163 \pm 0.0007$ mag, $V_{b,\text{OGLEIV}} = 16.23 \pm 0.08$ mag and $K_{b,\text{VVV}} = 13.61 \pm 0.03$ mag. Assuming that the blend suffers half the extinction, we found that the blend brightness is $I_{o,b,\text{OGLEIV}} = 14.7 \pm 0.1$ mag and the blend color is $(I_{\text{OGLEIV}} - K_{\text{VVV}})_{o,b} = 1.1 \pm 0.1$ mag, consistent with its being an early K-dwarf (Bessell & Brett 1988). This is in good agreement with the Gaia measurements.

5.3. Astrometry

Toward the Galactic Bulge and for stellar masses, microlensing occurs when the alignment between the lens and the source is less than a few mas. We therefore compared the position of centroids between the baseline object and the




Figure 5. Location of the target from the catalogs listed in the Table 4, centered at (RA = 271.6826°, DEC = -32.9076°) (J2000). North is up and East is left. Some uncertainties have been hidden for clarity. The source position measured from OGLE-IV is located at $\Delta(E, N) = (78, 78)$ mas (red square) from the OGLE-IV position. The grey cross indicates the prediction of the source, applying the offset to the Gaia position and assuming $\sigma(N, E) = (45, 35)$ mas uncertainties.

magnified source from the OGLE-IV images. In pixel coordinates, the magnified source has an offset of $\Delta(N, E) = (78 \pm 45, 78 \pm 35)$ mas from the bright blend centroid (the precision of the bright blend centroid is about 0.05 pixel, i.e. $\sigma(N, E) = (13, 13)$ mas). The two positions are different enough (i.e., 1.5σ) to assume that this is the third clue that the blend is likely not the lens.

5.4. The lens as a blend companion

In the following, we explore the possibility that the lens is a companion of the blend. From the astrometry offset derived in Section 5.3, we can derive the separation δ of the blend with its potential companion and found $\delta = 110$ mas, which corresponds to $a_{\text{proj}} \sim 110$ au at 1 kpc. If this potential companion is indeed a component of the lens system, then the mass ratio between the binary blend components is $q_b = (\theta_E/\theta_{E,b})^2 = (0.8/2.2)^2 \sim 0.13$, leading to a potential companion mass of $M_{b,2} = 0.13 \times 0.7 \sim 0.1 M_{\odot}$. Therefore, such a companion is not bright enough to have been significantly detected. Because the normalised separation between the putative companion and the bright blend is important $s_b = 110/2.2 \sim 50$, the hypothetical companion blend could have acted as a binary lens and left no signature of a triple-lens, as observed. However, this hypothetical companion would have a similar proper motion as the bright blend and the analysis on the relative proper motions in Section 5.1 also apply here. Therefore, the lens as a blend companion hypothesis is unlikely.

6. DISCUSSION AND POTENTIAL NEW CLUES

All available information seem to concur that the blend light is mainly due to a close K dwarf. Both astrometry and the constraint from finite-source effects reject the hypothesis that the bright blend is the lens. The light of the lens is not significantly detected and there are no constraints from the microlensing parallax: the distance and exact nature of the lens remains uncertain at the present time. However, considering a large mass range for the lens primary $M_{l,1} \in [0.1, 2.0] M_{\odot}$ (corresponding to $D_l \leq 5.5$ kpc according to Equation 4 and $\theta_E = 0.8 \pm 0.2$ mas), the companion mass range is $M_{l,2} \in [6, 130] M_{Jup}$. The lens companion is therefore a massive planet, a brown dwarf

Catalog	Source ID	Epoch	RA(J2000) °	DEC(J2000) °	Parallax mas	μ_α mas/yr	μ_δ mas/yr
Gaia	4042761215742767360	J2015.5	271.68268633(1)	-32.90760309(1)	0.96(7)	9.9(1)	2.0(1)
MOA	965	-	271.68263(4)	-32.90764(3)	-	-	-
OGLE-III	90793	J2002.46	271.68267(4)	-32.90761(3)	-	-	-
OGLE-IV (baseline)	58780	J2011.4	271.68267(4)	-32.90758(3)	-	-	-
OGLE-IV (source)	-	J2011.4	271.68269(4)	-32.90720(3)	-	-	-
VVV	2508	J2010	271.68262	-32.90763	-	-	-
PPMXL	4938889137283654706	J1991.21	271.68268(2)	-32.90760(2)	-	10.6(5.2)	6.3(5.2)

Table 4

Astrometry of the target in MOA, OGLE-III, OGLE-IV, VVV and Gaia catalogs. OGLE-III catalog of the field is from Szymański et al. (2011). OGLE-IV catalog is available online ^a. VVV catalog is from Beaulieu et al. (2016). The PPMXL catalog is from Roeser et al. (2010). Numbers in parenthesis are the 1σ uncertainties.

^a<http://ogle.astrouw.edu.pl/ogle4/ews/2018/ews.html>

or a low-mass M-dwarf if $D_l \leq 5.5$ kpc. If the lens is more distant, the primary is probably a stellar remnant, otherwise the lens light would have been detected. This indicates the need for supplementary observations to reveal the nature of the lens OGLE-2015-BLG-0232.

High-resolution imaging is an important tool for microlensing. Several planets have been confirmed using space or ground-based facilities and had their measured properties refined, see for example (Batista et al. 2015; Bennett et al. 2015; Beaulieu et al. 2017). High-resolution imaging is useful for two reasons. First, it is possible to estimate the source-lens proper motion μ from high-resolution images obtained several years after the microlensing event, when the source and the lens are well separated (Batista et al. 2015). High-resolution imaging can also provide measurements of source and, sometimes, lens fluxes and therefore tightly constrain the mass-distance relation of the lens (Ranc et al. 2015; Batista et al. 2015; Bennett et al. 2015; Beaulieu et al. 2017).

In the case of OGLE-2015-BLG-0232, high-resolution imaging will contribute to confirming/rejecting scenarios and possibly estimate the mass of the lens. The first step will be to challenge the assumption that the blend is a single star. This can be done immediately. Moreover, one can predict a more precise source position based on Gaia astrometry and the measured offset from the OGLE-IV photometry. The predicted position of the source is shown in the Figure 5, assuming 26 mas precision on OGLE-IV measurement (i.e., 0.1 pixel). The comparison of the flux at this position in high resolution images with the measured source fluxes from models could place constraints on the nature of the lens.

A second step will be to wait several years for the bright blend leaves the line of sight to obtain more information on the source/lens system. Because $\mu_b = 11 \pm 0.2$ mas/yr, the blend is separating faster than the lens/source system $\mu = 7.0 \pm 3$ mas/yr. In a decade, the blend should be about 11 pixels away from the line of sight while the source and the lens separation should be about 7 pixel (for a typical high-resolution pixel scale of 10 mas/pix).

Low-resolution spectroscopy could also confirm the spectral type of the bright blend. Similarly, the study of emission/absorption lines with high-resolution spectroscopy would allow a precise understanding of the blend. Finally, one could combine spectroscopic and photometric information to explore various scenarios in a Bayesian analysis (Santerne et al. 2016).

7. CONCLUSION

We presented an analysis of the binary microlensing event OGLE-2015-BLG-0232. Because the event occurred during full moon, the observations do not constrain much the deviations from the single-lens model. However, results from the modeling favor a close brown dwarf companion (i.e., $s \sim 0.55$ and $q \sim 0.06$). The source is estimated to be red and faint, probably a K dwarf in the Galactic Bulge. We also tested, and ultimately rejected, the hypothesis that the source belongs to the Sagittarius Dwarf Galaxy. Since the microlensing paral-

lax is not measured, we obtain only one (weak) constraint, from finite-source effects, on the mass and distance of the lens. Based on the recent Gaia DR2 release and OGLE-IV astrometry, we were able to infer that the bright blend is a K dwarf at 1 kpc and is most likely not the lens. We finally discuss the potential of additional observations to confirm the nature of the blend and ultimately to derive the exact nature of the lens.

ACKNOWLEDGEMENTS

The authors thank the anonymous referee for the constructive comments. This research has made use of NASA's Astrophysics Data System. Work by EB and RAS is supported by the NASA grant NNX15AC97G. Work by C. Han was supported by the grant (2017R1A4A1015178) of National Research Foundation of Korea. This work makes use of observations from the LCOGT network. The OGLE project has received funding from the National Science Centre, Poland, grant MAESTRO 2014/14/A/ST9/00121 to AU. The MOA project is supported by JSPS KAKENHI Grant Number JSPS24253004, JSPS26247023, JSPS23340064, JSPS15H00781, JP16H06287 and JP17H02871. The work by C.R. was supported by an appointment to the NASA Postdoctoral Program at the Goddard Space Flight Center, administered by USRA through a contract with NASA. This work has made use of data from the European Space Agency (ESA) mission *Gaia* (<https://www.cosmos.esa.int/Gaia>), processed by the *Gaia* Data Processing and Analysis Consortium (DPAC, <https://www.cosmos.esa.int/web/Gaia/dpac/consortium>). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the *Gaia* Multilateral Agreement. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. This research made use of Astropy, a community-developed core Python package for Astronomy (Astropy Collaboration, 2013). This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France. The DENIS project has been partly funded by the SCIENCE and the HCM plans of the European Commission under grants CT920791 and CT940627. It is supported by INSU, MEN and CNRS in France, by the State of Baden-Württemberg in Germany, by DGICYT in Spain, by CNR in Italy, by FFWF-BWF in Austria, by FAPESP in Brazil, by OTKA grants F-4239 and F-013990 in Hungary, and by the ESO C&EE grant A-04-046. Jean Claude Renault from IAP was the Project manager. Observations were carried out thanks to the contribution of numerous students and young scientists from all involved institutes, under the supervision of P. Fouqué, survey astronomer resident in Chile. DPB, AB, and CR were supported by NASA through grant NASA-80NSSC18K0274.

APPENDIX

COLOR TRANSFORMATIONS

In this work, we used several color transformations that we summarize here. First, we calibrated the MOA instrumental magnitudes to the OGLE-III catalog (Udalski 2003; Bond et al. 2017) using the relationships :

$$I_{\text{OGLE-III}} = R_{\text{MOA}} + (27.935 \pm 0.003) + (-0.244 \pm 0.003)(V_{\text{MOA}} - R_{\text{MOA}}) \pm 0.08 \quad (\text{A1})$$

$$V_{\text{OGLEIII}} = V_{\text{MOA}} + (28.556 \pm 0.002) + (-0.164 \pm 0.002)(V_{\text{MOA}} - R_{\text{MOA}}) \pm 0.08 \quad (\text{A2})$$

We also calibrated the MOA instrumental magnitudes to the OGLE-IV system using:

$$I_{\text{OGLEIV}} = R_{\text{MOA}} + (27.990 \pm 0.003) + (-0.247 \pm 0.009)(V_{\text{MOA}} - R_{\text{MOA}}) \pm 0.08 \quad (\text{A3})$$

$$V_{\text{OGLEIV}} = V_{\text{MOA}} + (28.425 \pm 0.005) + (-0.062 \pm 0.006)(V_{\text{MOA}} - R_{\text{MOA}}) \pm 0.08 \quad (\text{A4})$$

We also used the transformation of the 2MASS colors into the the VVV system (Soto et al. 2013):

$$J_{\text{VVV}} = J_{2\text{MASS}} - 0.077(J_{2\text{MASS}} - H_{2\text{MASS}}) \quad (\text{A5})$$

$$H_{\text{VVV}} = H_{2\text{MASS}} + 0.032(J_{2\text{MASS}} - H_{2\text{MASS}}) \quad (\text{A6})$$

$$K_{\text{VVV}} = K_{2\text{MASS}} + 0.010(J_{2\text{MASS}} - K_{2\text{MASS}}) \quad (\text{A7})$$

Transformations into the Bessell & Brett photometric system (Bessell & Brett 1988) are the revised version ⁶ of Carpenter (2001):

$$(K_s)_{2\text{MASS}} = K_{\text{BB}} + (-0.039 \pm 0.007) + (0.001 \pm 0.005)(J - K)_{\text{BB}} \quad (\text{A8})$$

$$(J - K_s)_{2\text{MASS}} = (-0.018 \pm 0.007) + (0.001 \pm 0.005)(J)_{\text{BB}} \quad (\text{A9})$$

Finally, the transformation of the Gaia DR2 to the Johnson-Cousins system is available online ⁷:

$$G - V_{\text{JC}} = -0.01760 - 0.006860(G_{\text{BP}} - G_{\text{RP}}) - 0.1732(G_{\text{BP}} - G_{\text{RP}})^2 \pm 0.045858 \quad (\text{A10})$$

$$G - I_{\text{JC}} = 0.02085 + 0.7419(G_{\text{BP}} - G_{\text{RP}}) - 0.096311(G_{\text{BP}} - G_{\text{RP}})^2 \pm 0.04956 \quad (\text{A11})$$

REFERENCES

- Albrow, M. D., Beaulieu, J.-P., Caldwell, J. A. R., et al. 2000, *ApJ*, 535, 176
 Albrow, M. D., Horne, K., Bramich, D. M., et al. 2009, *MNRAS*, 397, 2099
 Andrae, R., Fouesneau, M., Creevey, O., et al. 2018, *A&A*, 616, A8
 Andrae, R., Schulze-Hartung, T., & Melchior, P. 2010, *ArXiv e-prints*
 Bachelet, E., Beaulieu, J.-P., Boisse, I., Santerne, A., & Street, R. A. 2018, *ArXiv e-prints*
 Bachelet, E., Bramich, D. M., Han, C., et al. 2015, *ApJ*, 812, 136
 Bachelet, E., Fouqué, P., Han, C., et al. 2012a, *A&A*, 547, A55
 Bachelet, E., Norbury, M., Bozza, V., & Street, R. 2017, *AJ*, 154, 203
 Bachelet, E., Shin, I.-G., Han, C., et al. 2012b, *ApJ*, 754, 73
 Bailer-Jones, C. A. L., Rybizki, J., Fouesneau, M., Mantelet, G., & Andrae, R. 2018, *AJ*, 156, 58
 Batista, V., Beaulieu, J.-P., Bennett, D. P., et al. 2015, *ApJ*, 808, 170
 Beaulieu, J.-P., Batista, V., Bennett, D. P., et al. 2017, *ArXiv e-prints*
 Beaulieu, J.-P., Bennett, D. P., Batista, V., et al. 2016, *ApJ*, 824, 83
 Bennett, D. P., Bhattacharya, A., Anderson, J., et al. 2015, *ApJ*, 808, 169
 Bessell, M. S. & Brett, J. M. 1988, *PASP*, 100, 1134
 Binney, J. & Merrifield, M. 1998, *Galactic Astronomy*
 Bond, I. A., Abe, F., Dodd, R. J., et al. 2001, *MNRAS*, 327, 868
 Bond, I. A., Bennett, D. P., Sumi, T., et al. 2017, *MNRAS*, 469, 2434
 Bonfils, X., Delfosse, X., Udry, S., et al. 2013, *A&A*, 549, A109
 Boyajian, T. S., von Braun, K., van Belle, G., et al. 2012, *ApJ*, 757, 112
 Bozza, V. 2000, *A&A*, 359, 1
 Bozza, V. 2010, *MNRAS*, 408, 2188
 Bozza, V., Dominik, M., Rattenbury, N. J., et al. 2012, *MNRAS*, 424, 902
 Bramich, D. M. 2008, *MNRAS*, 386, L77
 Bramich, D. M., Horne, K., Albrow, M. D., et al. 2013, *MNRAS*, 428, 2275
 Bryan, M. L., Benneke, B., Knutson, H. A., Batygin, K., & Bowler, B. P. 2018, *Nature Astronomy*, 2, 138
 Buchhave, L. A., Latham, D. W., Johansen, A., et al. 2012, *Nature*, 486, 375
 Burrows, A., Hubbard, W. B., Lunine, J. I., & Liebert, J. 2001, *Reviews of Modern Physics*, 73, 719
 Burrows, A. & Liebert, J. 1993, *Reviews of Modern Physics*, 65, 301
 Carpenter, J. M. 2001, *AJ*, 121, 2851
 Casagrande, L., Ramírez, I., Meléndez, J., Bessell, M., & Asplund, M. 2010, *A&A*, 512, A54
 Cassan, A. 2008, *A&A*, 491, 587
 Cassan, A., Kubas, D., Beaulieu, J.-P., et al. 2012, *Nature*, 481, 167
 Chung, S.-J., Zhu, W., Udalski, A., et al. 2017, *ApJ*, 838, 154
 Clanton, C. & Gaudi, B. S. 2016, *ApJ*, 819, 125
 Cseresnes, P. & Alard, C. 2001, *A&A*, 369, 778
 Cutri, R. M., Skrutskie, M. F., van Dyk, S., et al. 2003, *VizieR Online Data Catalog*, 2246
 Dominik, M. 1998, *A&A*, 329, 361
 Dominik, M. 1999, *A&A*, 349, 108
 Dominik, M. 2009, *MNRAS*, 393, 816
 Dominik, M., Bachelet, E., Bozza, V., et al. 2018, *ArXiv e-prints*
 Dominik, M., Rattenbury, N. J., Allan, A., et al. 2007, *MNRAS*, 380, 792
 Dong, S., DePoy, D. L., Gaudi, B. S., et al. 2006, *ApJ*, 642, 842
 Erdl, H. & Schneider, P. 1993, *A&A*, 268, 453
 Fich, M., Blitz, L., & Stark, A. A. 1989, *ApJ*, 342, 272
 Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, *PASP*, 125, 306
 Fressin, F., Torres, G., Charbonneau, D., et al. 2013, *ApJ*, 766, 81
 Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, *ArXiv e-prints*
 Gaia Collaboration, Prusti, T., de Bruijne, J. H. J., et al. 2016, *A&A*, 595, A1
 Giuffrida, G., Sbordone, L., Zaggia, S., et al. 2010, *A&A*, 513, A62
 Gonzalez, O. A., Rejkuba, M., Zoccali, M., et al. 2012, *A&A*, 543, A13
 Gould, A. 2000, *ApJ*, 542, 785
 Gould, A. 2004, *ApJ*, 606, 319
 Gould, A. & Loeb, A. 1992, *ApJ*, 396, 104
 Griest, K. & Safizadeh, N. 1998, *ApJ*, 500, 37
 Han, C. 2009, *ApJ*, 691, L9
 Han, C., Jung, Y. K., Udalski, A., et al. 2016, *ApJ*, 822, 75
 Hundertmark, M., Street, R. A., Tsapras, Y., et al. 2017, *ArXiv e-prints*
 Ibata, R. A., Gilmore, G., & Irwin, M. J. 1994, *Nature*, 370, 194
 Kervella, P. & Fouqué, P. 2008, *A&A*, 491, 855
 Kozłowski, S., Woźniak, P. R., Mao, S., et al. 2006, *MNRAS*, 370, 435
 Kuijken, K. & Rich, R. M. 2002, *AJ*, 124, 2054
 Latham, D. W., Stefanik, R. P., Torres, G., et al. 2002, *AJ*, 124, 1144
 Liebig, C., D'Agó, G., Bozza, V., & Dominik, M. 2015, *MNRAS*, 450, 1565
 Lindegren, L., Hernández, J., Bombrun, A., et al. 2018, *A&A*, 616, A2
 Luri, X., Brown, A. G. A., Sarro, L. M., et al. 2018, *ArXiv e-prints*
 Majewski, S. R., Skrutskie, M. F., Weinberg, M. D., & Ostheimer, J. C. 2003, *ApJ*, 599, 1082
 Mao, S. & Di Stefano, R. 1995, *ApJ*, 440, 22
 Marconi, G., Buonanno, R., Castellani, M., et al. 1998, *A&A*, 330, 453
 Minniti, D., Lucas, P. W., Emerson, J. P., et al. 2010, *New A*, 15, 433
 Miyake, N., Udalski, A., Sumi, T., et al. 2012, *ApJ*, 752, 82
 Monaco, L., Bellazzini, M., Ferraro, F. R., & Pancino, E. 2004, *MNRAS*, 353, 874
 Monaco, L., Ferraro, F. R., Bellazzini, M., & Pancino, E. 2002, *ApJ*, 578, L47
 Mordasini, C., Alibert, Y., Benz, W., Klahr, H., & Henning, T. 2012, *A&A*, 541, A97
 Mortier, A., Santos, N. C., Sozzetti, A., et al. 2012, *A&A*, 543, A45
 Mróz, P., Han, C., and, et al. 2017, *AJ*, 153, 143
 Nataf, D. M., Gould, A., Fouqué, P., et al. 2013, *ApJ*, 769, 88

⁶ <https://www.ipac.caltech.edu/2mass/releases/allsky/doc/sec6.4b.html>

⁷ <https://gea.esac.esa.int/archive/documentation/GDR2/>

- Nishiyama, S., Tamura, M., Hatano, H., et al. 2009, *ApJ*, 696, 1407
- Penny, M. T. 2014, *ApJ*, 790, 142
- Poleski, R. 2013, *ArXiv e-prints*
- Poleski, R., Udalski, A., Bond, I. A., et al. 2017, *A&A*, 604, A103
- Ranc, C., Cassan, A., Albrow, M. D., et al. 2015, *A&A*, 580, A125
- Riess, A. G., Casertano, S., Yuan, W., et al. 2018, *ApJ*, 861, 126
- Roeser, S., Demleitner, M., & Schilbach, E. 2010, *AJ*, 139, 2440
- Santerne, A., Beaulieu, J.-P., Rojas Ayala, B., et al. 2016, *A&A*, 595, L11
- Schlaufman, K. C. 2018, *ApJ*, 853, 37
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Schneider, J., Dedieu, C., Le Sidaner, P., Savalle, R., & Zolotukhin, I. 2011, *A&A*, 532, A79
- Schönrich, R., Binney, J., & Dehnen, W. 2010, *MNRAS*, 403, 1829
- Schultheis, M., Chen, B. Q., Jiang, B. W., et al. 2014, *A&A*, 566, A120
- Shvartzvald, Y., Yee, J. C., Skowron, J., et al. 2018, *ArXiv e-prints*
- Skowron, J., Udalski, A., Kozłowski, S., et al. 2016, *Acta Astron.*, 66, 1
- Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, *AJ*, 131, 1163
- Soto, M., Barbá, R., Gunthardt, G., et al. 2013, *A&A*, 552, A101
- Spiegel, D. S., Burrows, A., & Milsom, J. A. 2011, *ApJ*, 727, 57
- Storn, R. & Price, K. 1997, *J. of Global Optimization*, 11, 341
- Suzuki, D., Bennett, D. P., Sumi, T., et al. 2016, *ApJ*, 833, 145
- Szymański, M. K., Udalski, A., Soszyński, I., et al. 2011, *Acta Astron.*, 61, 83
- Tsapras, Y., Street, R., Horne, K., et al. 2009, *Astronomische Nachrichten*, 330, 4
- Udalski, A. 2003, *Acta Astron.*, 53, 291
- Udalski, A., Szymański, M. K., & Szymański, G. 2015, *Acta Astron.*, 65, 1
- Wright, J. T., Fakhouri, O., Marcy, G. W., et al. 2011, *PASP*, 123, 412
- Yee, J. C., Hung, L.-W., Bond, I. A., et al. 2013, *ApJ*, 769, 77
- Zapolsky, H. S. & Salpeter, E. E. 1969, *ApJ*, 158, 809
- Zhu, W., Calchi Novati, S., Gould, A., et al. 2016, *ApJ*, 825, 60
- Zinn, J. C., Pinsonneault, M. H., Huber, D., & Stello, D. 2018, *ArXiv e-prints*