



## LJMU Research Online

Sun, M-Y, Wang, S-J, Li, X-Q, Shen, Y-L, Lu, J-R, Tian, X-H, Rahman, K, Zhang, L-J, Nian, H and Zhang, H

**CXCL6 Promotes Renal Interstitial Fibrosis in Diabetic Nephropathy by Activating JAK/STAT3 Signaling Pathway**

<http://researchonline.ljmu.ac.uk/id/eprint/10554/>

### Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Sun, M-Y, Wang, S-J, Li, X-Q, Shen, Y-L, Lu, J-R, Tian, X-H, Rahman, K, Zhang, L-J, Nian, H and Zhang, H (2019) CXCL6 Promotes Renal Interstitial Fibrosis in Diabetic Nephropathy by Activating JAK/STAT3 Signaling Pathway. *Frontiers in Pharmacology*. 10. ISSN 1663-9812**

LJMU has developed [LJMU Research Online](http://researchonline.ljmu.ac.uk/) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)

<http://researchonline.ljmu.ac.uk/>



# CXCL6 Promotes Renal Interstitial Fibrosis in Diabetic Nephropathy by Activating JAK/STAT3 Signaling Pathway

Meng-Yao Sun<sup>1,2†</sup>, Su-Juan Wang<sup>1,3†</sup>, Xiao-Qin Li<sup>1</sup>, Yu-Li Shen<sup>1</sup>, Jian-Rao Lu<sup>4</sup>, Xin-Hui Tian<sup>1</sup>, Khalid Rahman<sup>5</sup>, Li-Jun Zhang<sup>1\*</sup>, Hua Nian<sup>6\*</sup> and Hong Zhang<sup>1\*</sup>

<sup>1</sup> Institute of Interdisciplinary Integrative Medicine Research, Shanghai University of Traditional Chinese Medicine, Shanghai, China, <sup>2</sup> Central Laboratory, Putuo Hospital, Shanghai University of Traditional Chinese Medicine, Shanghai, China, <sup>3</sup> Department of Drug Preparation, Hospital of TCM and Hui Nationality Medicine, Ningxia Medical University, Wuzhong, China, <sup>4</sup> Department of Nephrology, Seventh People's Hospital, Shanghai University of Traditional Chinese Medicine, Shanghai, China, <sup>5</sup> School of Pharmacy and Biomolecular Sciences, Faculty of Science, Liverpool John Moores University, Liverpool, United Kingdom, <sup>6</sup> Department of Pharmacy, Yueyang Hospital of Integrated Traditional Chinese and Western Medicine, Shanghai University of Traditional Chinese Medicine, Shanghai, China

## OPEN ACCESS

### Edited by:

Matthew Griffin,  
National University of Ireland Galway,  
Ireland

### Reviewed by:

Peter Jon Nelson,  
Ludwig Maximilian University  
of Munich, Germany  
Ana Belen Sanz,  
Instituto de Investigación Sanitaria  
Fundación Jiménez Díaz, Spain

### \*Correspondence:

Li-Jun Zhang  
zhanglijun0407@163.com  
Hua Nian  
jackynian@126.com  
Hong Zhang  
hqzhang51@126.com

† These authors have contributed  
equally to this work

### Specialty section:

This article was submitted to  
Renal Pharmacology,  
a section of the journal  
Frontiers in Pharmacology

**Received:** 25 September 2018

**Accepted:** 22 February 2019

**Published:** 25 March 2019

### Citation:

Sun M-Y, Wang S-J, Li X-Q,  
Shen Y-L, Lu J-R, Tian X-H,  
Rahman K, Zhang L-J, Nian H and  
Zhang H (2019) CXCL6 Promotes  
Renal Interstitial Fibrosis in Diabetic  
Nephropathy by Activating  
JAK/STAT3 Signaling Pathway.  
*Front. Pharmacol.* 10:224.  
doi: 10.3389/fphar.2019.00224

In this study the role of CXCL6 in diabetic nephropathy (DN) was investigated. It was found to be overexpression in DN patients and DN rat model. And the expression of fibrosis-related cytokines was consistent with the expression of CXCL6. High glucose significantly increased the proliferation of rat renal fibroblasts NRK-49F cell and the expression of CXCL6. Knockdown of CXCL6 ameliorated the pro-proliferation effect of high glucose and decreased the expression of fibrosis-related cytokines, while CXCL6 overexpression exhibited the opposite phenomenon. Gene set enrichment analysis, Western blot and ELISA showed that Janus kinase-signal transducer and activator of transcription (JAK-STAT) and CYTOKINE\_CYTOKINE\_RECEPTOR\_INTERACTION signaling pathways were correlative with CXCL6. This data indicates that CXCL6 may promote fibrosis-related factors to accelerate the development of DN renal interstitial fibrosis by activating JAK/STAT3 signaling pathway. CXCL6 is promising to be a potential novel therapeutic target and candidate biomarker for JAK/STAT3 signaling for the treatment of DN.

**Keywords:** CXCL6, kidney, diabetics, fibrosis, pathway

## INTRODUCTION

Diabetic nephropathy (DN) is a well-known micro-vascular complication of diabetes, and the leading cause of ESRD (Gin et al., 2000). Over-deposition of the ECM in the renal interstitium is the main cause of renal interstitial fibrosis, which is the final common pathway of the majority of chronic progressive renal diseases, including DN (Strutz and Zeisberg, 2006). Fibroblasts are

**Abbreviations:** CCK-8, Counting Kit-8; CXCL6, C-X-C chemokine ligand 6; DMEM, Dulbecco's modified Eagle medium; DMSO, Dimethyl sulfoxide; DN, Diabetic Nephropathy; ECM, Extracellular matrix; ESRD, End-stage renal disease; FBS, Fetal bovine serum; GSEA, Gene Set Enrichment Analysis; JAK, Janus kinase; MMP, Matrix metalloproteinase; STAT, Signal transducer and activator of transcription; TGF- $\beta$ 1, Transforming Growth Factor  $\beta$ 1; TNF- $\alpha$ , Tumor Necrosis Factor  $\alpha$ .

the major cells that produce ECM in the renal interstitium and play an important role in the process of renal interstitial fibrosis (Grande and López-Novoa, 2009).

Many inflammatory factors are involved in the process of renal interstitial fibrosis in DN, such as chemokine, TGF- $\beta_1$ , interleukin family, adhesion molecule, etc., which can stimulate a series of inflammatory signaling pathways, resulting in ECM thickening, interstitial fibrosis, eventually triggering renal dysfunction (Wada and Makino, 2013).

Chemokines are a class of chemotactic and inducible small molecule peptides that are ubiquitous and play an important role in acute and chronic inflammation (Wasmuth et al., 2009). Among the chemokines, Soluble CXCL6 is an important inflammatory cytokine that recruits inflammatory cells to the site of inflammation by binding to the receptors CXCR1 and CXCR2 (Sadik et al., 2011). In the cigarette-induced CXCL6-deficient mouse inflammation model, macrophage recruitment in the lungs was significantly reduced (Balamayooran et al., 2012). CXCL6 is not only significantly increased in inflammatory bowel disease, but is also higher expressed in patients with periodontitis. Immunohistochemistry has confirmed that CXCL6 is mainly distributed in gingival vascular endothelial cells (Kebschull et al., 2009). Ionizing radiation can induce a significant inflammatory reaction in the skin, this can also promote the secretion of CXCL6 in fetal foreskin and adult human dermal fibroblasts, and mast cells mediate the secretion of CXCL6 (Müller and Meineke, 2011). Microparticles, a small vesicle that accumulates in the synovial fluid of patients with rheumatoid arthritis, not only promotes the secretion of inflammatory cytokines by synovial fibroblasts but also significantly induces the secretion of the chemokine CXCL6, thereby increasing the number of vascular endothelial cell migration and thus promoting angiogenesis and aggravation of joint inflammation (Reich et al., 2011). It has also been reported that CXCL6 can also promote lung fibrosis (Besnard et al., 2013) and it is significantly increased in patients with idiopathic pulmonary fibrosis.

Anti-CXCL6 treatment of 2 weeks is reported to significantly decrease the levels of IL-1 $\beta$ , collagen and Human Tissue Inhibitor Of Matrix Metalloprotease-1 (TIMP-1) in Bleomycin-induced lung inflammation and fibrosis mouse model together with the degree of lung inflammation and fibrosis suggesting that CXCL6 plays an important role in the process of fibrosis, however, the mechanism of action is not clearly defined (Besnard et al., 2013). The aim of this study was to investigate the potential functions and the underlying mechanism of CXCL6 in DN renal interstitial fibrosis.

## MATERIALS AND METHODS

### Patient Samples

All the tissue and plasma samples were obtained from The Seventh People's Hospital of Shanghai University of Traditional Chinese Medicine between February 2015 and December 2015. 40 DN and 20 normal kidney tissue samples were harvested from DN patients with kidney interstitial fibrosis at the time of biopsy or surgical resection. Plasma samples were collected from DN

patients ( $n = 40$ ) with kidney interstitial fibrosis and apparently healthy subjects ( $n = 40$ ) and the tissue and plasma samples were stored at  $-80^\circ\text{C}$  until used. The following data was recorded for all participants: sex, age, and risk factors for DN (hyperlipidemia, and glycosuria), showed in **Supplementary Table S1**.

This study was approved by the Medical Ethics Committee of The Seventh People's Hospital of Shanghai University of Traditional Chinese Medicine and written informed consent was obtained from all participants involved in this study.

### Bioinformatics Analysis

CXCL6 mRNA levels detected in human DN kidney samples ( $n = 19$ ) and healthy kidney tissue ( $n = 50$ ) were obtained from GEO DataSets (Access ID: GSE30122). Student's *t*-test was used to determine the statistical significance of CXCL6 expression between human diabetic kidney samples and healthy kidney tissue.

Gene Set Enrichment Analysis gene expression profiles about fibrosis was obtained from GEO DataSets (Access ID: GSE28221) was analyzed by GSEA. GSEA was performed using the GSEA software, Version 3.0, obtained from the Broad Institute<sup>1</sup>. GSEA version 3.0 software was from the Broad Institute at MIT. The nominal *P*-value and normalized enrichment score (NES) was used to sort the pathways associated with CXCL6-higher expression.

### Animal Model Preparation

Male SD rats (160–200 g) were obtained from Shanghai Sippr BK Laboratory Animals Ltd, (Shanghai, China). After feeding high-sugar-fat diet for 8 weeks, the rats were given a single intraperitoneal injection of 35 mg/kg STZ. After 72 h, the rats with fasting and random blood glucose reading of above 13.8 mmol/L, 16.7 mmol/L, respectively, meanwhile glycosuria showing ++++ were considered as diabetes. Four weeks after STZ injection, the DN model rat was established when the uromicroprotein exceeded 30 mg/day.

The animal experiments were carried out in accordance with the guidelines issued by Medical Ethics Committee of The Seventh People's Hospital of Shanghai University of Traditional Chinese Medicine.

### Sample Collection

After the model was successfully established, urine specimen, blood samples from rat heart, and renal tissues were collected and kept for corresponding examination. The urinary albumin concentration (U-Alb), urine albumin excretion (UAE), serum creatinine (Scr), serum urea nitrogen (BUN), uric acid (UA), and albumin (Alb) were measured by automatic biochemical analyzer. The level of IL-6, TNF- $\alpha$ , TGF- $\beta_1$ , CollagenI, CollagenIII, MMP2 and MMP9 in blood were detected by enzyme linked immunosorbent assay (Elisa). The expression of CXCL6, CXCR1, MMP2/9, CollagenI/III, TGF- $\beta_1$  in kidney tissues was determined by quantitative real-time reverse transcription polymerase chain reaction

<sup>1</sup><http://software.broadinstitute.org/gsea/downloads.jsp>

(qRT-PCR) and Western blotting. General histology of renal tissues was visualized by hematoxylin and eosin (H&E) and Masson staining.

## Cell Culture Experiments

A normal rat kidney fibroblast cell line NRK-49F cells were provided by the Shanghai Cell Bank, Chinese Academy of Sciences (Shanghai, China) and cultured in DMEM, which were supplemented with 10% FBS, 100 units/mL penicillin and 100  $\mu$ g/mL streptomycin at 37°C and 5% CO<sub>2</sub>.

## Cell Viability Analysis

Cells were seeded in 96-well plates at a density of 2000 cells per well and incubated overnight. The cell proliferation was quantitated using the Cell CCK-8; Dojindo Laboratories, Japan) according to the manufacturer's recommendations. Control cells were treated with cultured media containing 0.15% (v/v) DMSO.

## Lentivirus Transfection

The NRK-49F cells were seeded in 6-well plates, grown overnight and transfected with CXCL6 shRNA viral or overexpression lentivirus (both from GeneChem, Shanghai, China) to investigate the regulatory role of CXCL6. All of the transfections were performed using the Lipofectamine 3000 reagent (Invitrogen®;


Thermo Fisher Scientific, Inc.) according to the manufacturer's instructions. A blank vector lentivirus was used as a negative control (NC). CXCL6 expression in cells following transfection was confirmed by western blot analysis and quantitative real-time PCR.


## Quantitative Real-Time PCR Analysis (qRT-PCR)

Total RNA from the cells was extracted with TRIzol (Invitrogen, Carlsbad, CA, United States) according to the manufacturer's protocol. The SYBR Green-based qRT-PCR (Thermo Fisher) was performed to examine the relative CXCL6 and CXCR1 mRNA level which was normalized with GAPDH. The CXCL6 primers were: Primer F 5'-CTTAGCTCCAAGAATTAACC-3'; Primer R 5'-GGTCAAGACAAACATTATCC-3'. CXCR1 primers were: Primer F 5'-TCTTCCGCCAGGCATATAAAC-3'; Primer R 5'-TAGCAGACCAGCATGATGAAC-3'. The GAPDH primers were: Primer F 5'-CACCCACTCCTCCACCTTTG-3'; Primer R 5'-CCACCACCCTGTTGCTGTAG-3'.

## Western Blots

The bands were determined by western blotting using the enhanced chemiluminescent substrate (ECL, BioRad,





**FIGURE 2 |** Fibrosis related cytokines in DN rat model. **(A)** H&E and Masson staining of renal tissues. **(a)** H&E staining of renal tissues in control group **(b)** H&E staining of renal tissues model group **(c)** masson staining of renal tissues in control group **(d)** masson staining of renal tissues model group **(B)** the levels of several fibrosis related cytokines in peripheral blood were measured by Elisa. **(C)** Relative mRNA expressions of several fibrosis related cytokines were determined by qRT-PCR **(D)** the protein levels of several fibrosis related cytokines were examined by Western blotting. \*\*\* $p < 0.001$  vs. model group.

Richmond, CA, United States) in accordance with the manufacturer's instruction.

### Cytokine Detection

The fibrosis-related cytokines IL-6, TNF- $\alpha$ , TGF- $\beta$ 1, CollagenI, CollagenIII, MMP2 and MMP9 were detected in cell culture

supernatants and plasma samples. The following were used to measure the concentrations of cytokines: The Rat IL-6 ELISA kit, the Rat TNF- $\alpha$ ELISA kit, the Rat TGF- $\beta$ 1 ELISA Kit, the Rat CollagenIII ELISA kit, the Rat Collagen ELISA kit, the Rat MMP2 ELISA kit, and the Rat MMP9 ELISA kit (Xinyu Biological Technology, Shanghai, China).

## Histology

The tissue samples were fixed with 4% paraformaldehyde, dehydration and embedded in paraffin. General histology was visualized by hematoxylin and eosin (H&E) and Masson staining. The stained tissues were observed under an optical microscope, the renal fibrosis and collagen deposition were estimated, respectively.

## Statistical Analysis

The data are presented as the mean  $\pm$  SD, and all the experiments were performance three times independently. Unpaired two-tailed student's *t*-test was used to compare the difference between

**TABLE 1** | The detail data of biochemical indicators in each group (Mean  $\pm$  SD, *n* = 6).

Indicator	NC	DN rat model
Blood glucose (mM)	3.98 $\pm$ 0.21	16.42 $\pm$ 0.76***
Scr ( $\mu$ mol/L)	129.21 $\pm$ 9.04	457.75 $\pm$ 16.06***
BUN (mmol/L)	4.44 $\pm$ 0.35	10.80 $\pm$ 0.30***
UA ( $\mu$ mol/L)	101.09 $\pm$ 3.59	203 $\pm$ 5.19***
Alb(g/L)	39.5 $\pm$ 0.98	23.37 $\pm$ 0.76***
U-Alb( $\mu$ g/ml)	350.83 $\pm$ 44.24	1036.37 $\pm$ 25.94***
UAE(mg/L-h)	14.62 $\pm$ 1.23	43.18 $\pm$ 1.18***

\*\*\*compared with control: *P* < 0.001.


the two groups. The one-way analysis of variance (ANOVA) test with Dunnett *t* was performed for comparisons in the multiple groups and *P* < 0.05 was considered statistically significant.

## RESULTS

### CXCL6 Is Overexpression in DN

In order to define the CXCLs expression patterns in DN, the CXCLs (CXCL1, CXCL2, CXCL3, CXCL5, CXCL6, CXCL8, CXCL9, CXCL10, CXCL11, CXCL12, CXCL13, and CXCL14) mRNA levels in GEO DataSets (Asscess ID: GSE30122) were analyzed (**Figure 1A**). It was found that CXCL6 (3.08 fold, *P* < 0.001) was significantly higher in DN human kidney than in control (**Figure 1B**). Then Elisa and qRT-PCR were used to confirm the transcript of CXCL6 and it was found that CXCL6 was higher in plasma (**Figure 1C**) and kidney tissue (**Figure 1D**) of DN than in the control samples (*P* < 0.001).

The DN rat model was established by high-sugar-fat diet combined with a single intraperitoneal injection of streptozocin (STZ). As shown in **Table 1**, the level of blood glucose, Scr, BUN, UA, U-Alb, and UAE in the DN model group was drastically increased and Alb was decreased compared with the control group. Histology staining with H&E and Masson revealed typical damage and severe collagen fibrosis of kidney tissues in the DN model group (**Figure 2A**). The results of qRT-PCR and Western



blot were showed in **Figures 2C,D**. The expression of CXCL6 and its receptor CXCR1 of kidney tissues in the DN model group was significantly higher in DN rat kidney tissue than in the control group.

### Fibrosis-Related Cytokines Is High Expression in DN Rat Model


The level of fibrosis related cytokines (IL-6, TNF- $\alpha$ , TGF- $\beta$ 1, CollagenI/III, MMP2/9) in serum and the expression of CollagenI/III, MMP2/9 in kidney tissues were detected by Elisa, qRT-PCR and Western blot. As displayed in **Figure 2B**, the levels of IL-6, TNF- $\alpha$ , TGF- $\beta$ 1, CollagenI, CollagenIII, MMP2 and MMP9 were increased significantly when compared with the control group. And the expression of CollagenI/III, MMP2/9 of kidney tissues were significantly upregulated at both mRNA and protein levels in the DN model group of animals compared with the control group (**Figures 2C,D**).


### Glucose Increases the Expression of CXCL6 in NRK-49F Cells

Then effects of glucose on the expression of CXCL6 and its receptor CXCR1 in rat renal interstitial fibroblasts NRK-49F cells were detected by qRT-PCR and Western blotting after treatment with 5 mM, 10 mM, 15 mM, 25 mM, and 50 mM glucose, and further detected by ELISA. The results showed that 10 mM, 15 mM, 25 mM, 50 mM glucose significantly increased the expression of CXCL6 and CXCR1 both at mRNA (**Figure 3A**) and protein level (**Figure 3C**) in a concentration-dependent manner, furthermore the expression of CXCL6 increased with the treatment time of glucose (**Figure 3B**).

### CXCL6 Promoted the Proliferation of NRK-49F Cell

To further investigate the function of CXCL6 in high glucose-induced NRK-49F cells, lentivirus were transfected into high glucose (25 mM) induced NRK-49F, then the expression of





CXCR1 and CXCL6 were measured by RT-PCR and Western blotting. As shown in **Figures 4A,B**, compared with control, CXCL6 gene interference and over-expression resulted in a significant downregulation or upregulation of CXCL6 both at the mRNA and protein levels.

Then the effects of CXCL6 on high glucose-induced NRK-49F cells viability were investigated by CCK-8 assay. As shown in **Figure 4C**, high glucose (25 mM) increased the proliferation of NRK-49F cells strikingly. However, the proliferation of NRK-49F cell induced by high glucose was decreased markedly by knockdown CXCL6. Overexpression of CXCL6 significantly promoted the proliferation of high glucose (25 mM) induced NRK-49F cell and the addition of 30  $\mu$ M JAK inhibitor AG490 obviously reduced the proliferation of NRK-49F induced by CXCL6 overexpression (**Figure 4D**).


### CXCL6-Associated Signaling Pathways

To probe the CXCL6-associated pathways in fibrosis, GSEA were performed using RNA-sequencing data from the GEO DataSets (Asscess ID: GSE28221). JAK-STAT and CYTOKINE\_CYTOKINE\_RECEPTOR\_INTERACTION signaling

pathways were found to be strongly associated with CXCL6 expression (**Figure 6A**).

### CXCL6 Increased the Expression of Fibrosis-Related Cytokines in NRK-49F Cells

The expression of IL-6, TNF- $\alpha$ , TGF- $\beta$ 1, CollagenI, CollageIII, MMP2 and MMP9 in the cell supernatant was detected by ELISA. As shown in **Figure 5**, the expression of these cytokines was increased by high glucose concentration (25 mM) and overexpression of CXCL6 showed the same effect. On the contrary, the expression of these cytokines was decreased sharply by knockdown of CXCL6and 30  $\mu$ M JAK inhibitor AG490 significantly decreased CXCL6 overexpression-induced fibrosis-related cytokines expression. The expression of CollagenI/III, MMP2/9, PCNA, and p-STAT3 intracellular were evaluated by western blot. In **Figures 6B,C**, the results were consistent with that from ELISA. 25 mM glucose and overexpression of CXCL6 markedly promoted the expression of CollagenI/III, MMP2/9, PCNA, and p-STAT3. Knockdown of CXCL6 and treatment with 30  $\mu$ M JAK inhibitor AG490 drastically decreased their expression. The expression of STAT3 was



**FIGURE 6 |** CXCL6-associated pathways in DN. **(A)** GSEA was performed using GEO DataSets (Asscess ID: GSE28221). The JAK/STAT and CYTOKINE \_CYTOKINE \_RECEPTOR \_INTERACTION pathways were identified with the strongest association with CXCL6 expression. **(B,C)** Key moderators were determined by western blotting. Data were based on at least three independent experiments, and shown as mean  $\pm$  SD. \*\*\* $P < 0.001$ .

also measured, and no significant change was observed in all the groups.

## DISCUSSION

CXCL6 is a member of the chemokine family and plays a crucial role in experimental bleomycin-induced pulmonary fibrosis (Besnard et al., 2013). However, previous studies did not clarify the roles of CXCL6 in DN renal interstitial fibrosis. In this study, we revealed the overexpression of CXCL6 in DN and explored the underlying mechanism of CXCL6 in DN renal interstitial fibrosis.

There is emerging evidence that renal interstitial fibrosis is the most important pathologic feature of DN (He et al., 2016). Excessive deposition of ECM is the main reason of renal interstitial fibrosis (Picard et al., 2008) and it is widely believed that renal interstitial fibroblasts are the main cells

which play an important role in renal interstitial fibrosis due to producing excessive ECM (Faulkner et al., 2005). High glucose concentration, a distinctive risk factor in diabetes, is a potential stimulus for the proliferation, differentiation, and survival of certain types of cells (Lee and Han, 2010; Yuan et al., 2011; Le et al., 2012). Hence we explored the impacts of glucose on renal interstitial fibroblasts NRK-49F cell proliferation. The results from cck8 assay indicated that (10, 15, 25, 50 mM) glucose significantly promoted the proliferation of NRK-49F cells. Meanwhile, CXCL6 and CXCR1 expression were also dramatically induced by high glucose in a dose-dependent manner, demonstrating that CXCL6 may be a key factor in high glucose-induced renal fibrosis.

Then we examined the functions of CXCL6 in renal interstitial fibrosis. CXCL6 shRNA lentivirus was transfected into NRK-49F and we confirmed that CXCL6 shRNA viral transfection markedly reduced CXCL6 expression at protein

levels and partially attenuated the pro-proliferation effects of high glucose. On the contrary, ectopic expression of CXCL6 in NRK-49F significantly promoted the proliferation of NRK-49F. These data suggest that increasing the expression of CXCL6 contributes to high glucose-induced renal fibroblasts proliferation and activation.

In addition to cellular phenotype, we explored the molecular mechanism by which CXCL6 functions as a pro-fibrotic factor in DN. Overexpression of CXCL6 remarkably up-regulated the expression of pro-inflammatory cytokine (such as TNF- $\alpha$  and IL-6) and pro-fibrotic markers (such as TGF- $\beta$ , and collagen). Moreover, The JAK/STAT signaling pathway has previously been shown to implicate in the pathophysiology of fibrotic renal disease (Matsui and Meldrum, 2012; Matsui et al., 2017). Increasing the activity of genes involved in JAK/STAT signaling pathway may mediate renal fibrosis. In the present study, GSEA data indicated that higher CXCL6 expression was positively correlated with JAK/STAT pathways in fibrosis (Figure 5A). The expression of JAK/STAT and CYTOKINE\_CYTOKINE\_RECEPTOR\_INTERACTION pathway related factors was significantly suppressed by CXCL6 knockdown, while overexpression of CXCL6 in NRK-49F cells caused an inverse effect. With the addition of JAK inhibitor AG490, the proliferation of NRK-49F and the expression of these related factors induced by overexpression CXCL6 were inhibited significantly. These data suggest the roles of CXCL6 on JAK/STAT3, and may account for renal fibroblasts proliferation and renal interstitial fibrosis.

Approximately 30–40% of patients with diabetes develop nephropathy (Tesch, 2017) and the exact pathogenesis of DN is complex. Pathogenetic factors include hyperglycaemia, increased intraglomerular, the immune system, and systemic blood pressure, etc., Moreover, several cytokines and growth factors, which have complex mutual interactions, have been identified and may cause damaging effects on the kidney (Kim and Park, 2017). Renal response on these noxious effects further enhances renal damage. Although a number of therapeutic interventions have been shown to postpone the development or slow down the progression of DN, the proportion of ESRD due to DN remains high (Jin, 2015). Therefore, future interventions of other potential pathogenic factors were needed. Moreover, the use of representative animal models to defined clinical trial inclusion criteria may help to identify appropriate DN therapies.

## CONCLUSION

In conclusion, this study found that CXCL6 is overexpression in DN patients and the DN rat model, and the expression of fibrosis-related cytokines was consistent with the expression of

CXCL6. The results of *in vitro* experiment showed that high glucose concentration promotes renal fibroblasts proliferation and CXCL6 expression in NRK-49F. Knockdown of CXCL6 significantly inhibited the proliferation of NRK-49F and decreased the expression of pro-inflammatory and pro-fibrotic cytokines, while overexpression of CXCL6 exhibited the opposite phenomenon. In addition JAK inhibitor AG490 showed the same effect as knockdown of CXCL6 and it is likely that CXCL6 may promote fibrosis-related factors to accelerate the development of DN renal interstitial fibrosis by activating JAK/STAT3 signaling pathway. These findings not only broaden our understanding of the pathogenesis of renal DN but also provide a new attractive strategy in treating this disease. Due to the complexity and heterogeneity of the DN, further investigations of the molecular mechanisms of CXCL6 in DN are required to verify the function of CXCL6 in DN.

## AUTHOR CONTRIBUTIONS

M-YS and S-JW performed qRT-PCR and western blot assays and wrote the manuscript. X-QL contributed to the animal experiments. Y-LS and X-HT contributed to cell culture and data analyses. KR revised the manuscript. J-RL and HN collected clinical samples and implemented bioinformatics analysis. HZ and L-JZ applied for grants, designed the experimental protocols, and directed manuscript writing.

## FUNDING

This work was supported by funds from the National Natural Science Foundation of China (81502515, 81503332, 81703755, and 81773941), Shanghai Municipal Science and Technology Commission (15401902700 and 15401971800), Shanghai Municipal Health and Family Planning Commission (20154Y0063), Shanghai University of Traditional Chinese Medicine—Gaofeng Clinical Medicine Grant Support, Outstanding Leaders Training Program of Pudong Health Bureau of Shanghai (PWR12015-05), and Pudong New Area Science and Technology Commission (PKJ2015-Y13). Summit Discipline of Clinical Traditional Chinese Medicine in Pudong New Area of Shanghai (PDZY-2018-0601).

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fphar.2019.00224/full#supplementary-material>

## REFERENCES

- Balamayooran, G., Batra, S., Cai, S., Mei, J., Worthen, G. S., Penn, A. L., et al. (2012). Role of CXCL5 in leukocyte recruitment to the lungs during secondhand smoke exposure. *Am. J. Respir. Cell Mol. Biol.* 47, 104–111. doi: 10.1165/rcmb.2011-0260OC
- Besnard, A. G., Struyf, S., Guabiraba, R., Fauconnier, L., Rouxel, N., Proost, P., et al. (2013). CXCL6 antibody neutralization prevents lung inflammation and fibrosis in mice in the bleomycin model. *J. Leukoc. Biol.* 94, 1317–1323. doi: 10.1189/jlb.0313140
- Faulkner, J. L., Szykalski, L. M., Springer, F., and Barnes, J. L. (2005). Origin of interstitial fibroblasts in an accelerated model of angiotensin II-induced

- renal fibrosis. *Am. J. Pathol.* 167, 1193–1205. doi: 10.1016/S0002-9440(10)61208-4
- Gin, H., Rigalleau, V., and Aparicio, M. (2000). Lipids, protein intake, and diabetic nephropathy. *Diabetes Metab.* 26, 45–53.
- Grande, M. T., and López-Novoa, J. M. (2009). Fibroblast activation and myofibroblast generation in obstructive nephropathy. *Nat. Rev. Nephrol.* 5, 319–328. doi: 10.1038/nrneph.2009.74
- He, T., Xiong, J., Nie, L., Yu, Y., Guan, X., Xu, X., et al. (2016). Resveratrol inhibits renal interstitial fibrosis in diabetic nephropathy by regulating AMPK/NOX4/ROS pathway. *J. Mol. Med.* 94, 1359–1371. doi: 10.1007/s00109-016-1451-y
- Jin, D. C. (2015). Major changes and improvements of dialysis therapy in Korea: review of end-stage renal disease registry. *Korean J. Intern. Med.* 30, 17–22. doi: 10.3904/kjim.2015.30.1.17
- Kebschull, M., Demmer, R., Behle, J. H., Pollreis, A., Heidemann, J., Belusko, P. B., et al. (2009). Granulocyte chemotactic protein 2 (gcp-2/cxcl6) complements interleukin-8 in periodontal disease. *J. Periodontal. Res.* 44, 465–471. doi: 10.1111/j.1600-0765.2008.01134.x
- Kim, Y., and Park, C. W. (2017). New therapeutic agents in diabetic nephropathy. *Korean J. Intern. Med.* 32, 11–25. doi: 10.3904/kjim.2016.174
- Le, A., Lane, A. N., Hamaker, M., Bose, S., Gouw, A., Barbi, J., et al. (2012). Glucose-independent glutamine metabolism via TCA cycling for proliferation and survival in B cells. *Cell Metab.* 15, 110–121. doi: 10.1016/j.cmet.2011.12.009
- Lee, Y. J., and Han, H. J. (2010). Troglitazone ameliorates high glucose-induced EMT and dysfunction of SGLTs through PI3K/Akt, GSK-3 $\beta$ , Snail1, and  $\beta$ -catenin in renal proximal tubule cells. *Am. J. Physiol. Renal. Physiol.* 298, F1263–F1275. doi: 10.1152/ajprenal.00475.2009
- Matsui, F., Babitz, S. A., Rhee, A., Hile, K. L., Zhang, H., and Meldrum, K. K. (2017). Mesenchymal stem cells protect against obstruction-induced renal fibrosis by decreasing STAT3 activation and STAT3-dependent MMP-9 production. *Am. J. Physiol. Renal. Physiol.* 312, F25–F32. doi: 10.1152/ajprenal.00311.2016
- Matsui, F., and Meldrum, K. K. (2012). The role of the Janus kinase family/signal transducer and activator of transcription signaling pathway in fibrotic renal disease. *J. Surg. Res.* 178, 339–345. doi: 10.1016/j.jss.2012.06.050
- Müller, K., and Meineke, V. (2011). Radiation-induced mast cell mediators differentially modulate chemokine release from dermal fibroblasts. *J. Dermatol. Sci.* 61, 199–205. doi: 10.1016/j.jdermsci.2011.01.003
- Picard, N., Baum, O., Vogetseder, A., Kaissling, B., and Le Hir, M. (2008). Origin of renal myofibroblasts in the model of unilateral ureter obstruction in the rat. *Histochem. Cell Biol.* 130, 141–155. doi: 10.1007/s00418-008-0433-8
- Reich, N., Beyer, C., Gelse, K., Akhmetshina, A., Dees, C., and Zwerina, J. (2011). Microparticles stimulate angiogenesis by inducing ELR(+) CXC-chemokines in synovial fibroblasts. *J. Cell Mol. Med.* 15, 756–762. doi: 10.1111/j.1582-4934.2010.01051.x
- Sadik, C. D., Kim, N. D., and Luster, A. D. (2011). Neutrophils cascading their way to inflammation. *Trends Immunol.* 32, 452–460. doi: 10.1016/j.it.2011.06.008
- Strutz, F., and Zeisberg, M. (2006). Renal fibroblasts and myofibroblasts in chronic kidney disease. *J. Am. Soc. Nephrol.* 17, 2992–2998. doi: 10.1681/ASN.2006050420
- Tesch, G. H. (2017). Diabetic nephropathy - is this an immune disorder? *Clin. Sci.* 131, 2183–2199. doi: 10.1042/CS20160636
- Wada, J., and Makino, H. (2013). Inflammation and the pathogenesis of diabetic nephropathy. *Clin. Sci.* 124, 139–152. doi: 10.1042/CS20120198
- Wasmuth, H. E., Lammert, F., Zaldivar, M. M., Weiskirchen, R., Hellerbrand, C., Scholten, D., et al. (2009). Antifibrotic effects of CXCL9 and its receptor CXCR3 in livers of mice and humans. *Gastroenterology* 137, 309–319. doi: 10.1053/j.gastro.2009.03.053
- Yuan, P., Xue, H., Zhou, L., Qu, L., Li, C., Wang, Z., et al. (2011). Rescue of mesangial cells from high glucose-induced over-proliferation and extracellular matrix secretion by hydrogen sulfide. *Nephrol. Dial. Transplant.* 26, 2119–2126. doi: 10.1093/ndt/gfq749

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Sun, Wang, Li, Shen, Lu, Tian, Rahman, Zhang, Nian and Zhang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.