



LJMU Research Online

Afroz, M, Zihad, SMNK, Uddin, SJ, Rouf, R, Rahman, MS, Islam, MT, Khan, IN, Ali, ES, Aziz, S, Shilpi, JA, Nahar, L and Sarker, SD

A systematic review on antioxidant and anti-inflammatory activity of Sesame (Sesamum indicum L.) oil and further confirmation of anti-inflammatory activity by chemical profiling and molecular docking

<http://researchonline.ljmu.ac.uk/id/eprint/10871/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Afroz, M, Zihad, SMNK, Uddin, SJ, Rouf, R, Rahman, MS, Islam, MT, Khan, IN, Ali, ES, Aziz, S, Shilpi, JA, Nahar, L and Sarker, SD (2019) A systematic review on antioxidant and anti-inflammatory activity of Sesame (Sesamum indicum L.) oil and further confirmation of anti-inflammatory activity by

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

A systematic review on antioxidant and anti-inflammatory activity of Sesame (*Sesamum indicum* L.) oil and further confirmation of anti-inflammatory activity by chemical profiling and molecular docking

Running Head: A systematic review on antioxidant and anti-inflammatory activity of Sesame (*Sesamum indicum* L.) oil

Mohasana Afroz^{1,γ}, S. M. Neamul Kabir Zihad^{1,γ}, Shaikh Jamal Uddin^{1,γ*}, Razina Rouf², Md. Shamim Rahman³, Muhammad Torequl Islam^{4,5,*}, Ishaq N. Khan⁶, Eunüs S. Ali⁷, Shahin Aziz⁸, Jamil A. Shilpi¹, Lutfun Nahar⁹, Satyajit D. Sarker⁹

¹*Pharmacy Discipline, Life Science School, Khulna University, Khulna 9208, Bangladesh*

²*Department of Pharmacy, Faculty of Life Science, Bangabandhu Sheikh Mujibur Rahman Science & Technology University, Gopalganj, Bangladesh*

³*Biotechnology and Genetic Engineering Discipline, Life Science School, Khulna University, Khulna 9208, Bangladesh*

⁴*Department for Management of Science and Technology Development, Ton Duc Thang University, Ho Chi Minh City-700000, Vietnam*

⁵*Faculty of Pharmacy, Ton Duc Thang University, Ho Chi Minh City-700000, Vietnam*

⁶*PK-NeuroOncology Research Group, Institute of Basic Medical Sciences, Khyber Medical University, Peshawar, 25000, Pakistan*

⁷*Department of Biochemistry and Molecular Genetics, Northwestern University Feinberg School of Medicine, Chicago, IL, United States*

⁸*Chemical Research Division, Bangladesh Council of Scientific and Industrial Research, Dhaka, Bangladesh*

⁹*Medicinal Chemistry and Natural Products Research Group, School of Pharmacy and Biomolecular Sciences, Liverpool John Moores University, Liverpool, United Kingdom*

^γAuthors contributed equally

ABSTRACT

Traditionally sesame oil has been used as a popular food and medicine. The review aims to summarize the antioxidant and anti-inflammatory effects of sesame oil (SO) and its identified compounds as well as further fatty acid profiling and molecular docking study to correlate the interaction of its identified constituents with COX-2. For this, a literature study was made using Google Scholar, Pubmed and SciFinder databases. Literature study demonstrated that SO has potential antioxidant and anti-inflammatory effects in various test systems, including humans, animals and cultured cells through various pathways such as inhibition of COX, non-enzymatic defense mechanism, inhibition of pro-inflammatory cytokines, NF-kB or MAPK signaling and prostaglandin synthesis pathway. Fatty acid analysis of SO using gas chromatography identified known 9 fatty acids. *In-silico* study revealed that sesamin, sesaminol, sesamol, stigmasterol, Δ^5 -avenasterol, and Δ^7 -avenasterol (-9.6 to -10.7 kcal/mol) were the most efficient ligand for interaction and binding with COX-2. The known fatty acid were also showed binding efficiency with COX-2 to some extent (-6.0 to -8.4 kcal/mol). In summary, it is evident that sesame oil may be one of promising traditional medicine that we could use in the prevention and management of diseases associated with oxidative stress and inflammation.

Key words: *Sesamum indicum*, sesame oil, anti-inflammatory activity, antioxidant activity, molecular docking

1. INTRODUCTION

Sesame (*Sesamum indicum* L., Family: Pedaliaceae) is one of the most archaic crops cultivated since the civilizations of Indus valley, 2500 BC, Babylonia, 1750 BC, in Syria and Palestine around 3000 BC (Bedigian, 2004; Ricci, Groth, & Lago, 1999). From the ancient time sesame oil, derived from sesame seed, is used in various purposes including food, salve and medicine. The Assyrians community used sesame oil as food and medicine whereas it was a popular ingredient of massage in Ayurvedic medicine and interestingly in India it has been used as sacred oil (Shah, 2016). Ethnomedicinally sesame oil has been used against a number of ailments including pain, fever, inflammation, constipation, diuretic, healing burns and wounds and others (**Table 1**).

Sesame seed contains about 50% oil of its weight and has a distinguishable nutty taste and aroma. Literature study found that sesame oil consists of saturated and unsaturated fatty acids, lignans, phytosterols, tocopherols and vitamins. It is reported that sesame oil contains linoleic (39 to 59%), oleic acid (35 to 54%), palmitic acid (10%) and stearic acid (5%) (Gunstone, 2006). It also reported some antioxidant compounds such as tocopherol and lignans like sesamol, sesamin and sesamol (Hwang, 2005; Namiki, 1995) (**Figure 1 (a-d) and Table 1**). Several studies have been reported that sesame oil contributes to prevent a number of disorders including hypertension, hypercholesterolemia, cancer and aging (Kanu, Bahsoon, Kanu, & Kandeh, 2010). Moreover, sesame oil exhibits multiple physiological functions such as decreasing plasma triacylglycerol and arachidonic acid levels, imparting anti-inflammatory and estrogenic activities (Hemalatha, 2007; Shahidi, Wanasundara, & Wanasundara, 1997). Sesame oil showed an exceptionally high oxidative stability compared to soybean, corn and most other popular vegetable oils (Carrasco-Pancorbo et al., 2005; Miniotti & Georgiou, 2010; Shahidi et al., 1997). Some studies have shown that sesame seeds can reduce oxidative stress by modulating the concentration of antioxidant enzymes [superoxide dismutase (SOD), glutathione (GSH), glutathione peroxidase (GPx), catalase (CAT)], as well as of oxidative stress markers [thiobarbituric acid reactive substance (TBARS) and malondialdehyde (MDA)] (Alipoor, Haghghian, Sadat, & Asghari, 2012; Chen et al., 2005; Sankar, Ali, Sambandam, & Rao, 2011; Sankar, Rao, Sambandam, & Pugalendi, 2006a, 2006b; Sankar, Sambandam, Rao, & Pugalendi, 2005; Wichitsranoi et al., 2011). However, there is no chemical profiling and further interaction docking study of sesame oil constituents to COX enzyme to correlate its ethnopharmacological

and reported anti-inflammatory activity. As a part of our continuous search for natural lead molecules for antioxidant and anti-inflammatory activity (Rahman, Khatun, Uddin, & Shilpi, 2016; Uddin, Grice, & Tiralongo, 2011; Zihad et al., 2018; Zilani et al., 2018), here the aim of this review is to draw antioxidant and anti-inflammatory mechanisms of sesame oil on the basis of literature findings using the above-mentioned databases and further molecular docking study of its identified constituents with CoX-2 enzyme to confirm its anti-inflammatory activity.

2. MATERIALS AND METHODS

2.1 Literature review

2.1.1 Search strategy

To begin this review, a comprehensive literature search was conducted by using several database platforms such as PubMed, Google Scholar and SciFinder with the term ‘sesame oil’ with ‘antioxidant’, ‘oxidative stress’, ‘inflammation’ and ‘anti-inflammation’. We focused on reports that were only in English due to the language barrier, time efficiency and non-feasible costs of translation.

2.1.2 Selection of studies for inclusion in the systematic review

The following types of studies and investigations were included in this review: a) *in-vivo* animal studies, b) *in-vitro* studies, c) clinical studies, d) studies that include the effects of sesame oil or aqueous extract of sesame oil or isolated compounds of sesame oil on oxidation and inflammation, e) studies that indicate the concentrations or doses employed and the form of administration and f) studies that point out to the mechanisms of action associated with the sesame oil treatment and its isolated derivatives.

2.1.3 Data extraction

All the searched articles were assessed for the information according to surname of first author, year of publication, sesame oil and its isolated compound, test system, observation, result, suggested mechanism of action, concentrations tested and molecular mechanism involved. The general steps of the data search, the exclusion-inclusion data and other relevant information are presented in **Figure 2**.

2.2 Chemical profiling of sesame oil using gas chromatography

2.2.1 Extraction of sesame oil

The fresh sesame seeds were collected from local markets of Khulna, Bangladesh and identified by the expert of Forestry Discipline, Khulna University. A sample specimen of sesame seeds were deposited in Pharmacy Discipline, Khulna University for future reference. The collected samples were washed by water to remove any dust materials and shed dried. Finally the sesame oil was extracted using expeller pressing (also called oil pressing) method under mild conditions to preserve the natural characteristics of the fresh oil.

2.2.2 Preparation of methyl ester (FAMES)

The fatty acid composition was determined by analysis of their methyl esters. The fatty acid methyl esters (FAMES) were prepared by esterification reaction using $\text{BF}_3\text{-MeOH}$ complex according to an established method (Wirasmita et al., 2013). Briefly, sesame oil was taken in a screw capped glass tube and 1 mL of $\text{BF}_3\text{-MeOH}$ complex was added. The oil mixture was heated at 100°C for 1 hr in a water bath and then cooled at room temperature. 1 mL of deionized water and 2 mL of hexane were added in the cooled mixture and vortexed and centrifuged at low RPM for two minutes. The upper layer was collected by means of syringe and kept in refrigerator with air tight glass vial. The prepared FAMES were ready to analyze.

2.2.3 Gas Chromatography (GC) analysis of sesame oil

GC analysis of the fatty acids of sesame oil was carried out on Agilent 7890 system that equipped with flame ionization detector and split less injection system. The GC was fitted with a HP-5MS capillary column ($60\text{ m} \times 0.25\text{ mm}$). The temperature program was as follows: initial oven temperature at 50°C , then increased at $25^\circ\text{C}/\text{rpm}$ to 175°C for 6 min., then $4^\circ\text{C}/\text{rpm}$ to 230°C for 10 min and total run time was for 30 min. Helium was used as the carrier gas at 17.69 psi pressure with flow 0.6 mL/min. Samples were dissolved in methanol and 1 μL aliquot was injected automatically.

2.3 Molecular docking study of identified compounds in sesame oil

Docking is a method which predicts the preferred orientation of one molecule to a second when bound to each other to form a stable complex (Lengauer & Rarey, 1996). It's a popular method to perform virtual screening on large libraries of compounds and propose structural hypotheses of how the ligands inhibit the target, which is invaluable in lead optimization (Grouleff, Irudayam, Skeby, & Schiøtt, 2015). The goal of ligand-protein docking is to predict the predominant binding mode(s) of a ligand with a protein of known three-dimensional structure. Therefore, further confirmation of binding efficiency of identified 31 constituents in sesame oil with the COX-2 enzyme, a key protein responsible for inflammatory pain, a docking study was conducted.

2.3.1 Computational Study

Computational studies of the identified compounds in sesame oil were conducted against human COX-2 enzyme, which is the inducible enzyme in human body contributing to inflammatory pain. Human COX-2 or Prostaglandin G/H Synthase 2 (PGH2) is a homodimer protein comprising 604 amino acids in each monomeric unit (UniProtKB ID: P35354). Homology model of single monomeric unit of the human COX-2 was built using Modeller 9.19 (Zihad et al., 2018) taking three reported models as templates (PDB ID: 5F1A_A, 5IKQ_A, and 5F19_A) and dimeric structure predicted using Galaxy Gemini (Shin, Lee, Heo, Lee, & Seok, 2014) following energy minimization step. The initial model was selected based on dope (Discrete Optimized Protein Energy) score and further validated by analyzing Ramachandran plot (Using procheck and rampage) (Lovell et al., 2003). Knowledge based energy calculation was performed in ProSA (Wiederstein & Sippl, 2007). All the selected ligands were prepared using Autodock tools and the binding site in the enzyme was predicted through ProBiS server (Konc & Janežič, 2014) and further arachidonic acid was re-docked using Autodock Vina. Initial docking grid (40×40×40, 0.8 Å) enclosing the active binding site was generated depending on the binding of arachidonic acid. Finally all the prepared ligands were docked into the predicted binding pocket of the validated homology model of COX-2 utilizing Autodock Vina (Trott & Olson, 2010).

3. RESULTS AND DISCUSSION

3.1 LITERATURE REVIEW

The databases searched for antioxidant and anti-inflammatory activity of sesame oil were revealed 5039 records from PubMed (2096), Google Scholar (2000) and SciFinder (943) in which 87 studies were meeting the inclusion criteria. The exclusion of 4952 documents were due to absence of sesame oil mentioning in the title or the abstract, duplication of information and non-relevance of the study.

3.1.1 Antioxidant effect of sesame oil and its isolated compounds

Sesame is a popular ethnomedicinal plant and its oil has been reported to use against a number of diseases including oxidative induced different ailments. Experimental studies have been reported that sesame seeds and its oil can reduce oxidative stress by modulating the antioxidant enzymes as well as of oxidative stress markers. **Table 2** and **4** summarized the anti-oxidative effects of sesame oil and its reported constituents either *in-vivo*, *in-vitro* or in clinical test model. A number of possible mechanisms for antioxidant effect of sesame oil has been highlighted in different reports (**Figure 3**). One of the most common antioxidant effects of SO is to reduce oxidative stress in lipopolysaccharide (LPS) induced oxidative stress rat models (Chiang et al., 2014; Hsu, Chiang, Chien, Huang, & Liu, 2004a; Hsu & Liu, 2004b). Hsuet al., (2004) reported that the level of, lipid peroxidation, hydroxyl radical and nitrite levels reduced while the antioxidant enzyme superoxide dismutase (SOD) and catalase increased in SO treated LPS induced oxidative stress rat models (Hsu et al., 2004a). In addition, parenterally SO-treated animal groups showed attenuation of hepatic disorder, although the symptoms were different when the SO was given orally (Ahmad et al., 2006; Hsu et al., 2004a; Saleem, Chetty, & Kavimani, 2013). However, superoxide anion decreased and glutathione increased in the oral SO groups (Ahmad et al., 2006; Saleem et al., 2013). Ahmad S. et al. (2006) showed that dietary SO behave as an antioxidant, resulting in increased enzymatic and non-enzymatic antioxidants activity in the middle cerebral artery occlusion (MCAO rat model against MCAO-induced cerebral ischemia injury (Ahmad et al., 2006). Furthermore, giving SO orally could reduce oxidative myocardial damage induced by isoproterenol rat model (Saleem et al., 2013). According to Abdou et al., (2012), 5 mL/kg b.w, orally given sesame oil plays a protective role on cypermethrin-induced oxidative stress, biochemical changes, histopathological damage and genomic DNA fragmentation (Abdou, Hussien, & Yousef, 2012). SO not only protect rats from oxidative stress induced by chemical but also showed its protective effect against oxidative stress induced by chronic exposure to

electromagnetic radiation (EMR)(Marzook, El Moneim, & Elhadary, 2014). Majority of the reported protective effects of crude SO demonstrated in oxidative induced rat model within the dose of oral 5-10 mL/kg b.w(**Table 2**). However, only one clinical trial was conducted by Sankar et al., (2006) to investigate the effect of SO in 40 hypertensive patients (32 male and 18 female, aged 35-60 yrs) who were on antihypertensive therapy either with diuretics (hydrochlorothiazide) or Beta-blockers (atenolol) and the results demonstrated that sesame oil as edible oil lowered blood pressure, decreased lipid peroxidation, and increased antioxidant status (increased SOD, CAT and GSH activity) without changing lipid profile of the hypertensive patients(Sankar et al., 2006a).

Phytochemical constituents from SO have also been evaluated for their protective effects against oxidative stress both *in-vivo* and *in-vitro*. Among these sesamol, sesamin sesamol, sesaminol, ligning and their derivatives are the prominent constituents that reported for antioxidant effects (**Table 4**).Sesamol is one the main constituent that has been reported to have antioxidant effect (in both *in-vivo* and *in-vitro* model) through reduction of lipid peroxidation, superoxide and nitric oxide production as well as increased antioxidative enzyme (SOD, CAT, GSH) level at a different doses ranging 10-50 mg/kg b.w.(Chiang et al., 2014; Kang, Naito, Sakai, Uchida, & Osawa, 1999; Khamphio, Barusrux, & Weerapreeyakul, 2016; Lv, Zhu, & Liu, 2015; Yashaswini, Sadashivaiah, Ramaprasad, & Singh, 2017b). Its antioxidant effect has a strong correlation with its anti-inflammatory action. Sesamin another prominent compound reported in SO that possesses significant protective effects against oxidative stress in animal model including through ameliorate SOD, GPx, reduction of malondialdehyde and elevated different liver marker, TBARS and lipid peroxidation as well as reduction of superoxide production(Ahmad et al., 2016; Chen, Ying, Chen, Zhang, & Zhang, 2015; Hou, Chang, & Jeng, 2015; Hsieh et al., 2011; Jeng & Hou, 2005; Jnaneshwari et al., 2014; Nakai et al., 2003; Tian & Guo, 2017a; Zhang et al., 2016b). *In-vitro* studies of sesamin especially in neuronal cell line (PC12), it showed reduction of ROS generation especially NO production(Cao et al., 2013; Duarte, Chenet, de Almeida, Andrade, & de Oliveira, 2018; Lee et al., 2009; Yashaswini, Rao, & Singh, 2017a). However, Hou et al., (2015) reported that 10 μ M sesamin derivative, 3-bis (3-methoxybenzyl) butane-1, 4-diol inhibits LDH, lipid peroxidation and apoptosis as well as increases ACh release and also prevents cell damage, scavenges ROS, and attenuates the elevation of intracellular free Ca ion on A β -stressed PC12 cells(Hou et al., 2015).Both the

compounds sesamol and sesaminol reported to exert their antioxidant effect through the reduction of nitric oxide production and inhibit peroxidation although there is only an oxygen bridge difference (**Figure 1**) in their structure (Hou, Chen, Tzen, & Jeng, 2003; Kong et al., 2016; Lee et al., 2009). However, there is another report stated that sesaminol has the ability to increase antioxidant enzyme (such as SOD, CAT, GSH) in H₂O₂ exposed cell line (Wu et al., 2015).

3.1.3 Anti-inflammatory effect of sesame oil and its isolated compounds

Inflammation is a biological protective response induced by tissue injury or infection by harmful stimuli (microorganisms, damaged cell, irritants and non-self cells) involving an increase of permeability of endothelial lining cells, influxes of blood leukocytes into the interstitium, oxidative burst, release of kinins or cytokines, activation of several enzymes (oxygenases, nitric oxide synthases, peroxidases) as well as the arachidonic acid metabolism (Miguel, 2010). Uncontrolled chronic inflammation may cause a number of diseases and therefore anti-inflammatory agents either from natural source or synthetic are considered as therapeutic agents to prevent these ailments. Sesame oil has been reported to be used against a number of diseases including inflammation and a number of reports have been published on its anti-inflammatory action with highlighting a number of possible mechanisms for anti-inflammatory effects (**Figure 3**). **Table 3 and 4** summarize the anti-inflammatory effects of sesame oil and its reported constituents either *in-vivo*, *in-vitro* or in clinical test model. A number of reports showed that the anti-inflammatory activity of SO is associated with the reduction of pro-inflammatory cytokines and related factors in animal model (**Table 3**). A number of reports stated that SO showed the anti-inflammatory effect through reduction of pro-inflammatory cytokines (TNF- α , IL-6, IL-1 α , IL-1 β) as well as anti-oxidative effect either reduction of ROS or induction of anti-oxidative enzymes (GSH, NADPH oxidase, GPx) (Hsu, Chu, & Jou, 2016c; Hsu, Chu, Li, & Liu, 2008b; Kang, Naito, Tsujihara, & Osawa, 1998; Nakano et al., 2002; Narasimhulu, Riad, & Parthasarathy, 2018b; Narasimhulu et al., 2016) (**Figure 3**). Ali et al. (2017) showed that SO significantly decreased the levels of inflammatory mediator TNF- α and IL-6, although the levels of anti-inflammatory mediator IL-10 were significantly amplified in groups treated with combination of SO with MTX (first line anti-arthritis agent) compared to group treated with SO only or MTX only in adjuvant-induced arthritis rat model (Ali, Atia, El Allawy, & Alla, 2017). Later, Ismail et al., (2018) reported that daily oral administration of Δ^9 -THC/sesame

oil combination, over a period of 21 days, attenuated erythrocyte sedimentation rate (ESR) scores and pro-inflammatory cytokines, including TNF- α , IL-1 β , and IL-6 levels, to normal values and MTX treated values on adjuvant-induced arthritis (AIA) rat model (Ismail, Hasan, El-Orfali, Ismail, & Khawaja, 2018). Some *in-vitro* studies also showed that SO has potent anti-inflammatory properties by down regulating inflammatory markers as seen with LPS-induced inflammation in RAW macrophages or mouse peritoneal macrophages (Deme, Narasimhulu, & Parthasarathy, 2018; Narasimhulu, Burge, Doomra, Riad, & Parthasarathy, 2018a; Selvarajan, Narasimhulu, Bapputty, & Parthasarathy, 2015). The atherosclerosis PCR array revealed down regulation of a number of inflammatory markers such as Ccl2 or MCP-1, Ccl5 or RANTES, IL-1 α , IL-1 β , and TNF in the presence of SO (Selvarajan et al., 2015). Selvarajan et al., (2015) proved that sesame oil play a role in the prevention of atherosclerotic disease by suppression of colony stimulating factor 2 (Csf2), NF-kB1, an important pro-inflammatory transcription factor (Selvarajan et al., 2015). Other reports also demonstrated that SO reduced inflammatory gene expression and induced genes involved in cholesterol metabolism (Narasimhulu et al., 2018b; Narasimhulu et al., 2016). Independent analysis revealed significant inhibition of IL-6 expression in the presence of SO (Selvarajan et al., 2015).

Phytochemical constituents mainly sesamol, sesamin sesamolin, sesaminol, and their derivatives from SO have been reported for anti-inflammatory activity both *in-vivo* and *in-vitro* model (Table 4). A number of constituents (sesamol, sesamolin, sesamin, sesamin derivative and sesaminol) identified from SO showed their anti-inflammatory activity by down regulating the COX-2 activity or PGE2 synthesis (Chen et al., 2015; Hemshekhar et al., 2013; Hsieh et al., 2011; Hsu et al., 2008b; Hung, Chen, & Hou, 2017; Jeng & Hou, 2005; Kong et al., 2016; Monteiro et al., 2014; Yashaswini et al., 2017b). Other reports also demonstrated that some of these constituents (sesamol, sesamolin, sesamin, sesaminol) inhibit the pro-inflammatory cytokines such as TNF- α , IL-1 β , IL-6 and NF-kB either *in-vivo* rat (Bournival, Plouffe, Renaud, Provencher, & Martinoli, 2012; Chen et al., 2015; Hemshekhar et al., 2013; Hsu, Chu, Chandrasekaran, & Liu, 2009a; Jeng & Hou, 2005; Yashaswini et al., 2017b) or *in-vitro* cell culture model (Bournival et al., 2012). Sesamol, sesamolin and sesamin also suppress the inflammatory cascade by down regulating the expression of caspase-3 (Baluchnejadmojarad, Mansouri, Ghalami, Mokhtari, & Roghani, 2017; Chopra, Tiwari, Arora, & Kuhad, 2010; Hsieh et al., 2011; Hung et al., 2017; Jeng & Hou, 2005). A recent clinical trial was reported on

sesamin supplementation on cardiovascular risk factors in women with rheumatoid arthritis and the results found that sesamin research had been conducted and evaluated by Helli et al., (2016), showed protective effect of sesame oil(Helli, Mowla, Mohammadshahi, & Jalali, 2016).

3.2 CHEMICAL PROFILING OF SESAME OIL USING GAS CHROMATOGRAPHY

Sesame oil is an edible vegetable oil derived from sesame seeds. The presence of saturated and unsaturated fatty acids justifies the use of this oil to treat inflammation and other diseases. However, many researches have been carried out on sesame (*Sesamum indicum*), but no report on their fatty acid profiling FAME analysis using GC. Fatty acid has a strong inhibitory effect on the production of NO (Ren & Chung, 2007). They inhibit inducible nitric oxide synthase (iNOS), cyclooxygenase-2 (COX-2), and tumor necrosis factor-alpha (TNF-alpha) gene expressions induced by LPS (Chavali, Zhong, & Forse, 1998).Fatty acids reduce a translocation of NF-kappaB subunit and NF-kappaB-dependent transcriptional activity. The activation of NF-kappaB is inhibited by prevention of the degradation of inhibitory factor-kappa B alpha (Simopoulos, 2002). They also inhibit LPS-induced phosphorylation of mitogen-activated protein kinases (MAPKs). Sesame oil (SO) is rich in both polyunsaturated fatty acids (PUFA) and monounsaturated fatty acids (MUFA), and has been shown to reduce high blood pressure and lower the amount of medication required to control hypertension (Narasimhulu et al., 2018a; Sankar et al., 2005) in humans. Therefore, the present study was undertaken for the first time with an objective to carry out a complete investigation of the compositions of fatty acids of sesame oil and correlate their role in its reported antioxidant and anti-inflammatory activity. The chromatogram of FAME analysis of sesame oil is shown on the **Supplementary Figure S1**and **Supplementary Table S1**includes the identified constituents with their retention time, quantity, peak height, area(%) of each constituent identified of the collected Bangladeshi sesame oil. The results demonstrated that a total of 9 major fatty acid components (namely octanoic acid, decanoic acid, lauric acid, myristic acid, palmitic acid, oleatic acid, linoleatic acid, arachidonic acid and behenic acid)were identified in FAME analysis from the samples by using GC. Among the identified constituents all were reported before from SO of Indian sample (Crews et al., 2006; Mondal, Bhat, & Srivastava, 2010; Sengupta & Roychoudhury, 1976; Shah, 2016).

3.3 MOLECULAR DOCKING STUDY OF IDENTIFIED COMPOUNDS IN SESAME OIL

3.3.1 Role of COX-2 in molecular docking

Cyclooxygenase (COX), officially known as prostaglandin-endoperoxide synthase (PTGS), is an enzyme that is responsible for formation of prostanoids, including thromboxane and prostaglandins such as prostacyclin, from arachidonic acid (Litalien & Beaulieu, 2011). The two isozymes of COX involved in prostaglandin biosynthesis are COX-1 and COX-2. Arachidonic acid (AA) Pathway is a major component of inflammatory pathway (Kuehl & Egan, 1980). COX-2 is increasingly expressed during inflammatory conditions by pro-inflammatory molecules such as IL-1, TNF- α , LPS (Beuck, 1999; Kumar, Aggarwal, Saini, Kaushik, & Prakash, 2016; Ryn, Trummlitz, & Pairet, 2000) and its expression is absent or low in healthy individuals (Dinarello, 2010; Zhang et al., 2015). The target for anti-inflammatory drugs is cyclooxygenase (COX), a rate-limiting enzyme involved in the conversion of arachidonic acid into inflammatory prostaglandins. Literature study of SO showed that the anti-inflammatory activity of SO and its identified constituents via a number of molecular mechanism including inhibiting the COX-2. However, there is no report found on the role of each identified constituents of SO in the molecular binding with COX-2.

3.3.2 Analysis and visualization of docking study of sesame oil compounds against COX-2

The initial homology model of human COX-2 enzyme was selected based on the lowest dope profile and used for the docking showed on **Figure 4**. The sequence alignment is given in the **Supplementary Figure S2**. The Ramachandran plot analysis, Z-score and knowledge based energy are presented in **Supplementary Figure S3**. The homo-dimer model of COX-2 was generated from its monomeric units using GalaxyGemini following energy minimization step.

The arachidonic acid binding pocket was identified with the following amino acid residues: Ala513, Val335, Gly512, Ser516, Tyr341, Phe343, Val102, Met99, Leu103, Val509, Leu338, Tyr371, and Ser339 presented in **Supplementary Figure S4**. Nine ligands obtained from GC analysis and other twenty one ligands were docked into the active site of the human COX-2 homology model and analyzed in the **Supplementary Figures S5-S32** and the results are presented in **Table 5**. Nineteen out of thirty one compounds showed higher affinity for COX-2 than that of arachidonic acid (-6.8 kcal/mol).

Molecular docking studies with the identified ligands against human COX-2 enzyme revealed that Stigmasterol, Δ 5-Avenasterol and Δ 7-Avenasterol shows the best binding affinity - 10.7kcal/mol for COX-2 presented in **Figure 5 - 6**.

Among the 30 phytoconstituents, 12 compounds interacted with a number of amino acid residues (Ala513, Val335, Gly512, Ser516, Tyr341, Phe343, Val102, Met99, Leu103, Val509, Leu338, Tyr371, Ser339) associated with the binding of COX-2 at the catalytic site, with binding affinity ranging between -5.4 and -9.6 kcal/mol. Among these 12 compounds, Behenic acid showed lower binding affinity (-6.2 kcal/mol) than that of sesamol, pinoselin and vitamin K phylloquinone, but with greater number (13) of interaction with the amino acid residues involved in the binding of COX-2 through hydrophobic interactions. Vitamin K (phylloquinone) interacted with Phe343, Ser339, Leu517, Val102, Leu103, Met99 and Val335 with an affinity of -9.6 kcal/mol. Eicosenoic acid interacted with 12 amino acid residues with an affinity of -6.2 kcal/mol. Pinoselin also exhibited promising results in this computational study due to its binding with considerable interactions with amino acid residues (11 amino acids) around the catalytic site as well as with significant binding affinity (8.4 kcal/mol). Stearic acid (Binding affinity: -5.9 kcal/mol), Palmitic acid (Binding affinity: -5.4kcal/mol), Tocopherol (Binding affinity: -7.1 kcal/mol) and Vitamin A (Retinol) (Binding affinity: -7.4 kcal/mol) showed interaction with ten, nine, six and five amino acid residues, respectively. The other 18 compounds also showed high affinity for COX-2 enzyme but the amino acid residues were not involved in the binding of COX-2. Among the 18 compounds Sesamol, Campesterol, Δ 7-Stigmasterol, Beta-sitosterol, Sesamol and sesamol showed binding affinity -10.6 kcal/mol, -10.5 kcal/mol, -10.5 kcal/mol, -10.2 kcal/mol, -10.4 Kcal/mol and -9.6 Kcal/mol respectively. In this study, all values are negative meaning binding will be spontaneous. The results of this docking study confirmed that the binding potential of the reported constituents of sesame oil with their reported significant COX-2 inhibition activity and have a major role in their anti-inflammatory activity. The reported components of sesame oil were the competitive inhibitors of cyclooxygenase (COX), the enzyme which mediates the bioconversion of arachidonic acid to inflammatory prostaglandins (PGs).

4.CONCLUSION

In this review we have considered antioxidant and anti-inflammatory research on SO over the last decade and a total of 87 studies were found which reported that sesame oil possesses prominent antioxidant and anti-inflammatory effect on various test systems. The antioxidant effect of SO and its constituents were exert through reduction of lipid peroxidation, superoxide and nitric oxide production as well as increased antioxidative enzyme (SOD, CAT, GSH) level which has strong correlation with its anti-inflammatory action. The anti-inflammatory action of SO and its constituents were due to their down regulation of COX and PGE2 as well as inhibition of pro-inflammatory cytokines. COX inhibition was a major pathway for its anti-inflammatory activity which further confirmed through the molecular docking study. Fatty acid profiling of SO identified known 9 fatty acid which also showed molecular binding with COX-2 to some extent as a confirmation of its contribution in anti-inflammatory activity of SO. However, sesamol, sesamin, sesaminol and sesamolol are the main active principle of SO that are responsible for its antioxidant and anti-inflammatory activity as we have showed in this review. Overall findings suggests that although a number of research on SO and its constituents have conducted over the last decade are mainly *in-vitro* and *in-vivo* studies and data related preclinical and clinical trials remain very low. Therefore, more clinical and pharmacokinetic investigation on SO will be needed for future use as a therapeutic agent. In summary, this review reveals that SO can be an important traditional medicine in the treatment of different inflammation.

CONFLICT OF INTEREST

None

References:

Abdel-Daim, M.M., Taha, R., Ghazy, E.W., & El-Sayed, Y.S. (2015). Synergistic ameliorative effects of sesame oil and alpha-lipoic acid against subacute diazinon toxicity in rats: hematological, biochemical, and antioxidant studies. *Canadian Journal of Physiology and Pharmacology*, 94(1), 81-88. <https://doi.org/10.1139/cjpp-2015-0131>

- Abdou, H.M., Hussien, H.M., & Yousef, M.I. (2012). Deleterious effects of cypermethrin on rat liver and kidney: Protective role of sesame oil. *Journal of Environmental Science and Health, Part B*, 47(4), 306-314. <https://doi.org/10.1080/03601234.2012.640913>
- Ahmad, S., ElSherbiny, N.M., Jamal, M.S., Alzahrani, F.A., Haque, R., Khan, R., . . . Bhatia, K. (2016). Anti-inflammatory role of sesamin in STZ induced mice model of diabetic retinopathy. *Journal of Neuroimmunology*, 295, 47-53. <https://doi.org/10.1016/j.jneuroim.2016.04.002>
- Ahmad, S., Khan, M.B., Hoda, M.N., Bhatia, K., Haque, R., Fazili, I.S., . . . Katare, D.P. (2012). Neuroprotective effect of sesame seed oil in 6-hydroxydopamine induced neurotoxicity in mice model: cellular, biochemical and neurochemical evidence. *Neurochemical Research*, 37(3), 516-526. <https://doi.org/10.1007/s11064-011-0638-4>
- Ahmad, S., Yousuf, S., Ishrat, T., Khan, M.B., Bhatia, K., Fazli, I.S., . . . Islam, F. (2006). Effect of dietary sesame oil as antioxidant on brain hippocampus of rat in focal cerebral ischemia. *Life Sciences*, 79(20), 1921-1928. <https://doi.org/10.1016/j.lfs.2006.06.017>
- Ali, Z.Y., Atia, H., El Allawy, R.M., & Alla, S.M.A. (2017). Anti-hepatotoxic and synergistic effects of sesame oil with methotrexate in adjuvant-induced arthritis. *Egyptian Journal of Biochemistry & Molecular Biology*, 35(1&2), 77-92.
- Alipoor, B., Haghhighian, M.K., Sadat, B.E., & Asghari, M. (2012). Effect of sesame seed on lipid profile and redox status in hyperlipidemic patients. *International Journal of Food Sciences and Nutrition*, 63(6), 674-678. <https://doi.org/10.3109/09637486.2011.652077>
- Arumugam, P., & Ramesh, S. (2011). Protective effects of sesame oil on 4-NQO-induced oxidative DNA damage and lipid peroxidation in rats. *Drug and Chemical Toxicology*, 34(2), 116-119. <https://doi.org/10.3109/01480541003782310>
- Azab, A. (2014). Hepatoprotective effect of sesame oil against lead induced liver damage in albino mice: Histological and biochemical studies. *American Journal of BioScience*, 2(6-2), 1-11. <https://doi.org/10.11648/j.ajbio.s.20140202.11>
- Balkrishna, A. (2008). *Secrets of Indian herbs for good health* (1st edn). Divya Prakashan: Uttarakhanda, India.
- Baluchnejadmojarad, T., Mansouri, M., Ghalami, J., Mokhtari, Z., & Roghani, M. (2017). Sesamin imparts neuroprotection against intrastriatal 6-hydroxydopamine toxicity by inhibition of astroglial activation, apoptosis, and oxidative stress. *Biomedicine & Pharmacotherapy*, 88, 754-761. <https://doi.org/10.1016/j.biopha.2017.01.123>
- Bedigian, D. (2004). History and lore of sesame in Southwest Asia. *Economic Botany*, 58(3), 329-353. [https://doi.org/10.1663/0013-0001\(2004\)058\[0329:AR\]2.0.CO;2](https://doi.org/10.1663/0013-0001(2004)058[0329:AR]2.0.CO;2)
- Beuck, M. (1999). Nonsteroidal antiinflammatory drugs: a new generation of cyclooxygenase inhibitors. *Angewandte Chemie International Edition*, 38(5), 631-633. [https://doi.org/10.1002/\(SICI\)1521-3773\(19990301\)38:5%3C631::AID-ANIE631%3E3.0.CO;2-S](https://doi.org/10.1002/(SICI)1521-3773(19990301)38:5%3C631::AID-ANIE631%3E3.0.CO;2-S)
- Bhaskaran, S., Santanam, N., Penumetcha, M., & Parthasarathy, S. (2006). Inhibition of atherosclerosis in low-density lipoprotein receptor-negative mice by sesame oil. *Journal of Medicinal Food*, 9(4), 487-490. <https://doi.org/10.1089/jmf.2006.9.487>
- Bopitiya, D., & Madhujith, T. (2013). Antioxidant activity and total phenolic content of sesame (*Sesamum indicum* L.) seed oil. *Tropical Agricultural Research*, 24(3), 296-302.
- Bournival, J., Plouffe, M., Renaud, J., Provencher, C., & Martinoli, M.-G. (2012). Quercetin and sesamin protect dopaminergic cells from MPP. *Oxidative Medicine and Cellular Longevity*, 2012, 11. <https://doi.org/10.1155/2012/921941>

- Burdock, G.A. (2016). *Fenaroli's handbook of flavor ingredients* (6th edn). CRC press: Boca Raton, FL.
- Cao, W., Dai, M., Wang, X., Yuan, F., Chen, F., & Zhang, W. (2013). Protective effect of sesaminol from *Sesamum indicum* Linn. against oxidative damage in PC12 cells. *Cell Biochemistry and Function*, 31(7), 560-565. <https://doi.org/10.1002/cbf.2935>
- Carrasco-Pancorbo, A., Cerretani, L., Bendini, A., Segura-Carretero, A., Del Carlo, M., Gallina-Toschi, T., . . . Fernandez-Gutierrez, A. (2005). Evaluation of the antioxidant capacity of individual phenolic compounds in virgin olive oil. *Journal of Agricultural and Food Chemistry*, 53(23), 8918-8925. <https://doi.org/10.1021/jf0515680>
- Cengiz, N., Kavak, S., Güzel, A., Özbek, H., Bektaş, H., Him, A., . . . Balahoroğlu, R. (2013). Investigation of the hepatoprotective effects of Sesame (*Sesamum indicum* L.) in carbon tetrachloride-induced liver toxicity. *The Journal of Membrane Biology*, 246(1), 1-6. <https://doi.org/10.1007/s00232-012-9494-7>
- Chandrasekaran, V.R.M., Chien, S.-P., Hsu, D.-Z., Chang, Y.-C., & Liu, M.-Y. (2010). Effects of sesame oil against after the onset of acetaminophen-induced acute hepatic injury in rats. *Journal of Parenteral and Enteral Nutrition*, 34(5), 567-573. <https://doi.org/10.1177/0148607110362584>
- Chavali, S., Zhong, W., & Forse, R. (1998). Dietary α -linolenic acid increases TNF- α , and decreases IL-6, IL-10 in response to LPS: effects of sesamin on the Δ -5 desaturation of ω 6 and ω 3 fatty acids in mice. *Prostaglandins, Leukotrienes and Essential Fatty Acids*, 58(3), 185-191. [https://doi.org/10.1016/S0952-3278\(98\)90112-0](https://doi.org/10.1016/S0952-3278(98)90112-0)
- Chavali, S.R., Utsunomiya, T., & Forse, R.A. (2001). Increased survival after cecal ligation and puncture in mice consuming diets enriched with sesame seed oil. *Critical Care Medicine*, 29(1), 140-143. <https://doi.org/10.1097/00003246-200101000-00028>
- Chen, P.R., Chien, K.L., Su, T.C., Chang, C.J., Liu, T.-L., Cheng, H., & Tsai, C. (2005). Dietary sesame reduces serum cholesterol and enhances antioxidant capacity in hypercholesterolemia. *Nutrition Research*, 25(6), 559-567. <https://doi.org/10.1016/j.nutres.2005.05.007>
- Chen, X., Ying, X., Chen, L., Zhang, W., & Zhang, Y. (2015). Protective effects of sesamin on liver fibrosis through antioxidative and anti-inflammatory activities in rats. *Immunopharmacology and Immunotoxicology*, 37(5), 465-472. <https://doi.org/10.3109/08923973.2015.1085064>
- Chiang, H.-M., Chang, H., Yao, P.-W., Chen, Y.-S., Jeng, K.-C., Wang, J.-S., & Hou, C.-W. (2014). Sesamin reduces acute hepatic injury induced by lead coupled with lipopolysaccharide. *Journal of the Chinese Medical Association*, 77(5), 227-233. <https://doi.org/10.1016/j.jcma.2014.02.010>
- Chopra, K., Tiwari, V., Arora, V., & Kuhad, A. (2010). Sesamol suppresses neuro-inflammatory cascade in experimental model of diabetic neuropathy. *The Journal of Pain*, 11(10), 950-957. <https://doi.org/10.1016/j.jpain.2010.01.006>
- Chopra, R.N., & Nayar, S.L. (1956). *Glossary of Indian medicinal plants* (3rd edn). Council of Scientific And Industrial Research: New Delhi, India.
- Crews, C., Hough, P., Brereton, P., Godward, J., Lees, M., Guiet, S., & Winkelmann, W. (2006). Quantitation of the main constituents of some authentic sesame seed oils of different origin. *Journal of Agricultural and Food Chemistry*, 54(17), 6266-6270. <https://doi.org/10.1021/jf0603578>

- Deme, P., Narasimhulu, C.A., & Parthasarathy, S. (2018). Identification and evaluation of anti-inflammatory properties of aqueous components extracted from sesame (*Sesamum indicum*) oil. *Journal of Chromatography B*, 1087-1088, 61-69. <https://doi.org/10.1016/j.jchromb.2018.04.029>
- Dhar, P., Chattopadhyaya, K., Bhattacharyya, D., Biswas, A., Roy, B., & Ghosh, S. (2007). Ameliorative influence of sesame lignans on lipid profile and lipid peroxidation in induced diabetic rats. *Journal of Agricultural and Food Chemistry*, 55(14), 5875-5880. <https://doi.org/10.1021/jf063721b>
- Dinarello, C.A. (2010). Anti-inflammatory agents: present and future. *Cell*, 140(6), 935-950. <https://doi.org/10.1016/j.cell.2010.02.043>
- Duarte, A.R., Chenet, A.L., de Almeida, F.J.S., Andrade, C.M.B., & de Oliveira, M.R. (2018). The inhibition of heme oxygenase-1 (HO-1) abolishes the mitochondrial protection induced by sesamol in LPS-treated RAW 264.7 cells. *Chemico-biological Interactions*, 296, 171-178. <https://doi.org/10.1016/j.cbi.2018.09.012>
- Ghani, A. (1998). *Medicinal plants of Bangladesh: chemical constituents and uses*. Asiatic society of Bangladesh.
- Gharby, S., Harhar, H., Bouzoubaa, Z., Asdadi, A., El Yadini, A., & Charrouf, Z. (2017). Chemical characterization and oxidative stability of seeds and oil of sesame grown in Morocco. *Journal of the Saudi Society of Agricultural Sciences*, 16(2), 105-111. <https://doi.org/10.1016/j.jssas.2015.03.004>
- Grouleff, J., Irudayam, S.J., Skeby, K.K., & Schiøtt, B. (2015). The influence of cholesterol on membrane protein structure, function, and dynamics studied by molecular dynamics simulations. *Biochimica et Biophysica Acta (BBA)-Biomembranes*, 1848(9), 1783-1795. <https://doi.org/10.1016/j.bbamem.2015.03.029>
- Guimarães, R.d.C.A., Macedo, M.L.R., Munhoz, C.L., Filiu, W., Viana, L.H., Nozaki, V.T., & Hiane, P.A. (2013). Sesame and flaxseed oil: nutritional quality and effects on serum lipids and glucose in rats. *Food Science and Technology*, 33(1), 209-217. <https://dx.doi.org/10.1590/S0101-20612013005000029>
- Gülcan, A., BÜLBÜL, A., ÖZDEMİR, M., & EROL, A.Y.G. (2015). The investigation of the effects of omega 3 fatty acids and Sesame oil on Cyclosporine-A induced immunosuppressive and oxidant effects in rats. *Kocatepe Veterinary Journal*, 8(1), 45-50.
- Gunstone, F.D. (2006). Minor specialty oils. In *Nutraceutical and specialty lipids and their co-products*, Shahidi F (Ed). CRC Press: Boca Raton, FL; 99-133.
- Helli, B., Mowla, K., Mohammadshahi, M., & Jalali, M. (2016). Effect of sesamin supplementation on cardiovascular risk factors in women with rheumatoid arthritis. *Journal of the American College of Nutrition*, 35(4), 300-307. <https://doi.org/10.1080/07315724.2015.1005198>
- Hemalatha, S. (2007). Sesame lignans enhance the thermal stability of edible vegetable oils. *Food Chemistry*, 105(3), 1076-1085. <https://doi.org/10.1016/j.foodchem.2007.05.023>
- Hemalatha, S., & Raghunath, M. (2004). Dietary sesame (*Sesamum indicum* cultivar Linn) oil inhibits iron-induced oxidative stress in rats. *The British Journal of Nutrition*, 92(4), 581-587. <https://doi.org/10.1079/BJN20041239>
- Hemshekhhar, M., Thushara, R.M., Jnaneshwari, S., Devaraja, S., Kemparaju, K., & Girish, K.S. (2013). Attenuation of adjuvant-induced arthritis by dietary sesamol via modulation of inflammatory mediators, extracellular matrix degrading enzymes and antioxidant status.

- European Journal of Nutrition*, 52(7), 1787-1799. <https://doi.org/10.1007/s00394-012-0482-6>
- Hou, C.-W., Chang, S.-Y., & Jeng, K.-C. (2015). Protective effect of a sesamin derivative, 3-bis (3-methoxybenzyl) butane-1, 4-diol on A β -stressed PC12 cells. *Archives of Pharmacal Research*, 38(4), 543-548. <https://doi.org/10.1007/s12272-014-0426-1>
- Hou, R.C.-W., Chen, H.-L., Tzen, J.T., & Jeng, K.-C.G. (2003). Effect of sesame antioxidants on LPS-induced NO production by BV2 microglial cells. *Neuroreport*, 14(14), 1815-1819. <https://doi.org/10.1097/00001756-200310060-00011>
- Hsieh, P.F., Hou, C.-W., Yao, P.-W., Wu, S.-P., Peng, Y.-F., Shen, M.-L., . . . Jeng, K.-C. (2011). Sesamin ameliorates oxidative stress and mortality in kainic acid-induced status epilepticus by inhibition of MAPK and COX-2 activation. *Journal of Neuroinflammation*, 8(1), 10. <https://doi.org/10.1186/1742-2094-8-57>
- Hsu, C.-C., Huang, H.-C., Wu, P.-T., Tai, T.-W., & Jou, I.-M. (2016a). Sesame oil improves functional recovery by attenuating nerve oxidative stress in a mouse model of acute peripheral nerve injury: role of Nrf-2. *The Journal of Nutritional Biochemistry*, 38, 102-106. <https://doi.org/10.1016/j.jnutbio.2016.09.003>
- Hsu, D.-Z., Chen, S.-J., Chu, P.-Y., & Liu, M.-Y. (2013). Therapeutic effects of sesame oil on monosodium urate crystal-induced acute inflammatory response in rats. *SpringerPlus*, 2(1), 659. <https://doi.org/10.1186/2193-1801-2-659>
- Hsu, D.-Z., Chiang, P.-J., Chien, S.-P., Huang, B.-M., & Liu, M.-Y. (2004a). Parenteral sesame oil attenuates oxidative stress after endotoxin intoxication in rats. *Toxicology*, 196(1-2), 147-153. <https://doi.org/10.1016/j.tox.2003.12.001>
- Hsu, D.-Z., Chien, S.-P., Li, Y.-H., Chuang, Y.-C., Chang, Y.-C., & Liu, M.-Y. (2008a). Sesame oil attenuates hepatic lipid peroxidation by inhibiting nitric oxide and superoxide anion generation in septic rats. *Journal of Parenteral and Enteral Nutrition*, 32(2), 154-159. <https://doi.org/10.1177/0148607108314766>
- Hsu, D.-Z., Chu, P.-Y., Chandrasekaran, V.R.M., & Liu, M.-Y. (2009a). Sesame lignan sesamol protects against aspirin-induced gastric mucosal damage in rats. *Journal of Functional Foods*, 1(4), 349-355. <https://doi.org/10.1016/j.jff.2009.07.003>
- Hsu, D.-Z., Chu, P.-Y., & Jou, I.-M. (2016b). Daily sesame oil supplement attenuates joint pain by inhibiting muscular oxidative stress in osteoarthritis rat model. *The Journal of Nutritional Biochemistry*, 29, 36-40. <https://doi.org/10.1016/j.jnutbio.2015.10.007>
- Hsu, D.-Z., Chu, P.-Y., & Jou, I.-M. (2016c). Enteral sesame oil therapeutically relieves disease severity in rat experimental osteoarthritis. *Food & Nutrition Research*, 60(1), 29807. <https://doi.org/10.3402/fnr.v60.29807>
- Hsu, D.-Z., Chu, P.-Y., Li, Y.-H., & Liu, M.-Y. (2008b). Sesamol attenuates diclofenac-induced acute gastric mucosal injury via its cyclooxygenase-independent antioxidative effect in rats. *Shock*, 30(4), 456-462. <https://doi.org/10.1097/SHK.0b013e3181672185>
- Hsu, D.-Z., Li, Y.-H., Chien, S.-P., & Liu, M.-Y. (2004b). Effects of sesame oil on oxidative stress and hepatic injury after cecal ligation and puncture in rats. *Shock*, 21(5), 466-469. <https://doi.org/10.1097/01.shk.0000124028.92586.13>
- Hsu, D.-Z., Li, Y.-H., Chu, P.-Y., Periasamy, S., & Liu, M.-Y. (2011). Sesame oil prevents acute kidney injury induced by the synergistic action of aminoglycoside and iodinated contrast in rats. *Antimicrobial Agents and Chemotherapy*, 55(6), 2532-2536. <https://doi.org/10.1128/AAC.01597-10>

- Hsu, D.-Z., & Liu, M.-Y. (2004a). Effects of sesame oil on oxidative stress after the onset of sepsis in rats. *Shock*, 22(6), 582-585. <https://doi.org/10.1097/01.shk.0000135254.21699.45>
- Hsu, D.-Z., & Liu, M.-Y. (2004b). Sesame oil protects against lipopolysaccharide-stimulated oxidative stress in rats. *Critical Care Medicine*, 32(1), 227-231. <https://doi.org/10.1097/01.CCM.0000104947.16669.29>
- Hsu, D.-Z., Su, S.-B., Chien, S.-P., Chiang, P.-J., Li, Y.-H., Lo, Y.-J., & Liu, M.-Y. (2005). Effect of sesame oil on oxidative-stress-associated renal injury in endotoxemic rats: involvement of nitric oxide and proinflammatory cytokines. *Shock*, 24(3), 276-280. <https://doi.org/10.1097/01.shk.0000172366.73881.c7>
- Hsu, D.Z., Chu, P.Y., & Liu, M.Y. (2009b). Effect of sesame oil on acidified ethanol-induced gastric mucosal injury in rats. *Journal of Parenteral and Enteral Nutrition*, 33(4), 423-427. <https://doi.org/10.1177/0148607108327046>
- Hung, C.-T., Chen, L.-D., & Hou, C.-W. (2017). Neuroprotection of a sesamin derivative, 1, 2-bis [(3-methoxyphenyl) methyl] ethane-1, 2-dicarboxylic acid (MMEDA) against ischemic and hypoxic neuronal injury. *Iranian Journal of Basic Medical Sciences*, 20(12), 1324. <https://doi.org/10.22038/2FIJBMS.2017.9543>
- Hwang, L.S. (2005). Sesame oil. In *Bailey's industrial oil and fat products*, Shahidi F (Ed). John Wiley & Sons: Hoboken, New Jersey; 547-552.
- Ismail, M., Hasan, H., El-Orfali, Y., Ismail, H., & Khawaja, G. (2018). Anti-inflammatory, antioxidative, and hepatoprotective effects of trans 9-tetrahydrocannabinol/sesame oil on adjuvant-induced arthritis in rats. *Evidence-Based Complementary and Alternative Medicine*, 2018, 13. <https://doi.org/10.1155/2018/9365464>
- Jamieson, G.S., & Baughman, W.F. (1924). The chemical composition of sesame oil. *Journal of the American Chemical Society*, 46(3), 775-778. <https://doi.org/10.1021/ja01668a032>
- Jeng, K., & Hou, R. (2005). Sesamin and sesamolin: nature's therapeutic lignans. *Current Enzyme Inhibition*, 1(1), 11-20. <https://doi.org/10.2174/1573408052952748>
- Jnaneshwari, S., Hemshekhar, M., Thushara, R., Shanmuga Sundaram, M., Sebastin Santhosh, M., Sunitha, K., . . . Girish, K. (2014). Sesamol ameliorates cyclophosphamide-induced hepatotoxicity by modulating oxidative stress and inflammatory mediators. *Anti-Cancer Agents in Medicinal Chemistry*, 14(7), 975-983.
- Kang, M.-H., Naito, M., Sakai, K., Uchida, K., & Osawa, T. (1999). Mode of action of sesame lignans in protecting lowdensity lipoprotein against oxidative damage in vitro. *Life Sciences*, 66(2), 161-171. [https://doi.org/10.1016/S0024-3205\(99\)00574-3](https://doi.org/10.1016/S0024-3205(99)00574-3)
- Kang, M.-H., Naito, M., Tsujihara, N., & Osawa, T. (1998). Sesamolin inhibits lipid peroxidation in rat liver and kidney. *The Journal of Nutrition*, 128(6), 1018-1022. <https://doi.org/10.1093/jn/128.6.1018>
- Kanu, P.J., Bahsoon, J.Z., Kanu, J.B., & Kandeh, J.B. (2010). Nutraceutical importance of sesame seed and oil: A review of the contribution of their lignans. *Sierra Leone Journal of Biomedical Research*, 2(1), 4-16. <http://dx.doi.org/10.4314/sljbr.v2i1.56583>
- Kapadia, G.J., Azuine, M.A., Tokuda, H., Takasaki, M., Mukainaka, T., Konoshima, T., & Nishino, H. (2002). Chemopreventive effect of resveratrol, sesamol, sesame oil and sunflower oil in the Epstein-Barr virus early antigen activation assay and the mouse skin two-stage carcinogenesis. *Pharmacological Research*, 45(6), 499-505. <https://doi.org/10.1006/phrs.2002.0992>

- Kaur, I.P., & Saini, A. (2000). Sesamol exhibits antimutagenic activity against oxygen species mediated mutagenicity. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*, 470(1), 71-76. [https://doi.org/10.1016/S1383-5718\(00\)00096-6](https://doi.org/10.1016/S1383-5718(00)00096-6)
- Khamphio, M., Barusrux, S., & Weerapreeyakul, N. (2016). Sesamol induces mitochondrial apoptosis pathway in HCT116 human colon cancer cells via pro-oxidant effect. *Life Sciences*, 158, 46-56. <https://doi.org/10.1016/j.lfs.2016.06.017>
- Khan, M.M., Ishrat, T., Ahmad, A., Hoda, M.N., Khan, M.B., Khuwaja, G., . . . Ahmad, S. (2010). Sesamin attenuates behavioral, biochemical and histological alterations induced by reversible middle cerebral artery occlusion in the rats. *Chemico-biological Interactions*, 183(1), 255-263. <https://doi.org/10.1016/j.cbi.2009.10.003>
- Khaneshi, F., Nasrolahi, O., Azizi, S., & Nejati, V. (2013). Sesame effects on testicular damage in streptozotocin-induced diabetes rats. *Avicenna Journal of Phytomedicine*, 3(4), 347.
- Konc, J., & Janežič, D. (2014). ProBiS-ligands: a web server for prediction of ligands by examination of protein binding sites. *Nucleic Acids Research*, 42(W1), W215-W220. <https://doi.org/10.1093/nar/gku460>
- Kong, P., Chen, G., Jiang, A., Wang, Y., Song, C., Zhuang, J., . . . Yan, J. (2016). Sesamin inhibits IL-1 β -stimulated inflammatory response in human osteoarthritis chondrocytes by activating Nrf2 signaling pathway. *Oncotarget*, 7(50), 83720-83726. <https://doi.org/10.18632/oncotarget.13360>
- Kuehl, F.A., & Egan, R.W. (1980). Prostaglandins, arachidonic acid, and inflammation. *Science*, 210(4473), 978-984. <https://doi.org/10.1126/science.6254151>
- Kumar, A., Aggarwal, S., Saini, D., Kaushik, D., & Prakash, O. (2016). Synthesis and anti-inflammatory, analgesic activities of some new 6-aryl-4-(4-hydroxy-6-methyl-2-oxo-2H-pyran-3-yl)-2-S-benzylthopyrimidines. *Indian Journal of Heterocyclic Chemistry*, 25(3-4), 317-324.
- Kuo, P.-C., Lin, M.-C., Chen, G.-F., Yiu, T.-J., & Tzen, J.T. (2011). Identification of methanol-soluble compounds in sesame and evaluation of antioxidant potential of its lignans. *Journal of Agricultural and Food Chemistry*, 59(7), 3214-3219. <https://doi.org/10.1021/jf104311g>
- Lahaie-Collins, V., Bournival, J., Plouffe, M., Carange, J., & Martinoli, M.-G. (2008). Sesamin modulates tyrosine hydroxylase, superoxide dismutase, catalase, inducible no synthase and interleukin-6 expression in dopaminergic cells under mpp. *Oxidative Medicine and Cellular Longevity*, 1(1), 54-62.
- Lahorkar, P., Ramitha, K., Bansal, V., & Narayana, D.A. (2009). A comparative evaluation of medicated oils prepared using ayurvedic and modified processes. *Indian Journal of Pharmaceutical Sciences*, 71(6), 656. <https://doi.org/10.4103/2F0250-474X.59548>
- Lee, S.Y., Son, D.J., Lee, Y.K., Lee, J.W., Lee, H.J., Yun, Y.W., . . . Hong, J.T. (2006). Inhibitory effect of sesaminol glucosides on lipopolysaccharide-induced NF- κ B activation and target gene expression in cultured rat astrocytes. *Neuroscience Research*, 56(2), 204-212. <https://doi.org/10.1016/j.neures.2006.06.005>
- Lee, W.-J., Ou, H.-C., Wu, C.-M., Lee, I.-t., Lin, S.-Y., Lin, L.-Y., . . . Sheu, W.H.-H. (2009). Sesamin mitigates inflammation and oxidative stress in endothelial cells exposed to oxidized low-density lipoprotein. *Journal of Agricultural and Food Chemistry*, 57(23), 11406-11417. <https://doi.org/10.1021/jf902876p>

- Lengauer, T., & Rarey, M. (1996). Computational methods for biomolecular docking. *Current Opinion in Structural Biology*, 6(3), 402-406. [https://doi.org/10.1016/S0959-440X\(96\)80061-3](https://doi.org/10.1016/S0959-440X(96)80061-3)
- Lin, C.-H., Shen, M.-L., Zhou, N., Lee, C.-C., Kao, S.-T., & Wu, D.C. (2014). Protective effects of the polyphenol sesamin on allergen-induced TH2 responses and airway inflammation in mice. *PloS One*, 9(4), e96091. <https://doi.org/10.1371/journal.pone.0096091>
- Litalien, C., & Beaulieu, P. (2011). Molecular Mechanisms of Drug Actions: From Receptors to Effectors. In *Pediatric Critical Care* Fuhrman BP & Zimmerman JJ (Ed). Elsevier: Philadelphia, PA; 1553-1568.
- Liu, C.-T., & Liu, M.-Y. (2017). Daily sesame oil supplementation attenuates local renin-angiotensin system via inhibiting MAPK activation and oxidative stress in cardiac hypertrophy. *The Journal of Nutritional Biochemistry*, 42, 108-116. <https://doi.org/10.1016/j.jnutbio.2016.05.006>
- Lovell, S.C., Davis, I.W., Arendall III, W.B., De Bakker, P.I., Word, J.M., Prisant, M.G., . . . Richardson, D.C. (2003). Structure validation by C α geometry: ϕ , ψ and C β deviation. *Proteins: Structure, Function, and Bioinformatics*, 50(3), 437-450. <https://doi.org/10.1002/prot.10286>
- Lv, D., Zhu, C.-Q., & Liu, L. (2015). Sesamin ameliorates oxidative liver injury induced by carbon tetrachloride in rat. *International Journal of Clinical and Experimental Pathology*, 8(5), 5733.
- Ma, J.-Q., Ding, J., Zhang, L., & Liu, C.-M. (2014). Hepatoprotective properties of sesamin against CCl₄ induced oxidative stress-mediated apoptosis in mice via JNK pathway. *Food and Chemical Toxicology*, 64, 41-48. <https://doi.org/10.1016/j.fct.2013.11.017>
- Marzook, E.A., El Moneim, A.E.A., & Elhadary, A.A. (2014). Protective role of sesame oil against mobile base station-induced oxidative stress. *Journal of Radiation Research and Applied Sciences*, 7(1), 1-6. <https://doi.org/10.1016/j.jrras.2013.10.010>
- Matsumura, Y., Kita, S., Morimoto, S., Akimoto, K., Furuya, M., Oka, N., & Tanaka, T. (1995). Antihypertensive effect of sesamin. I. Protection against deoxycorticosterone acetate-salt-induced hypertension and cardiovascular hypertrophy. *Biological and Pharmaceutical Bulletin*, 18(7), 1016-1019. <https://doi.org/10.1248/bpb.18.1016>
- Miguel, M.G. (2010). Antioxidant and anti-inflammatory activities of essential oils: a short review. *Molecules*, 15(12), 9252-9287. <https://doi.org/10.3390/molecules15129252>
- Milder, I.E., Arts, I.C., van de Putte, B., Venema, D.P., & Hollman, P.C. (2005). Lignan contents of Dutch plant foods: a database including lariciresinol, pinoresinol, secoisolariciresinol and matairesinol. *British Journal of Nutrition*, 93(3), 393-402. <https://doi.org/10.1079/BJN20051371>
- Minioti, K.S., & Georgiou, C.A. (2010). Comparison of different tests used in mapping the Greek virgin olive oil production for the determination of its total antioxidant capacity. *Grasas y Aceites*, 61(1), 45-51. <https://doi.org/10.3989/gya.010508>
- Mondal, N., Bhat, K., & Srivastava, P. (2010). Variation in fatty acid composition in Indian germplasm of sesame. *Journal of the American Oil Chemists' Society*, 87(11), 1263-1269. <https://doi.org/10.1007/s11746-010-1615-9>
- Monteiro, É., Chibli, L., Yamamoto, C., Pereira, M., Vilela, F., Rodarte, M., . . . de Matos Araújo, A. (2014). Antinociceptive and anti-inflammatory activities of the sesame oil and sesamin. *Nutrients*, 6(5), 1931-1944. <https://doi.org/10.3390/nu6051931>

- Mosayebi, G., Ghazavi, A., Salehi, H., Payani, M., & Khazae, M. (2007). Effect of sesame oil on the inhibition of experimental autoimmune encephalomyelitis in C57BL/6 mice. *Pakistan Journal of Biological Sciences*, 10(11), 1790-1796.
- Nakai, M., Harada, M., Nakahara, K., Akimoto, K., Shibata, H., Miki, W., & Kiso, Y. (2003). Novel antioxidative metabolites in rat liver with ingested sesamin. *Journal of Agricultural and Food Chemistry*, 51(6), 1666-1670. <https://doi.org/10.1021/jf0258961>
- Nakano, D., Itoh, C., Ishii, F., Kawanishi, H., Takaoka, M., Kiso, Y., . . . Matsumura, Y. (2003). Effects of sesamin on aortic oxidative stress and endothelial dysfunction in deoxycorticosterone acetate-salt hypertensive rats. *Biological and Pharmaceutical Bulletin*, 26(12), 1701-1705. <https://doi.org/10.1248/bpb.26.1701>
- Nakano, D., Itoh, C., Takaoka, M., Kiso, Y., Tanaka, T., & Matsumura, Y. (2002). Antihypertensive effect of sesamin. IV. Inhibition of vascular superoxide production by sesamin. *Biological and Pharmaceutical Bulletin*, 25(9), 1247-1249. <https://doi.org/10.1248/bpb.25.1247>
- Namiki, M. (1995). The chemistry and physiological functions of sesame. *Food Reviews International*, 11(2), 281-329. <https://doi.org/10.1080/87559129509541043>
- Narasimhulu, C.A., Burge, K.Y., Doomra, M., Riad, A., & Parthasarathy, S. (2018a). Primary prevention of atherosclerosis by pretreatment of low-density lipoprotein receptor knockout mice with sesame oil and its aqueous components. *Scientific Reports*, 8(1), 12270. <https://doi.org/10.1038/s41598-018-29849-x>
- Narasimhulu, C.A., Riad, A., & Parthasarathy, S. (2018b). Sesame oil and an aqueous extract derived from sesame oil enhance regression of preexisting atherosclerotic lesions in low-density lipoprotein receptor knockout mice. *Journal of Medicinal Food*, 21(7), 641-646. <https://doi.org/10.1089/jmf.2018.0032>
- Narasimhulu, C.A., Selvarajan, K., Burge, K.Y., Litvinov, D., Sengupta, B., & Parthasarathy, S. (2016). Water-soluble components of sesame oil reduce inflammation and atherosclerosis. *Journal of Medicinal Food*, 19(7), 629-637. <https://doi.org/10.1089/jmf.2015.0154>
- Periasamy, S., Hsu, D.-Z., Chandrasekaran, V.R.M., & Liu, M.-Y. (2013). Sesame oil accelerates healing of 2, 4, 6-trinitrobenzenesulfonic acid-induced acute colitis by attenuating inflammation and fibrosis. *Journal of Parenteral and Enteral Nutrition*, 37(5), 674-682. <https://doi.org/10.1177/0148607112468768>
- Periasamy, S., Yang, S.-S., Chen, S.-Y., Chang, C.-C., & Liu, M.-Y. (2013). Prophylactic sesame oil attenuates sinusoidal obstruction syndrome by inhibiting matrix metalloproteinase-9 and oxidative stress. *Journal of Parenteral and Enteral Nutrition*, 37(4), 529-537. <https://doi.org/10.1177/0148607112454299>
- Pole, S. (2006). *Ayurvedic medicine: The principles of traditional practice* (1st edn). Elsevier Health Sciences: London, UK.
- Prasad, N.R., Mahesh, T., Menon, V.P., Jeevanram, R., & Pugalendi, K.V. (2005). Photoprotective effect of sesamol on UVB-radiation induced oxidative stress in human blood lymphocytes *in vitro*. *Environmental Toxicology and Pharmacology*, 20(1), 1-5. <https://doi.org/10.1016/j.etap.2004.09.009>
- Prasanthi, K., & Rajini, P. (2005). Fenvalerate-induced oxidative damage in rat tissues and its attenuation by dietary sesame oil. *Food and Chemical Toxicology*, 43(2), 299-306. <https://doi.org/10.1016/j.fct.2004.10.005>

- Rahman, M., Khatun, A., Uddin, S.J., & Shilpi, J.A. (2016). Comparative effect of *Lannea coromandelica* (Houtt.) Merr. leaves and stem barks on acetic acid induced pain model in mice and chromagenic reagents: exploring the analgesic potential and phytochemical groups. *Pharmacologyonline*, 1, 146-152.
- Ramesh, B., Saravanan, R., & Pugalendi, K. (2005). Influence of sesame oil on blood glucose, lipid peroxidation, and antioxidant status in streptozotocin diabetic rats. *Journal of Medicinal Food*, 8(3), 377-381. <https://doi.org/10.1089/jmf.2005.8.377>
- Ren, J., & Chung, S.H. (2007). Anti-inflammatory effect of α -linolenic acid and its mode of action through the inhibition of nitric oxide production and inducible nitric oxide synthase gene expression via NF- κ B and mitogen-activated protein kinase pathways. *Journal of Agricultural and Food Chemistry*, 55(13), 5073-5080. <https://doi.org/10.1021/jf0702693>
- Ricci, A., Groth, D., & Lago, A. (1999). Densidade de plantas, secagem e produção de sementes de gergelim cv. IAC-CHINA. *Revista Brasileira de Sementes*, 21(1), 82-86. <http://doi.org/10.17801/0101-3122/rbs>
- Ryn, J.v., Trummlitz, G., & Pairet, M. (2000). COX-2 selectivity and inflammatory processes. *Current Medicinal Chemistry*, 7(11), 1145-1161. <https://doi.org/10.2174/0929867003374255>
- Saleem, M.T., Chetty, M.C., & Kavimani, S. (2013). Putative antioxidant property of sesame oil in an oxidative stress model of myocardial injury. *Journal of Cardiovascular Disease Research*, 4(3), 177-181. <https://doi.org/10.1016/j.jcdr.2013.07.001>
- Saleem, M.T., Chetty, M.C., & Kavimani, S. (2014). Antioxidants and tumor necrosis factor alpha-inhibiting activity of sesame oil against doxorubicin-induced cardiotoxicity. *Therapeutic Advances in Cardiovascular Disease*, 8(1), 4-11. <https://doi.org/10.1177%2F1753944713516532>
- Saleem, T.M., Basha, S.D., Mahesh, G., Rani, P.S., Kumar, N.S., & Chetty, C.M. (2011). Analgesic, anti-pyretic and anti-inflammatory activity of dietary sesame oil in experimental animal models. *Pharmacologia*, 2(6), 172-177.
- Sankar, D., Ali, A., Sambandam, G., & Rao, R. (2011). Sesame oil exhibits synergistic effect with anti-diabetic medication in patients with type 2 diabetes mellitus. *Clinical Nutrition*, 30(3), 351-358. <https://doi.org/10.1016/j.clnu.2010.11.005>
- Sankar, D., Rao, M.R., Sambandam, G., & Pugalendi, K. (2006a). Effect of sesame oil on diuretics or β -blockers in the modulation of blood pressure, anthropometry, lipid profile, and redox status. *The Yale Journal of Biology and Medicine*, 79(1), 19-26.
- Sankar, D., Rao, M.R., Sambandam, G., & Pugalendi, K. (2006b). A pilot study of open label sesame oil in hypertensive diabetics. *Journal of Medicinal Food*, 9(3), 408-412. <https://doi.org/10.1089/jmf.2006.9.408>
- Sankar, D., Sambandam, G., Rao, M.R., & Pugalendi, K. (2005). Modulation of blood pressure, lipid profiles and redox status in hypertensive patients taking different edible oils. *Clinica Chimica Acta*, 355(1-2), 97-104. <https://doi.org/10.1016/j.cccn.2004.12.009>
- Selvarajan, K., Narasimhulu, C.A., Bapputty, R., & Parthasarathy, S. (2015). Anti-inflammatory and antioxidant activities of the nonlipid (aqueous) components of sesame oil: potential use in atherosclerosis. *Journal of Medicinal Food*, 18(4), 393-402. <https://doi.org/10.1089/jmf.2014.0139>

- Sengupta, A., & Roychoudhury, S. (1976). Triglyceride composition of *Sesamum indicum* seed oil. *Journal of the Science of Food and Agriculture*, 27(2), 165-169. <https://doi.org/10.1002/jsfa.2740270214>
- Shafipour, S.Z., Mosayebi, N., Asgari, F., Atrkarroushan, Z., & Pasdaran, A. (2017). Effect of external use of sesame oil in the prevention of chemotherapy-induced phlebitis in children with acute lymphoblastic leukemia (ALL). *Iranian Journal of Pediatric Hematology and Oncology*, 7(4), 198-206.
- Shah, N. (2016). *Sesamum indicum* (Sesame or Til): Seeds and oil-a historical and scientific evaluation from indian perspective. *Asian Agri-History*, 20(1), 3-19.
- Shahidi, F., Wanasundara, P., & Wanasundara, U. (1997). Changes in edible fats and oils during processing. *Journal of Food Lipids*, 4(3), 199-231. <https://doi.org/10.1111/j.1745-4522.1997.tb00093.x>
- Shin, W.-H., Lee, G.R., Heo, L., Lee, H., & Seok, C. (2014). Prediction of protein structure and interaction by GALAXY protein modeling programs. *Bio Design*, 2(1), 1-11.
- Simon, J.E., Chadwick, A.F., & Craker, L.E. (1984). *Herbs, an indexed bibliography, 1971-1980* (1st edn). Archon Books: Hamden, CT.
- Simopoulos, A.P. (2002). Omega-3 fatty acids in inflammation and autoimmune diseases. *Journal of the American College of nutrition*, 21(6), 495-505. <https://doi.org/10.1080/07315724.2002.10719248>
- Smith, D.E., & Salerno, J. (1992). Selective growth inhibition of a human malignant melanoma cell line by sesame oil *in vitro*. *Prostaglandins, Leukotrienes and Essential Fatty Acids*, 46(2), 145-150. [https://doi.org/10.1016/0952-3278\(92\)90221-4](https://doi.org/10.1016/0952-3278(92)90221-4)
- Sotnikova, R., Ponist, S., Navarova, J., Mihalova, D., Tomekova, V., Strosova, M., & Bauerova, K. (2009). Effects of sesame oil in the model of adjuvant arthritis. *Neuroendocrinology Letters*, 30(Suppl. 1), 22-24.
- Tian, H., & Guo, R. (2017a). Cardioprotective potential of sesamol against ischemia/reperfusion injury induced oxidative myocardial damage. *Biomed Res*, 28(5), 2156-2163.
- Tian, H., & Guo, R. (2017b). Cardioprotective potential of sesamol against ischemia/reperfusion injury induced oxidative myocardial damage. *Biomedical Research*, 28(5).
- Trott, O., & Olson, A.J. (2010). AutoDock Vina: improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading. *Journal of Computational Chemistry*, 31(2), 455-461. <https://doi.org/10.1002/jcc.21334>
- Uddin, S.J., Grice, I.D., & Tiralongo, E. (2011). Cytotoxic effects of Bangladeshi medicinal plant extracts. *Evidence-Based Complementary and Alternative Medicine*, 2011, 7. <http://doi.org/10.1093/ecam/nep111>
- Wan, Y., Li, H., Fu, G., Chen, X., Chen, F., & Xie, M. (2015). The relationship of antioxidant components and antioxidant activity of sesame seed oil. *Journal of the Science of Food and Agriculture*, 95(13), 2571-2578. <https://doi.org/10.1002/jsfa.7035>
- Wichitsranoi, J., Weerapreeyakul, N., Boonsiri, P., Settasatian, C., Settasatian, N., Komanasin, N., . . . Leelayuwat, N. (2011). Antihypertensive and antioxidant effects of dietary black sesame meal in pre-hypertensive humans. *Nutrition Journal*, 10(1), 82. <https://doi.org/10.1186/1475-2891-10-82>
- Wiederstein, M., & Sippl, M.J. (2007). ProSA-web: interactive web service for the recognition of errors in three-dimensional structures of proteins. *Nucleic Acids Research*, 35(suppl_2), W407-W410. <https://doi.org/10.1093/nar/gkm290>

- Wirasnita, R., Hadibarata, T., Novelina, Y.M., Yusoff, A.R.M., & Yusop, Z. (2013). A modified methylation method to determine fatty acid content by gas chromatography. *Bulletin of the Korean Chemical Society*, 34(11), 3239-3242. <http://doi.org/10.5012/bkcs.2013.34.11.3239>
- Wu, X.-L., Liou, C.-J., Li, Z.-Y., Lai, X.-Y., Fang, L.-W., & Huang, W.-C. (2015). Sesamol suppresses the inflammatory response by inhibiting NF- κ B/MAPK activation and upregulating AMP kinase signaling in RAW 264.7 macrophages. *Inflammation Research*, 64(8), 577-588. <https://doi.org/10.1007/s00011-015-0836-7>
- Yalagala, P.C., Sugasini, D., Ramaprasad, T.R., & Lokesh, B.R. (2017). Minor constituents in Rice bran oil and sesame oil play a significant role in modulating lipid homeostasis and inflammatory markers in rats. *Journal of Medicinal Food*, 20(7), 709-719. <https://doi.org/10.1089/jmf.2016.3780>
- Yashaswini, P., Rao, A.A., & Singh, S.A. (2017a). Inhibition of lipoxygenase by sesamol corroborates its potential anti-inflammatory activity. *International Journal of Biological Macromolecules*, 94(Part B), 781-787. <https://doi.org/10.1016/j.ijbiomac.2016.06.048>
- Yashaswini, P.S., Sadashivaiah, B., Ramaprasad, T.R., & Singh, S.A. (2017b). *In vivo* modulation of LPS induced leukotrienes generation and oxidative stress by sesame lignans. *The Journal of Nutritional Biochemistry*, 41, 151-157. <https://doi.org/10.1016/j.jnutbio.2016.12.010>
- Ying, Z., Kherada, N., Kampfrath, T., Mihai, G., Simonetti, O., Desikan, R., . . . Parthasarathy, S. (2011). A modified sesamol derivative inhibits progression of atherosclerosis. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 31(3), 536-542. <https://doi.org/10.1161/ATVBAHA.110.219287>
- Zhang, M.-M., Xie, X., Ma, Y.-T., Zheng, Y.-Y., Yang, Y.-N., Li, X.-M., . . . Chen, B.-D. (2015). Association of COX-2-765G> C genetic polymorphism with coronary artery disease: a meta-analysis. *International Journal of Clinical and Experimental Medicine*, 8(5), 7412-7418.
- Zhang, R., Yu, Y., Deng, J., Zhang, C., Zhang, J., Cheng, Y., . . . Yang, H. (2016a). Sesamin ameliorates high-fat diet-induced dyslipidemia and kidney injury by reducing oxidative stress. *Nutrients*, 8(5), 276. <https://doi.org/10.3390/nu8050276>
- Zhang, R., Yu, Y., Hu, S., Zhang, J., Yang, H., Han, B., . . . Luo, X. (2016b). Sesamin ameliorates hepatic steatosis and inflammation in rats on a high-fat diet via LXR α and PPAR α . *Nutrition Research*, 36(9), 1022-1030. <https://doi.org/10.1016/j.nutres.2016.06.015>
- Zihad, S.N.K., Bhowmick, N., Uddin, S.J., Sifat, N., Rahman, M.S., Rouf, R., . . . Aziz, S. (2018). Analgesic Activity, Chemical Profiling and Computational Study on *Chrysopogon aciculatus*. *Frontiers in Pharmacology*, 9, 1164. <https://doi.org/10.3389/fphar.2018.01164>
- Zilani, M.N.H., Uddin, S.J., Hossain, H., Hazni, H., Shilpi, J.A., & Hossain, M.G. (2018). Chemical characterization and bioactivity of *Trichosanthes dioica* edible shoot extract. *Oriental Pharmacy and Experimental Medicine*, 18(2), 167-175. <https://doi.org/10.1007/s13596-018-0310-5>

Table 1. Reported traditional uses, isolated compounds and pharmacological properties of Sesame oil (*Sesamum indicum* L.)

Scientific name	Family	Traditional uses	Isolated compounds	Reported pharmacological activities	References
<i>Sesamum indicum</i> L.	Pedaliaceae	<ul style="list-style-type: none"> ➤ Reduce body heat ➤ Relief pain ➤ Flavor enhancers ➤ Laxative ➤ Swish strong teeth and gums ➤ Healthy vaginal balance ➤ Emollient ➤ Tonic ➤ Diuretic ➤ Emmenagogue ➤ Aphrodisiac ➤ Amenorrhoea ➤ Dysmenorrhea ➤ Piles ➤ Demulcent ➤ Dysentary ➤ Urinary complaints ➤ Solvent for injection 	<p>1)Fatty acids</p> <p>a)Saturated- Palmitic acid, Stearic acid, Arachidic acid, Myristic acid, Behenic acid, Octanoic acid, Decanoic acid, Lauric acid</p> <p>b)Monounsaturated- Oleic acid, Eicosenoic acid</p> <p>c)Polyunsaturated- Linoleic acid, Linolenic acid</p> <p>2)Lignans Sesamin, Sesamol, Sesamol, Sesaminol, Sesamol, Pinoresinol, Lariciresinol, Asarinin</p> <p>3)Phytosterols Campesterol, Stigmasterol, Beta-sitosterol, Δ5-Avenasterol, Δ7-Stigmasterol and Δ7-Avenasterol</p> <p>4)Vitamins Alpha tocopherol, Beta tocopherol, Gamma tocopherol, Vitamin A, Vitamin B, Vitamin K</p>	<ul style="list-style-type: none"> ➤ Antioxidant ➤ Anti-inflammatory ➤ Antinociceptive ➤ Hepatoprotective ➤ Anti-arthritic ➤ Anti-hypertensive ➤ Chemo-protective ➤ Anti-cancer 	(Ahmad et al., 2006; Azab, 2014; Balkrishna, 2008; Burdock, 2016; Cengiz et al., 2013; Chopra & Nayyar, 1956; Ghani, 1998; Gharby et al., 2017; Guimarães et al., 2013; Hsu, Chen, Chu, & Liu, 2013; Jamieson & Baughman, 1924; Kapadia et al., 2002; Kuo, Lin, Chen, Yiu, & Tzen, 2011; Lahorkar, Ramitha, Bansal, & Narayana, 2009; Ma, Ding, Zhang, & Liu, 2014; Matsumura et al., 1995; Milder, Arts, van de Putte, Venema, & Hollman, 2005; Monteiro et al., 2014; Pole, 2006; Ramesh, Saravanan, & Pugalendi, 2005; Ricci et al., 1999; Saleem et al., 2011; Sankar et al., 2006b; Shafipour, Mosayebi, Asgari, Atrkarroushan, & Pasdaran, 2017; Simon, Chadwick, & Craker, 1984; Smith & Salerno, 1992; Sotnikova et al., 2009; Wan et al., 2015)

Table 2. Antioxidant activity of sesame oil

Test model	Dose/concentration	Route of administration	Mechanism of action	References
Myocardial injury rat model	5 and 10 mL/kg b.w	Orally	Reduction of thiobarbituric acid reactive substance (TBARS) and enhancement of endogenous antioxidant enzymes, lactate dehydrogenase (LDH), creatine kinase (CK) and aspartate transaminase (AST).	(Saleem et al., 2013)
Doxorubicin-induced rat model	5 and 10 mL/kg b.w	Orally	Enhancement of cardiac endogenous antioxidants and reduction of myocardial TNF- α expression.	(Saleem, Chetty, & Kavimani, 2014)
Streptozotocin (STZ) induced rat model	2% oil supplemented with 6%	With diet	Reduction of the levels of blood glucose, glycosylated hemoglobin, TBARS, lipid hydroperoxides and glucose-6-phosphatase, fructose-1,6-bisphosphatase activities; and elevation of the levels of hemoglobin, vitamin E, GSH and hexokinase activity.	(Ramesh et al., 2005)
Lipopolysaccharide-induced rat model	8 mL/kg	Parenterally	Reduction of lipid peroxidation, hydroxyl radical and nitrite level; and a boost in superoxide dismutase and catalase activity.	(Hsu et al., 2004a)
Rat (cecal ligation and puncture) model	4 mL/kg	Orally	Depletion of lipid peroxidation and serum nitrite levels.	(Hsu, Li, Chien, & Liu, 2004b)
Iron-induced rat model	0.4 g/kg sesame oil	With diet	Attenuation of the level of hepatic thiobarbituric acid-reactive substances, serum glutamate:oxaloacetate transaminase activities and serum glutamate pyruvate transaminase activities.	(Hemalatha & Raghunath, 2004)
Lipopolysaccharide induced rat model	8 mL/kg	Orally	Reduction of lipid peroxidation and superoxide anion counts; and increase in glutathione levels, and activities of superoxide dismutase and catalase.	(Hsu & Liu, 2004a)
Fenvalerate-induced rat model	10% sesame oil	With diet	Reduction of lipid peroxidation; and elevation of GSH levels and antioxidant enzymes.	(Prasanthi & Rajini,

model				2005)
(MCAO)-induced rat model	20% SO	With diet	Reduction of glutathione (GSH), glutathione-S-transferase (GST), glutathione peroxide (GPx), glutathione reductase (GR), catalase (CAT), superoxide dismutase (SOD), thiobarbituric acid reactive substance (TBARS) and lipid peroxidation (LPO).	(Ahmad et al., 2006)
Mice model	4mL /kg/day	Intraperitoneally	Significant increase in TAC, but non-significant elevation of serum nitrite production.	(Mosayebi, Ghazavi, Salehi, Payani, & Khazae, 2007)
Acetaminophen-induced rat model	8 mL/kg	orally	Significant depletion of glutathione levels and mitochondrial aconitase activity; but contradictory increase in superoxide anion, hydroxyl radical, and lipid peroxidation levels.	(Chandrasekaran, Chien, Hsu, Chang, & Liu, 2010).
Gentamicin-plus-iodinated contrast-induced rat model	0.5 mL/kg	Orally	Prevention of renal lipid peroxidation, myeloperoxidase, hydroxyl radicals, superoxide anion, nitrite/nitrate, and inducible nitric oxide synthase.	(Hsu, Li, Chu, Periasamy, & Liu, 2011)
Lipopolysaccharide-induced rat model	8 mL/kg	Subcutaneously	Reduction of lipid peroxidation, hydroxyl radical, nitric oxide, TNF- α , and IL-1 β along with considerable increase in the activities of superoxide dismutase, catalase and glutathione peroxidase in LPS intoxication.	(Hsu et al., 2005)
Cypermethrin-induced rat model	5 mL/kg b.w	Orally gavage	Elevation of the level of glutathione (GSH) and the activities of the antioxidant enzymes along with a decrease in the level of TBARS.	(Abdou et al., 2012)
Monocrotaline-induced rat model	0.5, 1, 2, or 4 mL/kg	Orally gavage	Protection against SOS by down-regulation of MMP-9 expression, up-regulation of TIMP-1 expression, and inhibition of oxidative stress.	(Periasamy, Yang, Chen, Chang, & Liu, 2013)
Cyclosporine-A induced rat model	1mL/kg	Orally gavage	. Significant restoration of the level of MDA, GSH, SOD and CAT compared to CsA group, but changes in the IL-1 α , IL-1 β levels	(Gülcan, BÜLBÜL,

model				were insignificant	ÖZDEMİR, & EROL, 2015)
Streptozotocin-induced rat model	100 mg/kg b.w	Intraperitoneally.		Up-regulation of testosterone, LH, and FSH.	(Khaneshi, Nasrolahi, Azizi, & Nejati, 2013)
Mice model	0.5, 1 and 2 mL/kg	Ingestion		Reduction of lipid peroxidation along with increased nuclear factor erythroid 2-related factor 2 and GAP43 expression in sciatic nerve.	(Hsu, Huang, Wu, Tai, & Jou, 2016a)
Rat model	0.5 or 1 mL/kg/day	Orally gavaged		Down-regulation the expression of angiotensin type 1 receptor, JNK and p38 MAPK and apoptosis signal regulating kinase 1, c-Fos and c-Jun with diminished production of superoxide anion, hydroxyl radical and lipid peroxidation.	(Liu & Liu, 2017)
Adjuvant arthritis rat model	1 mL/kg	Orally		Decreased plasma TBARS and GGT activity in the joint and spleen tissues.	(Sotnikova et al., 2009)
(4-NQO) induced rat model	- 1, 2, 4, and 8 mL/kg	Orally		Reduction of LPO and DNA damage from 4-NQO.	(Arumugam & Ramesh, 2011)
Septic rat model	4 mL/kg/day	Orally		Significant decline in hepatic lipid peroxidation, hydroxyl radical, superoxide anion, nitrite levels, xanthine oxidase activity and inducible nitric oxide synthase expression.	(Hsu et al., 2008a)
LPS induced rat model	8 mL/kg	Orally		Reduction of lipid peroxidation with increased activities of superoxide dismutase (SOD) and catalase (CAT).	(Hsu & Liu, 2004b)
Ethanol induced rat model	8 mL/kg	Orally		Reduction of mucosal lipid peroxidation, glutathione and nitric oxide production.	(Hsu, Chu, & Liu, 2009b)
Diazinon induced rat model	5 mL/kg body weight	Orally		Improvement in hematology and serum parameters, endogenous antioxidant status, and reduction of lipid peroxidation.	(Abdel-Daim, Taha, Ghazy, &

				El-Sayed, 2015)
Osteoarthritis rat model	1, 2, 4 mL/kg	Ingestion	Reduction of lipid peroxidation, muscular superoxide anion and peroxynitrite generations along with increased muscular glutathione and glutathione peroxidase levels, nuclear factor erythroid-2-related factor (Nrf2) expression	(Hsu, Chu, & Jou, 2016b)
EMR induced rat	1.5-3 mL	Orally	Reduction of cholesterol level with elevated SOD and CAT activities.	(Marzook et al., 2014)
DPPH and ABTS radical scavenging assays and β -carotene/linoleate model system.	0.1 – 0.5 g	<i>In-vitro</i>	Neutralization of DPPH and ABTS radical and the degradation of β -carotene.	(Bopitiya & Madhujith, 2013)
Human (clinical trial)	Use sesame oil as the only edible oil for 45 days.	With diet	Enhancement of superoxide dismutase (SOD), catalase (CAT), and the levels of vitamin C, vitamin E, beta-carotene, and reduced glutathione (GSH) along with decreased lipid peroxidation.	(Sankar et al., 2006a)

Table 3. Anti-inflammatory activity of sesame oil

Test model	Dose/concentration	Route of administration	Mechanism of action	References
LDLR -/- mice model	17% sesame oil	With diet	Reduction of plasma cholesterol, triglyceride, and LDL cholesterol levels in LDLR -/- mice.	(Bhaskaran, Santanam, Penumetcha, & Parthasarathy, 2006)
Mice (cecal ligation and puncture) model	5wt% SSO	With diet	Inhibition of delta-5 desaturase activity, resulting in an increase in the accumulation of dihomo-gamma-linolenic acid and subsequent decrease in the production of pro-inflammatory dienoic eicosanoids.	(Chavali, Utsunomiya, & Forse, 2001)
2,4,6-tri-nitro benzene sulfonic acid-induced rat model.	4 mL/kg	Orally gavaged	Reduction of the degree of inflammation, fibrosis, and acidic mucin with increased neutral mucin.	(Periasamy, Hsu, Chandrasekaran, & Liu, 2013)
(MSU) crystal-induced rat model.	0, 1, 2, or 4 mL/kg	Orally	Reduction of MSU crystal-induced total cell counts, tumor necrosis factor- α , interleukin (IL)-1 β , and IL-6 levels in lavage and pouch tissue.	(Hsu et al., 2013)
Osteoarthritis Rat model	0, 1, 2, or 4 mL/kg/day	Orally	Significant decrease in muscular interleukin-6, muscular lipid peroxidation, nuclear Nrf2 protein expression, and reactive oxygen species generations with increased citrate synthase activity and myosin heavy chain IL mRNA expression, glutathione production and glutathione peroxidase activity.	(Hsu et al., 2016c)
Both <i>in-vitro</i> and <i>in-vivo</i> (Atherosclerosis mice model)	340 mg/kg	With diet	Reduction of atherosclerotic lesions, plasma cholesterol, gene expression involved in inflammation and induction of genes involved in cholesterol metabolism and reverse cholesterol transport.	(Narasimhan et al., 2016).
Adjuvant-	2.5mg/kg/day	Orally	Attenuation of erythrocyte sedimentation rate (ESR) scores and pro-	(Ismail et

induced arthritis (AIA) rat model				inflammatory cytokines, including tumor necrosis factor-alpha (TNF- α), interleukin 1-beta (IL-1 β), and interleukin-6 (IL-6) levels.	al., 2018)
Rat model	10 wt% of GNO, RBO, and Sesame oil	With diet		Significant reduction in serum and liver lipids, 8-hydroxy-2-deoxyguanosine, cytokines in liver, and eicosanoids in leukocytes along with an up-regulation of sterol regulatory element-binding protein (SREBP)-2 and peroxisome proliferator-activated receptor gamma (PPAR γ) and down-regulation of nuclear factor-kappa B (NF- κ B) p65.	(Yalagala, Sugasini, Ramaprasad, & Lokesh, 2017).
Atherosclerosis Mice model	.75 mg/day	With diet		Reduction of the expression of inflammatory genes and induction of genes involved in cholesterol metabolism and RCT.	(Narasimhulu et al., 2018b)
Rheumatoid arthritis rat model	1mL/kg	Orally		Significantly decrease in the levels of TNF- α , IL-6 with an increase in the levels of IL-10 in groups treated with combination of SO and MTX compared to group treated with SO only or MTX only.	(Ali et al., 2017)
6-OHDA induced mice model	200 mL SO (20%)	With diet		Inhibition of activation of NADPH oxidase dependent inflammatory mechanism along with decreased TBARS level and increased GSH activity.	(Ahmad et al., 2012)
RAW macrophages (<i>in-vitro</i>) & mice (<i>in-vivo</i>)	264.7 50 and 250 μ g/mL	Cell line and with diet (<i>in-vivo</i>)		Prevention of Ox-LDL uptake by RAW macrophages and further inflammation <i>in-vitro</i> with reduction in plasma levels of TNF- α , IL-6, MCP-1 and VCAM1 in the SOAE pre-treated animals.	(Narasimhulu et al., 2018a)
Mice model and RAW macrophage cell.	264.7 10 μ g/mL-500 μ g/mL	Injected intraperitoneally		Down-regulation of the number of inflammatory markers such as Ccl2 or MCP-1, Ccl5 or RANTES, IL-1alpha, IL-1beta, TNF, colony stimulating factor 2 (Csf2), Nfkb1 and matrix metallo-proteinase 3 (Mmp3).	(Selvarajan et al., 2015)
MDMs /RAW cells	264.7 50 and 250 μ g/mL for MDMs, 5 and 25 μ g/mL for RAW macrophages	Cell line		Inhibition of IL-6 expression, TNF alpha or TLR4 or NFkB signaling pathways.	(Deme et al., 2018)

Table 4. Antioxidant and anti-inflammatory activity of pure compound

Isolated compound	Test model	Dose/conc.	Route of administration	Proposed mechanism	References
Sesamol	DLF-induced gastric rat model	0, 1, 3, or 10mg/kg	Subcutaneously.	Reduction of mucosal lipid peroxidation, hydroxyl radical levels, DLF induced mucosal prostaglandin E2 generation and cyclo-oxygenase activity.	(Hsu et al., 2008b)
	Aspirin induced rat model	0 to 30 mg/kg/day	Orally	Reduction of gastric mucosal lipid peroxidation, nitric oxide production, gastric mucosal proinflammatory cytokines (tumor necrosis factor- α and interleukin 1- β levels), and the activity of gastric mucosal myeloperoxidase.	(Hsu et al., 2009a)
	Cyclophosphamide-induced rat model	50 mg/kg	Orally	Attenuation of the levels of endogenous reactive oxygen species, lipid peroxidation, tumor necrosis factor - α , interleukin (IL)-1 β , IL-6 and cyclooxygenase-2 as well as increase in the levels of glutathione, total thiols, superoxide dismutase, catalase, glutathione-transferase and glutathione peroxidase.	(Jnaneshwari et al., 2014)
	M injured rat model	I/R Rat 50 mg/kg b.wt	Orally	Diminished infarct size, resorted the cardiac markers, reduces the lipid peroxidation, neutrophil infiltration and increases the antioxidants level. Downregulates inflammatory genes, Bax, Caspase-3 apoptotic proteins and upregulates anti-apoptotic Bcl-2 protein.	(Tian & Guo, 2017b)
	Blood lymphocytes (<i>in-vitro</i>)	1, 5 and 10 μ g/mL	Cell line	Reduction of lipid peroxidation marker, thiobarbituric acid reactive substances and elevation of glutathione (GSH), superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase.	(Prasad, Mahesh, Menon, Jeevanram, & Pugalendi, 2005)

Sesamol	<i>In-vitro</i> colorectal carcinoma cells (HCT116).	Low (0.05 and 0.25 mM) and high (0.5, 2, 5, and 10 mM) concentrations	Cell line	Suppression of cell viability via disruption of cell cycle progression at high concentrations.	(Khamphio et al., 2016)
	STZ-induced rat model	2, 4, and 8 mg/kg body weight	Orally	Attenuation of release of inflammatory mediators (TNF- α , IL-1 β , TGF-1 β), and caspase 3.	(Chopra et al., 2010)
	Mice model	2, 4, and 8 mg/kg	Orally gavaged	Reduction of immobility period, lipid peroxidation and nitrite levels and elevation of sucrose preference, glutathione levels, superoxide dismutase and catalase activities.	(Saleem et al., 2011)
	Adjuvant-induced arthritis rat model	25 and 50 mg/kg	Orally	Reduction of the non-enzymatic inflammatory markers (TNF, IL-1 β , IL-6, COX-2, PGE2, ROS, and H ₂ O ₂), withincreasedsuperoxide dismutase, catalase and glutathione-s-transferase activity.	(Hemshekh ar et al., 2013)
	RAW 264.7 cells	1–100 μ M	Cell line	Reduction of the levels of lipid peroxidation,protein nitration in the membranes of mitochondria,production of superoxide anion radical (O ₂ ^{-•}) and nitric oxide (NO [•]).	(Duarte et al., 2018)
	RAW 264.7 cells	3, 10, 30, 100 μ M	Cell line	Inhibition of production of nitric oxide, prostaglandin E2 (PGE2), pro-inflammatory cytokines,the epression of iNOS and COX-2 and NF- κ B and MAPK signaling pathwaysas wll as Upregulation of AMPK activation and Nrf2 signalling pathway..	(Wu et al., 2015)
	<i>In-vitro</i>	0-60 Mm	<i>In-vitro</i> strain	Inhibition ofCOX-1 andprolongation of the lag phase.	(Yashaswini et al., 2017a)
	Various strains of salmonella typhimurium	100 μ g/plate dose		Neutralization of oxygen free radicals and inhibition of the mutagenicity of H ₂ O ₂ .	(Kaur & Saini, 2000)

Modified form of Sesamol (INV-403) Rabbit and *in-vitro* (endothelial cell) 20 mg/kg per day, Dietary supplementat ion and 100 μmol/L With diet and cell line Improvement of vascular function (decreases constriction to angiotensin II and increases relaxation to acetylcholine), attenuation of systemic and plaque oxidative stress, and inhibition of nuclear factor-κB activation. (Ying et al., 2011)

	(DOCA)-salt induced rat model.	0.1% and 1% sesamin	With diet	Inhibition of enhanced vascular O ²⁻ production.	(Nakano et al., 2002)
	Middle cerebral artery occlusion (MCAO) rat model	30 mg/kg	Orally	Reduction of TBARS and protein carbonyl (PC) maintenance of healthy level of glutathione and its dependent enzymes (glutathione peroxidase [GPx] and glutathione reductase [GR]) in MCAO group.	(Khan et al., 2010)
	Lipid-induced rat model	40, 80, or 160 mg/(kg/day)	Orally	Improvement of the serum total cholesterol, triglyceride, low-density lipoprotein cholesterol, apolipoprotein-B, oxidized-low-density lipoprotein, and serum creatinine as well as amelioration of the superoxide dismutase activity and reduction of malondialdehyde levels in kidney tissue.	(Zhang et al., 2016a)
Sesamin	6-OHDA-lesioned rat model	10 or 20 mg/kg/day	Orally	Attenuation of motor imbalance in narrow beam test, striatal level of malondialdehyde (MDA) and reactive oxygen species (ROS), striatal caspase 3 activity and α-synuclein expression and improvement in superoxide dismutase (SOD) activity.	(Baluchnej admojarad et al., 2017)
	(CCl ₄)-induced rat model	100 mg/kg/day	Orally	Reduction of the release of liver enzymes (ALT, AST, and TBIL), protein carbonyls and the levels IL-6 and COX-2 in the liver by inhibition of NF-κB activation with improved SOD and GSH-Px activities.	(Chen et al., 2015)
	Kainic acid-induced rat model	(15 mg/kg or 30 mg/kg) and (0.1, 0.5,	Orally gavaged and cell	Significant decrease in the release of Ca ²⁺ , reactive oxygen species, and MDA from PC12 cells, expression of ERK1/2, p38 mitogen-activated protein kinases, Caspase-3, and COX-2	(Hsieh et al., 2011)

	and PC12 cell and BV-2 cell	1.0, or 2.0 μ M) in cell model.	line	in BV-2 cells and PGE ₂ production.	
	CCl ₄ induced rat model	5 and 10 mL/kg bw	Intraperitoneally	Reduction of the elevated serum liver marker enzymes, TBARS and improvement of GSH, SOD and catalase activities.	(Lv et al., 2015)
	STZ induced mice model	30 mg/kg BW	Intraperitoneally	Diminished progression of diabetic retinal injury by: 1) decreasing blood glucose level, 2) suppressing microglia activation, 3) reducing retinal TNF- α and ICAM-1 levels and 4) quenching iNOS expression.	(Ahmad et al., 2016)
	Pb and LPS induced rat model	10 mg/kg	Orally	Reduction of the serum levels of AST, ALT, CRP, TNF- α , IL-1, IL-6, NO, ROS generation, liver tissue expressions of c-Jun N-terminal kinase (JNK), p38 MAPK, GADD45 β , COX-2, and iNOS.	(Chiang et al., 2014)
	Ovalbumin (OVA)-induced mice model	1 mg/kg, 10 mg/kg, and 20 mg/kg body weight	Intraperitoneally	Inhibition of expression levels of interleukin-4 (IL-4), IL-5, IL-13, and serum IgE with reduced numbers of total inflammatory cells and eosinophils in BALF.	(Lin et al., 2014)
	<i>In-vitro</i> and <i>in-vivo</i> mice model	1 μ mol and 1 mg/kg	Cell line and orally	Significant decrease in lipid peroxidation and neutralization of free radicals.	(Nakai et al., 2003)
	DOCA-salt induced rat model	0.1% sesamin	With diet	Inhibition of the enhancement of aortic O ² -production.	(Nakano et al., 2003)
Sesamin	CCl ₄ induced mice model	10 mg/kg	Orally	Enhancement of the expression levels of phosphorylated Jun N-terminal kinases (JNK) with diminished release of mitochondrial cytochrome c in liver.	(Ma et al., 2014)
	Hepatic steatosis rat model	40, 80, and 160 mg/kg bw	Intragastrically	Reduction of the serum levels of total cholesterol, triacylglycerols, low-density lipoprotein cholesterol, free fatty acid, malonaldehyde and cytochrome P450 2E1 as well as potentiation of hepatic glutathione peroxidase and superoxide dismutase activities.	(Zhang et al., 2016b)
	Neuronal	1 Pm	Cell line	Significant depletion of MPP ⁺ -Induced IL-6, IL-1 β , and TNF- α	(Bournival

	PC12 cells cocultured with N9 microglial cells			mRNA and protein concentrations.	et al., 2012)
Sesamin	Primary chondrocytes (cultured cell)	2.5 and 5µM	Cell line	Inhibition of IL-1β induced production of PGE ₂ and NO, production of MMP1, MMP3 and MMP13 in IL-1β-stimulated chondrocytes, and IL-1β-induced phosphorylation of NF-κB p65 and IκBa.	(Kong et al., 2016)
	Human (clinical trial)	200 mg/day	With diet	Significant decrease in serum levels of MDA with increased TAC and high-density lipoprotein cholesterol (HDL-c) levels.	(Helli et al., 2016)
	Neuronal PC12 cells	1 pM	Cell line	Reduction of intracellular ROS production and modulation of tyrosine hydroxylase, superoxide dismutase, catalase, inducible NO synthase and interleukin-6 expression.	(Lahaie-Collins, Bournival, Plouffe, Carange, & Martinoli, 2008)
	Endothelial cell	12.5-100 µM	Cell line	Amelioration of oxLDL-induced ROS generation, SOD-1 inactivation, and activation of NF-κB.	(Lee et al., 2009)
Sesamin derivative-1, 2-bis(3-methoxybenzyl) ethane-1, 2-dicarboxylic acid (MMEDA)	Rat model, BV-2 microglia or PC12 cell	10 mg/kg, and 1, 10, 50 µM	Intraperitoneally and cell line	Attenuation of ROS, PGE ₂ release, hypoxia-induced JNK and caspase-3.	(Hung et al., 2017)
Sesamin derivative,	PC12 cells	10 µM	Cell line	Inhibition of LDH, lipid peroxidation, cell damage, ROS, elevation of intracellular free Ca ion and apoptosis activity	(Hou et al., 2015)

3-bis (3-methoxybenzyl) butane-1,4-diol					with increased ACh release.	
Sesamin and sesame oil	Paw edema mice model	sesame oil (100, 200, and 400 mg/kg) and sesamin (50, 100, and 200 mg/kg)	Orally		Inhibition of prostaglandin synthesis, pleural exudate formation and the leucocyte migration.	(Monteiro et al., 2014)
Sesamin and sesamol	Mice model	(0%, 0.1%, 0.2%, and 0.5%)	With diet		Inhibition of caspase-3 and MAPK activation, IL-6, PGE2, and tumor necrosis factor (TNF)- α and nitric oxide (NO) productions from microglia as well as elevation of IL-10..	(Jeng & Hou, 2005)
	BV-2 microglial cell line	50–100 μ M	Cell line		Inhibition of NO production, iNOS, mRNA and protein expression.	(Hou et al., 2003)
Sesamin and sesamol	LPS induced rat model	Oral supplement of sesamol and sesamin	Orally		Lowers LPS induced expression of cPLA ₂ , 5-LOX, BLT-1 and LTC ₄ synthase. The serum levels of TNF- α , MCP-1 and IL-1 β are found to be reduced in sesamol and sesamin group,	(Yashaswini et al., 2017b)
Sesamol	Rat model	1% sesamol	With diet		Reduction of lipid peroxidation, measured as 2-thiobarbituric acid reactive substances.	(Kang et al., 1998)
Sesaminol	Cultured rat astrocytes	sesaminol glucosides (10–100 μ g/ml)	Cell line		Inhibition of LPS-induced generation of nitric oxide (NO), reactive oxygen species (ROS) and expression of cytosolic phospholipase A ₂ (cPLA ₂), cyclooxygenase 2 (COX-2) and inducible nitric oxide synthase (iNOS) as well as prevention of LPS-induced DNA binding and transcriptional activity of nuclear factor KappaB (NF)- κ B.	(Lee et al., 2006)
	PC12 cells	25, 50 and 100 μ mol	Cell line		Elevation of cell survival rate and, SOD, CAT and GSH-Px activity with significant decline in the secreted LDH level,	(Cao et al., 2013)

apoptosis rate and ROS level of H₂O₂ exposed cells.


Sesame lignans and tocopherols	DM model	rat	0.5% sesame lignin + 0.25% alpha tocopherol	With diet	Amelioration of the alteration in lipid profile and the adverse free radical generation.	(Dhar et al., 2007)
Sesame lignans	<i>In-vitro</i>		0.1 µM	Cell line	Inhibition of TBARS formation.	(Kang et al., 1999)

Table 5. Binding affinity and interaction of sesame oil ligands with COX-2 after docking.



Compounds	Binding affinity to COX2 (kcal/mol)	Interacting Amino acid
Linoleic Acid	-7.0	Pro528, Lys346, Ile331, Gly324, Met99 , Phe347, Leu352, Leu351, Asp348, Ile327, Leu103 , Pro349, Lys328, Cys526, Trp531
Linolenic Acid	-7.3	Phe353, Phe347, Ile331, Asp348, Lys346, Ile327, Leu352, Cys526, Lys328, Pro528, Pro349, Trp531
Oleic Acid	-6.5	Ile331, Arg46, Leu352, Phe347, Ile327, Cys526, Pro528, Lys328, Pro349, Trp531
Palmitic Acid	-5.4	Gln336, Val102 , Leu517 , Ala513 , Leu345, Val335 , Val509 , Leu338 , Phe343 , Ser339 , Tyr341
Pinoresinol	-8.4	Ser339 , Tyr371 , Leu517 , Ala513 , Gly512 , Val335 , Met99 , Met508, Val509 , Phe343 , Tyr341 , Ser516, Val102
Sesamin	-10.4	Val524, Asn523, Arg362, Arg362, Phe128, Gln360, Asp215, Gly211, Asn361
Sesaminol	-10.6	Arg362, Asp215, Gly211, Ser129, Phe128, Gln360, Phe128, Arg362, Asn361
Sesamol	-6.4	Tyr359, Gln356, Pro528, Leu352, Ser527, Gln358, Phe357
Sesamolin	-9.6	Ser527, Lys355, Leu352, Phe357, Lys355, Ser112, Tyr359, Phe357, Phe353, Pro528, Thr104, Ser107, Gln358, Gln356

Sesamolinol	-8.8	Lys518, Lys355, Leu352, Pro528, Gln358, Thr104, Phe357, Gln356, Ser112, Leu103 , Ser107
Stearic Acid	-5.9	Met508, Leu345, Phe343 , Val335 , Ala513 , Gly512 , Val102 , Phe504, Leu338 , Leu517 , Val509 , Ser339 , Tyr341
Tocopherol	-7.1	Ser105, Ala513 , Leu345, Phe343 , Arg106, Ile98, Trp85, Phe84, Ile77, Leu517 , Met99 , Val74, Leu78, Val102 , Tyr341
Vitamin A Retinol	-7.4	Ser339 , Ser105, Tyr101, Leu78, Pro69, Lys68, Ala513 , Val335 , Glu510, Arg106, Val74, Val102 , Pro71, Tyr341
Vitamin K Phylloquinone	-9.6	Cys526, Leu352, Asp348, Lys346, Gly324, Met521, Phe343 , Ser339 , Val335 , Leu345, Ile327, Val102 , Tyr341 , Trp531, Leu103 , Met99 , Leu517 , Ile331, Lys328
Arachidic acid	-6.5	Met521, Asp348, Gly324, Met99 , Phe346, Lys346, Leu352, Ile327, Pro528, Phe351, Ile331, Cys526, Lys328, Trp531, Pro349
Myristic acid	-6.2	Cys526, Lys346, Gly324, Phe347, Leu352, Pro349, Ile331, Ile327, Lys328, Trp531
Behenic acid	-6.2	Gly512 , Tyr371 , Leu345, Leu103 , Met99 , Phe343 , Val102 , Leu517 , Leu338 , Ser516, Ala513 , Tyr341 , Ser339 , Val335 , Val509
Octanoic acid	-5.2	Asp348, Ile327, Trp531, Lys346, Phe347, Lys328
Decanoic acid	-5.8	Phe347, Gly324, Leu352, Ile327, Pro528, Cys526, Lys328, Pro349, Trp531
Lauric acid	-5.8	Cys526, Leu352, Pro349, Met99 , Asp348, Lys346, Ile327, Ile331, Phe347, Lys328, Trp531
Eicosenoic acid	-6.2	Phe343 , Gly512 , Leu338 , Leu103 , Val335 , Leu517 , Ala513 , Leu345, Met99 , Val509 , Tyr341 , Val102 , Ser516, Ser339
Lariciresinol	-8.8	Asp215, Tyr359, Gly211, Leu210, Trp125, Val524, Arg362, Gly211, Phe128, Gln360, Phe128, Arg362, Asn523, Asn361
Asarinin	-10.4	Val524, Asn523, Arg362, Asp215, Arg362, Phe128, Gly211, Phe128, Gln360, Asn361
Campesterol	-10.5	Asn523, Gly211, Trp125, Val524, Val524, Asn361, Leu131, Arg362, Gln360, Phe128, Trp125, Arg362, Gly211, Asn523, Asp215
Stigmasterol	-10.7	Val524, Gly211, Gln360, Leu210, Val524, Asn361, Leu131, Phe128, Arg362, Phe128, Trp125, Arg362, Gly211, Asn523, Asp215
Beta-sitosterol	-10.2	Val524, Arg362, Asn361, Phe128, Val524, Asn361, Leu131, Ser129, Gln360, Phe128, Trp125, Arg362, Gly211, Asn523, Asp215


Δ5-Avenasterol	-10.7	Val524, Gly211, Leu210, Ser129, Gln360, Val524, Asn361, Leu131, Phe128, Arg362, Phe128, Trp125, Arg362, Gly211, Asn523, Asp215
Δ7-Stigmasterol	-10.5	Val524, Leu131, Gln360, Gly211, Leu210, Asn361, Phe128, Trp125, Phe128, Gly211, Trp125, Val524, Gln360, Arg362, Asn523, Asp215
Δ7-Avenasterol	-10.7	Arg362, Leu210, Asn361, Leu131, Trp125, Val524, Asn523, Gln360, Phe128, Val524, Ser129, Phe128, Gly211, Trp125, Gly211, Arg362, Asp215, Asn523
Vitamin B(Riboflavin)	-8.5	Arg362, Trp125, Asn523, Asp215, Leu131, Ser129, Leu210, Arg362, Gln360, Gly211, Asn361
Control		
Arachidonic Acid	-6.8	Met99, Gly512, Tyr371, Val335, Leu103, Leu517, Ala513, Phe343, Ser339, Leu338, Val102, Val509, Tyr341



(a)



(b)



(c)




Figure 1: Structures of constituents reported in sesame oil. (a) Structures of fatty acids; (b) Structures of lignans; (c) Structures of phytosterols; (d) Structure of vitamins




Figure 2: Workflow of database searches with literature inclusion and exclusion criteria.




Figure3: Proposed mechanism of antioxidant and anti-inflammatory activity of sesame oil




Figure 4: Homology model of Human COX-2 enzyme.




Figure 5: Docking of Stigmasterol with human COX-2 enzyme




Figure 6: Docking of Δ^5 Avenasterol with human COX-2 enzyme