

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/266210195>

Marked eburnation among prehispanic individuals from La Gomera (Canary Islands)

Article · January 2014

CITATIONS

0

READS

86

4 authors, including:



Maria Castañeyra-Ruiz

Universidad de La Laguna

12 PUBLICATIONS 17 CITATIONS

[SEE PROFILE](#)



Matilde Arny

Universidad de La Laguna

116 PUBLICATIONS 782 CITATIONS

[SEE PROFILE](#)



Emilio Gonzalez-Reimers

Universidad de La Laguna

330 PUBLICATIONS 3,230 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



"Caracterización de las poblaciones prehispanicas de las Islas Canarias Occidentales a través del estudio de Isótopos estables en restos óseos" [View project](#)



Retinal occlusive disease [View project](#)

Marked eburnation among prehispanic individuals from La Gomera (Canary Islands)

María Castañeyra-Ruiz ¹, Aioze Trujillo-Mederos ¹, Matilde Arnay-de-la-Rosa ¹, Emilio González-Reimers ²

¹Departamento de Prehistoria, Arqueología, Antropología e Historia Antigua. Universidad de La Laguna. Tenerife. Canary Islands (Spain)

² Departamento de Medicina Interna. Hospital Universitario de Canarias. 38320. Universidad de La Laguna.

Author for correspondence. E-mail= egonrey@ull.es

Recibido: 24-04-2014, revisado: 21-05-2014, aceptado 10-06-2014

Summary

Marked eburnation among prehispanic individuals from La Gomera (Canary Islands)

Eburnation of the articular surfaces is an advanced feature of osteoarthritis. We here report some striking cases of such a condition in the proximal epiphysis of tibiae of prehispanic inhabitants of La Gomera, in the Canary Archipelago.

Key words

Osteoarthritis; Eburnation; Rheumatic diseases; Prehispanic Canary Islands; paleopathology.

Resumen

Marcada eburnación entre individuos prehistóricos de La Gomera (Islas Canarias)

La eburnación de las superficies articular es una característica avanzada de la osteoartritis. Presentamos aquí algunos casos llamativos de tal condición en la epífisis proximal de la tibia de los habitantes prehistóricos de la isla Gomera, en el Archipiélago Canario.

Palabras clave

Artrosis; eburnación; enfermedades reumáticas; Canarias Prehistóricas; paleopatología

Introduction

Osteoarthritis is a multi-factorial disease with a final common pathway of cartilage degeneration and bone eburnation. Articular cartilage and subchondral bone form a functional unit which become early altered during the osteoarthritic process [1,5]. Altered biomechanical forces, macrotrauma or repeated microtrauma are able to induce the release of proteolytic enzymes by chondrocytes, which contribute to destroy normal cartilage. In rare instances, defective cartilage may also become destroyed by normal loading forces.

Although subchondral bone sclerosis may precede detectable changes in articular cartilage [3], eburnation is a late event of the disease. For instance, it was present in the knee in about 40% of individuals undergoing tibial osteotomy and/or

replacement [7]. The presence of eburnation in an ancient bone implies that the destruction of the cartilage was complete [9]. Eburnation has been explained as the result of mechanical attrition and polishing of the exposed bony articular surface. However, the mechanisms underlying eburnation are by far more complex. Articular cartilage and subchondral bone respond in a coupled manner to loading forces that ultimately trigger the osteoarthritic process. The same forces which may destroy the cartilage are also involved in an osteosclerotic bone reaction which affects subchondral bone. Recent research supports a role of the Wnt β catenin /SOST system in increased sclerotic bone response [11]. Sclerostin, a protein derived from SOST gene transcription, inhibits bone formation [4,8], counteracting the action of WNT β catenin, which becomes activated in response to mechanical loading, and which is involved in increased bone synthesis. Indeed, some researchers have found that expression of SOST in osteocytes is decreased in osteoarthritic experimental models [10]. In addition, there is a complex response of the articular cartilage itself and the subchondral trabecular bone: a progressive thickening of the transverse trabeculae, together with a thinning of the vertical trabeculae have been described. Articular cartilage calcifies immediately adjacent to subchondral bone, and is then traversed by blood vessels [6]

Description of the cases

We report the cases of two right and one left tibiae probably belonging to adult men, affected by massive eburnation and polishing of the medial part of the proximal tibial epiphysis in two cases, and the lateral one in the third case (Figs. 1-3). As we can see, in the three cases, the tibial epiphyses show eburnated, polished surfaces. Marked osteophyte formation also surround the tibial articular surfaces. The intensity of osteoarthritic changes, at least in the two right tibiae, is more marked in the left (medial) epiphyseal plates, suggesting a misbalanced loading force acting on the medial side of the bone.

Fig. 1



Fig. 2



Fig.3



Fig.1,2,3: Cranial view of affected tibiae, showing the polished areas of eburnation (E). Intense osteophyte formation is also evident in the three cases.

Possibly, these individuals suffered genu varum deformity. The bones belong to prehispanic individual of La Gomera, and are preserved at the Museo Insular del Cabildo de San Sebastian (the capital of the island). Antiquity and precise burial sites are not known, but C14 dating of other samples from La Gomera have yielded antiquity ranging 1000-1500 years BP [2].

Discussion

We here report some advanced cases of tibial osteoarthritis in prehispanic inhabitants of La Gomera. This island is the second smallest (373 km²) of the seven “big “ islands of the Canary Archipelago, and was inhabited by people of North African origin, who arrived about 2500 years ago. The island, volcanic in origin, is round shaped, and in its central part there are remains of ancient volcanoes; indeed, no eruptions have happened in the last three millions years. All the central part, at an altitude over 800 m, is covered by dense forest, partly due to the contribution of the moisty north eastern winds which continuously blow during late spring and summer, converting the usually hot and dry months in moisty and mild ones. The economy of the inhabitants of the island was based on consumption of some forest species, coastal

fishing, and goat herding [2]. Probably the population had to climb the marked slopes of the island, and perhaps this activity contributed to the intense osteoarthritic changes. In any case, inspection of other bone remains suggest that osteoarthritis was a common disease among these individuals, and, in some cases, it was very intense, as illustrated in this report.

Bibliography

1. Ali SY Apatite-type crystal deposition in arthritic cartilage. *Scanning Electron Microsc.* 1985;(Pt 4):1555-66.
2. Arnay-de-la-Rosa M, Gámez-Mendoza A, Navarro-Mederos JF, Hernández-Marrero JC, Fregel R, Galindo-Martín et al. Dietary pattern during the early prehispanic settlement in La Gomera (Canary Islands). *J Archaeol Sci* 2009; 36:1972-1981.
3. Buckland-Wright C. Subchondral bone changes in hand and knee osteoarthritis detected by radiography. *Osteoarthritis cartilage* 2004; 12 (Suppl.A): S10-S19.
4. Chan BY, Fuller ES, Russell AK, Smith SM, Smith MM, Jackson MT, et al. Increased chondrocyte sclerostin may protect against

- cartilage degradation in osteoarthritis. Osteoarthritis cartilage 2011; 19:874-885.
5. Goldring MB, Goldring SR. Osteoarthritis. J Cell Physiol 2007; 213:626-634.
 6. Goldring SR. Alterations in periarticular bone and cross talk between subchondral bone and articular cartilage in osteoarthritis. Ther Adv Musculoskeletal Dis 2012; 4 :249-258.
 7. Koshino T, Machida J Grading system of articular cartilage degeneration in osteoarthritis of the knee. Bull Hosp Jt Dis. 1993
 8. Li X, Zhang Y, Kang H, Liu W, Liu P, Zhang J, Harris Se, Wu D (2005) Sclerostin binds to LRP5/6 and antagonizes canonical Wnt signaling. J Biol Chem 280:19883-19887.
 9. Ortner DJ, Putschar WGJ. Identification of Pathological Conditions in Human Skeletal Remains. Paleopathology. Smithsonian Institution Press 1981. Washington.
 10. Robling AG, Niziolek PJ, Baldridge LA, Condon KW, Allen MR, Alam I, et al. Mechanical stimulation of bone in vivo reduces osteocyte expression of SOST/sclerostin. J Biol Chem 2008; 283: 5866-5875.
 11. Williams BO, Isogna KL (2009) Where Wnts went: the exploding field of Lrp5 and Lrp6 signaling in bone. J Bone Miner Res 24:171-178.