



LJMU Research Online

Punt, A, Firman, J, Boobis, A, Cronin, MTD, Gosling, JP, Wilks, MF, Hepburn, PA, Thiel, A and Fussell, K

Potential of ToxCast data in the safety assessment of food chemicals

<http://researchonline.ljmu.ac.uk/id/eprint/12270/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Punt, A, Firman, J, Boobis, A, Cronin, MTD, Gosling, JP, Wilks, MF, Hepburn, PA, Thiel, A and Fussell, K (2020) Potential of ToxCast data in the safety assessment of food chemicals. Toxicological Sciences. ISSN 1096-6080

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

Potential of ToxCast data in the safety assessment of food chemicals

Ans Punt^{1*}, James Firman², Alan Boobis³, Mark Cronin², John Paul Gosling⁴, Martin F. Wilks⁵, Paul A. Hepburn⁶, Anette Thiel⁷, Karma Fussell⁸

¹Wageningen Food Safety Research, PO Box 230, 6700 AE Wageningen, The Netherlands,

²School of Pharmacy and Biomolecular Sciences, Liverpool John Moores University, Byrom Street, Liverpool, L3 3AF, UK

³National Heart & Lung Institute, Imperial College London, London W12 0NN, UK

⁴School of Mathematics, University of Leeds, Leeds, LS2 9JT, UK,

⁵Swiss Centre for Applied Human Toxicology, University of Basel, Switzerland

⁶Unilever, Safety & Environmental Assurance Centre, Colworth Science Park, Sharnbrook. MK44 1LQ, UK.

⁷DSM Nutritional Products, Wurmisweg 576, 4303-Kaiseraugst, Switzerland

⁸Nestlé Research, Case Postale 44, CH-1000 Lausanne 26, Switzerland.

*Corresponding author: publications@ilsieurope.be

© The Author(s) 2020. Published by Oxford University Press on behalf of the Society of Toxicology. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Abstract

Tox21 and ToxCast are high-throughput *in vitro* screening (HTS) programmes coordinated by the U.S. National Toxicology Program and the U.S. Environmental Protection Agency, respectively, with the goal of forecasting biological effects *in vivo* based on bioactivity profiling. The present study investigated whether mechanistic insights in the biological targets of food-relevant chemicals can be obtained from ToxCast results, when the chemicals are grouped according to structural similarity. Starting from the 556 direct additives that have been identified in the ToxCast database by Karmaus *et al.* (2017), the results showed that, despite the limited number of assays in which the chemical groups have been tested, sufficient results are available within so-called "DNA binding" and "nuclear receptor" target families to profile the biological activities of the defined chemical groups for these targets. The most obvious activity identified was the estrogen receptor (ER)-mediated actions of the chemical group containing parabens and structurally related gallates, as well the chemical group containing genistein and daidzein (the latter particularly towards ER β as potential health beneficial target). These group effects, as well as the biological activities of other chemical groups, was evaluated in a series of case studies. Overall, the results of the present study suggest HTS data could add to the evidence considered for regulatory risk assessments for food chemicals and to the evaluation of desirable effects of nutrients and phytonutrients. The data will be particularly useful for providing mechanistic information and to fill data gaps with read-across.

1. Introduction

Automatic high-throughput screening (HTS) of chemicals across a wide range of biological targets is an emerging practice in many chemical sectors. HTS plays a crucial role in the prioritization of chemicals based on toxicological mode of action as well as finding lead actives based on intended biological activity (Hartman *et al.*, 2018; Brunner *et al.*, 2019; Mayr and Fuerst, 2008; Olker *et al.*, 2019). Within next-generation (non-animal) risk assessment strategies, HTS will be one of the key technologies to characterize the ability of chemicals perturb biological pathways associated with an adverse outcome pathway (Villeneuve *et al.*, 2019). Much effort has been devoted to the application of HTS to various sectors and regulatory environments and strategies to achieve a broader acceptance of HTS and computational approaches in regulatory decision making have recently been laid out by Thomas *et al.* (2019). However, little has been done to relate these approaches to the assessment of foods and food ingredients, which are often assumed to be harmless, although a variety of toxicological or beneficial biological effects can be elicited. The aim of the present study was to explore the potential of ToxCast HTS data to be integrated into regulatory safety assessment of food chemicals.

The Tox21 and ToxCast programmes are high-throughput *in vitro* screening programmes co-ordinated by the U.S. National Toxicology Program and the U.S. Environmental Protection Agency, respectively, with the goal to forecast biological effects *in vivo*, especially toxicity, based on bioactivity profiling (Kavlock *et al.*, 2012). Tox21/ToxCast results (together referred to as ToxCast) have been evaluated by several groups in various publications using clustering algorithms and self-organizing maps (Karmaus *et al.*, 2016; Kleinstreuer *et al.*, 2014), hierarchical clustering techniques (Sipes *et al.*, 2013) or through links with chemical fingerprinting (Richard *et al.*, 2016). Specific to food relevant chemicals, Karmaus *et al.* (2016, 2017) identified and evaluated the activity patterns of 1211 food-use compounds within ToxCast, comprising 556 direct food additives, 371 food contact substances, and 543 pesticides.

A challenge with applying such non-directed, quantitative approaches on food-relevant chemicals is that an observed uneven coverage of chemical-endpoint combinations within this class of compounds leads to a significant bias in the results (i.e. chemicals with a high biological activity are those that have a broader test coverage). A second challenge of hierarchical clustering and self-organizing heatmaps is that they do not provide any direct mechanistic insights in the biological targets of a chemical relative to an adverse outcome pathway. The acquisition of such qualitative mechanistic insights is just as crucial to the consideration of ToxCast data in risk assessments of food chemicals and can be an important resource to evaluate nutrients and phytonutrients and their corresponding desirable effects.

The present study investigated whether insights in the biological targets of food-relevant chemicals can be obtained from the results of the ToxCast assays when the chemicals are grouped according to structural similarity (for example, homologous series), exploring those targets that are induced by multiple chemicals in the group. The current study focussed only on the 556 direct food additives that have been identified by Karmaus et al. (2017) (chemicals that are for example added to foods to preserve, colour and stabilise food as well as flavourings), and not on the 371 food contact substances and 543 pesticides that were identified by Karmaus et al. (2017). Both food contact substances and pesticides may lead to indirect exposures via food ingredients, but these compounds are not intended to be added to foods. The 556 direct food additives were supplemented with seven chemicals from the original non-curated list of food-use chemicals published by Karmaus et al. (2016) to also include natural food constituents (safrole, quercetin, resveratrol, genistein, daidzein, coumarin) as well as heptyl paraben, a non-approved food contact material that is structurally related to the approved methyl and ethyl parabens (EFSA, 2004), to give 563 reference compounds. The compounds within the dataset are clustered based on their chemical structural characteristics (e.g. alcohol, aldehydes and carboxylic acids, and ketones) as well as their functional uses in food (e.g. flavouring agents, nutrients, additives and regulatory restricted). Whereas the clustering into structurally similar chemicals was used to explore the relationship

between chemical homology and biological activity, the clustering into functional use categories related biological activities to current food uses. A method was set-up that allows to scroll through the activities of the groups of structurally related chemicals towards different targets. Relevant biological targets of a chemical group are considered those towards which a high percentage of chemicals within a group are active. Overall, the results of the present study offer insights into the possible integration of HTS data in the safety and risk assessments of food chemical.

2. Methods

2.1 Grouping of chemicals

2.1.1 Grouping of the chemicals according to functional use classes

The chemical names and CAS numbers were obtained from the appendices as published by Karmaus et al. (2016, 2017). To obtain a link between the 563 selected direct food additives and their use in foods, particularly within the EU, the compounds were subdivided into different use categories. To this end, the CAS numbers were first matched with the European Union list of flavourings (Annex I of Regulation 1334/2008) using the R script provided https://git.wur.nl/Punt001/ilsi_toxcast.git. There were 449 compounds that matched and were categorised as EU flavourings. The majority of the remaining 114 compounds were manually categorised into “novel foods”, “nutrients”, “polyphenols”, “E-numbers” (subdivided into “sweeteners”, “antioxidants”, “preservatives”, “colours”, and “remaining E-numbers”), and flavouring oils (which were merged with the EU flavourings use class), based on an online search using particularly the EU food additives database (EU) (DG SANTE, 2011), EFSA’s OpenFoodTox database (Dorne *et al.*, 2017; EFSA, 2017) and PubMed. The final 17 compounds that could not be linked to any known food use in the EU were categorized as “other”.

2.1.2 Grouping of the chemicals according to chemical structure

From the list of 563 compounds, 552 were found to correspond to discrete chemical entities with defined molecular structure. The SMILES (simplified molecular-input line-

entry specification) strings of these compounds were extracted from the ToxCast Data Spreadsheets (U.S. EPA, 2018a). The remaining 11 entities correspond to mixtures (for example, peppermint oil, clover leaf oil, polysorbate 80) and were either grouped together as structurally undefined or, in case of the flavouring oils, were assigned to the chemical group of the major constituent of oil. To this end, cornmint oil and peppermint oil are grouped in the same chemical group as menthol, whereas clove leaf oil is grouped with eugenol, anise oil with anethole, nutmeg oil with alpha-pinene, petitgrain oil with limonene, and cananga oil with beta-caryophyllene (Jelen, 2012; Han *et al.*, 2017). Using ChemoTyper software (Molecular Networks, Erlangen, Germany) and the SMILES strings of the chemicals, the compounds were classified through application of chemical knowledge, focussing on shared structural features and, where applicable, with their known physiological roles (Mellor *et al.*, 2019; Yang *et al.*, 2015). During the course of this undertaking, a three-tier system of grouping was adopted, in which larger primary clusters (for example, alcohols) were further subdivided as appropriate into secondary (for example, alcohol, alkyl) and tertiary (for example, alcohol, alkyl, primary, straight chain) groups. The final groupings can be found in Supporting information 1, along with additional information on the log P, log D, and the Henry's law constants of the chemical (estimated with ACD/Labs software). In addition, an estimate of the mean similarity of the chemicals within each group was made using the ChemmineR (Cao *et al.*, 2008) and fmcsR (Wang *et al.*, 2013) packages in R, to calculate the Maximum Common Substructures (MCSs) between the chemicals within a group and the Tanimoto coefficients based on these MCSs. The average of the calculated Tanimoto coefficients (excluding the Tanimoto coefficient of the chemicals with themselves) is taken as marker for group similarity. The R codes for these calculations have been made available at https://git.wur.nl/Punt001/ilsi_toxcast.git.

The grouping according to the functional use classes (section 2.1.1) and the chemical groups were combined in a so-called circle pack plot using the igraph and ggraph libraries in R (Csardi G and Nepusz T, 2006; Pedersen, 2017). The R code for the

circle pack graphic of the chemical groups has been made available at https://git.wur.nl/Punt001/ilsi_toxcast.git.

2.2 ToxCast data

The ToxCast activity data of the chemicals were derived from the data spreadsheet "ac50_Matrix_180918.csv" (U.S. EPA, 2018a) containing results from 1410 different assays. These crude ToxCast data for the 563 individual chemicals provided several positive hit-calls i.e. assays for which the concentration producing fifty percent of maximum activity (AC50), with a value less than 1,000,000 (the value used to indicate negative results), could be derived. Starting from this dataset, all assays were excluded that did not directly relate to a specific biological activity. These included all assays for which the 'assay_function_type' was 'background control', the 'assay_design_type' was either 'background reporter' or 'viability reporter', the 'intended_target_family' was 'background measurement' and the 'biological_process_target' was 'cell death', 'cell proliferation', or 'cytotoxicity'. These results are already taken into account during data analysis steps, e.g. through the production of Z-scores, and do not represent a specific activity of toxicological interest. In addition, only assays relevant to humans were extracted by setting 'species' to 'human'. The remaining assay endpoints were annotated according to the targets (for example ESR1 and ESR2, being the estrogen receptor (ER) alpha and beta, respectively), target family (for example, nuclear assays, DNA binding, or cytokines), and target subfamily (for example, nuclear assays-steroidal, nuclear assays-non steroidal, nuclear assays-orphan) parameters as provided in the "Assay_Summary_180918.csv" file (U.S. EPA, 2018a). By filtering out the assays that did not relate to a specific biological activity, 559 of the 1410 ToxCast assays were excluded, leaving 851 assays in the dataset.

Z-scores are utilised within the ToxCast dataset to filter out the AC50 results that were potentially affected by non-specific effects such as cytotoxicity. Z-scores represent the number of standard deviations (on a standardised scale) that separate the potency for the specified assay from the median potency of a range of cytotoxicity assays (Judson

et al., 2015; Houck *et al.*, 2017). Assay results with a large Z-score are more likely to reflect a target specific effect that is not caused by cell stress or cytotoxicity-related processes (Kleinstreuer *et al.*, 2014). For the present study, the available Z-scores from the "zscore_Matrix_180918.csv" (U.S. EPA, 2018a) file were used. These Z-scores were derived for chemicals with two or more positive responses in cytotoxicity assays. AC50 results with Z-scores lower than 3 were removed from the dataset as potential activity data that were affected by non-specific effects like cytotoxicity (Judson *et al.*, 2015). For 261 compounds out of the 563 food-relevant chemicals this filtering based on Z-score<3 resulted in a more than 75% reduction in positive hit-calls. For example, retinol expressed activity in 101 out of the 851 evaluated ToxCast assays, but 84 (83%) of these assay results had Z-scores<3. The mean AC50 for retinol in the 84 assays with Z-scores<3 was 57±41 µM, whereas the mean AC50 was 6.5±4.3 µM for the 17 assays with Z-score>3. A similar result can be seen for quercetin that was active in 91 assays of which 84 assay-results had a Z-scores<3. The mean AC50 value in the assays with Z-scores<3 was 31±33 µM whereas this was 2.8±2.0 µM for the assays that had Z-scores>3. These results suggest that the specificity increases after filtering for Z-scores. On average, for all chemicals, the mean AC50 values were 10-fold lower for the results with Z-scores >3 compared with the result with Z-scores<3.

Warning signs ("flags") are used in ToxCast data files to provide an indication of any unwanted influence of the method of data collection or automatic data processing on the obtained AC50 values. Possible flags include: i) "only highest concentration above baseline, active", ii) "only one concentration above baseline, active", iii) "multiple points above baseline, inactive", iv) "noisy data", v) "borderline active", vi) "borderline inactive", vii) "gain AC50 < lowest concentration & loss AC50 < mean concentration", viii) "hit-call potentially confounded by overfitting", and ix) "biochemical assay with < 50% efficacy". Flagged results were not filtered out from the ToxCast dataset in the present study but were considered in the different case studies to interpret the relevance of certain assays. The available flags were derived from the

"AllResults_flags_180918.csv" file (U.S. EPA, 2018a). For more information please see the US EPA documentation on the data analysis steps (U.S. EPA, 2018b).

In Supporting information 2, the background information on the different ToxCast assays is provided, including, per assay, the number of food-relevant chemicals that were tested, the fraction of the tested food-relevant chemicals that tested positive (AC50 value less than 1,000,000), the fraction of the tested food-relevant chemicals that contained flags (specified for each of the different flags). In addition, the targets, target families, and target subfamilies to which the assays belong, are provided in Supporting information 2.

2.3 Defining the biological activities of the chemical groups towards different ToxCast targets

Within the Assay_Summary_180918.csv file, the intended biological target of each assay is defined under "technological_target_official_symbol". For each of the tertiary chemical groups as identified with ChemoTyper (see Section 2.2.2. and Table 1 of the Results section), the biological activities towards the different biological targets were defined by calculating the percentage of chemicals (per tertiary chemical group) that were active in that assay of that target. To this end, the number of chemicals per tertiary chemical group that were tested in the assays of a specific target and the number of chemicals for which AC50 values (that is, the chemical tested positively) were defined, based on which the percentage activity could be calculated. For this evaluation, only those assays in which at least three chemicals of a group had been tested were considered. As a result, all chemical groups with less than 3 chemicals were removed for further analysis. For larger groups, this means that only chemical group-assay endpoint combinations with an $n > 3$ were included in the dataset.

For the different target families within ToxCast (for example, DNA binding and nuclear receptor targets) and the target subfamilies of the nuclear receptor target family (being steroidal, non-steroidal and orphan), the percent activities of each tertiary chemical grouping per individual biological target were plotted as a heatmap using the

ggplot package in R. In addition, the percentage of positive hits per target (sub)family was calculated and plotted along with the circle pack of the chemical groups. The R codes for the calculation of the biological activities per chemical group and the resulting heatmaps, and circle pack have been made available through https://git.wur.nl/Punt001/ilsi_toxcast.git. In addition, the R workflow has been made available as a web application using R Shiny (Rstudio). This web application is available through: <https://ilsi.eu/exploitation-of-toxcast-data-on-food-chemicals-for-safety-risk-assessment/>. The R codes for the calculation of the biological activities per chemical group and the resulting heatmaps, and circle pack have been made available through https://git.wur.nl/Punt001/ilsi_toxcast.git. In addition, the R workflow has been made available as a web application using R Shiny (Rstudio). This web application can be accessed through: <https://ilsi.eu/exploitation-of-toxcast-data-on-food-chemicals-for-safety-risk-assessment/>.

3. Results

3.1 Grouping of the ToxCast chemicals based on functional use and chemical structure.

The 563 food-relevant chemicals were clustered according to their chemical structure as well as their functional use classes. The obtained groups are displayed in Figure 1 as a so-called circle pack, which displays the hierarchical architecture of the defined functional and chemical groups. Note that a given chemical group may be split across more than one functional group and vice versa. The first layer within Figure 1 displays the functional-use classes of which the largest group consists of food flavourings (obtained after matching the CAS numbers with the EU food flavourings regulation). A total of 455 chemicals fell into the flavouring's category. Other relevant functional use classes included the group of European E-numbers (43 chemicals, food additives that perform a certain technological function in food, subdivided into sweeteners, antioxidants, preservatives, colours, and remaining E-numbers) nutrients (31 chemicals)

and regulatory restricted chemicals (19 chemicals). Chemicals that fell into multiple categories are counted multiple times, once for each category. For example, ascorbic acid is included as both an E-number (as a preservative) and nutrient (being also a vitamin). Chemicals for which no clear food use could be defined are grouped as "Other".

The clusters that were obtained based on the chemical structure are shown alongside the functional use classes in Figure 1 using different colours, representing the primary chemical clusters to which the chemical groups belong. Details on how the large primary clusters (e.g. alcohols) are further subdivided as appropriate into secondary (e.g. alcohol, alkyl) and tertiary (e.g. alcohol, alkyl, primary, straight chain) groups can be found in Table 1. Chemicals that fell into multiple chemical groups are counted multiple times, once for each group. For example, ascorbic acid falls into the "Ester, Lactone, Ascorbic acid and derivatives" group as well as the "Vitamines and derivatives" chemical group. Overall, 102 tertiary groups were defined for which the biological activity was explored. These tertiary groups consist of at least three closely related chemicals, with the largest chemical group consisting of seventeen (group 12, alkyl substituted phenols) chemicals. The majority of the defined chemical groups has a mean Tanimoto coefficient that is higher than 0.6. Some of the chemical groups are more structurally diverse and are atypical of the wider set, holding as they do compounds which exhibit unique characteristics. For example, the "amino acids and derivatives" grouping consists of a series of complex, often natural products, whereas "metallic salts organic" is founded solely upon the possession of an inorganic counter ion. Azo dyes furthermore represent a collection of compounds which may exhibit variation in wider structure despite unification by a distinctive functional group."

3.2 Global evaluation of the biological activity of the tertiary homologous chemical groups

3.2.1. ToxCast biological activities plotted as a heatmap

Figure 2 displays different heatmaps demonstrating the activity of the 102 defined tertiary structural groupings towards different biological targets of the different target families and Figure 3 of the "DNA binding" and "nuclear receptor target families" in detail. White spots in the heatmaps represent chemical group-biological target combinations for which insufficient data are available ($n < 3$ in all assays that cover that biological target). Grey means that all the chemicals within the group were inactive in the assays for that target, while the colours ranging from orange to red represents an increasing percentage of chemicals within the chemical group that responded in the assays of that target.

It is clear from the number of white areas in the heatmaps of Figure 2 that there are significant data gaps in ToxCast for the food-relevant chemicals, indicating that the ToxCast data set is not yet comprehensive for some of these types of food chemicals. Therefore, it is important to note that a lack of observed activity in the summarized findings should not be construed as indicative of inactive food-relevant chemicals, but that this is often the consequence of insufficient data. This observation may not be unique to food-relevant chemicals; other test substances within the wider ToxCast dataset beyond the scope of this inquiry may be as yet insufficiently tested for any broad conclusions to be made regarding their biological activities. Among the different target families, most of the food-relevant chemicals were tested in assays that are linked to the "DNA binding" and "nuclear receptor" target families (i.e. most grey/colour). The food-relevant chemicals have also been tested in assays that are linked to "cell cycle", "growth factor", "hydrolase", and "steroidal hormone". However, these latter target families consist of only one to three targets each (few y-axis tick marks), whereas the "DNA binding" and "nuclear receptor" target families consist of thirty-six and forty targets, respectively. Further evaluations therefore focus on the activities within these latter two target families.

The activities within the "DNA binding" and "nuclear receptor" target families are further highlighted in Figure 3, in which the "nuclear receptor" target family is also subdivided into its three distinctive subfamilies ("steroidal", "non-steroidal", and

“orphan”). Figure 3 reveals that most tertiary chemical groups are not active towards most of biological targets (i.e. 0% of the chemicals in the tertiary group showed activity in the assays of the target) within the “DNA binding” and “nuclear receptor” target families. One notable exception is chemical group 55 (containing parabens and gallates) that stands out in Figure 3 for its high activity towards ESR1|ESR2 (corresponding to ER α and ER β), ESR1 (ER α) and ESR2 (ER β). Other relevant chemical groups in Figure 3 are group 93 (containing thiols), which has a relatively high group activity at a variety of DNA binding targets, group 79 (containing genistein and daidzein), which has a high activity towards ESR1, ESR/ESR2, and ESR2, and group 101 (retinoids), which has a relatively high group activity towards RXRA, RAXRB, NR1I2 and NRF2.

Within the “DNA binding” and “nuclear receptor” target families there are a few targets for which almost all the chemical groups appear to be active. Examples are the NFE2L2, RXRA, RXRB, NR1I2, and ESR1 targets (horizontal orange stripes in Figure 3). Particularly, the frequent responses towards NFE2L2, RXRA, RXRB, and NR1I2 are likely because these are relatively general endpoints that are involved in increasing metabolic capacity or oxidative stress response (Mazaira *et al.*, 2019; Lousse *et al.*, 2018). However, to some extent this frequent activity also appears to be due to a proportionately high number of chemicals that are active in certain individual assays that fall under these targets (Ryan, 2017). For example, among the different assays that measure effects on ESR1, 10% of the food-relevant chemicals were active in the ATG_ERE_CIS_up assay and 7% in the TOX21_ERa_LUC_BG1_Agonist assay, whereas only 0.4 to 3% of the chemicals were active in other assays that measure ESR1 (see Supporting information 2), suggesting that the high positive rate in some assays might be an artefact. In addition, 12% of the food relevant chemicals were active in the ATG_NRF2_ARE_CIS_up assay (NFE2L2 target), 25% in the ATG_PXRE_CIS_up assay (NR1I2 target), TOX21_RXR_BLA_Agonist_ratio assay (RXRA target), and 7% in the ATG_RXRb_TRANS_up (RXRB target), with much lower rates in other assays for these targets. This suggests that care should be taken in the interpretation of the ToxCast results when activity towards a biological target is due to activity in one of these specific

assays that generate a high number of positive results. Supporting information 2 provides a list of the ToxCast assays, the percentage of the food relevant chemicals that were active in each assay, and the percentage of the results that contained flags. Based on these data the specificity of the different assays can be assessed, which is highly relevant for the interpretation of the test results for the individual chemical groups.

3.2.2. Biological activities of the chemical groups in the context of their functional uses

Figure 4 combines the ToxCast activity data with the circle pack of Figure 1, providing an indication of the biological activities of the groups of food-relevant chemicals in the context of their functional uses. To this end, for each chemical group the percent activities in the assays that belong to a specific target family was calculated. For example, the distinct activities of group 55 (parabens-gallates), 79 (containing genistein and daidzein), group 98 (retinoids), and group 90 (thiols) as was observed in Figure 3, result in an overall high activity of these groups in the "steroidal nuclear receptor" (group 55 and 79), "nonsteroidal" (group 101) and "dna binding" target (sub)families (group 93) in Figure 4. Figure 4 also reveals that many chemical groups are slightly active within the steroidal and non-steroidal nuclear receptor target families. These activities generally relate to activities in the assays with a disproportionately high number of positive hits and/or assays that capture general response mechanism to chemical exposure, as discussed above.

Some of the tertiary structural groups in Figure 4 consist of chemicals that fall into different functional use classes. In those cases, the activity that is displayed in Figure 4 corresponds to percent activity of the chemicals that fall into the same use class and not the activity of the whole group. For example, Figure 4 reveals that the group of parabens (group 55) contains both regulatory restricted compounds and compounds that are used as antioxidants and preservatives. Particularly the restricted parabens and parabens used as preservative appear to have activity in the steroidal nuclear receptor target family of assays. In comparison, group 79 consists of polyphenols (genistein,

daidzein, and quercetin) and of different flavourings (for example coumarin). Figure 4 reveals that the high steroidal-nuclear receptor activity of group 79 comes only from the polyphenols of group 79 and not from the flavourings.

The results from the heat map (Figure 3) and circle pack (Figure 4) reveal that some of the key biological targets of the food-relevant chemicals can be defined by focussing on ToxCast activities of predefined homologous chemical groups. The circle pack plot (Figure 4) provides insight into the overall biological activities of the tertiary chemical groups within the "DNA binding" and "(steroidal, non-steroidal, and orphan) nuclear receptor" target (sub)families and places the results in the context of the functional uses. The heatmap of Figure 3 provides insights into the specific targets within these target (sub)families that are affected. Based on these results several case studies were defined to explore how the ToxCast data can be used in food safety risk evaluations and for the evaluation of desirable effects of nutrients and phytonutrients. The case studies are used to check whether the mechanistic information that is obtained from ToxCast matches with what is expected from the chemical group. To this purpose, case studies were selected around chemical groups that express a high biological activity towards a specific target (parabens), chemicals that are restricted for food use due a specific activity (some parabens and genotoxic and carcinogenic compounds like estragole, methyleugenol and safrole), and chemical groups are related to specific health benefits (e.g. flavonoids and fatty acid). For the selected case studies sufficient literature data is available on the mechanisms of action of the compounds. The comparison of the observed target(s) with expected target(s) is considered a crucial step to find potential caveats in the HTS data that need to be considered for future use of the data on chemicals for which little animal experimental or *in vitro* reference data is available.

3.4.1 Case study on regulatory restricted chemicals

The group of regulatory restricted chemicals provides an interesting group of food-relevant chemicals for the evaluation of ToxCast activities. Several compounds that are

restricted for food use in regulations can be found within the ToxCast data set of food-relevant compounds. Most of these compounds have an E-number (EU codes for substances that are permitted as food additive) yet have been discontinued for food use in the EU. The exact reasons for the discontinuation are not always clear but do not necessarily relate to the demonstrable toxicity of the chemical. For example, ethoxyquin (E324) was suspended from its authorisation as a feed additive in (EU, 2017) because of a lack of data on some aspects of its safety, but it is currently being re-evaluated by EFSA again for this use (EFSA, 2019). However, for two chemical groups within the regulatory restricted group, demonstrable toxicological findings have played an important role in their restriction for food use. These are group 55 (containing parabens and gallates, which have estrogenic activities) and groups 70 and 77 (containing estragole, methyleugenol, and safrole, which are genotoxic and carcinogenic (Phillips *et al.*, 1984). It is of interest to observe whether for these groups a perturbation of the underlying biological target responsible for the restrictions can be detected with the goal of determining the potential contribution of ToxCast data in such evaluations of food safety risk.

3.4.1.1 Restricted and non-restricted parabens and gallates

The ToxCast evaluation of the biological activities of structural groups detected a relative high activity of the paraben-gallate (group 55) towards ER alpha and beta activation (ESR1, ESR2, ESR1|ESR2 in Figure 3). This structural group consists of the approved E-numbers methylparaben (preservative), dodecyl gallate, octyl gallate and propyl gallate (used as anti-oxidants); two parabens that are not used in foods within the EU; butyl- and heptylparaben; and the restricted propylparaben (EFSA, 2004). Figure 4 reveals that the biological activity within the paraben-gallate mainly comes from the restricted and non-restricted parabens (preservatives) rather than the gallates that are used as anti-oxidants. Particularly the percentage of positive assays towards the steroidal nuclear receptors (Fig. 4A) was higher for parabens than gallates. Positives were largely for estrogenic assays. The differences within the group of parabens-gallates is also reflected

in the relative estrogenic potencies of these two groups; the mean AC50 of the positive ER alpha assay results for the restricted/non-approved parabens (with Z-scores > 3) is 10.9 ± 4.7 μM for propylparaben, 5.2 ± 1.4 μM for butylparaben, and 3.7 ± 0.6 for heptylparaben, whereas the non-restricted methylparaben has a lower potency with a mean AC50 of 53 ± 18 μM . Dodecyl and propyl gallate were not active in the ER receptor assays in the final dataset and octyl gallate is active in only one out of the eight ER receptor assays. It should, however, be noted that many of the positive ER hit-calls of the gallates appeared to have been filtered out as a result of their low Z-scores. This suggests a potential influence of, for example, cytotoxicity, on the estrogen receptor results of the different gallates.

Altogether, the results obtained for group 55 reveal an interesting potency difference between the parabens and gallates within this group. These results provide relevant information that can be considered within the regulatory risk assessments of these compounds, particularly to perform a read-across. The estrogenic hazard potential of the parabens has long been included in their risk assessment (EFSA, 2004; SCCS, 2013; EMA, 2015). In case of gallates, the potential these compounds to interfere with the human estrogen receptor *in vitro* has recently been mentioned by EFSA in re-evaluations of dodecyl, octyl and propyl gallate, but has not been included in their final risk evaluation as confirmatory *in vivo* data are lacking (EFSA, 2015a; 2015b; 2015c). Based on the results of the present study the association between gallates and the estrogen receptor was considered low, but follow-up *in vitro* studies may be needed to better understand the origin of the low Z-scores for the gallates in the ER-related assays that had to be dismissed on the basis of these low Z-scores.

3.4.1.3 Regulatory restricted genotoxic carcinogens

Evaluation of the ToxCast activities of chemical group 70 and 77 that contain the known genotoxic and carcinogenic compounds estragole (group 70), methyleugenol (group 70), and safrole (group 77) (SCF, 2001) shows that the hazard of these type of compounds cannot be adequately identified in ToxCast. In Figure 4, groups 70 and 77 can be found

to have a slight activity within the non-steroidal nuclear receptor target family. However, based on Figure 4, it can be concluded that this activity relates to activation of RXRB and NR1I2 (also called PXR), targets that induce xenobiotic metabolism enzyme synthesis and for which many non-carcinogenic chemical groups are active. Hence this slight activity towards RXRB and NR1I2 is not considered diagnostic for the genotoxic hazard of these compounds nor for potential other specific mechanisms of actions. Given the genotoxic mechanisms of estragole, methyleugenol and safrole, activity in assays that include p53 tumor suppressor gene activity might be expected (Paini *et al.*, 2011). Such activity forms part of the "DNA binding" target family. However, no such activity was found, nor was there any other indication of genotoxicity for these chemicals within the ToxCast dataset.

Since estragole, methyleugenol, and safrole require bioactivation for their genotoxic and carcinogenic effects (SCF, 2001; Punt *et al.*, 2007), this lack of detectable activity could be due to the lack of metabolic capacity within the ToxCast assays (DeGroot *et al.*, 2018). Moreover, genotoxicity can be difficult to detect without very tight concentration spacing, as the high-throughput assays often quantify the upregulation of DNA-repair pathways (Iyer *et al.*, 2019). While these assays quantify the cellular mechanisms evolved to fix low-level DNA damage, when the damage-levels are great, cells instead die without attempting repair, resulting in false-negative tests. Therefore, it is also possible that the lack of detectable genotoxic activity could be due to the large concentration-spacing used for ToxCast testing and the subsequent misclassification of genotoxicity as cytotoxicity. Taken together, these findings suggest that currently genotoxicants and/or carcinogens cannot always be adequately detected within the ToxCast activity data, which is supported by other literature findings (Becker *et al.*, 2017).

3.4.2 Use of ToxCast data in assessment of beneficial effects

The flavonoids genistein, daidzein, and quercetin; unsaturated fatty acids like linolenic, linoleic and oleic acid are all examples of compounds with health-beneficial effects that

can be found within the set of food-relevant chemicals used in the present study. For these substances the relationship between beneficial and adverse biological effects is of particular interest. Therefore, the ToxCast data from these two substance-groups was examined to characterize the biological targets of each group of health beneficial chemicals and evaluate how the ToxCast information might be used to inform a risk-benefit assessment of the compounds.

3.4.2.1 Flavonoids

Flavonoids have been extensively studied for their biological effects against cancer, cardiovascular diseases, obesity and diabetes, as well as neurodegenerative disorders (Williamson *et al.*, 2018). Within the set of food-relevant chemicals of the present study, three flavonoids are included: quercetin and the isoflavones daidzein and genistein. All are part of the “heterocycles and polycycles- oxygen heterocycles-pyranone”-chemical group (group 79). Other chemicals that are part of this group are maltol, 2-ethyl-3-hydroxy-4-pyrone, maltol isobutyrate, coumarin, and 6-methyl coumarin. The molecular targets that have been suggested to play a predominant role in the health beneficial effects of flavonoids are displayed in Table 2.

In contrast to what was expected, no activity of group 79 was found for most of the targets described in Table 2, except for activity in ER-related assays and NRF2. Similarly, when examining the data for each individual flavonoid in the group it is clear that this group is made up of diverse substances; each has relatively few identifiable activities with little overlap with those of other members of the structural group.

Interestingly, the apparent biological activities of flavonoids are significantly affected by the filtering out of results with Z-scores less than three. Without this filtering, quercetin, genistein, and daidzein are active in 19.4%, 31.3%, and 23.6% of the assays, respectively, that are part of the target (sub)families of Figure 2, whereas they are active in only 1%, 11%, and 10% of these assays after filtering. Judson *et al.* (2016) also identified quercetin as a highly active chemical within ToxCast, with low specificity (low

Z-scores). Many of the effects of the chemicals in group 79 are thus filtered out as being non-specific. Though results with a Z-score lower than three may reflect an indirect influence of cytotoxicity or other non-specific mechanisms of action, the low Z-scores may also point to a non-specific interference with the assays. For example, flavonoids are capable of stabilization of luciferase, frequently used in reporter gene assays (Prinsloo *et al.*, 2017). The high number of assay-results with low Z-scores indicate that challenges exist in using HTS to explore the biological activities of certain compound such as flavonoids.

Although the ToxCast activities of flavonoids seem uncertain due to possible non-specific effects, group 79 does show a distinct activity within the “nuclear receptor-steroidal” target (sub)family towards the estrogen receptor alpha and beta. This result is predominantly due to activity of the isoflavones genistein and daidzein (see Figure 4). The interaction of genistein and daidzein with the ER receptor has been linked to both beneficial health effects (for example, lowering menopausal symptoms, lowering cancer risks and risk for cardiovascular diseases) and adverse effects (endocrine disruption, increased hormone cancer risk) (Rietjens *et al.*, 2017). A key hypothesis behind the benefits and risks of isoflavonoids is the differences between the activation of ER α and ER β . ER α activation enhances cell proliferation, whereas ER β counteracts the ER α -mediated stimulation of cell proliferation (Rietjens *et al.*, 2017). Though many estrogenic compounds within ToxCast interact with both ER α and ER β (including the parabens as described above), the AC50 values for genistein and daidzein were 20 and 11-fold lower for ER β compared with ER α (based on the OT_ER_ER α ER α _0480/1440 and OT_ER_ER β ER β _0480/1440 assays), respectively, suggesting a predominantly ER β -mediated effect at low concentrations (This is not the case for the parabens which have comparable AC50 values towards ER α and ER β for example). This selective estrogen receptor modulation suggests that the risk-benefit profile of genistein and daidzein is probably dose-dependent but must be extrapolated to *in vivo* dose-response or potency information to identify whether the effective concentrations *in vitro* are capable of being attained *in vivo*. For genistein, this has for example been done by Boonpawa *et al.*

(2017), revealing that both Asian dietary intake levels and the use of genistein-containing supplements are sufficient for ER β activation, but not for ER α modulation. Thus, the *in vitro* potency information over a range of ToxCast targets can be used to prioritize measurement and evaluation of *in vivo* biological effects within the context of risk-benefit assessments.

3.4.2.2 Fatty acids

Fatty acids, particularly unsaturated fatty acids, play a key role in reducing cardiovascular risks and anti-inflammatory effects (Williams 2000). The set of food-relevant chemicals of the present study contains a series of both unsaturated (group 22) and saturated (group 24) fatty acids. Group 22 consists of 2-butenoic-, sorbic-, 10-undecenoic-, oleic-, linolenic-, and linoleic-acid. Group 24 consists of acetic acid, butanoic-, pentanoic-, hexanoic-, heptanoic-, octanoic-, decanoic-, dodecanoic-, tetradecanoic-, hexadecenoic (palmitic)-, and octadecanoic(stearic)- acid.

An important mode of action of fatty acids is the regulation of lipid metabolism (Varga *et al.*, 2011). For example, Popeijus *et al.* (2014) have shown that fatty acid chain length and saturation influences PPAR α transcriptional activation and repression in HepG2 cells, and specifically the saturated fatty acids palmitic acid (C16:0) and stearic acid (C18:0) both repress PPAR α activation, whilst their unsaturated metabolites palmitoleic acid (C16:1(*n*-7)) and oleic acid (C18:1(*n*-9)) activate PPAR transcription. Other potentially relevant targets of fatty acids within lipid homeostasis are SREBPs, LXR, and HNF4 (Müller and Kersten, 2003). Table 3 provides an overview of the ToxCast activity of the chemical groups containing fatty acids towards these different targets.

The ToxCast activity of both groups 22 and 24 towards the expected targets of Table 3 appears to be very limited. For example, within group 22 only 10-undecenoic acid is active in two PPAR α -related assays, and within group 24 the PPAR α activity mainly comes from decanoic acid. The low responses seem to be partly due to the filtering based on Z-scores lower than three. Without filtering, all the long chain fatty acids of the unsaturated

fatty acid group (oleic acid, 10-undecenoic acid, linoleic acid and linolenic acid) are active in the ATG_PPARa_TRANS_up assay, which is in line with what is expected (Popeijus *et al.*, 2014). This raises questions as to whether AC50s with low Z-scores should indeed be considered to be the result of non-specific activities and as to what causes these low Z-scores. The low observed biological activity of the saturated fatty acids group towards PPARa (either up or down regulation) was not affected by the Z-score filtering.

The activity towards the other potential molecular targets of saturated and unsaturated fatty acids are also limited but does not seem to be caused by the filtering based on Z-scores. In the case of SREBPs, the limited activity might be the result of to the fact that unsaturated fatty acids are down regulators (Hannah *et al.*, 2001), whereas ToxCast only contains the ATG_SREBP_CIS_up assay. The LXR receptor, which is involved in the regulation of cholesterol and fatty acid homeostasis was not active as a Toxcast assay target for group 22 and 25, which may be a reflection of it being responsive to HNF4A intracellular cholesterol alterations (Lund *et al.*, 2006). Overall, the results indicate that the ToxCast dataset is at present not yet adequate to obtain insights into the biological activities of fatty acids or, for example, for the extrapolation of the potential effects over different chain-lengths of fatty acids.

3.5 Remaining relevant groups

Table 4 provides a list of remaining relevant chemical groups that display a relatively high activity as displayed in Figures 3 and 4, but which were not assessed further as case studies. The observed activities of these groups generally relate to endpoints such as increased metabolic capacity or oxidative stress response. However, the targets that are affected by the Thiol group (group 93) within the "DNA binding" target family may point at a specific activity of this chemical group that is potentially relevant for the safety evaluation of this chemical group.

4. Discussion

This investigation aimed to i) explore how HTS can be leveraged to obtain chemical-specific insights into the biological targets that may be affected by different food-relevant chemicals, and ii) assess the utility of the data for the safety assessment of food chemicals as well as the evaluation of health beneficial effects of chemicals in a few case studies. A method was set-up to group the chemicals according to functional use and structural similarity. For each of the tertiary chemical groups of homologous chemicals, the percent of chemicals that were active in the assays for different targets and target families were calculated. The targets that are elicited (and directionality thereof, e.g. activation vs down-regulation), but also the inactivity towards certain targets can provide key information in characterizing biological patterns.

A general challenge in the use of HTS data in chemical safety evaluations is the uncertainty around the individual assay results. The diverse assay space and challenges with automatic processing contribute to this uncertainty (Watt and Judson, 2018; Ryan, 2017; Cox *et al.*, 2014). The approach of the present study, in which the focus is not on the individual chemical results but on the activity of homologues within chemical groups may contribute to reducing the uncertainty and improving the specificity when defining biological targets of chemicals based on HTS data. Taking the example of genistein, this compound is active in 114 assays within the crude ToxCast data set, with 49% of these results containing flags. Removing all data with Z-scores lower than three as a cut-off value for non-specific effects, the number of positive hit calls is reduced to 16, of which 30% contain flags. Combining the results with the other flavonoids within the list of food-relevant chemicals, including daidzein, points towards the expected ER activation as the most predominant biological effect. Both genistein and daidzein tested positive in 11 different ESR1 and/or ESR2 assays with one flagged result for genistein.

Though these results indicate that the specificity increases by filtering the ToxCast data for Z-scores and focussing on targets that are induced by homologous chemical groups, such filtering may also result in a potential loss of information. In the different

case studies, filtering of Z-scores lower than 3, sometimes appeared to eliminate valuable information. It is therefore also important to go back to the crude data to evaluate the filtering process prior to the use of ToxCast results for risk evaluations.

The (tertiary) chemical groups for which the ToxCast activities were assessed were obtained with ChemoTyper software (Yang *et al.*, 2015; Mellor *et al.*, 2019). Other methods for chemical grouping exist as well, including for example AMBIT (<http://cefic-iri.org/toolbox/ambit/>), the OECD QSAR toolbox (Dimitrov *et al.*, 2016) and ToxMatch (<https://ec.europa.eu/jrc/en/scientific-tool/toxmatch>). Key to the chemical grouping is that groups should become neither too large nor too small. The highest number of chemicals within one group that was obtained using ChemoTyper contained seventeen analogues. Most of the groups consist of three to five chemicals. Sixty-six of the originally 168 defined groups could not be used in the present study as these contained only one or two chemicals.

The fact that the number of chemicals within a group varies may pose some bias in the evaluation of the biological activity group targets. If two compounds in a group of three chemicals are active towards a specific target, this corresponds to 66% activity, whereas an activity of two compounds in a group of six would correspond to 33% activity. A similar bias in the results occurs due to the varying number of assays per target in the ToxCast data set. Many of the biological targets as displayed in Figure 4 are covered by only one or two assays, whereas for the ESR1 target there are 16 assays of which 11 measure estrogen receptor agonism. A high percentage activity of a chemical group in the case of ESR1 will therefore occur only when the chemicals of that group are active across a wide range of assays for ESR1. Even though these results indicate that for larger chemical groups and for targets that are covered by multiple assays it will be more difficult to pick up group activities, the results are expected to become more specific. For example, the activity of the compounds of the paraben-gallate (group 55) and combined activity of genistein and daidzein within the pyrole group (group 79) in multiple ESR1, ESR2 and ESR1|2-related assays give confidence that estrogen receptor activation is an important target for these groups of chemicals. This effect of enrichment was less

apparent within fatty acid and genotoxic carcinogen-case studies, due to the overall limited activity of the individual chemicals in the expected assays.

Whereas the ToxCast data may add to the evidence considered in food chemical safety evaluations (that can be used for read-across and risk assessments), the results should serve primarily as a screening tool to set priorities for further evaluations relative to hypothesized biological targets. Examples of the use of ToxCast data, particularly for read-across, can also increasingly be found in the literature (Blackburn *et al.*, 2019; Lizarraga *et al.*, 2019). Some care however should be taken to avoid overinterpretation of the data. Not all available chemicals within ToxCast have been tested in all available assays and not all toxicity endpoints are covered by the available assays. In addition, better understanding of relevance of data with low Z-scores may minimize loss of potentially relevant information. For individual cases where low Z-scores are found, follow-up analyses may be needed to identify what causes these low Z-scores. These facts are critical aspects of the dataset which we found crucial for proper data interpretation during the present study.

A more general aspect that needs to be kept in mind is the fact that metabolic activation of chemicals is not accounted for in the ToxCast assays. This probably contributed to the observed poor prediction of genotoxicity in the case studies. Research that focuses on enhancing the metabolic capacity in HTS assays is therefore important (DeGroot *et al.* 2018).

Finally, it should be noted that *in vitro* activity data do not directly reflect *in vivo* biological potencies (in vivo effects will for example also depend on the availability of a chemical in the body). Extrapolation of the concentration-response curves to *in vivo* potency information is an important next step. There are an increasing number of publications that focus on establishing such an extrapolation (Becker *et al.*, 2014; Punt *et al.*, 2019; Fabian *et al.*, 2019; Boonpawa *et al.*, 2017; Wetmore *et al.*, 2015; Dent *et al.*, 2019). One approach is to use kinetic modelling or human biomonitoring data to compare the AC50 values with internal plasma concentrations reached during daily exposures in so-called exposure:activity ratios, EARs. EARs of different compounds can subsequently

be compared in a so-called 'dietary comparator ratio' (DCR) approach to prioritise exposure-activity data relative a known reference compound (Becker *et al.*, 2014; Dent *et al.*, 2019).

To increase regulatory use of HTS it will be important to tackle the different challenges related to HTS and quantitative *in vitro* to *in vivo* extrapolations. Recently Thomas *et al.*, 2019 published a blueprint to systematically address these key challenges, which can be expected to move the field forward.

Overall, the results of the present study suggest HTS data could add to the evidence considered for regulatory risk assessments for food chemicals and to the evaluation of desirable effects of nutrients and phytonutrients. The data will be particularly useful for providing mechanistic information and to fill data gaps with read-across. Whereas the current study mainly focussed on setting up a method to find key biological targets of chemical groups and the qualitative interpretation thereof, the key next step for use in risk evaluations or follow-up research is to also focus on the quantitative aspects of the results. This includes, for example, the evaluation of the (differences in) potencies of the chemicals towards targets of interest and placing the potencies in the context of human *in vivo*-relevant exposure.

Acknowledgements

This work was conducted by an expert group of the European branch of the International Life Sciences Institute (ILSI Europe). We appreciate the contribution of Dr. Agnes Karmaus who gave us insights in the pre-existing work done on food-relevant chemicals in the ToxCast data set. We also appreciate the contribution of Sylvain Etter (Firmenich) and Heinz Traussnig (Mayr-Melnhof Karton) to the discussions on the work of the expert group. The expert group received funding from the ILSI Europe New Approaches to Chemical Risk Assessment for Food and Food Ingredients Task Force. Industry members of this task force are listed on the ILSI Europe website at www.ilsieurope.eu. For further information about ILSI Europe, please email info@ilsieurope.be or call +32 2771 00 14. The opinions expressed herein, and the conclusions of this publication are those of the authors and do not necessarily represent the views of ILSI Europe nor those of its member companies.

References

- Becker, R.A., Hays, S.M., Kirman, C.R., Aylward, L.L., and Wise, K. (2014) Interpreting Estrogen Screening Assays in the Context of Potency and Human Exposure Relative to Natural Exposures to Phytoestrogens. *Birth Defects Res. Part B Dev. Reprod. Toxicol.*, **101**, 114–124.
- Blackburn, K.L., Ellison, C.A., Stuard, S.B., and Wu, S. (2019) Dosimetry considerations for in vivo and in vitro test data and a novel surrogate iTTC approach for read-across based on metabolites. *Comput. Toxicol.*, **10**, 145–157.
- Boonpawa, R., Spenkelink, A., Punt, A., and Rietjens, I.M.C.M. (2017a) In vitro-in silico-based analysis of the dose-dependent in vivo oestrogenicity of the soy phytoestrogen genistein in humans. *Br. J. Pharmacol.*, **174**, 2739–2757.
- Boonpawa, R., Spenkelink, A., Punt, A., and Rietjens, I.M.C.M. (2017b) *In vitro-in silico* -based analysis of the dose-dependent *in vivo* oestrogenicity of the soy phytoestrogen genistein in humans. *Br. J. Pharmacol.*, **174**, 2739–2757.
- Brunner, A.M., Dingemans, M.M.L., Baken, K.A., and van Wezel, A.P. (2019) Prioritizing anthropogenic chemicals in drinking water and sources through combined use of mass spectrometry and ToxCast toxicity data. *J. Hazard. Mater.*, **364**, 332–338.
- Cao, Y., Charisi, A., Cheng, L.-C., Jiang, T., and Girke, T. (2008) ChemmineR: a compound mining framework for R. *Bioinformatics*, **24**, 1733–1734.
- Cox, L.A. (Tony), Popken, D., Marty, M.S., Rowlands, J.C., Patlewicz, G., Goyak, K.O., and Becker, R.A. (2014) Developing scientific confidence in HTS-derived prediction models: Lessons learned from an endocrine case study. *Regul. Toxicol. Pharmacol.*, **69**, 443–450.
- Csardi G and Nepusz T (2006) The igraph software package for complex network research, InterJournal, Complex Systems. *InterJournal*, **Complex Systems**, 1695.
- DeGroot, D.E., Swank, A., Thomas, R.S., Strynar, M., Lee, M.-Y., Carmichael, P.L., and Simmons, S.O. (2018) mRNA transfection retrofits cell-based assays with xenobiotic metabolism. *J. Pharmacol. Toxicol. Methods*, **92**, 77–94.
- Dent, M.P., Li, H., Carmichael, P.L., and Martin, F.L. (2019) Employing Dietary

Comparators to Perform Risk Assessments for Anti-Androgens Without Using Animal Data. *Toxicol. Sci.*, **167**, 375–384.

DG SANTE (2011) EU database on Food Additives.

https://webgate.ec.europa.eu/foods_system/main/?sector=FAD&auth=SANCAS
(accessed 12 Sept 2019).

Dimitrov, S.D., Diderich, R., Sobanski, T., Pavlov, T.S., Chankov, G. V., Chapkanov, A.S., Karakolev, Y.H., Temelkov, S.G., Vasilev, R.A., Gerova, K.D., Kuseva, C.D., Todorova, N.D., Mehmed, A.M., Rasenberg, M., and Mekenyan, O.G. (2016) QSAR Toolbox – workflow and major functionalities. *SAR QSAR Environ. Res.*, **27**, 203–219.

Dorne, J. Lou, Richardson, J., Kass, G., Georgiadis, N., Monguidi, M., Pasinato, L., Cappe, S., Verhagen, H., and Robinson, T. (2017) Editorial: OpenFoodTox: EFSA’s open source toxicological database on chemical hazards in food and feed. *EFSA J.*, **15**.

EFSA (2017) OpenFoodTox database.

<https://www.efsa.europa.eu/en/microstrategy/openfoodtox> (accessed 12 Sept 2019).

EFSA (2004) Opinion of the Scientific Panel on food additives, flavourings, processing aids and materials in contact with food (AFC) related to para hydroxybenzoates (E 214-219). *EFSA J.*, **2**, 83.

EFSA (2019) Scientific panel on additives and products or substances used in animal feed. Minutes of the 106th meeting of the working group on technological additives. <https://www.efsa.europa.eu/sites/default/files/wgs/animal-feed/wg-technological-additives-2018-2021>.

EMA (2015) Reflection paper on the use of methyl- and propylparaben as excipients in human medicinal products for oral use.

<https://www.ema.europa.eu/en/documents/scientific-guideline/reflection-paper-use-methyl-propylparaben-excipients-human-medicinal-products-oral>.

EU (European Union) (2017) Commission Implementing Regulation (EU) 2017/962 of 7 June 2017 suspending the authorisation of ethoxyquin as a feed additive for all animal

- species and categories. *Off J Eur Comm L* 145: 13–17 (accessed 12 Sept 2019).
- Fabian, E., Gomes, C., Birk, B., Williford, T., Hernandez, T.R., Haase, C., Zbranek, R., van Ravenzwaay, B., and Landsiedel, R. (2019) In vitro-to-in vivo extrapolation (IVIVE) by PBTK modeling for animal-free risk assessment approaches of potential endocrine-disrupting compounds. *Arch. Toxicol.*, **93**, 401–416.
- Han, X., Beaumont, C., and Stevens, N. (2017) Chemical composition analysis and in vitro biological activities of ten essential oils in human skin cells. *Biochim. Open*, **5**, 1–7.
- Hannah, V.C., Ou, J., Luong, A., Goldstein, J.L., and Brown, M.S. (2001) Unsaturated fatty acids down-regulate srebp isoforms 1a and 1c by two mechanisms in HEK-293 cells. *J. Biol. Chem.*, **276**, 4365–72.
- Hartman, J.K., Beames, T., Parks, B., Doheny, D., Song, G., Efremenko, A., Yoon, M., Foley, B., Deisenroth, C., McMullen, P.D., and Clewell, R.A. (2018) An in vitro approach for prioritization and evaluation of chemical effects on glucocorticoid receptor mediated adipogenesis. *Toxicol. Appl. Pharmacol.*, **355**, 112–126.
- Houck, K.A., Judson, R.S., Knudsen, T.B., Martin, M.T., Richard, A.M., Crofton, K.M., Simeonov, A., Paules, R.S., Bucher, J.R., and Thomas, R.S. (2017) Comment on “On the Utility of ToxCast™ and ToxPi as Methods for Identifying New Obesogens”;. *Environ. Health Perspect.*, **7425**, A8–A11.
- Iyer, S., Pham, N., Marty, M., Sandy, M., Solomon, G., and Zeise, L. (2019) An Integrated Approach Using Publicly Available Resources for Identifying and Characterizing Chemicals of Potential Toxicity Concern: Proof-of-Concept With Chemicals That Affect Cancer Pathways. *Toxicol. Sci.*, **169**, 14–24.
- Jelen, H. (2012) Food flavors : chemical, sensory and technological properties CRC Press.
- Judson, R., Houck, K., Martin, M., Richard, A.M., *et al.* (2016) Analysis of the Effects of Cell Stress and Cytotoxicity on *In Vitro* Assay Activity Across a Diverse Chemical and Assay Space. *Toxicol. Sci.*, **153**, 409–409.
- Judson, R.S., Magpantay, F.M., Chickarmane, V., Haskell, C., *et al.* (2015) Integrated Model of Chemical Perturbations of a Biological Pathway Using 18 In Vitro High-

- Throughput Screening Assays for the Estrogen Receptor. *Toxicol. Sci.*, **148**, 137–54.
- Karmaus, A.L., Trautman, T.D., Krishan, M., Filer, D.L., and Fix, L.A. (2017) Curation of food-relevant chemicals in ToxCast. *Food Chem. Toxicol.*, **103**, 174–182.
- Karmaus, A.L., Filer, D.L., Martin, M.T., and Houck, K.A. (2016) Evaluation of food-relevant chemicals in the ToxCast high-throughput screening program. *Food Chem. Toxicol.*, **92**, 188–196.
- Kavlock, R., Chandler, K., Houck, K., Hunter, S., Judson, R., Kleinstreuer, N., Knudsen, T., Martin, M., Padilla, S., Reif, D., Richard, A., Rotroff, D., Sipes, N., and Dix, D. (2012) Update on EPA's ToxCast Program: Providing High Throughput Decision Support Tools for Chemical Risk Management. *Chem. Res. Toxicol.*, **25**, 1287–1302.
- Kleinstreuer, N.C., Yang, J., Berg, E.L., Knudsen, T.B., Richard, A.M., Martin, M.T., Reif, D.M., Judson, R.S., Polokoff, M., Dix, D.J., Kavlock, R.J., and Houck, K.A. (2014) Phenotypic screening of the toxcast chemical library to classify toxic and therapeutic mechanisms. *Nat. Biotechnol. Vol.*, **32**.
- Lizarraga, L.E., Dean, J.L., Kaiser, J.P., Wesselkamper, S.C., Lambert, J.C., and Zhao, Q.J. (2019) A case study on the application of an expert-driven read-across approach in support of quantitative risk assessment of p,p'-dichlorodiphenyldichloroethane. *Regul. Toxicol. Pharmacol.*, **103**, 301–313.
- Louisse, J., Dingemans, M.M.L., Baken, K.A., van Wezel, A.P., and Schriks, M. (2018) Exploration of ToxCast/Tox21 bioassays as candidate bioanalytical tools for measuring groups of chemicals in water. *Chemosphere*, **209**, 373–380.
- Lund, E.G., Peterson, L.B., Adams, A.D., Lam, M.-H.N., Burton, C.A., Chin, J., Guo, Q., Huang, S., Latham, M., Lopez, J.C., Menke, J.G., Milot, D.P., Mitnaul, L.J., Rex-Rabe, S.E., Rosa, R.L., Tian, J.Y., Wright, S.D., and Sparrow, C.P. (2006) Different roles of liver X receptor α and β in lipid metabolism: Effects of an α -selective and a dual agonist in mice deficient in each subtype. *Biochem. Pharmacol.*, **71**, 453–463.
- Mayr, L.M. and Fuerst, P. (2008) The Future of High-Throughput Screening. *J. Biomol. Screen.*, **13**, 443–448.
- Mazaira, G.I., Zgajnar, N.R., Lotufo, C.M., Daneri-Becerra, C., Sivils, J.C., Soto, O.B.,

- Cox, M.B., and Galigniana, M.D. (2019) Nuclear Receptors: A Historical Perspective. Humana, New York, NY, pp. 1–5.
- Mellor, C.L., Marchese Robinson, R.L., Benigni, R., Ebbrell, D., Enoch, S.J., Firman, J.W., Madden, J.C., Pawar, G., Yang, C., and Cronin, M.T.D. (2019) Molecular fingerprint-derived similarity measures for toxicological read-across: Recommendations for optimal use. *Regul. Toxicol. Pharmacol.*, **101**, 121–134.
- Müller, M. and Kersten, S. (2003) Nutrigenomics: goals and strategies. *Nat. Rev. Genet.*, **4**, 315–322.
- Olker, J.H., Korte, J.J., Denny, J.S., Hartig, P.C., Cardon, M.C., Knutsen, C.N., Kent, P.M., Christensen, J.P., Degitz, S.J., and Hornung, M.W. (2019) Screening the ToxCast Phase 1, Phase 2, and e1k Chemical Libraries for Inhibitors of Iodothyronine Deiodinases. *Toxicol. Sci.*, **168**, 430–442.
- Paini, A., Guignard, G., Bezencon, C., Latado, H., Schilter, B., van Bladeren, P.J., Rietjens, I.M.C.M., and Marin-Kuan, M. (2011) DNA damage response induced by 1'-hydroxyestragole in rat primary hepatocytes. In, *Generation of in vitro data to model dose dependent in vivo DNA binding of genotoxic carcinogens and its consequences: the case of estragole*. <http://edepot.wur.nl/200999> (accessed 12 Sept 2019).
- Pedersen, T.L. (2017) ggraph: An Implementation of Grammar of Graphics for Graphs and Networks. <https://cran.r-project.org/package=ggraph> (accessed 12 Sept 2019).
- Phillips, D.H., Reddy, M.V., and Randerath, K. (1984) ³²P-Post-labelling analysis of DNA adducts formed in the livers of animals treated with safrole, estragole and other naturally-occurring alkenylbenzenes. II. Newborn male B6C3F₁ mice. *Carcinogenesis*, **5**, 1623–1628.
- Popeijus, H.E., van Otterdijk, S.D., van der Krieken, S.E., Konings, M., Serbonij, K., Plat, J., and Mensink, R.P. (2014) Fatty acid chain length and saturation influences PPARα transcriptional activation and repression in HepG2 cells. *Mol. Nutr. Food Res.*, **58**, 2342–2349.
- Prinsloo, G., Papadi, G., Hiben, M.G., de Haan, L., Lousse, J., Beekmann, K., Vervoort,

- J., and Rietjens, I.M.C.M. (2017) In vitro bioassays to evaluate beneficial and adverse health effects of botanicals: promises and pitfalls. *Drug Discov. Today*, **22**, 1187–1200.
- Punt, A., Aartse, A., Bovee, T.F.H., Gerssen, A., van Leeuwen, S.P.J., Hoogenboom, R.L.A.P., and Peijnenburg, A.A.C.M. (2019) Quantitative in vitro-to-in vivo extrapolation (QIVIVE) of estrogenic and anti-androgenic potencies of BPA and BADGE analogues. *Arch. Toxicol.*, **93**, 1941–1953.
- Richard, A.M., Judson, R.S., Houck, K.A., Grulke, C.M., Volarath, P., Thillainadarajah, I., Yang, C., Rathman, J., Martin, M.T., Wambaugh, J.F., Knudsen, T.B., Kancherla, J., Mansouri, K., Patlewicz, G., Williams, A.J., Little, S.B., Crofton, K.M., and Thomas, R.S. (2016) ToxCast Chemical Landscape: Paving the Road to 21st Century Toxicology. *Chem. Res. Toxicol.*, **29**, 1225–1251.
- Rietjens, I.M.C.M., Louisse, J., and Beekmann, K. (2017) The potential health effects of dietary phytoestrogens. *Br. J. Pharmacol.*, **174**, 1263–1280.
- Ryan, N. (2017) A User's Guide for Accessing and Interpreting ToxCast Data.
- SCCS (2013) Scientific Committee on Consumer Safety (SCCS) Opinion on parabens. Updated request for a scientific opinion on propyl- and butylparaben. https://ec.europa.eu/health/scientific_committees/consumer_safety/docs/sccs_o_132.pdf (accessed 12 Sept 2019).
- Sipes, N.S., Martin, M.T., Kothiya, P., Reif, D.M., Judson, R.S., Richard, A.M., Houck, K.A., Dix, D.J., Kavlock, R.J., and Knudsen, T.B. (2013) Profiling 976 ToxCast chemicals across 331 enzymatic and receptor signaling assays. *Chem. Res. Toxicol.*, **26**, 878–95.
- Thomas, R.S., Bahadori, T., Buckley, T.J., Cowden, J., *et al.* (2019) The Next Generation Blueprint of Computational Toxicology at the U.S. Environmental Protection Agency. *Toxicol. Sci.*, **169**, 317–332.
- U.S. EPA (2018a) ToxCast & Tox21 summary files for invitrodb_v2. *ToxCast Tox21 Summ. Files from Invit.* https://epa.figshare.com/articles/Previously_Published_ToxCast_Data/6062551/2

(accessed 12 Sept 2019).

U.S. EPA (2018b) ToxCast Owner's Manual - Guidance for Exploring Data.

Varga, T., Czimmerer, Z., and Nagy, L. (2011) PPARs are a unique set of fatty acid regulated transcription factors controlling both lipid metabolism and inflammation. *Biochim. Biophys. Acta - Mol. Basis Dis.*, **1812**, 1007–1022.

Villeneuve, D.L., Coady, K., Escher, B.I., Mihaich, E., Murphy, C.A., Schlekat, T., and Garcia-Reyero, N. (2019) High-throughput screening and environmental risk assessment: State of the science and emerging applications. *Environ. Toxicol. Chem.*, **38**, 12–26.

Wang, Y., Backman, T.W.H., Horan, K., and Girke, T. (2013) fmcsR: mismatch tolerant maximum common substructure searching in R. *Bioinformatics*, **29**, 2792–2794.

Watt, E.D. and Judson, R.S. (2018) Uncertainty quantification in ToxCast high throughput screening. *PLoS One*, **13**.

Wetmore, B.A., Wambaugh, J.F., Allen, B., Ferguson, S.S., Sochaski, M.A., Setzer, R.W., Houck, K.A., Strobe, C.L., Cantwell, K., Judson, R.S., LeCluyse, E., Clewell, H.J., Thomas, R.S., and Andersen, M.E. (2015) Incorporating High-Throughput Exposure Predictions With Dosimetry-Adjusted *In Vitro* Bioactivity to Inform Chemical Toxicity Testing. *Toxicol. Sci.*, **148**, 121–136.

Williamson, G., Kay, C.D., and Crozier, A. (2018) The Bioavailability, Transport, and Bioactivity of Dietary Flavonoids: A Review from a Historical Perspective. *Compr. Rev. Food Sci. Food Saf.*, **17**, 1054–1112.

Yang, C., Tarkhov, A., Maruszyk, J., Bienfait, B., Gasteiger, J., Kleinoeder, T., Magdziarz, T., Sacher, O., Schwab, C.H., Schwoebel, J., Terfloth, L., Arvidson, K., Richard, A., Worth, A., and Rathman, J. (2015) New Publicly Available Chemical Query Language, CSRML, To Support Chemotype Representations for Application to Data Mining and Modeling. *J. Chem. Inf. Model.*, **55**, 510–528.

Figure 1. Circle pack plot of the defined functional groups with the colouring highlighting the different defined chemical groups. The larger a circle, the more chemicals fall into the group and closely related chemicals are packed more closely together. Tertiary groups (closest related chemicals) are labelled and coloured according to the primary chemical group to which they belong. Details about the composition of the groups can be found in Table 1.

Figure 2. Heatmaps showing coverage of biological activity for the 102 tertiary chemical groups within the different ToxCast target families. The targets are displayed on the y-axes with ticks, one per target. The range from orange to red corresponds to an increasing percentage of chemicals within the chemical group that was active in the different assays of that target. White spots mean that less than 3 chemicals were tested in all assays of that target. Grey spots mean that none of the chemicals in the chemical group was active in the assays of that target.

Figure 3. Heatmaps of the biological activity of the 102 tertiary chemical groups within "DNA binding" and "nuclear receptor" target families. Each target (displayed on the y-axes with labels) is covered by 1 to 11 assays. The range from orange to red corresponds to an increasing percentage of chemicals within the chemical group that showed activity in the different assays of that target. White spots mean that less than 3 chemicals were tested in all assays of that target. Grey spots mean that none of the chemicals in the chemical group was active in the assays of that target. The results for all target families can be interactively viewed through [www. https://ilsi.eu/exploitation-of-toxcast-data-on-food-chemicals-for-safety-risk-assessment/](https://ilsi.eu/exploitation-of-toxcast-data-on-food-chemicals-for-safety-risk-assessment/).

Figure 4. Biological activity of the tertiary chemical groups within the "Nuclear receptor" (A, B, C) and "DNA binding" (D) target families. For each tertiary chemical group, the percent of chemicals that were active in the assays for different target families were

calculated and displayed in the colours indicated. The results for all target families can be interactively viewed through [www. https://ilsi.eu/exploitation-of-toxcast-data-on-food-chemicals-for-safety-risk-assessment/](https://ilsi.eu/exploitation-of-toxcast-data-on-food-chemicals-for-safety-risk-assessment/).

Table 1. Defined chemical groups

Primary groups^a	Secondary groups^a	Tertiary groups ^a (including the mean Tanimoto coefficient) ^b
Alcohol	Alkenyl; Alkyl; Hydroxybenzene; Phenylalkanol	1. Alkenyl, primary (0.44); 2. Alkenyl, secondary (0.42); 3. Alkenyl, tertiary (0.7); 4. Alkyl, diol (0.42); 5. Alkyl, primary, branched-chain (0.58); 6. Alkyl, primary, straight chain (0.63); 7. Alkyl, secondary, cyclic (0.68); 8. Sugar alcohol (0.59); 9. Alkoxy phenol ether, substituted (0.57); 11. Hydroxy benzyl ketones (0.71); 12. Phenol, aliphatic substituted(0.67); 13. Salicyclic acid and derivatives (0.68); 14. Phenalkyl/alkenyl (0.60)
Aldehyde	Alkenyl; Alkyl; Aromatic	15. Alkenyl, acyclic (0.51); 16. Alkyl, branched-chain (0.50); 17. Alkyl, straight chain (0.76); 18. Benzaldehyde derivatives (0.64); 19. Phenylalkenyl (0.67); 20. Phenylalkyl (0.66)
Carboxylic acid	Alkenyl; Alkyl; Amino acids and derivatives; Aryl; Hydroxy acid; Keto acid; Polycarboxylic acid	21. Alkenyl, branched-chain (0.48); 22. Alkenyl, straight chain (0.37); 23. Alkyl, branched-chain (0.72); 24. Alkyl, straight chain (0.59); 25. Amino acids and derivatives (0.26); 26. Benzoic acid (0.90); 27. Phenylaliphatic carboxylic acid (0.41); 28. Lactic acids; 29. Keto acid (0.53); 30. Polycarboxylic acid, alkyl and alkenyl (0.52)
Dyes	Azo; Triarylmethane	31. Azo (0.38); 32. Triarylmethane (0.76)
Ester	Aliphatic alcohol diester/triester; Alkenyl alcohol; Alkyl alcohol; Aromatic acid ester; Aromatic alcohol; Lactone	33. Aliphatic alcohol diester/triester (0.37); 34. (3Z)-Hex-3-en-1-yl alcohol; 35. Allyl alcohol; 36. Citronellol; 37. Geraniol; 38. Linalool; 39. Branched-chain alcohol, aliphatic (0.54); 40. Branched-chain alcohol, aryl (0.69); 41. Butanol (0.51); 42. Ethanol, aliphatic (0.58); 43. Ethanol, aryl (0.55); 44. Hexanol (0.76); 45. Isobutanol (0.79); 46. Methanol, aliphatic (0.43); 47. Methanol, aryl (0.79); 48. Pentanol (0.79); 49. Propanol (0.67); 50. Straight chain (7 +) alcohol, aliphatic (0.80); 51. Straight chain (7 +) alcohol, aryl (0.80); 52. 2-Aminobenzoate (0.62); 53. Benzoate (0.69); 54. Cinnamate (0.57); 55. Parabengallate (0.72); 56. Phenylacetate (0.73); 57. Salicylate (0.67); 58. 3-Phenylpropen-2-enyl alcohol (0.76); 59. Anisyl (0.90); 60. Benzyl alcohol, aliphatic (0.85); 61. Benzyl alcohol, aryl (0.83); 62. Phenylethyl alcohol, aliphatic (0.77); 63. Phenylethyl alcohol, aryl (0.77); 64. Ascorbic acid and derivatives (0.41); 65. Lactone, five-membered (0.66); 66. Lactone, six-membered (0.64)
Ether	Alkenyl; Alkyl; Aromatic	67. Alkenyl, acyclic (0.59); 68. Alkyl, cyclic (0.39); 69. Aryl methoxy (0.67); 70. Aryl methoxy, aliphatic substituted(0.65)

Heterocycles and polycycles	Hydrocarbon polycycles; Nitrogen heterocycles; Oxygen heterocycles; Sulfur-nitrogen heterocycles	71. Bicycloheptanes and derivatives (0.67); 72. Biphenyl derivatives (0.90); 73. Naphthalene derivatives (0.50); 74. Pyrazine derivatives (0.57); 75. Pyridine derivatives (0.48); 76. Quinoline derivatives (0.56); 77. Benzodiazole (0.48); 78. Furan derivatives (0.49); 79. Pyranone (0.45); 80. Thiazole and thiazoline (0.28)
Hydrocarbon	Terpene	81. Terpene (0.57)
Inorganic	Inorganic	82. Inorganic (0.08)
Ketone	Alkenyl; Alkyl; Aryl; Jasmone derivatives	83. Alkenyl, acyclic (0.55); 84. Cyclohexenyl (0.58); 85. Ionone/irone (0.62); 86. Alkyl, acyclic (0.61); 87. Alkyl, cyclic (0.61); 88. Benzyl (0.71); 89. Jasmone derivatives (0.42)
Metallic salts organic	Metallic salts organic	90. Metallic salts organic (0.21)
Organosulfur	Alkyl thioether; Disulfide; Thiol	91. Aliphatic thioether (0.44); 92. Disulfide (0.37); 93. Thiol (0.33)
Structure undefined	Structure undefined	94. Structure undefined (NA)
Sugars and derivatives	Sugars and derivatives	95. Sugars and derivatives (0.49)
Terpene and Terpenoid derivatives	Carvone derivatives; Citronellol derivatives; Farnesene derivatives; Geraniol derivatives; Linalool derivatives; Retinol derivatives	96. Carvone derivatives (0.67); 97. Citronellol derivatives (0.70); 98. Farnesene derivatives (0.62); 99. Geraniol derivatives (0.75); 100. Linalool derivatives (0.75); 101. Retinol derivatives (0.69)
Vitamins and derivatives	Vitamins and derivatives	102. Vitamins and derivatives (0.25)

^a Only those chemicals chemical groups that contain at least three chemicals are displayed. The full list of chemicals and their grouping is provided in appendi

^b Mean Tanimoto coefficient, calculated based on the Maximum Common Substructures of the chemicals within a group (see Materials and Methods).

Table 2. Expected important biological targets of different flavonoids and the percentage active in the chemical group containing flavonoids (group 79) towards these targets.

Biological target^a	ToxCast targets (% active in group 79)	Function	ToxCast Target family	ToxCast Target subfamily
NRF2	NFE2L2 (15)	antioxidant	DNA binding	basic leucine zipper
NF- κ B	NFKB1 (0)	free-radical scavenging	DNA binding	NF-kappa B
VEGF	KDR (0), FLT1 (0), FLT4 (0)	regulation of vascular cell development	kinase	receptor tyrosine kinase
PPAR	PPARA (0), PPARD (0), PPARG (0), PPARA PPARD PPARG (0), PPARG SRC (0)	lipid metabolism and glucose homeostasis	nuclear receptor	non-steroidal
VCAM-1	VCMA1 (0)	vascular cell adhesion	cell adhesion molecules	immunoglobulin CAM
ER	ESR1 (14), ESR2 (29), ESR1 ESR2 (40)	estrogen-dependent proliferation and differentiation	nuclear receptor	steroidal

^aBeekmann et al. 2012; Williamson et al. 2018; Rietjens et al. 2017

Table 3. Expected important biological targets of different fatty acids within lipid homeostasis and the percentage of actives in the chemical group containing unsaturated fatty acids (groups 22) and saturated fatty acids (group 24) towards these targets.

Biological target^a	ToxCast target (% active in groups 22 and 24)	Function	ToxCast Target family	ToxCast Target subfamily
PPARs	PPARA (11)(10), PPARD (0)(0), PPARG (8)(4), PPARA PPARD PPARG (13)(5) PPARG SRC (13)(0)	lipid metabolism and homeostasis, glucose utilization	nuclear receptor	non-steroidal
SREBPs	SREBF1 (0)(6)	lipid metabolism and homeostasis	DNA binding	basic helix-loop-helix leucine zipper
LXR	NR1H2 (0)(0), NR1H3 (0)(0), NR1H2 NR1H3 (0)(0) SRC NR1H4)(10),	lipid metabolism and homeostasis	nuclear receptor	non-steroidal
HNF4	HNF4A (0)(0)	lipid metabolism and homeostasis	nuclear receptor	orphan

^aMüller and Kersten, 2003

Table 4. List of chemical groups that displayed a relatively high activity in Figure 3 and 4 and the targets that are affected.

Chemical group	Target family and key targets ^a
Ester-Aliphatic alcohol diester/triester (group 33, n =11 of which 1 chemical is part of the regulatory restricted group)	Non-steroidal nuclear receptor: NR1I2 (33), PPARG (22), NR1H4 (20)
Metallic salts organic (group 87, n =7 of which 1 chemical is part of the regulatory restricted group)	Non-steroidal nuclear receptor: NR1I2 (36), RXRA (33), PPARG (30)
Retinol derivatives (101, n=3)	Non-steroidal nuclear receptor: NR1I2 (45), RXRA (40), RXRB (30), VDR (25), NR1H2 NR1H3 (20)
Organosulfur.Thiol (group 93, n = 6)	DNA binding: TCF7 TCF7L2 LEF1 TCF7L1 (60) IRF1 (40), FOS JUN, (33), SMAD1 (33), USF1 (33), NFKB1 (33), NFE2L2 (33), POU2F1 (33), TP53 (33), AHR (28), SREBF1 (20), HSF1 (20), XBP1ng (20)

^a>Only the targets with more than 20% activity are displayed




Figure 1. Circle pack plot of the defined functional groups with the colouring highlighting the different defined chemical groups. The larger a circle, the more chemicals fall into the group and closely related chemicals are packed more closely together. Tertiary groups (closest related chemicals) are labelled and coloured according to the primary chemical group to which they belong. Details about the composition of the groups can be found in Table 1.

269x209mm (150 x 150 DPI)




Figure 2. Heatmaps showing coverage of biological activity for the 102 tertiary chemical groups within the different ToxCast target families. The targets are displayed on the y-axis with ticks, one per target. The range from orange to red corresponds to an increasing percentage of chemicals within the chemical group that was active in the different assays of that target. White spots mean that less than 3 chemicals were tested in all assays of that target. Grey spots mean that none of the chemicals in the chemical group was active in the assays of that target.

350x249mm (150 x 150 DPI)




Figure 3. Heatmaps of the biological activity of the 102 tertiary chemical groups within “DNA binding” and “nuclear receptor” target families. Each target (displayed on the y-axes with labels) is covered by 1 to 11 assays. The range from orange to red corresponds to an increasing percentage of chemicals within the chemical group that showed activity in the different assays of that target. White spots mean that less than 3 chemicals were tested in all assays of that target. Grey spots mean that none of the chemicals in the chemical group was active in the assays of that target.

499x299mm (150 x 150 DPI)




Figure 4. Biological activity of the tertiary chemical groups within the “Nuclear receptor” (A, B, C) and “DNA binding” (D) target families. For each tertiary chemical group, the percent of chemicals that were active in the assays for different target families were calculated and displayed in the colours indicated. The results for all target families can be interactively viewed through [www. https://ils.eu/exploitation-of-toxcast-data-on-food-chemicals-for-safety-risk-assessment/](https://ils.eu/exploitation-of-toxcast-data-on-food-chemicals-for-safety-risk-assessment/).

402x399mm (150 x 150 DPI)