



## LJMU Research Online

Ho, AYQ, Perley, DA, Kulkarni, SR, Dong, DZJ, De, K, Chandra, P, Andreoni, I, Bellm, EC, Burdge, KB, Coughlin, M, Dekany, R, Feeney, M, Frederiks, DD, Fremling, C, Golkhou, VZ, Graham, MJ, Hale, D, Helou, G, Horesh, A, Kasliwal, MM, Laher, RR, Masci, FJ, Miller, AA, Porter, M, Ridnaia, A, Ben, R, Shupe, DL, Soumagnac, MT and Svinkin, DS

**The Koala: A Fast Blue Optical Transient with Luminous Radio Emission from a Starburst Dwarf Galaxy at  $z=0.27$**

<http://researchonline.ljmu.ac.uk/id/eprint/13218/>

### Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Ho, AYQ, Perley, DA, Kulkarni, SR, Dong, DZJ, De, K, Chandra, P, Andreoni, I, Bellm, EC, Burdge, KB, Coughlin, M, Dekany, R, Feeney, M, Frederiks, DD, Fremling, C, Golkhou, VZ, Graham, MJ, Hale, D, Helou, G, Horesh, A, Kasliwal, MM, Laher, RR, Masci, FJ, Miller, AA, Porter, M, Ridnaia, A, Ben, R.**

LJMU has developed [LJMU Research Online](http://researchonline.ljmu.ac.uk/) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

<http://researchonline.ljmu.ac.uk/>

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)

<http://researchonline.ljmu.ac.uk/>

## The Koala: A Fast Blue Optical Transient with Luminous Radio Emission from a Starburst Dwarf Galaxy at $z = 0.27$

ANNA Y. Q. HO,<sup>1</sup> DANIEL A. PERLEY,<sup>2</sup> S. R. KULKARNI,<sup>1</sup> DILLON Z. J. DONG,<sup>1</sup> KISHALAY DE,<sup>1</sup> POONAM CHANDRA,<sup>3</sup> IGOR ANDREONI,<sup>1</sup> ERIC C. BELLM,<sup>4</sup> KEVIN B. BURDGE,<sup>1</sup> MICHAEL COUGHLIN,<sup>5</sup> RICHARD DEKANY,<sup>6</sup> MICHAEL FEENEY,<sup>6</sup> DMITRY D. FREDERIKS,<sup>7</sup> CHRISTOFFER FREMLING,<sup>1</sup> V. ZACH GOLKHOV,<sup>4,8</sup> MATTHEW J. GRAHAM,<sup>1</sup> DAVID HALE,<sup>6</sup> GEORGE HELOU,<sup>9</sup> ASSAF HORESH,<sup>10</sup> MANSI M. KASLIWAL,<sup>1</sup> RUSS R. LAHER,<sup>9</sup> FRANK J. MASCI,<sup>9</sup> A. A. MILLER,<sup>11,12</sup> MICHAEL PORTER,<sup>6</sup> ANNA RIDNAIA,<sup>7</sup> BEN RUSHOLME,<sup>9</sup> DAVID L. SHUPE,<sup>9</sup> MAAYANE T. SOUMAGNAC,<sup>13,14</sup> AND DMITRY S. SVINKIN<sup>7</sup>

<sup>1</sup>*Cahill Center for Astrophysics, California Institute of Technology, MC 249-17, 1200 E California Boulevard, Pasadena, CA, 91125, USA*

<sup>2</sup>*Astrophysics Research Institute, Liverpool John Moores University, IC2, Liverpool Science Park, 146 Brownlow Hill, Liverpool L3 5RF, UK*

<sup>3</sup>*National Centre for Radio Astrophysics, Tata Institute of Fundamental Research, PO Box 3, Pune, 411007, India*

<sup>4</sup>*DIRAC Institute, Department of Astronomy, University of Washington, 3910 15th Avenue NE, Seattle, WA 98195, USA*

<sup>5</sup>*School of Physics and Astronomy, University of Minnesota, Minneapolis, Minnesota 55455, USA*

<sup>6</sup>*Caltech Optical Observatories, California Institute of Technology, Pasadena, CA 91125*

<sup>7</sup>*Ioffe Institute, Politekhnicheskaya 26, St. Petersburg 194021, Russia*

<sup>8</sup>*The eScience Institute, University of Washington, Seattle, WA 98195, USA\**

<sup>9</sup>*IPAC, California Institute of Technology, 1200 E. California Blvd, Pasadena, CA 91125, USA*

<sup>10</sup>*Racah Institute of Physics, The Hebrew University of Jerusalem, Jerusalem 91904, Israel*

<sup>11</sup>*Center for Interdisciplinary Exploration and Research in Astrophysics and Department of Physics and Astronomy, Northwestern University, 1800 Sherman Ave, Evanston, IL 60201, USA*

<sup>12</sup>*The Adler Planetarium, Chicago, IL 60605, USA*

<sup>13</sup>*Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, CA 94720, USA*

<sup>14</sup>*Department of Particle Physics and Astrophysics, Weizmann Institute of Science, Rehovot 76100, Israel*

(Received 2 March 2020; Revised 13 April 2020)

Submitted to The Astrophysical Journal

### Abstract

We present ZTF18abvkwla (the “Koala”), a fast blue optical transient discovered in the Zwicky Transient Facility (ZTF) One-Day Cadence (1DC) Survey. ZTF18abvkwla has a number of features in common with the groundbreaking transient AT2018cow: blue colors at peak ( $g - r \approx -0.5$  mag), a short rise time from half-max of under two days, a decay time to half-max of only three days, a high optical luminosity ( $M_{g,\text{peak}} \approx -20.6$  mag), a hot ( $\gtrsim 40,000$  K) featureless spectrum at peak light, and a luminous radio counterpart. At late times ( $\Delta t > 80$  d) the radio luminosity of ZTF18abvkwla

( $\nu L_\nu \gtrsim 10^{40} \text{ erg s}^{-1}$  at 10 GHz, observer-frame) is most similar to that of long-duration gamma-ray bursts (GRBs). The host galaxy is a dwarf starburst galaxy ( $M \approx 5 \times 10^8 M_\odot$ ,  $\text{SFR} \approx 7 M_\odot \text{ yr}^{-1}$ ) that is moderately metal-enriched ( $\log [\text{O}/\text{H}] \approx 8.5$ ), similar to the hosts of GRBs and superluminous supernovae. As in AT2018cow, the radio and optical emission in ZTF18abvkwla likely arise from two separate components: the radio from fast-moving ejecta ( $\Gamma\beta c > 0.38c$ ) and the optical from shock-interaction with confined dense material ( $< 0.07 M_\odot$  in  $\sim 10^{15} \text{ cm}$ ). Compiling transients in the literature with  $t_{\text{rise}} < 5 \text{ d}$  and  $M_{\text{peak}} < -20 \text{ mag}$ , we find that a significant number are engine-powered, and suggest that the high peak optical luminosity is directly related to the presence of this engine. From 18 months of the 1DC survey, we find that transients in this rise-luminosity phase space are at least two to three orders of magnitude less common than CCSNe. Finally, we discuss strategies for identifying such events with future facilities like the Large Synoptic Survey Telescope, and prospects for detecting accompanying X-ray and radio emission.

## 1. INTRODUCTION

Historically, the cadence of optical time-domain surveys was tuned to detecting Type Ia supernovae (SNe), whose optical light curves rise from first light to peak in 15–20 days (Miller et al. 2020). Recognizing that this observing strategy resulted in “gaps” in timescale-luminosity phase-space, surveys such as the Palomar Transient Factory (Law et al. 2009; Rau et al. 2009) and the Pan-STARRS1 Medium Deep Survey (Drout et al. 2014) sought to systematically chart the landscape of short-timescale ( $< 10 \text{ day}$ ) phenomena. These efforts delineated populations of fast transients spanning many orders of magnitude in peak luminosity, from faint calcium-rich transients (Kasliwal et al. 2012) to luminous relativistic explosions (Cenko et al. 2013).

A population of particular recent interest is “fast evolving luminous transients” (Rest et al. 2018) or “fast blue optical transients” (Margutti et al. 2019). A consistent definition of this “class” does not yet exist; these terms typically refer to transients with rise times and peak luminosities too fast and too luminous, respectively, to be explained by the radioactive decay

of  $^{56}\text{Ni}$ . Although they likely arise from a variety of progenitors, fast-luminous transients are primarily found in star-forming galaxies (Drout et al. 2014; Pursiainen et al. 2018) and therefore are thought to represent a variety of poorly understood endpoints of massive-star evolution. As summarized in Kasen (2017), fast and luminous light curves may be powered by shock breakout or shock-cooling emission from material that is closely confined to the progenitor star at the time of explosion, or alternatively by a “central engine:” accretion onto a black hole, or the rotational spindown of a magnetar.

Most fast-luminous optical transients have been found in archival searches of optical-survey data, including PS1 (Drout et al. 2014), the Dark Energy Survey (Pursiainen et al. 2018), Kepler (Rest et al. 2018), and the Supernova Legacy Survey (Arcavi et al. 2016). A handful have been discovered while the transient was still active, enabling prompt follow-up observations. For example, spectroscopic monitoring of the fast-luminous transients iPTF16asu and ZTF18abukavn (SN 2018gcp) revealed that as the optical emission faded, the spectrum developed features typical of broad-lined Ic SNe (Whitesides et al. 2017; Wang et al. 2019; Ho et al. 2019b).

\* Moore-Sloan, WRF Innovation in Data Science, and DIRAC Fellow

The discovery of the fast-luminous transient AT2018cow (Prentice et al. 2018) generated considerable excitement because of its proximity ( $z = 0.0141$ ) and therefore the opportunity for detailed observations. AT2018cow had several remarkable features: (1) near-relativistic ejecta velocities at early times, from optical spectroscopy (Perley et al. 2019a); (2) luminous and fast-varying X-ray emission suggesting an exposed central engine (Rivera Sandoval et al. 2018; Ho et al. 2019a; Margutti et al. 2019); (3) high-velocity emission lines of hydrogen and helium emerging at late times (Perley et al. 2019a); (4) no second peak that would indicate a significant role for radioactive ejecta in powering the light curve (Perley et al. 2019a); and (5) luminous submillimeter emission indicating a large explosion energy injected into a shell of very dense material (Ho et al. 2019a; Huang et al. 2019). Despite extensive observations across the electromagnetic spectrum, the progenitor of AT2018cow is unknown. One suggestion is a massive-star explosion that resulted in the formation of an accreting black hole or magnetar, which drove a mildly relativistic jet or wind (Perley et al. 2019a; Margutti et al. 2019; Ho et al. 2019a). Other suggestions include an electron-capture SN (Lyutikov, & Toonen 2019) and a tidal disruption event (TDE; Vinkó et al. 2015; Perley et al. 2019a; Kuin et al. 2019). If AT2018cow was a massive-star explosion, the dense confined CSM points to eruptive mass-loss shortly before core-collapse (Ho et al. 2019a), and indeed Fox & Smith (2019) pointed out the similarity between AT2018cow and interaction-powered Type Ibn SNe.

Here we report the discovery in Zwicky Transient Facility (ZTF) data of ZTF18abvkwla<sup>1</sup>, a fast-rising luminous optical transient at  $z =$

0.27.<sup>2</sup> In §2 we present the key observational features of ZTF18abvkwla—a rest-frame  $g$ -band light curve similar to that of AT2018cow, a luminous radio counterpart similar to gamma-ray burst (GRB) afterglows, and a starburst dwarf host galaxy. In §3 we compare ZTF18abvkwla to transients in the literature that have  $t_{\text{rise}} < 5$  d and  $M < -20$  mag, where  $t_{\text{rise}}$  is defined from 0.75 mag below peak to peak (half-max to max in flux space). We use a cut of  $M < -20$  mag to exclude “normal” Type Ibn SNe (Hosseinzadeh et al. 2017) and we exclude the hundreds of optical afterglows discovered in GRB follow-up observations (Kann et al. 2010). The comparison sample is shown in Table 1 and Figure 1. Note that the Table 1 transients have thermal spectra at peak, unlike GRB afterglows which arise from synchrotron radiation.

In §4.1 we model the optical emission from ZTF18abvkwla as thermal emission from shock breakout in dense confined material, and in §4.2 we use the radio emission to estimate properties of the forward shock (velocity, shock energy) and the ambient medium. In §4.3, we discuss possible progenitor systems. Finally, in §5 we use 18 months of survey observations to estimate the rate of transients in the phase-space of Figure 1, and find that the rate is 2–3 times smaller than the CC SN rate.

Throughout this paper, we use a standard  $\Lambda$ CDM cosmology (Planck Collaboration et al. 2016) and times are reported in UT. Optical magnitudes are reported in the AB system (Oke, & Gunn 1983), and corrected for foreground Galactic extinction using reddening measurements in Schlafly & Finkbeiner (2011) and the extinction law from Fitzpatrick (1999).

<sup>1</sup> nicknamed “Koala” on account of the last four letters of its ZTF ID

<sup>2</sup> After the submission of our paper, Coppejans et al. (2020) published radio and X-ray observations of CSS161010, another transient in a dwarf galaxy with properties similar to AT2018cow.

**Table 1.** Transients in the literature with  $t_{\text{rise}} < 5$  d and  $M < -20$  mag. Timescales are presented in rest-frame and measured using the light curve that most closely matches rest-frame  $g$ . Luminosity is corrected for Galactic extinction, assuming zero host-galaxy extinction in all cases except for iPTF15ul and SN 2011kl. SN 2011kl was associated with GRB 111209A, and the afterglow emission has been subtracted.

Name	Redshift	$M_{g,\text{max}}$	$t_{\text{rise}}$ days	$t_{\text{fade}}$ days	Ref
Dougie	0.19	$-23.03 \pm 0.13$	$3.92 \pm 0.14$	$9.69 \pm 1.19$	[1]
SN 2011kl	0.677	$-20.31 \pm 0.13$	$4.97 \pm 1.20$	$17.70 \pm 5.82$	[2,3]
SNLS04D4ec	0.593	$-20.26 \pm 0.03$	$< 3.81$	$8.60 \pm 0.43$	[4]
SNLS05D2bk	0.699	$-20.39 \pm 0.02$	$2.90 \pm 0.06$	$12.75 \pm 0.78$	[4]
SNLS06D1hc	0.555	$-20.28 \pm 0.03$	$4.59 \pm 0.06$	$12.35 \pm 0.45$	[4]
iPTF15ul	0.066	$-21.2 \pm 0.3$	$1.53 \pm 0.05$	$3.72 \pm 0.08$	[5]
DES16X1eho	0.76	$-20.39 \pm 0.09$	1.28–2.53	$1.01 \pm 0.27$	[6]
iPTF16asu	0.187	$-20.3 \pm 0.1$	$1.14 \pm 0.13$	$10.62 \pm 0.55$	[7]
AT2018cow	0.0141	$-20.89 \pm 0.06$	$1.43 \pm 0.08$	$1.95 \pm 0.06$	[8,9]

**References**— [1] Vinkó et al. (2015), [2] Greiner et al. (2015), [3] Kann et al. (2019), [4] Arcavi et al. (2016), [5] Hosseinzadeh et al. (2017), [6] Pursiainen et al. (2018) [7] Whitesides et al. (2017), [8] Prentice et al. (2018) [9] Perley et al. (2019a)

## 2. DISCOVERY AND BASIC ANALYSIS

### 2.1. Optical

#### 2.1.1. Photometry

Since April 2018, ZTF (Bellm et al. 2019a; Graham et al. 2019) has been conducting a wide-area (2000–3000 deg<sup>2</sup>) one-day cadence (1DC) survey in  $g$  and  $r$  (Bellm et al. 2019). The sky coverage of the 1DC survey is shown in Figure 2 and a histogram of the typical time between exposures is shown in Figure 3.


The IPAC ZTF pipeline (Masci et al. 2019) uses the method described in Zackay et al. (2016) to generate difference images using a coadded reference image. Every  $5\sigma$  point-source detection is assigned a score based on a machine learning real-bogus metric (Mahabal et al. 2019; Duev et al. 2019), and is cross-matched against external catalogs to search for resolved and extended counterparts (Tachibana & Miller 2018). Alerts are distributed in Avro format (Patterson et al. 2019) and are filtered by the ZTF collaboration using a web-based sys-

tem called the GROWTH Marshal (Kasliwal et al. 2019).

ZTF18abvkwla was discovered in an image obtained on 12 Sept 2018. The alert passed a filter designed to look for rapidly-evolving transients, and as a result we obtained a follow-up spectrum 24 hours later (§2.1.2). The discovery magnitude was  $g = 19.73 \pm 0.16$  mag and the last non-detection was one day prior, with a limiting magnitude  $g > 20.74$ .

The source position was measured to be  $\alpha = 02^{\text{h}}00^{\text{m}}15.19^{\text{s}}$ ,  $\delta = +16^{\text{d}}47^{\text{m}}57.3^{\text{s}}$  (J2000), which is  $0'.28 \pm 0'.13$  from the nucleus of a blue ( $g - r = 0.32$  mag) extended source that has a photometric redshift of 0.11 (68 percentile confidence interval 0.08–0.29) in the eighth data release of LegacySurvey (DR8; Dey et al. 2019). At  $z = 0.2714$  (§2.1.2) this offset corresponds to  $1.9 \pm 0.9$  kpc. The host is approximately  $2''$  (14 kpc) across.

The light curve (Figure 4; Table 2) has a similar timescale and peak luminosity to that of AT2018cow. In rest-frame  $g$ -band, the rise time




**Figure 1.** Phase-space of luminosity and rise time considered in this paper; see Table 1 for data sources. We do not show the transient Dougie (Vinkó et al. 2015), which had a peak absolute magnitude of  $-23$ . Note that the peak mag of iPTF15ul includes a large host-galaxy extinction correction, whereas the other sources have zero host extinction correction. Also note that SN 2011kl was associated with an ultra-long duration GRB 111209A (Kann et al. 2018), and the light-curve properties shown here reflect the afterglow-subtracted light curve (Kann et al. 2019).

is  $1.83 \pm 0.05$  d, the fade time is  $3.12 \pm 0.22$  d, and the peak magnitude is  $-20.59 \pm 0.07$  mag.


We estimate that the onset of the optical emission was around the time of the last non-detection ( $t_0 = 2458372.9206$  JD) and use this as a reference epoch for the remainder of the paper.

### 2.1.2. Spectroscopy and Host Galaxy Properties

One day after discovery, we obtained a spectrum of ZTF18abvkwla using the Double Beam Spectrograph (DBSP; Oke & Gunn 1982) on the 200-inch Hale telescope at Palomar Observatory. We used the D55 dichroic, a slit width of 1.5 arcseconds, the 600/4000 blue grating, and the 316/7500 red grating. The spectrum was reduced using a PyRAF-based pipeline (Bellm & Sesar 2016). As shown in Figure 5, the spectrum




**Figure 2.** Number of epochs obtained by the ZTF one-day cadence survey from 3 April 2018 to 18 October 2019



**Figure 3.** Histogram of times between successive observations of a field in the same filter for the ZTF one-day cadence survey. Intervals greater than 10 days are not shown.

shows a hot blue continuum with no broad features in emission or absorption. Superimposed



**Figure 4.** Light curve of ZTF18abvkwla in P48  $g$  (filled green squares) and  $r$  (open orange circles) with a comparison to AT2018cow at similar rest wavelengths, both corrected for Galactic extinction. The ‘S’ at the top of the inset indicates the epoch of our DBSP spectrum. Dashed lines show  $^{56}\text{Ni}$ -powered light curves for two different nickel masses.

on the spectrum are a variety of narrow emission lines typical of a star-forming galaxy ( $\text{H}\alpha$ ,  $\text{H}\beta$ , O III, S II, O II) at a redshift of  $z = 0.2714$  plus the Mg II UV doublet in absorption at consistent redshift.


A blackbody fit to the continuum (after subtracting a host-galaxy continuum model, discussed later in this section) indicates an effective temperature  $T \gtrsim 40,000\text{K}$ , although we caution that it could be significantly higher as the bulk of the energy was clearly emitted in the UV ( $< 2750\text{\AA}$  in the rest frame) and we have no firm constraint on the host-galaxy extinction. Together with the peak absolute magnitude of the  $g$ -band light curve, we derive a bolometric luminosity of  $L_{\text{bol}} > \nu L_{\nu} \sim 10^{44}\text{ erg s}^{-1}$ . Assuming  $T = 40,000\text{K}$ , the photospheric radius is  $R > 2 \times 10^{14}\text{ cm}$ . Since the peak is 2 d after first light, assuming  $R(t = t_0) = 0$  gives  $V > 0.04c$ .

On 4 Jan 2019 (+115 d), we obtained a spectrum of the host galaxy of ZTF18abvkwla using the Low Resolution Imaging Spectrometer (Oke et al. 1995; McCarthy et al. 1998) on the Keck I 10-m telescope, with the 400/3400 grism in the

blue camera and the 400/8500 grating in the red camera. Exposure times were 940 and 900 seconds for the blue and red camera respectively. The spectrum was reduced and extracted using `Lpipe` (Perley 2019). The absolute calibration was established independently for each camera (red vs. blue) by calculating synthetic photometry of the output spectra in the blue and red cameras in the  $g$  and  $r$  bands, respectively, and rescaling to match the  $g$  and  $r$  photometry from SDSS DR14 (Abolfathi et al. 2018). The SDSS magnitudes (AB, converted to Pogson) are  $u = 21.74 \pm 0.20$ ,  $g = 21.20 \pm 0.04$ ,  $r = 20.81 \pm 0.05$ ,  $i = 20.92 \pm 0.09$ , and  $z = 20.52 \pm 0.20$ .

The host-galaxy spectrum (Figure 6) consists of a weak continuum and a series of very strong emission lines. Line fluxes were extracted using an identical procedure as in Perley et al. (2016). We first fit a model to the spectral energy distribution (SED). We used a custom IDL routine based on the templates of Bruzual & Charlot (2003) to fit the SDSS  $ugriz$  photometry, including the contribution of nebular lines. As only SDSS  $ugriz$  photometry is available to fit





**Figure 5.** The spectrum of ZTF18abvkwla at the peak of the  $g$ -band optical light curve (black), which was 1 day after the first detection. The source is extremely hot and blue with no spectral features except those associated with the host galaxy. Overplotted in pink is a rescaled late-time spectrum of the host galaxy with a 40,000 K blackbody added.

the host-galaxy SED it is difficult to constrain the nature of the stellar population of the host galaxy in detail, and we were only able to fit the simplest possible model (a continuous star-formation history). However, the stellar mass is unambiguously low ( $\sim 5 \times 10^8 M_{\odot}$ , comparable to the SMC),

This model was then used to produce a synthetic galaxy continuum spectrum, which was subtracted from the observed one (this correction is significant only for higher-order Balmer lines, which overlay strong galaxy absorption features). Emission line fluxes were then measured by fitting a Gaussian function to each emission line (plus a linear baseline to fit any continuum residuals). Lines that were blended or very nearby were fit in groups, and lines whose ratios are fixed from theory were tied together in fitting. A list of all measured line fluxes is given in Table 3.


The SED fitting and the emission-line analysis produce consistent estimates of  $7 M_{\odot} \text{ yr}^{-1}$  for the star-formation rate, and a very high specific star-formation rate of  $\sim 10^{-8} \text{ yr}^{-1}$ . This implies a stellar population dominated by young

stars formed in a recent triggered star-formation burst episode.

We used the host galaxy spectrum (Figure 6) to calculate standard emission-line diagnostics, including metallicity estimates on a variety of scales using the Monte-Carlo code of Bianco et al. (2016). These metallicity measurements are provided in Table 4. The basic properties of the host galaxy are listed in Table 5.

## 2.2. Radio Observations

We obtained four epochs of observations of ZTF18abvkwla using the Karl G. Jansky Very Large Array (VLA; Perley et al. 2011) under the program VLA/18B-242 (PI: D. Perley), listed in Table 6. The first epoch was at  $\Delta t \approx 81$  d at X-band, while the VLA was in C configuration. We used 3C138 as our flux density and band-pass calibrator, and J0204+1514 as our complex gain calibrator. The next three epochs were at  $\Delta t \approx 310$  d,  $\Delta t \approx 350$  d, and  $\Delta t \approx 400$  d, all while the VLA was in A configuration. We continued to use 3C138 but switched to J0238+1636 as our complex gain calibrator. For each observation, we ran the standard VLA calibration pipeline available in the Common Astronomy Software



**Figure 6.** Spectrum of the host galaxy of ZTF18abvkwla. The scale on the bottom half has been zoomed in to show the galaxy continuum and weak emission lines. The feature at 9500 Å is a sky-subtraction residual.

Applications (CASA; McMullin et al. 2007). After calibration, we inspected the data manually for further flagging. We imaged the data using the CLEAN algorithm (Högbom 1974) available in CASA, using a cell size that was 1/5 of the synthesized beamwidth. The field size was set to be the smallest magic number ( $10 \times 2^n$ ) larger than the number of cells needed to cover the primary beam.

In addition, the position of ZTF18abvkwla was serendipitously covered by the VLA Sky Survey (VLASS; Lacy et al. 2019), which has been mapping the entire sky visible to the VLA at low frequencies (2–4 GHz) in three epochs at a cadence of 32 months. The Quicklook images are now available for the first epoch ( $17,000 \text{ deg}^{-2}$ ). We searched the existing Quicklook data using code available on Github<sup>3</sup> that locates the appropriate VLASS tile and subtile for a given RA and Dec and extracts a cutout 12 arcsec on a side. Given a non-detection we estimated an upper limit on the flux density by taking the standard deviation of the pixel val-

ues in this cutout, after performing initial  $3\sigma$  clipping (removing pixels with a value greater than  $3 \times$  the standard deviation). The VLASS observation of ZTF18abvkwla is also listed in Table 6.

We obtained one epoch of observations with the upgraded Giant Metrewave Radio Telescope (GMRT; Gupta et al. 2017; Swarup et al. 1991) under a proposal for Director’s Discretionary Time (Proposal # ddtC086; PI: A. Ho). For our GMRT observations, we used 3C147 and 3C48 as our flux density and bandpass calibrators and 0238+166 for our phase calibrator. We calibrated the GMRT data manually using commands in CASA, with 6 rounds of phase-only self-calibration and 2 rounds of amplitude and phase self-calibration.

The radio light curve from the VLA is shown in Figure 7. At the time of our first observation, the 10 GHz (rest-frame 12 GHz) luminosity was  $10^{30} \text{ erg s}^{-1} \text{ Hz}^{-1}$ , and the in-band spectral index was  $\alpha = -0.16 \pm 0.05$  where  $F_\nu \propto \nu^{-\alpha}$ . At late times ( $\Delta t > 300 \text{ d}$ ) the decline is very steep: at 6 GHz we find  $F_\nu \propto t^{-6.8 \pm 0.9}$ , and at 10 GHz we find  $F_\nu \propto t^{-3.2 \pm 1.4}$ .

<sup>3</sup> [https://github.com/annayqho/Query\\_VLASS](https://github.com/annayqho/Query_VLASS)

**Table 2.** Optical photometry for ZTF18abvkwla from forced photometry on P48 images (Yao et al. 2019). Values have not been corrected for Galactic extinction. Phase  $\Delta t$  is defined from  $t_0$ , the last non-detection.

Date (MJD)	$\Delta t$	Filter	AB Mag
58372.39	-1.02	<i>r</i>	< 21.39
58372.42	-0.99	<i>g</i>	< 21.56
58373.41	0.00	<i>g</i>	19.71 ± 0.05
58373.45	0.04	<i>r</i>	20.18 ± 0.09
58374.39	0.98	<i>r</i>	20.00 ± 0.07
58374.41	1.00	<i>g</i>	19.53 ± 0.05
58375.37	1.96	<i>r</i>	19.92 ± 0.07
58375.37	1.96	<i>r</i>	20.02 ± 0.07
58375.43	2.03	<i>g</i>	19.65 ± 0.04
58375.43	2.03	<i>g</i>	19.72 ± 0.05
58376.42	3.01	<i>r</i>	20.15 ± 0.07
58376.44	3.04	<i>g</i>	19.77 ± 0.05
58377.39	3.98	<i>g</i>	20.10 ± 0.07
58377.43	4.02	<i>r</i>	20.29 ± 0.09
58378.40	4.99	<i>r</i>	20.50 ± 0.10
58378.40	4.99	<i>r</i>	20.62 ± 0.12
58378.45	5.04	<i>g</i>	20.64 ± 0.10
58378.45	5.05	<i>g</i>	20.42 ± 0.09
58379.42	6.02	<i>r</i>	20.85 ± 0.15
58379.44	6.04	<i>g</i>	20.72 ± 0.12
58380.43	7.03	<i>r</i>	21.04 ± 0.15
58382.34	8.93	<i>r</i>	21.06 ± 0.27
58382.34	8.93	<i>r</i>	21.22 ± 0.28
58382.43	9.03	<i>g</i>	21.35 ± 0.19
58382.43	9.03	<i>g</i>	21.56 ± 0.23
58383.48	10.07	<i>g</i>	21.51 ± 0.21


To estimate the contribution to the radio emission from the host galaxy, we use the relation in Greiner et al. (2016), adapted from Murphy et al. (2011):

$$\left(\frac{\text{SFR}_{\text{Radio}}}{M_{\odot} \text{ yr}^{-1}}\right) = 0.059 \left(\frac{F_{\nu}}{\mu\text{Jy}}\right) (1+z)^{-(\alpha+1)} \times \left(\frac{D_L}{\text{Gpc}}\right)^2 \left(\frac{\nu}{\text{GHz}}\right)^{-\alpha}. \quad (1)$$

In the final epoch of our radio observations, assuming  $\alpha = -0.75$  (Condon 1992) where  $F_{\nu} \propto \nu^{\alpha}$ , the 10 GHz flux density of  $0.031 \pm 0.003$  mJy

**Table 3.** Host emission line fluxes and equivalent widths

Species	Wavelength	Flux	Eq. Width
	(Å)	(erg cm <sup>-2</sup> s <sup>-1</sup> )	(Å)
H $\alpha$	6562.82	214.74 ± 2.71	205.9 ± 7.0
H $\beta$	4861.33	57.57 ± 1.07	41.3 ± 1.1
H $\gamma$	4340.47	26.98 ± 1.03	17.6 ± 0.8
H $\delta$	4101.74	13.92 ± 0.91	7.2 ± 0.5
H $\epsilon$	3970.08	11.44 ± 0.86	5.9 ± 0.4
H $\zeta$	3889.06	9.72 ± 0.88	5.0 ± 0.5
[O II]	3727	159.44 ± 1.72	89.0 ± 2.4
[Ne III]	3868.76	16.00 ± 0.94	8.3 ± 0.5
[O III]	4363.21	<3.31	<2.1
[O III]	4958.91	66.35 ± 1.37	47.6 ± 1.4
[O III]	5006.84	196.88 ± 1.60	141.3 ± 3.1
He I	5875.62	6.76 ± 0.72	5.7 ± 0.6
[N II]	6548.06	4.90 ± 0.69	4.7 ± 0.7
[N II]	6583.57	13.91 ± 0.82	13.3 ± 0.9
[S II]	6716.44	27.86 ± 0.95	29.3 ± 1.2
[S II]	6730.82	21.81 ± 0.71	22.9 ± 0.9
O I	6300.30	6.76 ± 0.71	7.2 ± 0.8
[Ar III]	7135.79	5.49 ± 0.53	6.8 ± 0.7



**Figure 7.** The radio light curve of ZTF18abvkwla with the spectral energy distribution at  $\Delta t \approx 350$  d (rest-frame  $\Delta t \approx 275$  d) shown inset. The upper limit at 3 GHz comes from a serendipitous observation by VLASS.

predicts a SFR of  $20 M_{\odot} \text{ yr}^{-1}$ . So, we conclude that during the final observation the radio emission is still dominated by the transient, but the host may contribute a non-trivial fraction of the flux.

**Table 4.** Host galaxy properties (metallicities, mainly) from PyMCZ.

SFR <sup>a</sup>	6.47 ± 1.3
E(B-V)	0.220 <sup>+0.023</sup> <sub>-0.022</sub>
logR23	0.903 <sup>+0.012</sup> <sub>-0.012</sub>
D02	8.253 <sup>+0.130</sup> <sub>-0.128</sub>
Z94	8.450 <sup>+0.016</sup> <sub>-0.010</sub>
M91	8.219 <sup>+0.026</sup> <sub>-0.026</sub>
PP04_N2Ha	8.200 <sup>+0.010</sup> <sub>-0.010</sub>
PP04_O3N2	8.187 <sup>+0.008</sup> <sub>-0.009</sub>
P10_ONS	8.708 <sup>+0.024</sup> <sub>-0.024</sub>
P10_ON	8.172 <sup>+0.046</sup> <sub>-0.047</sub>
M08_N2Ha	8.361 <sup>+0.020</sup> <sub>-0.021</sub>
M08_O3O2	8.521 <sup>+0.011</sup> <sub>-0.011</sub>
M13_O3N2	8.174 <sup>+0.009</sup> <sub>-0.009</sub>
M13_N2	8.194 <sup>+0.041</sup> <sub>-0.042</sub>
KD02_N2O2	7.567 <sup>+0.722</sup> <sub>-0.074</sub>
KK04_N2Ha	8.381 <sup>+0.028</sup> <sub>-0.029</sub>
KK04_R23	8.390 <sup>+0.021</sup> <sub>-0.021</sub>
KD02comb	8.304 <sup>+0.024</sup> <sub>-0.024</sub>

<sup>a</sup>SFR is not from PyMCZ but is calculated directly from the corrected Balmer-line fluxes based on the relation of Kennicutt et al. (1994).

### 3. COMPARISON WITH EXTRAGALACTIC EXPLOSIONS

#### 3.1. Optical Light Curve and Spectrum

As shown in §1, the fast rise time and high peak luminosity of ZTF18abvkwla is shared by only a handful of transients in the literature. In this section we compare the optical properties of ZTF18abvkwla to the transients in Table 1. We exclude Dougie because it resided in an old stellar population with no signs of enhanced star formation (Vinkó et al. 2015); the dominance of absorption features and much lower star-formation rate were confirmed by additional LRIS spectroscopy (Arcavi et al. 2016).

To compare light curves, we selected the light curve in a filter closest to rest-frame  $g$  (the same

**Table 5.** Properties of the host galaxy of ZTF18abvkwla. The stellar mass, star-formation rate, maximum age, and extinction are from a fit to the galaxy SED; the  $\chi^2$  refers to that fit. The metallicity [O/H] was measured using the host galaxy spectrum and is provided on the Z94 scale. This value corresponds to 0.6× solar.

Stellar mass	M	$5.1^{+3.4}_{-2.0} \times 10^8 M_{\odot}$
Star-formation rate	SFR	$6.8^{+3.7}_{-4.6} M_{\odot} \text{ yr}^{-1}$
Maximum age	age	$7.5^{+30}_{-4.5} \times 10^7 \text{ yr}$
Extinction	Av	$0.72^{+0.17}_{-0.54} \text{ mag}$
	$\chi^2/\text{dof}$	1.6 / 2
Metallicity	12+log[O/H]	8.5

The SFR listed here is derived from the photometry, while the SFR in Table 4 was derived from the spectrum. So, there is no expectation of identical values or errors.

**Table 6.** Radio observations of ZTF18abvkwla with the VLA and the GMRT. Upper limit is reported as 3× the image RMS.

$\Delta t$	Facility	Obs. Date	Config.	$\nu$	Flux Density
days		(UT)		(GHz)	(mJy)
81	VLA	2018-12-01	C	10	$0.364 \pm 0.006$
188	VLA <sup>a</sup>	2019-03-19	B	3	< 0.134
310	VLA	2019-07-19	BnA	10	$0.061 \pm 0.003$
343	VLA	2019-08-21	A	6	$0.089 \pm 0.003$
346	VLA	2019-08-24	A	3	$0.068 \pm 0.004$
351	VLA	2019-08-29	A	1.5	$0.146 \pm 0.013$
352	VLA	2019-08-30	A	10	$0.045 \pm 0.003$
364	GMRT	2019-09-11	–	0.6	< 0.105
396	VLA	2019-10-13	A	10	$0.031 \pm 0.003$
397	VLA	2019-10-14	A	6	$0.033 \pm 0.003$

<sup>a</sup>From VLASS

filters used in constructing Figure 1). Following Whitesides et al. (2017), we calculated absolute magnitudes using


$$M = m_{\text{obs}} - 5 \log_{10} \left( \frac{D_L}{10 \text{ pc}} \right) + 2.5 \log_{10}(1+z). \quad (2)$$

We cannot perform a true  $K$ -correction because most objects lack sufficient spectroscopic coverage. These equations will introduce systematic errors on the order of 0.1 mag.

In Figure 8 we show the rest-frame  $g$ -band light curve of ZTF18abvkwla compared to the light curves of transients in Table 1. The fast rise time of ZTF18abvkwla is most similar to that of iPTF15ul, AT2018cow, and perhaps iPTF16asu: it is faster than SN2011kl and the SNLS transients. ZTF18abvkwla fades much more quickly than iPTF16asu (which spectroscopically evolved into a Ic-BL SN) and in this sense is more similar to iPTF15ul and AT2018cow. In terms of peak luminosity, ZTF18abvkwla is close to iPTF15ul, AT2018cow, DES16X1eho, and iPTF16asu, and brighter than SN2011kl and the SNLS transients. However, we caution that the high peak luminosity of iPTF15ul results from a large host-galaxy extinction inferred in Hosseinzadeh et al. (2017), without which the peak magnitude would be  $-19.6$ .

Next we consider color evolution. ZTF18abvkwla showed tentative evidence for reddening over time, from  $g - r = -0.47 \pm 0.09$  mag at peak to  $g - r = -0.03 \pm 0.21$  mag in the final epoch a week later; however, this is only a  $2\sigma$  change. AT2018cow, iPTF15ul, and DES16X1eho remained very blue throughout the evolution of their optical light curves, whereas iPTF16asu reddened significantly as the SN became the dominant component.

Finally, we consider the spectral evolution of the transients in Table 1. Peak-light spectra were not obtained for DES16X1eho (Pursiainen et al. 2018) or the SNLS transients (Arcavi et al. 2016). The peak-light spectra of iPTF16asu, AT2018cow, and SN2011kl were featureless (Whitesides et al. 2017; Perley et al. 2019a;



**Figure 8.** The rest-frame  $g$ -band (observer-frame  $r$ -band) light curve of ZTF18abvkwla (black line), compared to light curves of other transients in the literature in as close to the same rest-frame filter as possible. Each panel shows one transient highlighted in orange for comparison, with the rest shown in grey in the background.

Greiner et al. 2015), and iPTF15ul<sup>4</sup> had a weak emission feature attributed to C III (Hosseinzadeh et al. 2017). After peak, iPTF16asu developed features of a Ic-BL SN (Whitesides et al. 2017), and AT2018cow had a complex spectral evolution, with a broad feature ( $v > 0.1c$ ) that appeared and disappeared over several days following peak light and a variety of emission lines that appeared one week later (Perley et al. 2019a). Unfortunately we do not have any spectra of ZTF18abvkwla after peak.

### 3.2. Radio Light Curve

In the previous section (§3.1) we compared the optical properties of ZTF18abvkwla to the transients in Table 1: the light curve shape, the color

<sup>4</sup> iPTF15ul was classified as Type Ibn in Hosseinzadeh et al. (2017), but the lack of distinct He I at peak make this classification uncertain.)

evolution, and the spectrum. In this section we compare the radio properties of ZTF18abvkwla to the same set of transients.

Of the transients in Table 1, only AT2018cow and GRB 111209A/SN 2011kl had a detected radio counterpart.<sup>5</sup> Prompt radio follow-up observations were also obtained for iPTF15ul<sup>6</sup> and iPTF16asu, but neither was detected. To our knowledge, Dougie, the SNLS transients, and DES16X1eho did not have deliberate radio follow-up observations; we searched the VLASS archive and found that all except SNLS04D4ec were observed but none were detected. In Figure 9 we show the radio measurements of the Table 1 transients compared to stellar explosions and tidal disruption events. For completeness, we also searched the positions of all of the transients in the two largest collections of unclassified fast-rising luminous optical transients reported to date, PS1 (Drout et al. 2014) and the Dark Energy Survey (Pursiainen et al. 2018). None were detected, and the limits are listed in Table 7.


As shown in Figure 9, ZTF18abvkwla is most similar in luminosity to long-duration GRB afterglows (Berger et al. 2003; Perley et al. 2014). The SED is also similar: in §2.2 we found that the SED of ZTF18abvkwla peaked near 10 GHz at  $\Delta t = 81$  d, while the SED of GRB 030329 ( $z = 0.1685$ ) peaked at 5 GHz (Berger et al. 2003) at 67 days post-explosion, and the SED of GRB 130427A ( $z = 0.340$ ) peaked at 10 GHz

<sup>5</sup> In the case of GRB 111209A/SN 2011kl, the radio emission was likely from the GRB afterglow itself (Kann et al. 2018).

<sup>6</sup> Observations of iPTF15ul were obtained within five days of the optical discovery, two observer-frame days after peak optical light, at 6 GHz and 22 GHz with the VLA, at 15 GHz with the Arcminute Microkelvin Imager (Zwart et al. 2008), and at 95 GHz with the Combined Array for Research in Millimeter-wave Astronomy (Bock et al. 2006). There was no detection at any frequency, with an RMS of 0.235 mJy with CARMA and an RMS of 0.03 mJy with AMI.

**Table 7.** Radio limits for rapidly evolving transients in Drout et al. (2014) and Pursiainen et al. (2018) The  $\Delta t$  is the number of days between the discovery date as listed in Drout et al. (2014) or the time of peak as listed in Pursiainen et al. (2018) and the epoch of the VLASS observation of that field.

ID	$z$	RA [hh:mm:ss]	Dec [dd:mm:ss]	$\Delta t$ (days)	Limit ( $\mu$ Jy)
PS1-10ah	0.074	10:48:15.784	+57:24:19.48	2836	102
PS1-11qr	0.324	09:56:41.767	+01:53:38.25	2467	130
PS1-12bb	0.101	09:57:23.866	+03:11:04.47	2174	149
PS1-12bv	0.405	12:25:34.602	+46:41:26.97	2642	129
PS1-12brf	0.275	22:16:06.892	-00:58:09.81	1892	124
PS1-11bbq	0.646	08:42:34.733	+42:55:49.61	2731	159
PS1-13duy	0.27	22:21:47.929	-00:14:34.94	1505	127
PS1-13dwm	0.245	22:20:12.081	+00:56:22.35	1422	155
PS1-10iu	–	16:11:34.886	+55:08:47.91	2689	103
PS1-13aea	–	12:18:14.320	+47:20:12.60	2199	88
PS1-13bit	–	16:12:00.765	+54:16:08.16	1618	104
PS1-13cgt	–	16:18:56.245	+54:19:33.71	1552	123
DES15S1fli	0.45	02:52:45.15	-00:53:10.21	826	150
DES13X3gms	0.65	02:23:12.27	-04:29:38.35	1520	139
DES15S1flf	0.23	02:51:09.36	-00:11:48.71	826	139
DES14S2anq	0.05	02:45:06.67	-00:44:42.77	1199	118
DES14X3pkl	0.3	02:28:50.64	-04:48:26.44	1100	105
DES15C3lpq	0.61	03:30:50.89	-28:36:47.08	849	145
DES16S1dxu	0.14	02:50:43.53	-00:42:33.29	385	154
DES15C3mgq	0.23	03:31:04.56	-28:12:31.74	835	99
DES16X1eho	0.76	02:21:22.87	-04:31:32.64	365	152
DES16X3cxn	0.58	02:27:19.32	-04:57:04.27	393	128
DES15C3lzm	0.33	03:28:41.86	-28:13:54.96	839	106
DES13C3bcok	0.35	03:32:06.47	-28:37:29.70	1513	98
DES15C3nat	0.84	03:31:32.44	-28:43:25.06	810	108
DES15C3opk	0.57	03:26:38.76	-28:20:50.12	777	125
DES15C3opp	0.44	03:26:57.53	-28:06:53.61	781	112
DES13X3npb	0.5	02:26:34.11	-04:08:01.96	1411	122
DES16C3axz	0.23	03:31:14.15	-28:40:00.25	523	100
DES16C3gin	0.35	03:31:03.06	-28:17:30.98	391	107
DES14X1bnh	0.83	02:14:59.79	-04:47:33.32	1172	145
DES16X3ega	0.26	02:28:23.71	-04:46:36.18	357	111
DES15C3mfu	–	03:28:36.08	-28:44:20.00	835	187
DES13C3abtt	–	03:30:28.91	-28:09:42.12	1513	107
DES15C3pbi	–	03:28:56.68	-28:00:07.98	772	182
DES15X3atd	–	02:23:21.64	-04:17:28.95	830	146
DES13C3nxi	–	03:27:51.22	-28:21:26.21	1559	75
DES13C3smn	–	03:27:53.08	-28:05:00.93	1564	124
DES13X3aakf	–	02:22:50.84	-04:41:57.01	1441	108
DES13X3afjd	–	02:28:00.31	-04:34:59.39	1411	123
DES13X3kgm	–	02:26:00.92	-04:51:59.29	1508	103
DES16S2fqu	–	02:47:05.94	-00:20:50.40	356	139
DES16X1ddm	–	02:15:18.88	-04:21:52.07	386	111
DES16X3ddi	–	02:21:45.39	-04:41:08.95	393	127
DES16X3erw	–	02:24:49.31	-04:30:51.45	357	117



**Figure 9.** The 10 GHz radio light curve of ZTF18abvkwla compared to low-frequency (1–10 GHz) light curves of different classes of energetic explosions: tidal disruption events (purple; Zauderer et al. 2011; Berger et al. 2012; Zauderer et al. 2013; Alexander et al. 2016; Eftekhari et al. 2018), supernovae exploding in dense CSM (blue lines,  $\gtrsim 10^{37}$  erg s $^{-1}$ ; Soderberg et al. 2005, 2006; Salas et al. 2013), relativistic Ic-BL supernovae (red lines; Kulkarni et al. 1998; Soderberg et al. 2010), AT2018cow (black line, small stars), long-duration gamma-ray bursts (orange lines; Berger et al. 2003; Hancock et al. 2012; Perley et al. 2014; van der Horst et al. 2014), and “ordinary” supernovae ( $\lesssim 10^{37}$  erg s $^{-1}$ ; Weiler et al. 1986, 2007; Horesh et al. 2013; Krauss et al. 2012). The CSS161010 light curve was taken from Coppejans et al. (2020). The AT2018cow light curve is at 9 GHz with data taken from Ho et al. (2019a), Margutti et al. (2019), and Bietenholz et al. (2020).

(Perley et al. 2014) at a similar epoch post-explosion.

### 3.3. A Starburst Host Galaxy

In Sections 3.1 and 3.2 we compared the optical and radio properties of ZTF18abvkwla, respectively, to other transients in the literature.

Here we put its host galaxy properties into context.

Galaxies with very high specific star-formation rates (e.g.,  $sSFR \gtrsim 10^{-8}$  yr $^{-1}$ , our operational definition of a “starburst”) contribute a small fraction of star-formation in the low-redshift Universe (Lee et al. 2009), so the appearance of ZTF18abvkwla in such a galaxy ( $sSFR \sim 1.4 \times 10^{-8}$  yr $^{-1}$ ) is notable. However, their contribution to low-metallicity star-formation is more significant, as they are typically low-mass and therefore low-metallicity (Tremonti et al. 2004). They are also promising candidates to experience a top-heavy IMF (Dabringhausen et al. 2009) and potential sites of enhanced binary or dynamical stellar interactions (van den Heuvel & Portegies Zwart 2013). Each of these mechanisms have been appealed to in attempts to interpret the relatively high abundance of exotic transients of other types found in these systems, including superluminous SNe (SLSNe; Lunnan et al. 2014; Leloudas et al. 2015; Perley et al. 2016; Schulze et al. 2018), broad-lined Ic SNe (Modjaz et al. 2019), GRBs (Fruchter et al. 2006; Krühler et al. 2015; Schulze et al. 2015; Vergani et al. 2015), and at least some fast radio bursts (Katz 2016; Tendulkar et al. 2017).

Based on our measurements in §2.1.2 we conclude the following about the host of ZTF18abvkwla:

*The host is not an AGN* — We confirm the lack of any evidence for an optical AGN based on the very weak [NII] emission. The host falls squarely in the star-forming locus of the BPT diagram (Figure 10a).

*The host metallicity is typical for its mass* — The host is relatively metal-poor: the precise number is of course scale-dependent, but using the Z94 scale we calculate [O/H] of 8.45, or about 0.6×Solar. This is a lower metallicity than the majority of star-formation in the local

Universe, but not an outlier and unexceptional for low-mass galaxies in particular (Figure 10b).

*The star-formation intensity is similar to extreme SLSN and GRB hosts* — The most striking nature of the host galaxy is its very high specific star-formation rate, which is evident in Figure 10c and 10d.

The host of AT2018cow was also a dwarf galaxy, although it was more massive than that of ZTF18abvkwla and not starbursting, with a mass and star-formation rate of  $1.4 \times 10^9 M_\odot$  and  $0.22 M_\odot \text{yr}^{-1}$  respectively (Perley et al. 2019a). The host galaxy of DES16X1eho had a stellar mass  $\log(M/M_\odot) = 9.96^{+0.14}_{-0.51}$  and a specific SFR of  $\log(\text{sSFR}/M_\odot \text{yr}^{-1}) = -9.25$  (Pursiainen et al. 2018). The host galaxy of iPTF16asu had a stellar mass  $M = 4.6^{+2.0}_{-2.3} \times 10^8 M_\odot$  and an H $\alpha$  SFR of  $0.7 M_\odot \text{yr}^{-1}$ , corresponding to a sSFR of  $1.4 \text{Gyr}^{-1}$  (Whitesides et al. 2017). Finally, the host galaxies of the SNLS transients harbored relatively evolved stellar populations, and were noted to be markedly different from starburst galaxies (Arcavi et al. 2016).

#### 4. INTERPRETATION

Even with the small number of events in the Table 1 menagerie, the diversity of optical and radio properties (§3.1, §3.2) suggests that there are several progenitor systems involved. In this section we model the optical and radio light curves of ZTF18abvkwla and discuss the implications for the progenitor.

##### 4.1. Modeling the Optical Light Curve

Shock-interaction with extended low-mass material is an efficient mechanism for producing a fast-peaking luminous optical light curve. Shock breakout occurs when the photon diffusion time drops below the shock crossing time ( $\tau < c/v_s$ , where  $\tau$  is the optical depth and  $v_s$  is the shock velocity). For normal stellar progenitors, this emission is primarily at X-ray and UV wavelengths and lasts for seconds to a fraction of an hour. In the wake of this shockwave, the

outer stellar material is heated to high temperatures, and as it cools it radiates on the timescale of a day (“cooling envelope” emission). See Waxman & Katz (2017) for a review.

Prior to core-collapse, massive stars can undergo mass-loss via steady winds or eruptive episodes (Smith 2014). As a result, a star can be surrounded by dense, recently-expelled material at the time of explosion. If this material is optically thick, it increases the effective radius of the star and prolongs the light curve from shock breakout. If the light curve of ZTF18abvkwla arises from shock breakout in a shell, we can estimate the radius of this extended material (CSM) assuming a rise to peak bolometric luminosity  $t_{\text{rise}} < 2 \text{d}$ , a peak luminosity  $L_{\text{bol}} > 10^{44} \text{erg s}^{-1}$  and a typical SN shock velocity of  $10^4 \text{km s}^{-1}$ . The rise timescale is

$$\begin{aligned} t_{\text{BO}} &\sim \frac{R_{\text{CSM}}}{v_s} \\ &= (1.3 \text{d}) \left( \frac{R_{\text{CSM}}}{10^{15} \text{cm}} \right) \left( \frac{v_s}{10^4 \text{km s}^{-1}} \right)^{-1}. \end{aligned} \quad (3)$$


For ZTF18abvkwla, we find  $R_{\text{CSM}} < 1.5 \times 10^{15} \text{cm}$ .

We can also estimate the mass in the shell, assuming that the shock deposits half its kinetic energy  $(1/2)\rho v_s^2$  and that this deposited energy is  $E_{\text{BO}} \sim 4\pi R^2 dR e_s$ , where the energy density reflects the amount of thermal energy in the layer. The luminosity scales as

$$\begin{aligned} L_{\text{BO}} &\sim \frac{E_{\text{BO}}}{t_{\text{cross}}} \sim \frac{v_s^3}{4} \frac{dM}{dR} = (2.2 \times 10^{45} \text{erg s}^{-1}) \\ &\times \left( \frac{v_s}{10^4 \text{km s}^{-1}} \right)^3 \left( \frac{dM}{M_\odot} \right) \left( \frac{dR}{10^{15} \text{cm}} \right)^{-1}. \end{aligned} \quad (4)$$

Assuming  $dR \sim R$ , we find  $M_{\text{CSM}} < 0.07 M_\odot$ . In this framework, the differences in the light curves of different objects corresponds to differences in the shell mass, shell radius, and shock velocity. The luminosity is most sensitive to





**Figure 10.** Comparison of the host galaxy of ZTF18abvkwla to <11 Mpc comparison galaxies (grey) and to the host galaxies of nearby hydrogen-poor SLSNe (diamonds), as in Perley et al. 2016. Light diamonds indicate mass-metallicity estimated metallicities. Comparison galaxies are weighted by their SFR; histograms show the SFR-weighted binned totals on each axis. ZTF18abvkwla is indicated by a yellow cross. From top left: (a) BPT diagram. (b) Mass–star-formation rate relation. (c) Mass–metallicity relation. (d) Specific star-formation-rate–metallicity relation. The host is a starbursting galaxy with no evidence of AGN activity, and while it is metal-poor it is not particularly so given its mass.

the velocity, so it is possible that the transients in Table 1 are distinguished by fast velocities, which would naturally explain the inclusion of a Ic-BL SN. For a fixed shock velocity, a fast rise time corresponds to a small shell radius, which in turn requires a large shell mass to produce a high luminosity.

Another possibility is that the light curve is powered not by shock breakout in a shell, but by post-shock envelope-cooling emission. For example, this was the model invoked for iPTF16asu (Whitesides et al. 2017), which led to an inferred shell mass of  $0.45 M_{\odot}$  and a shell

radius of  $1.7 \times 10^{12}$  cm. The light curve of ZTF18abvkwla has a similar rise time but a higher peak luminosity than that of iPTF16asu, and the effective temperature at peak is significantly higher. According to the one-zone analytic formalism in Nakar, & Piro (2014) and Piro (2015), a higher peak temperature for a fixed rise time and a fixed opacity arises from a larger shell radius. A larger shell radius can also explain the higher bolometric luminosity, although that could also arise from a larger explosion energy or faster ejecta velocity.

Another mechanism suggested to explain the optical light curve of AT2018cow was reprocessing by dense outer ejecta (Margutti et al. 2019). In this picture, a central source (such as an accretion disk or magnetar) emits high-energy (i.e. X-ray) emission, which is reprocessed by surrounding material to produce lower-energy (i.e. optical) radiation. This is one setup for tidal disruption events, in which case the surrounding material is unbound stellar debris (Strubbe & Quataert 2009). Indeed, several properties of ZTF18abvkwla and AT2018cow are similar to TDEs in the literature, such as the photospheric radius of  $10^{14}$ – $10^{15}$  cm, the effective temperature of  $10^4$  K, and high radio luminosities attributed to jets (for reviews of TDE observations, see Gezari (2012) and Komossa (2015)).

Regardless of the power source at peak, we also use the optical light curve to put an upper limit on the mass of  $^{56}\text{Ni}$  that could have been synthesized in the explosion. Using Equation (16) in Kasen (2017), the luminosity from the radioactive decay of  $^{56}\text{Ni}$  is


$$L(t) = 2 \times 10^{43} \left( \frac{M_{\text{Ni}}}{M_{\odot}} \right) \times \left[ 3.9e^{-t/\tau_{\text{Ni}}} + 0.678 \left( e^{-t/\tau_{\text{Co}}} - e^{-t/\tau_{\text{Ni}}} \right) \right] \text{ erg s}^{-1} \quad (5)$$

where  $\tau_{\text{Ni}} = 8.8$  d and  $\tau_{\text{Co}} = 113.6$  d. Using the final  $g$ -band measurement ( $g = 21.51 \pm 0.21$ ) at  $\Delta t = 10$  d ( $\Delta t = 8$  d rest-frame)  $L \approx \lambda F_{\lambda} \approx 1.4 \times 10^{43}$  erg s $^{-1}$ , so the amount of  $^{56}\text{Ni}$  that could power the light curve at this epoch is  $M_{\text{Ni}} \lesssim 0.36 M_{\odot}$  (Figure 4). From a compilation of CC SNe, Lyman et al. (2016) found nickel masses of  $0.11 \pm 0.04 M_{\odot}$  for Type IIb SNe,  $0.17 \pm 0.16 M_{\odot}$  for Type Ib SNe,  $0.22 \pm 0.16 M_{\odot}$  for Type Ic SNe, and  $0.32 \pm 0.15 M_{\odot}$  for Type Ic-BL SNe. So, we cannot rule out an underlying nickel-powered light curve for ZTF18abvkwla.

#### 4.2. Modeling the Radio Light Curve

The high luminosity and fast variability timescale of the 10 GHz light curve implies a high brightness temperature  $T_B \approx 10^{11}$  K, so we conclude that the emission is synchrotron radiation. In the first epoch, the 10 GHz observation is declining and has an in-band (8–12 GHz) spectral index of  $\alpha = -0.16 \pm 0.05$  where  $F_{\nu} \propto \nu^{-\alpha}$ . This is much shallower than the optically thick ( $\alpha = -2.5$ ) or the optically thin ( $\alpha = +0.7$ ) regimes of a synchrotron self-absorption (SSA) spectrum, which suggests that the peak of the SED is near 10 GHz (observer-frame) at this epoch. In what follows, we assume that the SSA spectrum has a rest-frame peak frequency  $\nu_p \lesssim 8$  GHz (the bottom of the band) and a rest-frame peak flux density  $F_p \gtrsim 0.364$  mJy.

When the SSA peak is known, the outer shock radius  $R_p$  and magnetic field strength  $B_p$  can be derived assuming that energy is equally partitioned into magnetic fields and relativistic electrons (Scott & Readhead 1977; Readhead 1994). We use the equations for  $R_p$  and  $B_p$  for radio SNe in Chevalier (1998) (Equations 11 and 12). Assuming an optically thin spectral index of  $\nu^{-1}$  and a filling factor  $f = 0.5$ , we find  $R_p \gtrsim 8.0 \times 10^{16}$  cm and  $B_p \lesssim 0.51$  G. So, the mean velocity until  $t_{\text{obs}} = 81$  d is  $\Gamma\beta c = R_p(1+z)/t_{\text{obs}} = 0.38c$ . Using Equations 12, 16, and 23 in Ho et al. (2019a), and assuming  $\epsilon_e = \epsilon_B = 1/3$ , we find that the shock has swept up energy  $U = 3.4 \times 10^{49}$  erg into an ambient medium of density  $n_e = 190$  cm $^{-3}$ , corresponding to a mass-loss rate of  $\dot{M} = 5.8 \times 10^{-4} M_{\odot} \text{ yr}^{-1}$  assuming a wind velocity  $v_w = 1000$  km s $^{-1}$ . In Figure 11 we show these quantities compared to those of other energetic explosions. The peak radio luminosity density is directly proportional to  $U/R$ , the energy swept up by the shock divided by the shock radius (right-hand side of Figure 11). So, the fact that ZTF18abvkwla, AT2018cow, and CSS161010 are distinguished by high radio luminosities is primarily a consequence of a large explosion energy.



**Figure 11.** Approximate luminosity and frequency of the SSA peak of ZTF18abvkwla at  $\Delta t = 81$  d (observer-frame), compared to other energetic explosions in the literature, including AT2018cow (Ho et al. 2019a; Margutti et al. 2019) and CSS161010 (Coppejans et al. 2020). Lines of constant mass-loss rate are shown in units of  $10^{-4} M_{\odot} \text{yr}^{-1}$  scaled to a wind velocity of  $1000 \text{ km s}^{-1}$ . The corresponding energy of the explosion (assuming equipartition) is shown on the right-hand side.

#### 4.3. Progenitor Systems and a Search for an Associated Gamma-ray Burst

The physical setups outlined in §4.1 — a shock driven through a shell, reprocessing of a high-energy compact source by optically thick material — could arise in a variety of different progenitor systems. An additional clue for ZTF18abvkwla is the host galaxy, which experienced a very recent burst of star-formation activity. In that sense, a massive-star origin seems most natural.

AT2018cow was suggested to have two distinct components: a shock driven through dense equatorial material (producing the optical emission), and a faster polar outflow (producing the radio emission; Margutti et al. 2019). As

shown by early millimeter observations (Ho et al. 2019a), later radio observations (Margutti et al. 2019), and Very Long Baseline Interferometry (Bietenholz et al. 2020; Mohan et al. 2020), the fast outflow was subrelativistic with a near-constant velocity of  $v = 0.1c$ . In ZTF18abvkwla, the radio-emitting ejecta is faster:  $> 0.38c$  at the same epochs when the outflow from AT2018cow was  $0.1c$ . As shown in Figure 11, the higher luminosity at late times arises from this faster velocity; the explosion energy of the two events appears to have been similar.

Because the late-time radio light curve is similar of that of GRBs, we searched for potential GRB counterparts to ZTF18abvkwla in the period between the last non-detection (MJD 58372.4206; 2018-09-11 10:05:39.84) and the first detection (MJD 58373.4075; 2018-09-12 09:46:48.00). There were two bursts detected by the interplanetary network (IPN; Hurley et al. 2010, 2016), one by the Gamma-ray Burst Monitor (GBM) aboard the *Fermi* spacecraft (Gruber et al. 2014; von Kienlin et al. 2014; Bhat et al. 2016) and one detected by the Konus-Wind experiment aboard the *Wind* spacecraft (Aptekar et al. 1995). The positions of both bursts are inconsistent with that of ZTF18abvkwla.

Due to the lack of detected GRB, we can set a limit on the fluence and corresponding isotropic equivalent energy of a prompt burst associated with ZTF18abvkwla. The IPN has essentially a 100% duty cycle across the sky, and detects GRBs with  $E_p > 20 \text{ keV}$  down to  $6 \times 10^{-7} \text{ erg cm}^{-2}$  at 50% efficiency (Hurley et al. 2010, 2016). At  $t_0$ , the estimated 20–1500 keV limiting peak flux at the position of ZTF18abvkwla was  $2 \times 10^{-7} \text{ erg cm}^{-2} \text{ s}^{-1}$  for a Band model that has  $E_{\text{pk}}$  in the 50–500 keV range. At the distance of ZTF18abvkwla, this corresponds to a limit on the isotropic peak luminosity of  $L_{\text{iso}} < 5 \times 10^{49} \text{ erg s}^{-1}$ . Therefore

we strongly disfavor an on-axis classical GRB (which is also consistent with the lack of observed optical afterglow emission).

Among GRBs, two events have shown evidence for a luminous optical blackbody component at early times: GRB 060218 ( $z = 0.033$ ; Soderberg et al. 2006; Mirabal et al. 2006; Pian et al. 2006; Sollerman et al. 2006; Ferrero et al. 2006) and GRB 101225A ( $z \approx 0.3$ ; Thöne et al. 2011). GRB 060218 was a very long-duration ( $T_{90} \approx 2100$  s) low-luminosity ( $L_{\text{iso}} = 2.6 \times 10^{46}$  erg s $^{-1}$ ) GRB associated with the ICBL SN 2006aj (Cano et al. 2017). A GRB with these properties cannot be ruled out by our limits. GRB 101225A also had a very long duration  $T_{90} > 2000$  sec, and a candidate faint ( $M \approx -16.7$ ) SN counterpart.

As in the case of AT2018cow, we cannot rule out a TDE origin. In that case, the similarity to the light curve of AT2018cow would suggest a similar kind of system, i.e. an intermediate-mass black hole ( $M \sim 10^4 M_{\odot}$ ; Perley et al. 2019a) with a white dwarf (Kuin et al. 2019) or a Solar-type (Perley et al. 2019a) stellar companion. In the case of AT2018cow, the main argument against a TDE hypothesis was the large ambient density ( $10^5$  cm $^{-3}$ ) from millimeter (Ho et al. 2019a) and radio (Margutti et al. 2019) observations. For ZTF18abvkwla, assuming that the flat spectral index indicates a 10 GHz peak at 81 d, we find a much lower density ( $10^2$  cm $^{-3}$ ). Among TDEs, the radio light curve of ZTF18abvkwla is most similar to that of the TDE candidate IGR J12580+0134 (Irwin et al. 2015), which had a nearly identical  $\nu L_{\nu}$  (and fade rate) one year post-discovery. The radio emission from IGR J12580+0134 has been attributed to an off-axis relativistic jet (Irwin et al. 2015; Lei et al. 2016) but interpretation is complicated by the coincidence of the source with a known AGN.

## 5. RATE ESTIMATE

An important clue to the progenitor of sources like ZTF18abvkwla is the cosmological rate. Furthermore, three fast-luminous transients – SN 2011kl (associated with GRB 111209A), AT2018cow, and ZTF18abvkwla – have detected luminous radio emission, although the radio emission from SN 2011kl likely arose from the GRB afterglow. Clearly, being able to recognize additional members of this phase-space in optical surveys would be valuable for radio follow-up observations. In this section, we conduct an archival search of 18 months of the 1DC survey (2018 Apr 3 – 2019 Oct 18 UT) to estimate the rate of transients in the phase-space of Figure 1 and delineate false positives.

First we selected field-nights in the survey for which the 1-night coverage was approximately maintained. Specifically, we require

- at least one observation the night before ( $0.5 < dt < 1.5$  days)
- at least one observation two nights before ( $2.5 < dt < 1.5$  days)
- at least three observations in the next five nights ( $dt < 5.5$  days)

We find 8064 fields satisfying these criteria. Of these, 6854 fields (85%) have limiting magnitude  $> 19.75$  mag and 4596 fields (57%) have limiting magnitude  $> 20.5$  mag. The dominant effect is lunation, with some night-to-night variations due to weather.

For each of the 8064 field-nights, we searched for fast transients. To detect a fast transient, we require that the peak of the light curve be “resolved:” that is, that there are measurements both before and after peak light that are  $> 0.75$  mag fainter than the peak magnitude. We then measure the time from 0.75 mag below peak to peak by linearly interpolating the light curve. If this rise time is  $< 5$  d, we include the transient in our sample. More specifically, we filtered sources as summarized in Table 8. We

scanned the remaining 659 sources by eye and removed sources with very noisy light curves or flaring behavior.

**Table 8.** Filtering criteria for sources similar to ZTF18abvkwla in the ZTF 1DC survey

Criteria	# sources remaining
Real <sup>a</sup> , bright <sup>b</sup> , pos. sub. <sup>c</sup> , not star <sup>d</sup>	758,528
Short duration <sup>e</sup> and peak resolved <sup>f</sup>	659

NOTE— <sup>a</sup>  $\text{drb} > 0.99$  <sup>b</sup>  $\text{magpsf} < 20$  <sup>c</sup>  $\text{isdiffpos}=\text{'t'}$  or  $\text{'1'}$  <sup>d</sup>  $\text{not}(\text{sgscore1} > 0.76 \text{ and } \text{distpsnr1} < 1)$  <sup>e</sup> Duration between 1 and 100 days <sup>f</sup> Peak has preceding or subsequent detection/non-detection in a  $\pm 5$  d window that is at least 0.75 mag fainter

In Table 9 we list all 27 sources with rise times faster than 5 d, including ZTF18abvkwla itself. Five sources are spectroscopically classified SNe: two Type II, two Type Ibn, and one Type IIb. Three sources are classified as CVs, two spectroscopically and one by cross-matching with the AAVSO International Variable Star Index VSX (Watson et al. 2017). Two are very likely flare stars based on previous detections in Pan-STARRS individual-epoch images, and a third is a likely flare star based on a GALEX counterpart. Nine sources are likely extragalactic (based on proximity to a host galaxy). When redshift estimates for these galaxies were not available, we attempted to obtain them using LRIS on 17 Feb 2020. Two sources remain without definitive redshift estimates, so we provide a photometric redshift from LegacySurvey DR8. One source (ZTF18abxxeai) has a very faint host classified as a PSF in LegacySurvey DR8, and the remaining five sources have no clear host counterpart.

Of the sources with a definitive host redshift measurement, ZTF18abvkwla is the only one that is more luminous than  $M = -20$  mag. Clearly, the primary interlopers in searches for transients like ZTF18abvkwla are CVs and less luminous SNe. CVs can be ruled out on the basis of repeated flaring, whereas less luminous

SNe can only be ruled out if the redshift of the host galaxy is known *a priori*. Aside from ZTF18abvkwla, eight transients in our sample remain as possibly having  $M_{\text{g,peak}} < -20$ , although the lack of an obvious host for six of them suggest that these may be CVs.

We take eight as an upper limit for the number of transients in ZTF that could fall within the phase-space of Figure 1. Of these, three peak brighter than 19 mag, and four have a peak between 19 and 19.75 mag. We now calculate two all-sky rates. First we assume that the transient peaks at  $< 19$  mag, in which case we discard field-nights with a limiting magnitude shallower than 19.75 mag. Then we assume that the transient peaks at  $< 19.75$  mag, in which case we discard field-nights with a limiting magnitude shallower than 20.5 mag.

Each ZTF field is  $47 \text{ deg}^2$ , but there is latitude-dependent overlap that has to be taken into account when converting this to a rate per square degrees in the sky. For the primary grid, a rough estimate of the fill factor is 87.5%. For the 1DC survey, the footprint is 10% smaller than the number of fields multiplied by 47 square degrees. So, taking fill factor and overlap into account, we estimate a typical area-per-field of  $37 \text{ deg}^2$ . So for transients brighter than 19 mag we have  $2.5 \times 10^5 \text{ deg}^2 \text{ d}$  and for transients brighter than 19.75 mag we have  $1.7 \times 10^5 \text{ deg}^2 \text{ d}$ . For transients peaking brighter than 19 mag we have a limiting all-sky rate

$$3 \times \frac{41253 \text{ deg}^2}{2.5 \times 10^5 \text{ deg}^2 \text{ d}} \times \frac{365 \text{ d}}{1 \text{ yr}} \approx 180 \text{ yr}^{-1}. \quad (6)$$

For transients peaking brighter than 19.5 mag we have a limiting all-sky rate

$$4 \times \frac{41253 \text{ deg}^2}{1.7 \times 10^5 \text{ deg}^2 \text{ d}} \times \frac{365 \text{ d}}{1 \text{ yr}} \approx 350 \text{ yr}^{-1}. \quad (7)$$

Now, we use the limiting magnitude to estimate a volumetric rate. Assuming a tran-

**Table 9.** Fast-rising transients in ZTF resulting from our archival search of the one-day cadence survey. In the redshift column, a range refers to the 68 percentile range on the photometric redshift from LegacySurvey DR8 (we provide a corresponding range of absolute magnitude) and a single value corresponds to a spectroscopic redshift. When the distance is known, the peak mag is an absolute magnitude, and when the distance is not known the peak mag is an apparent magnitude. These values correspond to the filter as close to rest-frame  $g$ -band as possible, and when the distance is not known they correspond to the observed  $g$ -band filter. Magnitudes are corrected for Galactic extinction and timescales are in rest-frame when the redshift is known, and in observer-frame when the redshift is not known.

ZTF Name (IAU Name)	Redshift	Peak Mag	$t_{\text{rise}}$	$t_{\text{fade}}$	Type	Notes
18abvkwla	0.2714	$-20.59 \pm 0.07$	$1.83 \pm 0.05$	$3.12 \pm 0.22$	FBOT	This paper
19aavbjfp (SN2019fkl)	0.028	$-17.4 \pm 0.4$	$3.2 \pm 0.9$	$21.8 \pm 6.1$	SN II	
19abgbdep (AT2019lbv)	0.0318	$-18.36 \pm 0.03$	$2.32 \pm 0.03$	$14.4 \pm 0.9$	SN II	
18aalrxas	0.0588	$-18.43 \pm 0.03$	$1.86 \pm 0.02$	$2.4 \pm 0.3$	SN I Ib	Fremling et al. (2019)
19abuvqgw (AT2019php)	0.087	$-18.9 \pm 0.1$	$3.6 \pm 0.1$	$4.5 \pm 0.3$	SN I bn	
19aapfinki (SN2019deh)	0.05469	$-19.90 \pm 0.01$	$4.38 \pm 0.03$	$7.2 \pm 0.4$	SN I bn	
18abskrix	Galactic	$17.78 \pm 0.02$	$1.26 \pm 0.03$	$2.5 \pm 0.2$	CV	Spectroscopic classification
18absrffm (AT2018ftw)	Galactic	$16.34 \pm 0.01$	$2.60 \pm 0.03$	$5.00 \pm 0.03$	CV	Spectroscopic classification
18abyzkeq	Galactic	$18.32 \pm 0.10$	$1.37 \pm 0.04$	$0.93 \pm 0.06$	CV	AAVSO Name: CSS 151114:224934+375554
18ablxawt	Galactic	$18.31 \pm 0.04$	$2.4 \pm 0.3$	$5.2 \pm 0.9$	Likely flare star	Previous detection in PS1 DR2 at $i = 19.4$
19abpwygn	-	$16.74 \pm 0.01$	$2.03 \pm 0.03$	$1.73 \pm 0.03$	Likely flare star	Previous detection in PS1 DR2 at $z = 18.75$
18abyjgaa	-	$18.39 \pm 0.03$	$0.82 \pm 0.03$	$2.05 \pm 0.08$	Likely flare star	GALEX source, possible flaring in PS1 DR2
18aasaiyp	0.104	$-19.13 \pm 0.05$	$1.9 \pm 0.6$	$17.1 \pm 0.6$	Unknown	
18abuvqgo	0.155	$-19.93 \pm 0.05$	$4.7 \pm 0.2$	$9.9 \pm 0.6$	Unknown	
18abydmfv (AT2018hkr)	0.042	$-18.66 \pm 0.03$	$3.15 \pm 0.04$	$7.7 \pm 2.5$	Unknown	
18acepuyx (AT2018kxh)	0.0711	$-19.1 \pm 0.2$	$1.4 \pm 0.3$	$10.8 \pm 1.2$	Unknown	
19aatoboa (AT2019esf)	0.0758	$-18.90 \pm 0.03$	$2.41 \pm 0.03$	$4.9 \pm 0.3$	Unknown	
19abgbbp (AT2019leo)	0.0625	$-18.83 \pm 0.03$	$4.2 \pm 0.3$	$> 5$	Unknown	
19abiyhd (AT2019lwj)	0.07	$-18.11 \pm 0.05$	$2.5 \pm 0.2$	$4.1 \pm 0.3$	Unknown	
19aaadfc	0.08–0.15	$19.04 \pm 0.04$	$2.44 \pm 0.15$	$5.86 \pm 0.15$	Unknown	
19aanvhyc (AT2019coi)	0.056–0.076	$18.41 \pm 0.04$	$4.39 \pm 0.04$	$12.1 \pm 2.1$	Unknown	
18abxxeai		$18.55 \pm 0.06$	$1.9 \pm 0.1$	$6.0 \pm 0.8$	Unknown	‘PSF’ host in LegacySurvey DR8
18acgnwpo	-	$18.90 \pm 0.05$	$0.52 \pm 0.03$	$6.5 \pm 0.5$	Unknown	No clear host
19aanqqzb	-	$16.63 \pm 0.04$	$1.91 \pm 0.07$	$1.2 \pm 0.1$	Unknown	No clear host
19aaqfdvu	-	$19.02 \pm 0.06$	$2.0 \pm 0.2$	$1.6 \pm 0.4$	Unknown	No clear host
19aaxfqyx	-	$18.76 \pm 0.03$	$0.98 \pm 0.03$	$4.68 \pm 0.27$	Unknown	No clear host
19abfzfs	-	$19.36 \pm 0.17$	$3.7 \pm 2.2$	$13.4 \pm 3.2$	Unknown	No clear host

sient that peaks at  $M = -20$  mag, requiring a peak apparent magnitude brighter than 19 mag restricts our sensitivity to 400 Mpc. So, we find a volumetric rate of  $7 \times 10^{-7} \text{ yr}^{-1} \text{ Mpc}^{-3}$ . Requiring a peak apparent magnitude brighter than 19.75 mag restricts our sensitivity to 560 Mpc, leading to a volumetric rate of  $4 \times 10^{-7} \text{ yr}^{-1} \text{ Mpc}^{-3}$ . For reference, we provide rates of core-collapse SNe and GRBs in Table 10. The rate of events like ZTF18abvkwla appears to be at least two orders of magnitude smaller than

the CC SN rate, and more similar to the rate of GRBs in the local universe.

## 6. PROSPECTS FOR DETECTING X-RAY EMISSION

Clearly, radio observations are an important avenue of follow-up for transients like ZTF18abvkwla. Another valuable avenue is X-ray observations, which were not obtained for ZTF18abvkwla. We can estimate what the predicted X-ray luminosity would be from inverse

**Table 10.** Local ( $z = 0$ ) Rates of core-collapse supernovae and GRBs. Approximately 30% of CC SNe arise from a progenitor stripped of its hydrogen envelope. Among these stripped events, there are roughly equal numbers of Ib, Ic, and IIb events. Of the Ic events,  $\sim 10\%$  are “broad-lined” with photospheric velocities  $\gtrsim 30,000$  km/s. The fraction of Ic-BL SNe with associated GRBs has been estimated to be 1/40 (Graham & Schady 2016) although the rate is highly uncertain. The fraction of Ic-BL SNe with associated LLGRBs remains uncertain. Note that the rate quoted for LLGRBs does not include a beaming correction.

Class	Rate/Fraction	References
SN II	$4.47 \pm 1.39 \times 10^{-5} \text{ yr}^{-1} \text{ Mpc}^{-3}$	[1]
SN Ibc	$2.58 \pm 0.72 \times 10^{-5} \text{ yr}^{-1} \text{ Mpc}^{-3}$	[1]
Frac. of Ibc SN that are Ic	$0.69 \pm 0.09$	[2,3]
Frac. of Ic SN that are Ic-BL	$0.21 \pm 0.05$	[2,3]
LLGRB	$2.3^{+4.9}_{-1.9} \times 10^{-7} \text{ yr}^{-1} \text{ Mpc}^{-3}$	[4]
	$3.3^{+3.5}_{-1.8} \times 10^{-7} \text{ yr}^{-1} \text{ Mpc}^{-3}$	[5]
$\ell$ GRB	$\mathcal{R}_{\text{obs}} = 4.2^{+9.0}_{-4.0} \times 10^{-10} \text{ yr}^{-1} \text{ Mpc}^{-3}$	[6]
	$f_b = 0.0019 \pm 0.0003$	[7]
	$f_b = 0.013 \pm 0.004$	[8]

**References**—[1] Li et al. (2011), [2] Kelly & Kirshner (2012), [3] Graham & Schady (2016), [4] Soderberg et al. (2006), [5] Liang et al. (2007), [6] Lien et al. (2014), [7] Frail et al. (2001), [8] Guetta et al. (2005)

Compton scattering, using the optical and radio luminosities:

$$\frac{L_X}{L_{\text{radio}}} = \frac{u_{ph}}{u_B}. \quad (8)$$

Taking  $L_{\text{radio}} = 10^{40} \text{ erg s}^{-1}$ ,  $u_{ph} = 10^{44} \text{ erg s}^{-1} / (4\pi R^3/3)$  where  $R = 10^{14} \text{ cm}$ , and  $u_B = B^2/8\pi$  where  $B = 0.6 \text{ G}$ , we find  $L_X \approx 10^{43} \text{ erg s}^{-1}$ . This is even more luminous than the X-ray emission observed accompanying AT2018cow, which had  $L_X \approx 10^{42} \text{ erg s}^{-1}$  (Rivera Sandoval et al. 2018; Ho et al. 2019a; Margutti et al. 2019). To our knowledge there were no X-ray follow-up observations of DES16X1eho, while observations of iPTF16asu resulted in an X-ray upper limit of  $10^{43} \text{ erg s}^{-1}$ . Hosseinzadeh et al. (2017) report pre-peak UV measurements from *Swift* for iPTF15ul, but to our knowledge X-ray observations have

not been reported. We measured an upper limit of  $0.005 \text{ count s}^{-1}$  in a single epoch from the publicly available *Swift* data. Assuming  $n_H = 1.7 \times 10^{20} \text{ cm}^{-2}$  and a power-law source model with a photon index  $\Gamma = 2$  we obtain an upper limit on the unabsorbed 0.3–10 keV luminosity of  $2 \times 10^{42} \text{ erg s}^{-1}$ .

## 7. SUMMARY AND CONCLUSIONS

ZTF18abvkwla is distinguished by two key characteristics: a fast-evolving optical light curve with a hot ( $T > 40,000 \text{ K}$ ) and featureless thermal spectrum at peak, and a long-lived, fast-fading radio light curve similar to those of jet-powered long-duration GRBs. The host galaxy underwent a recent starforming episode and has a very high specific star-formation rate, similar to that of extreme SLSN and GRB hosts. Events like ZTF18abvkwla are rare: from one

year of the ZTF 1DC survey, we estimate that the rate is at least 2–3 orders of magnitude smaller than the CC SN rate.

Due to the lack of late-time photometry, we cannot conclude whether the late-time light curve was powered by the same mechanism as the peak or whether another mechanism such as nickel decay became dominant, and we have only tentative evidence for color evolution (cooling) over time. Furthermore, we cannot determine whether this source developed supernova features and whether it most closely resembles a Ic-BL like iPTF16asu, a continuum with emission lines like the Ibn iPTF15ul or the SN/TDE candidate AT2018cow, or neither.

Among the fast-luminous optical transients in Table 1, only AT2018cow and SN 2011kl had detected radio emission. ZTF18abvkwla thus adds to the very small number of events in the literature established to have fast-blue optical light curves, as well as a separate fast ejecta component that produces luminous radio emission. Interestingly, most of the well-studied transients in Table 1 are associated with a candidate engine-powered explosion. AT2018cow had a long-lived central engine that powered a fast ( $0.1c$ ) outflow. The Koala likely had a central engine that powered an even faster ( $> 0.38c$ ) outflow, perhaps a relativistic jet. iPTF16asu was a Ic-BL SN, and therefore by definition had faster ejecta velocities than ordinary core-collapse supernovae, although there was no evidence for a jet. SN 2011kl had a burst of high-energy emission (GRB 111209A) and an

associated luminous afterglow. Given the sensitivity of the luminosity to the shock speed (Equation 7), perhaps this apparent relationship between engine-driven supernovae and luminous fast-luminous optical transients should not be surprising.

At  $z = 0.27$ , ZTF18abvkwla was much more distant than AT2018cow ( $z = 0.0141$ ), but the lesson from §2.2 and §5 is that we should not be deterred by cosmological distances in pursuing X-ray and radio follow-up observations. The radio emission from ZTF18abvkwla would be easily detectable by the VLA out to  $z = 0.5$  (assuming  $5 \mu\text{Jy}$  RMS in half an hour of integration time) or even out to  $z = 0.8$  (when it would be  $30 \mu\text{Jy}$ ). Assuming a *Swift*/XRT sensitivity limit of  $4 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$ , the X-ray emission from ZTF18abvkwla may have been on the detection threshold. For a *Chandra* sensitivity limit an order of magnitude deeper, this may be on the detection threshold at  $z = 0.7$ . At these larger distances ( $z = 0.5, z = 0.7$ ) the optical  $g$ -band magnitude would be 21.1 and 22.3 respectively. This is out of reach for current surveys like ZTF, but standard for LSST. The false positives in such a search are lower-luminosity explosions (Type IIb, II, and Ibn SNe) and CVs. These can be ruled out via knowledge of the host redshift (and therefore intrinsic luminosity), so we emphasize the need for extensive and reliable galaxy-redshift catalogs.

The code used to produce the results described in this paper was written in Python and is available online in an open-source repository<sup>7</sup>.

## APPENDIX

### A. LIGHT-CURVE MEASUREMENTS

To construct Table 1, we used observed bands as close as possible to rest-frame  $g$ :  $g$ -band for  $z < 0.15$ ,  $r$ -band for  $0.15 < z < 0.45$ ,  $i$ -band for  $0.45 < z < 0.78$ , and  $z$ -band for  $0.78 < z < 1.0$ . We excluded transients with  $z > 1.0$ . We measured rise and fade times to 0.75 mag below peak by

<sup>7</sup> <https://github.com/annayqho/Koala>



linearly interpolating the single-filter light curve, and measured uncertainties using a Monte Carlo with 1000 realizations of the light curve. Additional notes on each transient are below.

For iPTF15ul ( $z = 0.066$ ; Hosseinzadeh et al. 2017) the uncertainty on the peak magnitude was dominated by the uncertainty from the host-galaxy extinction estimate. For AT2018cow ( $z = 0.0141$ ; Prentice et al. 2018; Perley et al. 2019a) we used the time between the last non-detection and the first detection as an upper limit on the rise time, although we note that interpolation would give 0.4 d, much shorter than 3 d. We also corrected for 0.287 mag of Galactic extinction, which was not applied in Table 3 of Perley et al. (2019a). For a lower limit, we used the  $o$ -band detection before peak (dominated by  $r$ -band flux at this epoch), corrected for 0.198 mag of Galactic extinction. We assumed  $g - r = -0.4$  mag and  $g - i = -0.7$  mag.

For SN 2011kl ( $z = 0.677$ ) we used column  $M_{4556}$  in Table 2 of Kann et al. (2019). These values are corrected for rest-frame extinction, and the contributions from the GRB afterglow and host galaxy have been subtracted. For SNLS04D4ec ( $z = 0.593$ ), SNLS05D2bk ( $z = 0.699$ ), and SNLS06D1hc ( $z = 0.555$ ) we used the  $i$ -band light curve from Arcavi et al. (2016) and corrected for Milky Way extinction.

For Dougie ( $z = 0.19$ ; Vinkó et al. 2015) we added an additional 0.1 mag in quadrature to account for the zero-point uncertainty, and corrected for 0.031 mag of Milky Way extinction. For iPTF16asu ( $z = 0.187$ ; Whitesides et al. 2017) we could not measure the rise or peak magnitude in rest-frame  $g$  because observations in the appropriate filter ( $r$ ) began only 3 days after peak. We estimated an upper limit to the peak magnitude by assuming that the  $g - r$  color at peak was identical to the  $g - r$  color during the first  $r$ -band measurement. We used the first  $r$ -band measurement as a lower limit. For the time from half-max to max, we used the observed  $g$ -band light curve instead. We obtained the  $i$ -band light curve of DES16X1eho ( $z = 0.76$ ; Pursiainen et al. 2018) from M. Pursiainen (private communication).

## ACKNOWLEDGMENTS

A.Y.Q.H would like to thank the NRAO staff for their help with data calibration and imaging, particularly Steve Myers, Aaron Lawson, Drew Medlin, and Emmanuel Momjian. She is grateful for their support and hospitality during her visit to Socorro. She thanks Gregg Hallinan and Brad Cenko for their advice on reducing the radio and X-ray data, respectively, Jochen Greiner and Iair Arcavi for their assistance in obtaining the afterglow-subtracted light curve of SN 2011kl, Miika Pursiainen for sharing light curves of DES fast-luminous transients, Griffin Hosseinzadeh for useful discussions about iPTF15ul, Jesper Sollerman and Steve Schulze for carefully reading the manuscript, and Tony Piro and Ben Margalit for other productive conversations. This work made use of the IPN master burst list ([ssl.berkeley.edu/ipn3/masterli.html](http://ssl.berkeley.edu/ipn3/masterli.html)) maintained by Kevin Hurley. We would like to thank Raffaella Margutti for pointing out a typo in an earlier version of this paper, and to the anonymous referee for detailed comments that greatly improved the flow and clarity of the paper.

A.Y.Q.H. would like to acknowledge the support of a National Science Foundation Graduate Research Fellowship under Grant No. DGE-1144469, the GROWTH project funded by the National Science Foundation under PIRE Grant No. 1545949, and the Heising-Simons Foundation. P.C. acknowledges the support from the Department of Science and Technology via SwarnaJayanti Fellowship awards (DST/SJF/PSA-01/2014-15). A.H. is grateful for the support by grants from the Israel

Science Foundation, the US-Israel Binational Science Foundation, and the I-CORE Program of the Planning and Budgeting Committee and the Israel Science Foundation. This research was funded in part by a grant from the Heising-Simons Foundation and a grant from the Gordon and Betty Moore Foundation through Grant GBMF5076, and benefited from interactions with Daniel Kasen and David Khatami also funded by that grant. A.A.M. is funded by the Large Synoptic Survey Telescope Corporation, the Brinson Foundation, and the Moore Foundation in support of the LSSTC Data Science Fellowship Program; he also receives support as a CIERA Fellow by the CIERA Post-doctoral Fellowship Program (Center for Interdisciplinary Exploration and Research in Astrophysics, Northwestern University).

Based on observations obtained with the Samuel Oschin Telescope 48-inch and the 60-inch Telescope at the Palomar Observatory as part of the Zwicky Transient Facility project. ZTF is supported by the National Science Foundation under Grant No. AST-1440341 and a collaboration including Caltech, IPAC, the Weizmann Institute for Science, the Oskar Klein Center at Stockholm University, the University of Maryland, the University of Washington, Deutsches Elektronen-Synchrotron and Humboldt University, Los Alamos National Laboratories, the TANGO Consortium of Taiwan, the University of Wisconsin at Milwaukee, and Lawrence Berkeley National Laboratories. Operations are conducted by COO, IPAC, and UW. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc. We thank the staff of the GMRT that made these observations possible. GMRT is run by the National Centre for Radio Astrophysics of the Tata Institute of Fundamental Research. Some of the data presented herein were obtained at the W. M. Keck Observatory, which is operated as a scientific partnership among the California Institute of Technology, the University of California and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W. M. Keck Foundation. The authors wish to recognize and acknowledge the very significant cultural role and reverence that the summit of Maunakea has always had within the indigenous Hawaiian community. We are most fortunate to have the opportunity to conduct observations from this mountain.

*Facilities:* Keck:I (LRIS), Hale (DBSP), EVLA, GMRT

*Software:* `Astropy` (Astropy Collaboration et al. 2013, 2018), `IPython` (Pérez & Granger 2007), `matplotlib` (Hunter 2007), `numpy` (Oliphant 2006), `scipy` (Virtanen et al. 2019), `extinction` (Barbary et al. 2016), `LPipe` (Perley 2019), `PYRAF-DBSP` (Bellm & Sesar 2016), `IDLAstro`

## REFERENCES

- Abolfathi, B., Aguado, D. S., Aguilar, G., et al. 2018, *ApJS*, 235, 42
- Alexander, K. D., Berger, E., Guillochon, J., Zauderer, B. A., & Williams, P. K. G. 2016, *ApJL*, 819, L25
- Aptekar, R. L., Frederiks, D. D., Golenetskii, S. V., et al. 1995, *SSRv*, 71, 265
- Arcavi, I., Wolf, W. M., Howell, D. A., et al. 2016, *ApJ*, 819, 35
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, 558, A33
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, *AJ*, 156, 123
- Barbary, K. 2016, `extinction`, v0.3.0, Zenodo, doi: 10.5281/zenodo.804967, <https://github.com/kbarbary/extinction>
- Bellm, E. C., & Sesar, B. 2016, *Astrophysics Source Code Library*, ascl:1602.002

- Bellm, E. C., Kulkarni, S. R., Graham, M. J., et al. 2019, *PASP*, 131, 018002
- Bellm, E. C., Kulkarni, S. R., Barlow, T., et al. 2019, *PASP*, 131, 068003
- Berger, E., Kulkarni, S. R., Pooley, G., et al. 2003, *Nature*, 426, 154
- Berger, E., Zauderer, A., Pooley, G. G., et al. 2012, *ApJ*, 748, 36
- Narayana Bhat, P., Meegan, C. A., von Kienlin, A., et al. 2016, *ApJS*, 223, 28
- Bianco, F. B., Modjaz, M., Oh, S. M., et al. 2016, *Astronomy and Computing*, 16, 54
- Bietenholz, M. F., Margutti, R., Coppejans, D., et al. 2020, *MNRAS*, 491, 4735
- Bock, D. C.-J., Bolatto, A. D., Hawkins, D. W., et al. 2006, *Proc. SPIE*, 626713
- Bruzual, G., & Charlot, S. 2003, *MNRAS*, 344, 1000
- Campana, S., Mangano, V., Blustin, A. J., et al. 2006, *Nature*, 442, 1008
- Cano, Z., Wang, S.-Q., Dai, Z.-G., & Wu, X.-F. 2017, *Advances in Astronomy*, 2017, 8929054
- Cenko, S. B., Kulkarni, S. R., Horesh, A., et al. 2013, *ApJ*, 769, 130
- Cenko, S. B., Urban, A. L., Perley, D. A., et al. 2015, *ApJL*, 803, L24
- Chevalier, R. A. 1998, *ApJ*, 499, 810
- Chevalier, R. A., & Irwin, C. M. 2011, *ApJL*, 729, L6
- Condon, J. J. 1992, *ARA&A*, 30, 575
- Coppejans, D. L., Margutti, R., Terreran, G., et al. 2020, arXiv e-prints, arXiv:2003.10503
- Dabringhausen, J., Kroupa, P., & Baumgardt, H. 2009, *MNRAS*, 394, 1529
- De Colle, F., & Lu, W. 2019, arXiv e-prints, arXiv:1911.01442
- Dey, A., Schlegel, D. J., Lang, D., et al. 2019, *AJ*, 157, 168
- Drout, M. R., Chornock, R., Soderberg, A. M., et al. 2014, *ApJ*, 794, 23
- Duev, D. A., Mahabal, A., Masci, F. J., et al. 2019, *MNRAS*, 489, 3582
- Eftekhari, T., Berger, E., Zauderer, B. A., et al. 2018, *ApJ*, 854, 86
- Ferrero, P., Kann, D. A., Zeh, A., et al. 2006, *A&A*, 457, 857
- Firth, R. E., Sullivan, M., Gal-Yam, A., et al. 2015, *MNRAS*, 446, 3895
- Fitzpatrick, E. L. 1999, *PASP*, 111, 63
- Fox, O. D., & Smith, N. 2019, *MNRAS*, 488, 3772
- Frail, D. A., Kulkarni, S. R., Sari, R., et al. 2001, *ApJL*, 562, L55
- Fremming, C., Ko, H., Dugas, A., et al. 2019, *ApJL*, 878, L5
- Fruchter, A. S., Levan, A. J., Strolger, L., et al. 2006, *Nature*, 441, 463
- Gal-Yam, A. 2019, *ARA&A*, 57, 305
- Gezari, S., Heckman, T., Cenko, S. B., et al. 2009, *ApJ*, 698, 1367
- Gezari, S. 2012, *European Physical Journal Web of Conferences*, 03001
- Giannios, D., & Metzger, B. D. 2011, *MNRAS*, 416, 2102
- Graham, J. F., & Schady, P. 2016, *ApJ*, 823, 154
- Graham, M. J., Kulkarni, S. R., Bellm, E. C., et al. 2019, *PASP*, 131, 078001
- Greiner, J., Mazzali, P. A., Kann, D. A., et al. 2015, *Nature*, 523, 189
- Greiner, J., Michałowski, M. J., Klose, S., et al. 2016, *A&A*, 593, A17
- Gruber, D., Goldstein, A., Weller von Ahlefeld, V., et al. 2014, *ApJS*, 211, 12
- Guetta, D., Piran, T., & Waxman, E. 2005, *ApJ*, 619, 412
- Gupta Y., et al., 2017, *Current Science*, 113, 707
- Hancock, P. J., Murphy, T., Gaensler, B., et al. 2012, *GRB Coordinates Network* 12804, 1
- Ho, A. Y. Q., Phinney, E. S., Ravi, V., et al. 2019, *ApJ*, 871, 73
- Ho, A. Y. Q., Goldstein, D. A., Schulze, S., et al. 2019, *ApJ*, 887, 169
- Högbom, J. A. 1974, *A&AS*, 15, 417
- Horesh, A., Stockdale, C., Fox, D. B., et al. 2013, *MNRAS*, 436, 1258
- Hosseinzadeh, G., Arcavi, I., Valenti, S., et al. 2017, *ApJ*, 836, 158
- Huang, K., Shimoda, J., Urata, Y., et al. 2019, *ApJL*, 878, L25
- Hunter, J. D. 2007, *CISE*, 9(3), 90
- Hurley, K., Golenetskii, S., Aptekar, R., et al. 2010, *American Institute of Physics Conference Series*, 330
- Hurley, K., Svinikin, D. S., Aptekar, R. L., et al. 2016, *ApJL*, 829, L12
- Irwin, J. A., Henriksen, R. N., Krause, M., et al. 2015, *ApJ*, 809, 172
- Kann, D. A., Klose, S., Zhang, B., et al. 2010, *ApJ*, 720, 1513
- Kann, D. A., Schady, P., Olivares, E. F., et al. 2018, *A&A*, 617, A122

- Kann, D. A., Schady, P., Olivares E., F., et al. 2019, *A&A*, 624, A143
- Kasen, D. 2017, *Handbook of Supernovae*, ISBN 978-3-319-21845-8. Springer International Publishing AG, 2017, p. 939, 939
- Kasliwal, M. M., Kulkarni, S. R., Gal-Yam, A., et al. 2012, *ApJ*, 755, 161
- Kasliwal, M. M., Cannella, C., Bagdasaryan, A., et al. 2019, *PASP*, 131, 038003
- Katz, J. I. 2016, *Modern Physics Letters A*, 31, 1630013
- Kelly, P. L., & Kirshner, R. P. 2012, *ApJ*, 759, 107
- Kennicutt, R. C., Tamblyn, P. C., & Congdon, C. E. 1994, *ApJ*, 435, 22
- Komossa, S. 2015, *Journal of High Energy Astrophysics*, 7, 148
- Krauss, M. I., Soderberg, A. M., Chomiuk, L., et al. 2012, *ApJL*, 750, L40
- Krühler, T., Malesani, D., Fynbo, J. P. U., et al. 2015, *A&A*, 581, A125
- Kuin, N. P. M., Wu, K., Oates, S., et al. 2019, *MNRAS*, 487, 2505
- Kulkarni, S. R., Frail, D. A., Wieringa, M. H., et al. 1998, *Nature*, 395, 663
- Lacy, M., Baum, S. A., Chandler, C. J., et al. 2019, arXiv e-prints, arXiv:1907.01981
- Law, N. M., Kulkarni, S. R., Dekany, R. G., et al. 2009, *PASP*, 121, 1395
- Lee, J. C., Kennicutt, R. C., Funes, S. J. J. G., et al. 2009, *ApJ*, 692, 1305
- Lei, W.-H., Yuan, Q., Zhang, B., et al. 2016, *ApJ*, 816, 20
- Li, W., Chornock, R., Leaman, J., et al. 2011, *MNRAS*, 412, 1473
- Liang, E., Zhang, B., Virgili, F., et al. 2007, *ApJ*, 662, 1111
- Lien, A., Sakamoto, T., Gehrels, N., et al. 2014, *ApJ*, 783, 24
- Lu, W., & Bonnerot, C. 2020, *MNRAS*, 492, 686
- Lunnan, R., Chornock, R., Berger, E., et al. 2014, *ApJ*, 787, 138
- Leloudas, G., Schulze, S., Krühler, T., et al. 2015, *MNRAS*, 449, 917
- Lyman, J. D., Bersier, D., James, P. A., et al. 2016, *MNRAS*, 457, 328
- Lyutikov, M., & Toonen, S. 2019, *MNRAS*, 487, 5618
- Mahabal, A., Rebbapragada, U., Walters, R., et al. 2019, *PASP*, 131, 038002
- Margutti, R., Metzger, B. D., Chornock, R., et al. 2019, *ApJ*, 872, 18
- Masci, F. J., Laher, R. R., Rusholme, B., et al. 2019, *PASP*, 131, 018003
- Matheson, T., Filippenko, A. V., Chornock, R., et al. 2000, *AJ*, 119, 2303
- McCarthy, J. K., Cohen, J. G., Butcher, B., et al. 1998, *Proc. SPIE*, 81
- McMullin, J. P., Waters, B., Schiebel, D., Young, W., & Golap, K. 2007, *Astronomical Data Analysis Software and Systems XVI*, 376, 127
- Miller, A. A., Yao, Y., Bulla, M., et al. 2020, arXiv e-prints, arXiv:2001.00598
- Mirabal, N., Halpern, J. P., An, D., et al. 2006, *ApJL*, 643, L99
- Modjaz, M., Bianco, F. B., Siwek, M., et al. 2019, arXiv e-prints, arXiv:1901.00872
- Mohan, P., An, T., & Yang, J. 2020, *ApJL*, 888, L24
- Murphy, E. J., Condon, J. J., Schinnerer, E., et al. 2011, *ApJ*, 737, 67
- Nakar, E., & Piro, A. L. 2014, *ApJ*, 788, 193
- Nakar, E. 2015, *ApJ*, 807, 172
- Nakar, E., & Piran, T. 2017, *ApJ*, 834, 28
- Ofek, E. O., Rabinak, I., Neill, J. D., et al. 2010, *ApJ*, 724, 1396
- Oke, J. B., & Gunn, J. E. 1982, *PASP*, 94, 586
- Oke, J. B., & Gunn, J. E. 1983, *ApJ*, 266, 713
- Oke, J. B., Cohen, J. G., Carr, M., et al. 1995, *PASP*, 107, 375
- Oliphant, T. E. 2006, *A guide to NumPy*, Trelgol Publishing
- Patterson, M. T., Bellm, E. C., Rusholme, B., et al. 2019, *PASP*, 131, 018001
- Pérez, F. & Granger, B. E. 2007, *CISE*, 9(3), 29
- Perley, R. A., Chandler, C. J., Butler, B. J., et al. 2011, *ApJL*, 739, L1
- Perley, D. A., Levan, A. J., Tanvir, N. R., et al. 2013, *ApJ*, 778, 128
- Perley, D. A., Cenko, S. B., Corsi, A., et al. 2014, *ApJ*, 781, 37
- Perley, D. A., Quimby, R. M., Yan, L., et al. 2016, *ApJ*, 830, 13
- Perley, D. A., Mazzali, P. A., Yan, L., et al. 2019, *MNRAS*, 484, 1031
- Perley, D. A. 2019, *PASP*, 131, 084503
- Pian, E., Mazzali, P. A., Masetti, N., et al. 2006, *Nature*, 442, 1011
- Pietka, M., Fender, R. P., & Keane, E. F. 2015, *MNRAS*, 446, 3687

- Piro, A. L. 2015, *ApJL*, 808, L51
- Planck Collaboration, Ade, P. A. R., Aghanim, N., et al. 2016, *A&A*, 594, A13
- Prentice, S. J., Maguire, K., Smartt, S. J., et al. 2018, *ApJL*, 865, L3
- Pursiainen, M., Childress, M., Smith, M., et al. 2018, *MNRAS*, 481, 894
- Quimby, R. M., Kulkarni, S. R., Kasliwal, M. M., et al. 2011, *Nature*, 474, 487
- Rau, A., Kulkarni, S. R., Law, N. M., et al. 2009, *PASP*, 121, 1334
- Readhead, A. C. S. 1994, *ApJ*, 426, 51
- Rest, A., Garnavich, P. M., Khatami, D., et al. 2018, *Nature Astronomy*, 2, 307
- Rivera Sandoval, L. E., Maccarone, T. J., Corsi, A., et al. 2018, *MNRAS*, 480, L146
- Salas, P., Bauer, F. E., Stockdale, C., & Prieto, J. L. 2013, *MNRAS*, 428, 1207
- Schlafly, E. F., & Finkbeiner, D. P. 2011, *ApJ*, 737, 103
- Schulze, S., Chapman, R., Hjorth, J., et al. 2015, *ApJ*, 808, 73
- Schulze, S., Krühler, T., Leloudas, G., et al. 2018, *MNRAS*, 473, 1258
- Scott, M. A., & Readhead, A. C. S. 1977, *MNRAS*, 180, 539
- Shivvers, I., Modjaz, M., Zheng, W., et al. 2017, *PASP*, 129, 054201
- Slysh, V. I. 1990, *Soviet Astronomy Letters*, 16, 339
- Smith, N., & McCray, R. 2007, *ApJL*, 671, L17
- Smith, N. 2014, *ARA&A*, 52, 487
- Soderberg, A. M., Kulkarni, S. R., Berger, E., et al. 2004, *Nature*, 430, 648
- Soderberg, A. M., Kulkarni, S. R., Berger, E., et al. 2005, *ApJ*, 621, 908
- Soderberg, A. M., Kulkarni, S. R., Nakar, E., et al. 2006, *Nature*, 442, 1014
- Soderberg, A. M., Chakraborti, S., Pignata, G., et al. 2010, *Nature*, 463, 513
- Sollerman, J., Jaunsen, A. O., Fynbo, J. P. U., et al. 2006, *A&A*, 454, 503
- Stratta, G., Gendre, B., Atteia, J. L., et al. 2013, *ApJ*, 779, 66
- Strubbe, L. E., & Quataert, E. 2009, *MNRAS*, 400, 2070
- Swarup, G., Ananthakrishnan, S., Kapahi, V. K., et al. 1991, *Current Science*, 60, 95
- Tachibana, Y., & Miller, A. A. 2018, *PASP*, 130, 128001
- Taggart, K., & Perley, D. 2019, arXiv e-prints, arXiv:1911.09112
- Tendulkar, S. P., Bassa, C. G., Cordes, J. M., et al. 2017, *ApJL*, 834, L7
- Thöne, C. C., de Ugarte Postigo, A., Fryer, C. L., et al. 2011, *Nature*, 480, 72
- Tremonti, C. A., Heckman, T. M., Kauffmann, G., et al. 2004, *ApJ*, 613, 898
- van den Heuvel, E. P. J., & Portegies Zwart, S. F. 2013, *ApJ*, 779, 114
- van der Horst, A. J., Paragi, Z., de Bruyn, A. G., et al. 2014, *MNRAS*, 444, 3151
- Vergani, S. D., Salvaterra, R., Japelj, J., et al. 2015, *A&A*, 581, A102
- Vinkó, J., Yuan, F., Quimby, R. M., et al. 2015, *ApJ*, 798, 12
- Virtanen, P., Gommers, R., Oliphant, T. E., et al. 2019, arXiv e-prints, arXiv:1907.10121
- von Kienlin, A., Meegan, C. A., Paciesas, W. S., et al. 2014, *ApJS*, 211, 13
- Wang, L. J., Wang, X. F., Cano, Z., et al. 2019, *MNRAS*, 489, 1110
- Watson, C., Henden, A. A., & Price, A. 2017, *VizieR Online Data Catalog*, B/vsx
- Waxman, E., & Katz, B. 2017, *Handbook of Supernovae*, 967
- Weiler, K. W., Sramek, R. A., Panagia, N., van der Hulst, J. M., & Salvati, M. 1986, *ApJ*, 301, 790
- Weiler, K. W., van Dyk, S. D., Panagia, N., Sramek, R. A., & Discenna, J. L. 1991, *ApJ*, 380, 161
- Weiler, K. W., Williams, C. L., Panagia, N., et al. 2007, *ApJ*, 671, 1959
- Whitesides, L., Lunnan, R., Kasliwal, M. M., et al. 2017, *ApJ*, 851, 107
- Yao, Y., Miller, A. A., Kulkarni, S. R., et al. 2019, arXiv e-prints, arXiv:1910.02967
- Zackay, B., Ofek, E. O., & Gal-Yam, A. 2016, *ApJ*, 830, 27
- Zauderer, B. A., Berger, E., Soderberg, A. M., et al. 2011, *Nature*, 476, 425
- Zauderer, B. A., Berger, E., Margutti, R., et al. 2013, *ApJ*, 767, 152
- Zwart, J. T. L., Barker, R. W., Biddulph, P., et al. 2008, *MNRAS*, 391, 1545