



LJMU Research Online

Thomas, C, Jones, H, Whitworth-Turner, C and Louis, J

A Sleep Analysis of Elite Female Soccer Players During a Competition Week

<http://researchonline.ljmu.ac.uk/id/eprint/14431/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Thomas, C, Jones, H, Whitworth-Turner, C and Louis, J (2021) A Sleep Analysis of Elite Female Soccer Players During a Competition Week. International Journal of Sports Physiology and Performance. ISSN 1555-0265

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

1 **Title:** A Sleep Analysis of Elite Female Soccer Players During a Competition Week

2 **Submission type:** Original investigation

3 **Authors:** Craig Thomas¹, Helen Jones¹, Craig Whitworth-Turner² and Julien Louis^{1*}

4 **Institutional affiliations:** ¹Research Institute for Sport and Exercise Sciences (RISES),
5 Liverpool John Moores University, Liverpool, UK.

6 ²Philadelphia 76ers training complex, 1-99 S Front St, Camden, New Jersey 08103, USA.

7 ***Corresponding author**

8 Dr Julien Louis

9 Research Institute for Sport and Exercise Sciences

10 Liverpool John Moores University

11 Tom Reilly Building

12 Byrom St. Campus

13 Liverpool, L3 3AF, UK

14 E-mail: J.B.Louis@ljmu.ac.uk

15 Tel: 0151 904 6285

16 ORCID ref: orcid.org/0000-0002-9109-0958

17 **Running head:** Sleep of Elite Female Soccer Players

18 **Abstract word count:** 250

19 **Text-only word count:** 3,673

20 **Number of figures and tables:** 0 figures and 4 tables

21

22

23

24

25

26 **Abstract**

27 **Purpose:** (1) To compare the sleep of female players from a professional soccer team to non-
28 athlete controls across an in-season week and (2) to compare the sleep of core and fringe players
29 from the same team on the night after a match to training nights.

30 **Methods:** Using an observational design, 18 professional female soccer players and 18 female
31 non-athlete controls were monitored for their sleep via wristwatch actigraphy across one week.
32 Independent sample *t*-tests and Mann Whitney U tests were performed to compare sleep
33 between groups whilst an ANOVA compared sleep on training nights to the night after a match.

34 **Results:** Soccer players had significantly greater sleep duration than non-athlete controls (+38
35 min; $P = 0.009$; $d: 0.92$), which may have resulted from an earlier bedtime (-00:31 h: min; $P =$
36 0.047 ; $d: 0.70$). The soccer players also had less intraindividual variation in bedtime than non-
37 athletes (-00:08 h: min; $P = 0.023$; $r: 0.38$). Despite this, sleep onset latency was significantly
38 longer within soccer players (+8 min; $P = 0.032$; $d: 0.78$). On the night after a match, sleep
39 duration of core players was significantly lower than on training nights (-49 min; $P = 0.010$; $d:$
40 0.77). In fringe players, there was no significant difference between nights for any sleep
41 characteristic.

42 **Conclusions:** During the in-season period, sleep duration of professional female soccer players
43 is greater than non-athlete controls. However, the night after a match challenges the sleep of
44 players with more match involvement and warrant priority of sleep hygiene strategies.

45 **Keywords:** wristwatch actigraphy, team sports, training, in-season, recovery

46

47

48

49

50

51

52

53

54

55

56

57

58

59 **Introduction**

60 Sleep is considered the single best recovery strategy available for elite athletes following
61 training and competition.¹ This is attributed to the bodily processes that occur during sleep
62 thought to serve physical and psychological restoration.² Individuals are recommended to sleep
63 at least 7h per night with 85% or more sleep efficiency (time asleep as a percentage of time in
64 bed [sleep quality]).^{3,4} During the competitive soccer season, however, sleep may be impacted
65 by numerous factors such as scheduling, training stress, arousal and travel.⁵ For example, a
66 recent investigation of different match situations (i.e. home, away, day and night) revealed that
67 sleep duration (5:49 h: min) and sleep efficiency (79 %) were lower on matchdays compared
68 to training days (6:36 h: min and 85% for sleep duration and sleep efficiency respectively) in
69 players from a professional men's Portuguese team.⁶ There has also been no alteration to sleep
70 after a match (i.e. day and night) and evening high intensity training compared with a rest day
71 in male youth soccer players^{7,8}, though the sleep duration in these studies was highly variable
72 (5.7–7.5 h). Such findings coupled with other studies in semi-professional and professional
73 players showing a reduction in sleep duration after home, away and night-time fixtures (4.5-6
74 h)⁹⁻¹¹ highlight that although matchdays tend to impact sleep more, male soccer players may
75 not meet current sleep guidelines during the season.

76 Data concerning the sleep of female soccer players during the competitive season is somewhat
77 more limited. Sleep duration was lower following night-time training sessions (21:00 h) than
78 on rest day nights (7:17 vs. 7:51 h: min) in players from a semi-professional female Portuguese
79 team.¹² This was corroborated in a follow up study, as sleep duration after night-time training
80 was markedly lower (7:09 h: min) compared to matchdays (i.e. home and away matches; 8:44
81 h: min) and rest days (8:35 h: min), also in semi-professional female players.¹³ This was despite
82 sleep efficiency falling within guidelines on all days (88-91 %).¹³ Other than after night-time
83 training, these findings may indicate female soccer players attain optimal sleep during the
84 soccer season. That said, the players in the aforementioned studies were not full-time and thus,
85 would not have the same schedule or training load as professional players. Besides, in the study
86 by Costa et al¹³, it is unknown if matchday sleep would have differed from training nights
87 according to players match involvement. Fullagar et al¹⁰ showed that male soccer players who
88 started a night match had a smaller reduction in sleep efficiency from pre-match days than non-
89 players (-3.9 vs. -20.7 %). Consequently, it may be appropriate to compare the sleep
90 characteristics of professional female players between training and matchday nights based on
91 playing time.

92 There is also no study comparing the sleep characteristics of female soccer players with a non-
93 athlete, control group. Elite athletes are thought to have a greater sleep need than their non-
94 athlete counterparts due to the physical demands associated with training and competition.¹⁴
95 Within soccer players, studies employing objective measures of sleep have produced
96 conflicting findings on the comparison with non-athletes.^{15,16} This may be explained by the
97 length of study, as Whitworth-Turner et al¹⁶ is the only investigation to have monitored sleep
98 over consecutive nights, thereby capturing the variation in scheduled activities. In this study,
99 male youth soccer players had a longer sleep duration (486 vs. 422 min), but a lower sleep
100 efficiency than non-athletes (93 vs. 96 %), as measured using a wireless system. These findings
101 are in accordance with suggestions that elite athletes may have reduced sleep quality compared
102 with non-athlete controls.^{17,18} It would therefore seem important to first compare professional
103 female soccer players and non-athlete controls to provide an insight into the sleep differences
104 between these two populations. The primary aim of this study was to compare the sleep
105 characteristics of female players from a professional soccer team to non-athlete controls across
106 an in-season training week. Additionally, as it is unclear whether matchday alters sleep in

107 professional female players, a secondary aim was to compare the sleep of core and fringe
108 players on the night after a match to training nights. This type of analysis could further the
109 understanding of how professional female soccer players sleep during the season and where
110 their sleep may be challenged.

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140 **Methods**

141 **Subjects**

142 Eighteen female soccer players (age = 23.2±4.5yrs) and 18 female non-athlete controls (age =
143 24.9±2.8yrs) were recruited. The soccer players were contracted to a professional team in
144 England (competing in the FA Women's Super League, the top tier of English women's
145 football) and had playing experience of 6.7±4.7yrs. In the group, there were two goalkeepers,
146 five defenders, six midfielders and five forwards. The non-athlete controls were full time
147 university students and physically active. Inclusion criteria consisted of the following: 18-40
148 years of age; non-smoker for at least six months; not a night-shift worker and not engaged in
149 competitive sport. None of the participants were using sleep aids (e.g. medication) nor had they
150 travelled across different time zones in the month prior to participation. Before the study
151 commenced, written informed consent was obtained and participants filled in the morningness-
152 eveningness questionnaire for the assessment of their chronotype.¹⁹ The mean score for soccer
153 players and non-athletes was 58±6 (intermediate) and 59±8 (moderate morning) respectively.
154 This study was granted ethical approval from the Local University research ethics committee.

155 **Design**

156 Using an observational, descriptive study design, sleep characteristics were monitored across
157 a week via wristwatch actigraphy. Actigraphy was implemented as this has shown to have a
158 good overall agreement with polysomnography (gold standard measurement of sleep) and is
159 considered a valid alternative in field settings (90-91 %).²⁰ Female soccer players were
160 monitored across one of four in-season weeks (early to mid-season) that had similar training
161 and matchday schedules. This was important to prevent differences in sleep characteristics
162 between soccer players induced by the timing of training and competition in relation to the
163 amount of days that were recorded. Accordingly, from Tuesday to Saturday, players trained
164 and rested to prepare for matchday on Sunday, whilst on Monday they either rested or
165 participated in activities to aid post-match recovery (Table 1). All training sessions were
166 scheduled by the coaching staff at the club and the team played two home and two away
167 matches (travelled night before and on the day) across the four-week period. The non-athlete
168 controls were monitored during a week where they participated in their habitual routines,
169 consisting of typical lecture and study days. Rest days (i.e. days with no set schedule) were
170 excluded from data analysis in both groups. This was conducted to avoid unwanted variation
171 associated with non-work activities²¹ and this approach has previously been used within soccer
172 players.¹⁶ In addition, participants filled in a diary for the quantification of their internal
173 training load. Across the monitoring period, participants were encouraged not to alter their
174 sleep behaviour and they followed their usual dietary intake, which included caffeinated
175 products.

176 ***INSERT TABLE 1 HERE***

177 **Methodology**

178 **Sleep Assessment**

179 To monitor sleep characteristics, an actiwatch (Actiwatch 4, Cambridge Technology Ltd, UK)
180 was provided and set to an epoch length of 1 min at a medium sensitivity.²² On each night,
181 participants were asked to wear the actiwatch on their non-dominant wrist at least 30 min before
182 they retired to bed and then press the marker button upon their bedtime (lights out). The marker
183 button was used again the following morning to indicate their final awakening time (lights on)
184 before they were instructed to fill in the Consensus Sleep Diary²³ within an hour of getting out

185 of bed. The Consensus Sleep Diary asks questions relating to bedtime, sleep onset latency,
186 number of awakenings, final awakening time, get up time and sleep quality. Using the
187 actiwatch markers and the information from the Consensus Sleep Diary; bedtime, sleep onset,
188 final awakening time and get up time were determined so that sleep behaviour could be
189 automatically calculated using the appropriate actiwatch software (Actiwatch activity and sleep
190 analysis version 5.24, Cambridge Neurotechnology Ltd, UK). From the actiwatch analysis, the
191 following characteristics were chosen to describe sleep: time in bed (min), bedtime (h: min),
192 time of final awakening (h: min), sleep onset latency (min), sleep duration (min), sleep
193 efficiency (%) and wake after sleep onset (min) (Table 2).

194 ***INSERT TABLE 2 HERE***

195 **Quantification of Training Load**

196 After exercise, the duration and session rating of perceived exertion (sRPE) were recorded in
197 the diary to calculate internal training load (CR10 scale).²⁴ The soccer players were asked to
198 fill in the diary within an hour of finishing each training session and as soon as possible after a
199 match to account for post-match activities. The non-athlete controls were asked to fill in the
200 diary on days when they performed exercise and was also to be completed within an hour after
201 their sessions. From this information, daily training load (arbitrary units [AU]) was calculated
202 for both groups by multiplying the session duration in minutes by the sRPE.²⁴

203 **Statistical Analysis**

204 Statistical Package for the Social Sciences (SPSS v26) was used for data analysis. To compare
205 average sleep between soccer players and non-athletes, the mean of the characteristics across
206 five nights was calculated for each individual. The data of both groups were then assessed for
207 normality using the Shapiro-Wilk statistic and independent sample *t*-tests were conducted to
208 assess group differences. When assessing the average daily training load between groups, the
209 same process was used. Intraindividual variation of sleep characteristics was also calculated
210 for groups across five nights by obtaining the mean of individual standard deviations.^{25,26} For
211 intraindividual sleep characteristics that met normality (i.e. time in bed), independent sample
212 *t*-tests were performed, but for those that violated normality (i.e. bedtime, time of final
213 awakening, sleep duration, sleep efficiency, sleep onset latency and wake after sleep onset),
214 the Mann Whitney U test was chosen. A Linear Mixed Model ANOVA was used to assess if
215 sleep on training nights and the night after a match differed within core (involved in ≥ 60
216 minutes of a match) and fringe (unused substitute or involved in ≤ 45 minutes of a match)
217 players. The average of the characteristics from training days was calculated for each player
218 and assessed against the night after a match. Within the models, type of night (i.e. training or
219 match) was inputted as the fixed effect and individual player identification was included as the
220 random effect. Data are presented as mean \pm standard deviation, mean difference and 95%
221 confidence intervals (CI) or the median and interquartile range (IQR) for the reporting of non-
222 parametric tests. Cohens *d* was calculated for effect size where normality was met and was
223 subsequently assessed using the following thresholds: < 0.20 = trivial effect; $0.20-0.60$ = small
224 effect; $> 0.60-1.20$ = moderate effect; $> 1.20-2.00$ = large effect; $> 2.00-4.00$ = very large and
225 > 4.00 = extremely large effect.²⁷ In circumstances where normality was not displayed, the *r*
226 (uses *z* score from Mann Whitney U test) was used for effect size.²⁸ The *r* was interpreted from
227 Cohen's criteria, where: 0.1 = small effect; 0.3 = moderate effect and 0.5 = large effect.²⁹
228 Statistical significance was set at level $P < 0.05$.

229

230

231 **Results**

232 **Soccer Players vs. Non-Athlete Controls**

233 The soccer players had a significantly greater daily training load compared with non-athlete
234 controls (738±169 vs. 364±167 AU; $P < 0.001$; $d: 2.23$ [very large]; 95% CI: 260-488 AU).
235 Average sleep characteristics for soccer players and non-athletes are shown in Table 3. Soccer
236 players spent significantly more time in bed than non-athlete controls (+55 min; $P = 0.001$; $d:$
237 1.22 [large effect]). Bedtime was significantly earlier in soccer players (-00:31 h: min; $P =$
238 0.047; $d: 0.70$ [moderate effect]) but final awakening was similar between groups ($P = 0.167$;
239 $d: 0.47$ [small effect]). Soccer players had significantly greater sleep duration than non-athlete
240 controls (+38 min; $P = 0.009$; $d: 0.92$ [moderate effect]). Sleep onset latency of soccer players
241 was also significantly longer than non-athlete controls (+8 min; $P = 0.032$; $d: 0.78$ [moderate
242 effect]). There were no significant differences between groups for sleep efficiency ($P = 0.362$;
243 $d: 0.39$ [small effect]) and wake after sleep onset ($P = 0.733$; $d: 0.14$ [trivial effect]).

244 Intraindividual variation of sleep variables for both groups are also in Table 3. Soccer players
245 had significantly less intraindividual variation in bedtime compared with non-athlete controls
246 (-00:08 h: min; $P = 0.023$; $r: 0.38$ [moderate effect]). A marginal non-significant difference
247 between groups was shown for intraindividual variation of final awakening ($P = 0.050$; $r: 0.33$
248 [moderate effect]). There were no significant differences between groups for intraindividual
249 variation of time in bed ($P = 0.226$; $d: 0.44$ [small effect]), sleep duration ($P = 0.496$; $r: 0.11$
250 [small effect]), sleep efficiency ($P = 0.649$; $r: 0.08$ [trivial effect]), sleep onset latency ($P =$
251 0.178; $r: 0.23$ [small effect]) and wake after sleep onset ($P = 0.326$; $r: 0.16$ [small effect]).

252 ***INSERT TABLE 3 HERE***

253 **Night After Match vs. Training Nights**

254 The time played by core and fringe players on matchday was 84±11 min and 14±18 min
255 respectively. All core players started whilst one fringe player started, and three others were
256 used as substitutes. Table 4 displays sleep characteristics of these groups on the night after a
257 match and on training nights.

258 In core players, bedtime was significantly later on the night after a match compared with
259 training nights (+00:37 h: min; $P = 0.032$; $d: 0.76$ [moderate effect]) but there was no
260 significant difference for final awakening ($P = 0.359$; $d: 0.37$ [small effect]). Sleep duration
261 was significantly lower on the night after a match compared with training nights (-49 min; $P =$
262 0.010; $d: 0.77$ [moderate effect]). There were marginal non-significant differences for time in
263 bed ($P = 0.069$; $d: 0.68$ [moderate effect]), sleep efficiency ($P = 0.069$; $d: 0.60$ [small effect])
264 and wake after sleep onset ($P = 0.059$; $d: 0.54$ [small effect]). A small non-significant
265 difference was observed for sleep onset latency ($P = 0.208$; $d: 0.45$). In fringe players, there
266 was no significant difference for any of the sleep characteristics between the night after a match
267 and training nights ($P > 0.05$, $d < 0.60$).

268

269 ***INSERT TABLE 4 HERE***

270

271

272

273 Discussion

274 The aims of the current study were (1) to compare the sleep characteristics of female players
275 from a professional soccer team to non-athlete controls (2) to compare the sleep of core and
276 fringe players on the night after a match to training nights. In comparison to non-athlete
277 controls, the soccer players displayed a greater sleep duration, which was likely a result of an
278 earlier bedtime. Despite this, the soccer players had a longer sleep onset latency compared with
279 non-athlete controls. On the night after a match, bedtime was later and sleep duration was lower
280 within core players compared with training nights, whilst there was no alteration to the sleep
281 of fringe players. These findings may suggest sleep duration of professional female soccer
282 players is greater than non-athletes but that matchday challenges the sleep of players with more
283 match involvement.

284 Sleep Comparison Between Soccer Players and Non-Athlete Controls

285 This is the first study to compare the sleep characteristics of female soccer players with non-
286 athlete controls during the competitive season. In line with a previous investigation that
287 described the sleep of male youth soccer players and non-athletes¹⁶, the female soccer players
288 slept more than the non-athlete controls. Additionally, the soccer players sleep time was similar
289 to that of other elite female athletes monitored during multiple training days (7.2-7.6 h).³⁰ The
290 greater sleep duration observed within the soccer players was likely facilitated by an earlier
291 bedtime compared with non-athletes. After the completion of a training day, the soccer players
292 did not have any scheduled work commitments that may have delayed their time to bed. In
293 contrast, the bedtime of the non-athlete controls may have been constrained by independent
294 study activities in the evening (i.e. writing, reading and revision) or late-night socialising. The
295 timing of these activities are a frequent cause of poor sleep and sleepiness in students, and the
296 use of psycho-active stimulants may have added effects.³¹ In the current study, this notion is
297 also supported by the disruption to bedtime within non-athletes, as indicated by greater
298 intraindividual variability of this measure compared with soccer players. Thus, by having fewer
299 commitments in the late evening, it is possible professional female soccer players may be able
300 to increase sleep duration during the season.

301 Despite more time asleep, the female soccer players had a longer sleep onset latency
302 accompanied by a moderate effect size compared with non-athlete controls, which suggests the
303 soccer players had more difficulty falling asleep. It was also observed that the sleep onset
304 latency of the soccer players was within the range of 20-30min deemed long in duration.³² The
305 difference in time to sleep onset with the non-athlete controls was comparable to values
306 reported in previous studies (+10-13 min).^{16,18} Our findings may be attributed to excessive
307 electronic device use by players prior to bedtime, as this behaviour has been linked to delayed
308 sleep onset within senior athletes.³³ With some of the female soccer players living together, it
309 is also possible that noise outside of bedrooms or **pre-bedtime arousal** may have led to the
310 difficulty in falling asleep. The longer sleep onset latency, however, was not associated with a
311 lower sleep efficiency as found within Olympic athletes and male youth soccer players.^{16,18}
312 Indeed, the female soccer players had a similar sleep efficiency to the non-athlete controls.
313 Irrespective of this, it seems there may be a need to develop individual strategies to improve
314 the sleep onset latency of professional female soccer players.

315 Sleep Comparison Between Night After Match and Training Nights

316 Previous research has shown that sleep duration was lower following night-time training than
317 on matchdays (i.e. home and away) within semi-professional female soccer players.¹³
318 However, this study did not reflect the training schedule of professional soccer players and
319 there was no consideration that matchday sleep may be dependent upon players match
320 involvement. Within core players, we found that sleep duration was lower on the night after a
321 match compared with training nights, whereas there was no alteration to the sleep of fringe
322 players. These findings may suggest that sleep duration on the matchday night is lower
323 compared to training nights within professional female soccer players that have more match
324 involvement. This opposes Fullagar et al¹⁰, where starting players from a men's national team
325 had a smaller reduction in sleep efficiency after a night match compared with pre-match days
326 than non-players. In the current study, sleep duration of core players on the night after a match
327 was below the minimum 7h recommended to maintain optimal health and functioning.³ This
328 length of sleep may impact post-match recovery and subsequently, the training readiness of
329 core players.¹⁴ Taken together, these findings may provide a rationale to prioritise the sleep
330 hygiene of core players over fringe players post-match.

331 Given the bedtime of core players was later on the night after a match, a delayed time to bed
332 was likely responsible for the reduced sleep duration. Later bedtimes coupled with lower sleep
333 durations are common among soccer players, more so after evening matches (i.e. late kick
334 offs).^{5,6,11} Of note, our data demonstrates that core players still retired to bed later and slept less
335 on the night of an afternoon match (12-14:00 kick off). This broadly concurs with findings
336 from Australian football, where the night after a match, regardless of the match start time,
337 impacted bedtime and sleep duration compared with non-matchdays.³⁴ The reason for the later
338 bedtime is unclear but is probably related to match factors as there was no alteration to the
339 sleep of fringe players. It should also be mentioned more core players were involved in away
340 matches, however, those that travelled the shortest (derby match) had the later bedtimes,
341 indicating post-match travel was not influential. An obvious candidate from the match are the
342 physical effects of playing (i.e. increased muscle damage)², though such mechanisms have not
343 gained appraisal.³⁵ It is perhaps more plausible that the core players may have experienced a
344 change in their mood (i.e. elation or tension) as a result of the match outcome that meant they
345 adopted a later bedtime than usual. Male youth soccer players displayed poorer mood and sleep
346 quality the day after losing a match but better mood and sleep quality after winning.³⁶
347 Nonetheless, further research is warranted to provide a better understanding of the factors that
348 impact sleep following daytime matches in professional soccer players.

349 **Limitations**

350 Although the current study utilised female soccer players at the professional level, there are
351 some limitations that should be considered. There was no controlling for the menstrual cycle,
352 so habitual sleep may have been altered due to the participants being within different phases.
353 It is acknowledged other factors (e.g. napping, psychology, electronic device use, nutrition,
354 water immersion recovery strategies and sleep environment) were not monitored, which would
355 have been useful to explain sleep differences. Due to monitor availability, not all players were
356 assessed at the same time and thus, this may have increased variation in the sleep characteristics
357 reported in this article. As the data was collected from one team during five days of the season,
358 it may not reflect other teams or indeed, competitive schedules (i.e. two games per week). It is
359 also unknown from our data if core and fringe players sleep is affected by match location and
360 time. Future research should examine the sleep of professional female soccer players over a
361 longer period during the season that incorporates more match situations.

362 **Practical Applications**

- 363 • Professional female soccer players may have greater sleep duration during the season
364 than non-athlete controls due to earlier bedtimes. Practitioners could, therefore,
365 highlight the importance of having control over bedtime to achieve more sleep.
- 366 • Despite a greater sleep duration, professional female soccer players may also have
367 longer sleep onset latencies compared with non-athlete controls. Implementing
368 individual sleep strategies may be useful to assist soccer players with falling asleep.
- 369 • On the night after a match, core but not fringe players may sleep less than on training
370 nights. Practitioners could use this information to prioritise core players sleep following
371 a match as reduced sleep duration may impact their recovery from the match itself.

372 **Conclusion**

373 In conclusion, sleep duration of professional female soccer players is greater than non-athlete
374 controls. This is likely due to fewer evening commitments allowing for an earlier bedtime and
375 highlights the importance of professional female soccer players having control of bedtime to
376 achieve more sleep. However, the night after a match challenges the sleep of players with more
377 match involvement and as such, they warrant priority of sleep hygiene strategies.

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393 **Acknowledgements**

394 The authors thank the participants for their involvement and the performance coach of the
395 professional soccer team that wishes to remain anonymous.

396 **Conflicts of Interest**

397 The authors have no conflicts of interest to declare.

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

- 420 1. Halson SL. Nutrition, sleep and recovery. *Eur J Sport Sci.* 2008; 8(2): 119-126.
- 421 2. Fullagar H, Duffield R, Skorski S, Coutts AJ, Julian R, Meyer T. Sleep and recovery in
422 team sport: current sleep-related issues facing professional team sport athletes. *Int J*
423 *Sports Physiol Perform.* 2015; 10(8): 950-957.
- 424 3. Hirshkowitz M, Whiton K, Albert SM et al. National Sleep Foundation's sleep time
425 duration recommendations: methodology and results summary. *Sleep Health.* 2015;
426 1(1): 40-43.
- 427 4. Ohayon M, Wickwire EM, Hirshkowitz M et al. National Sleep Foundation's sleep
428 quality recommendations: first report. *Sleep Health.* 2017; 3(1): 6-19.
- 429 5. Nedelec M, Halson S, Abaidia AE, Ahmaidi S, Dupont G. Stress, sleep and recovery in
430 elite soccer: a critical review of the literature. *Sports Med.* 2015; 45(10): 1387-1400.
- 431 6. Carrico S, Skorski S, Duffield R, Mendes B, Calvete F, Meyer T. Post-match sleeping
432 behavior based on match scheduling over a season in elite football players. *Sci Med*
433 *Football.* 2018; 2(1): 9-15.
- 434 7. Roach GD, Schmidt WF, Aughey RJ et al. The sleep of elite athletes at sea level and
435 high altitude: a comparison of sea-level natives and high-altitude natives. *Br J Sports*
436 *Med.* 2013; 47(supp 1): 114-120.
- 437 8. Robey E, Dawson B, Halson S, Gregson W, Goodman C, Eastwood P. Sleep quantity
438 and quality in elite youth soccer players: a pilot study. *Eur J Sport Sci.* 2014; 14(5):
439 410-417.
- 440 9. Fowler P, Duffield R, Vaile J. Effects of domestic air travel on technical and tactical
441 performance and recovery in soccer. *Int J Sports Physiol Perform.* 2014; 9(3): 378-386.
- 442 10. Fullagar H, Duffield R, Skorski S, White D, Bloomfield J, Kolling S, Meyer T. Sleep,
443 travel, and recovery responses of national footballers during and after long-haul
444 international air travel. *Int J Sports Physiol Perform.* 2016a; 11(1): 86-95.
- 445 11. Fullagar H, Skorski S, Duffield R, Meyer T. The effect of an acute sleep hygiene
446 strategy following a late-night soccer match on recovery of players. *Chronobiol Int.*
447 2016b; 33(5): 490-505.
- 448 12. Costa JA, Brito J, Nakamura FY, Oliveira EM, Costa OP, Rebelo AN. Does night
449 training load affect sleep patterns and nocturnal cardiac autonomic activity in high-level
450 female soccer players? *Int J Sports Physiol Perform.* 2019a; 14(6): 779-787.
- 451 13. Costa JA, Brito J, Nakamura FY, Figueiredo P, Oliveira E, Rebelo A. Sleep patterns
452 and nocturnal cardiac autonomic activity in female athletes are affected by the timing
453 of exercise and match location. *Chronobiol Int.* 2019b; 36(3): 360-373.
- 454 14. Davenne D. Sleep of athletes – problems and possible solutions. *Biol Rhythm Res.* 2009;
455 40(1): 45-52.
- 456 15. Brand S, Beck J, Gerber M, Hatzinger M, Holsboer-Trachsler E. Evidence of favorable
457 sleep-EEG patterns in adolescent male vigorous football players compared to controls.
458 *World J Biol Psychiatry.* 2010; 11(2 Pt 2): 465-475.
- 459 16. Whitworth-Turner C, Di Michele R, Muir I, Gregson W, Drust B. A comparison of
460 sleep patterns in youth soccer players and non-athletes. *Sci Med Football.* 2017; 2(1):
461 3-8.
- 462 17. Bender AM, Van Dongen HPA, Samuels CH. Sleep quality and chronotype differences
463 between elite athletes and non-athlete controls. *Clocks and Sleep.* 2019; 1(1): 3-12.
- 464 18. Leeder J, Glaister M, Pizzoferro K, Dawson J, Pedlar C. Sleep duration and quality in
465 elite athletes measured using wristwatch actigraphy. *J Sports Sci.* 2012; 30(6): 541-545.
- 466 19. Horne JA, Ostberg O. A self-assessment questionnaire to determine morningness-
467 eveningness in human circadian rhythms. *Int J Chronobiol.* 1976; 4(2): 97-110.

- 468 20. Rupp TL, Balkin TJ. Comparison of Motionlogger Watch and Actiwatch actigraphs to
469 polysomnography for sleep/wake estimation in healthy young adults. *Behav Res*
470 *Methods*. 2011; 43(4): 1152-1160.
- 471 21. Wittmann M, Dinich J, Merrow M, Roenneberg T. Social jetlag: misalignment of
472 biological and social time. *Chronobiol Int*. 2006; 23(1&2): 497-509.
- 473 22. Fuller KL, Juliff L, Gore CJ, Peiffer JJ, Halson SL. Software thresholds alter the bias
474 of actigraphy for monitoring sleep in team-sport athletes. *J Sci Med Sport*. 2017; 20(8):
475 756-760.
- 476 23. Carney CE, Buysse DJ, Ancoli-Israel S, Edinger JD, Krystal AD, Lichstein KL, Morin,
477 CM. The Consensus Sleep Diary: standardising prospective sleep self-monitoring.
478 *Sleep*. 2012; 35(2): 287-302.
- 479 24. Foster C, Florhaug JA, Franklin J, Gottschall L, Hrovatin LA, Parker S, Doleshal P,
480 Dodge C. A new approach to monitoring exercise training. *J Strength Cond Res*. 2001;
481 15(1): 109-115.
- 482 25. Caia JP, Halson SL, Scott TJ, Kelly VG. Intra-individual variability in the sleep of
483 senior and junior rugby league athletes during the competitive season. *Chronobiol Int*.
484 2017; 34(9): 1239-1247.
- 485 26. Dillon HR, Lichstein KL, Dautovich ND, Taylor DJ, Riedel BW, Bush AJ. Variability
486 in self-reported normal sleep across the adult age span. *J Gerontol B Psychol Sci Soc*
487 *Sci*. 2015; 70(1): 46-56.
- 488 27. Hopkins W, Marshall SW, Batterham AM, Hanin J. Progressive statistics for studies in
489 sports medicine and exercise science. *Med Sci Sports Exerc*. 2009; 41(1): 3-13.
- 490 28. Fritz CO, Morris PE, Richler JJ. Effect size estimates: current use, calculations and
491 interpretation. *J Exp Psychol Gen*. 2012; 141(1): 2-18.
- 492 29. Coolican H. *Research Methods and Statistics in Psychology*. London, UK: Hodder;
493 2009.
- 494 30. Lastella M, Roach GD, Vincent GE, Scanlan AT, Halson SL, Sargent C. The impact of
495 training load on sleep during a 14-day training camp in elite, adolescent, female
496 basketball players. *Int J Sports Physiol Perform*. 2020; 15(5): 724-730.
- 497 31. Hershner SD, Chervin RD. Causes and consequences of sleepiness among college
498 students. *Nat Sci Sleep*. 2014; 6: 73-84.
- 499 32. McMahan DM, Burch JB, Wirth MD, Youngstedt SD, Hardin JW, Hurley TG, Blair
500 SN, Hand GA, Shook RP, Drenowatz C, Burgess S, Hebert JR. Persistence of social
501 jetlag and sleep disruption in healthy young adults. *Chronobiol Int*. 2018; 35(3): 312-
502 328.
- 503 33. Jones MJ, Dawson B, Gucciardi DF, Eastwood PR, Miller J, Halson SL, Dunican IC,
504 Peeling P. Evening electronic device use and sleep patterns in athletes. *J Sports Sci*.
505 2019; 37(8): 864-870.
- 506 34. Lalor BJ, Halson SL, Tran J, Kemp JG, Cormack SJ. No compromise of competition
507 sleep compared with habitual sleep in elite Australian footballers. *Int J Sports Physiol*
508 *Perform*. 2018; 13(1): 29-36.
- 509 35. Aloulou A, Duforez F, Bieuzen F, Nedelec M. The effect of night-time exercise on
510 sleep architecture among well-trained male endurance runners. *J Sleep Res*. 2019; Dec
511 10: 1-12.
- 512 36. Abbott W, Brownlee TE, Harper LD, Naughton RJ, Clifford T. The independent effects
513 of match location, match result and the quality of opposition on subjective wellbeing in
514 under 23 soccer players: a case study. *Res Sports Med*. 2018; 26(3): 262-275.

516 **Table 1.** Female soccer players training schedule during the week of sleep monitoring.

	Mon	Tue	Wed	Thurs	Fri	Sat	Sun
AM	Rest	S&C (10:00)	S&C (10:00)	Rest	S&C (10:00)	Tactical (11:00)	
		Technical (11:00)	Technical (11:00)		Tactical (11:00)		
PM	Rest or recovery		S&C (14:30)	Rest	S&C (14:30)		Match (12-14:00 KO)

517 *Monday and Thursday were excluded from data collection as these were rest days. Abbreviations: AM
518 = morning; PM = afternoon; S&C = strength and conditioning session; KO = kick off.

519

520 **Table 2.** Definitions of sleep variables taken from the wristwatch actigraphy analysis.

Sleep variable	Definition
Time in bed (min)	Time difference between bedtime and get up time
Bedtime (h: min)	Estimated clock time participant attempted to sleep
Time of final awakening (h: min)	Estimated clock time participant woke up for the final time
Sleep onset latency (min)	Time between bedtime and sleep onset
Sleep duration (min)	Time asleep between sleep onset and final awakening
Sleep efficiency (%)	Time asleep divided by time in bed multiplied by 100
Wake after sleep onset (min)	Time awake between sleep onset and final awakening

521

522 **Table 3.** Mean±SD or the median (IQR) where applicable, and effect size with 95% CI for comparisons
523 of average sleep and intraindividual sleep variability in female soccer players vs. non-athlete controls.

	Soccer players	Non-athletes	ES (95% CI)
Average sleep			
Time in bed (min)	550±38*	495±51	1.22 (24-85)
Bedtime (h: min)	22:59±00:38*	23:30±00:51	0.70 (-01:01-00:00)
Time of final awakening (h: min)	07:49±00:31	07:31±00:46	0.47 (-00:08-00:45)
Sleep onset latency (min)	21±12*	13±8	0.78 (0-15)
Sleep duration (min)	456±43*	418±40	0.92 (10-67)
Sleep efficiency (%)	83.0±5.9	84.6±4.0	0.39 (-5.0-1.9)
Wake after sleep onset (min)	51±25	48±19	0.14 (-12-18)
Intraindividual sleep variability			
Time in bed (min)	43±20	54±29	0.44 (-27-6)
Bedtime (h: min)	00:29(00:12)*	00:37(00:30)	0.38 (-00:25--00:01)
Time of final awakening (h: min)	00:38(00:26)	00:50(00:36)	0.33 (-00:33-00:00)
Sleep onset latency (min)	10(12)	6(10)	0.23 (-2-9)
Sleep duration (min)	43(22)	42(38)	0.11 (-22-10)
Sleep efficiency (%)	4.0(1.9)	3.0(2.5)	0.08 (-1.0-1.7)
Wake after sleep onset (min)	14(12)	10(17)	0.16 (-4-9)

524 *Indicates $P < 0.05$ compared to non-athlete controls.

525

526 **Table 4.** Mean±SD and effect size with 95% CI for comparisons of average sleep in core and fringe
 527 players on night after match vs. training nights (*n* = 10 core players and *n* = 8 fringe players).

	Night after match	Training nights	ES (95% CI)
Core players			
Time in bed (min)	517±81	557±37	0.68 (-85-6)
Bedtime (h: min)	23:26±01:03*	22:49±00:34	0.76 (00:01-01:12)
Time of final awakening (h: min)	07:31±00:49	07:44±00:21	0.37 (-00:45-00:19)
Sleep onset latency (min)	30±31	21±9	0.45 (-7-26)
Sleep duration (min)	414±70*	463±56	0.77 (-86--12)
Sleep efficiency (%)	80.0±5.3	82.9±5.1	0.60 (-6.3-0.5)
Wake after sleep onset (min)	40±17	50±20	0.54 (-20-1)
Fringe players			
Time in bed (min)	569±41	547±39	0.55 (-9-54)
Bedtime (h: min)	23:04±00:45	23:05±00:43	0.02 (-00:30-00:28)
Time of final awakening (h: min)	08:04±00:58	07:59±00:37	0.02 (-00:31-00:42)
Sleep onset latency (min)	16±15	20±12	0.30 (-12-6)
Sleep duration (min)	467±44	458±23	0.27 (-21-39)
Sleep efficiency (%)	82.6±10.0	84.2±6.7	0.12 (-5.4-2.1)
Wake after sleep onset (min)	55±29	54±35	0.03 (-18-22)

528 *Indicates *P* < 0.05 compared to training nights.

529

530