

Ivany, E, Ritchie, L, Lip, GYH, Lotto, R, Werring, D and Lane, DA

Effectiveness and safety of antithrombotic medication in patients with atrial fibrillation and intracranial haemorrhage: systematic review and meta-analysis

<http://researchonline.ljmu.ac.uk/id/eprint/16970/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Ivany, E, Ritchie, L, Lip, GYH, Lotto, R, Werring, D and Lane, DA (2022)
Effectiveness and safety of antithrombotic medication in patients with atrial fibrillation and intracranial haemorrhage: systematic review and meta-analysis. *Stroke*. 53 (10). pp. 3035-3046. ISSN 0039-2499**

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

Effectiveness and Safety of Antithrombotic Medication in Patients With Atrial Fibrillation and Intracranial Hemorrhage: Systematic Review and Meta-Analysis

Elena Ivany¹, MRes; Leona A. Ritchie¹, MPharm; Gregory Y.H. Lip¹, MD; Robyn R. Lotto¹, PhD; David J. Werring¹, PhD; Deirdre A. Lane¹, PhD

BACKGROUND: For patients with atrial fibrillation who survive an intracranial hemorrhage (ICrH), the decision to offer oral anticoagulation (OAC) is challenging and necessitates balancing risk of thromboembolic events with risk of recurrent ICrH.

METHODS: This systematic review assesses the effectiveness and safety of OAC and/or antiplatelets in patients with atrial fibrillation with nontraumatic ICrH. Bibliographic databases CENTRAL, MEDLINE, EMBASE, and CINAHL were searched. Articles on adults with atrial fibrillation with spontaneous ICrH (intracerebral, subdural, and subarachnoid), receiving antithrombotic therapy for stroke prevention were eligible for inclusion.

RESULTS: Twenty articles (50 470 participants) included 2 randomized controlled trials (n=304), 8 observational studies, 8 cohort studies, and 2 studies that meta-analyzed individual-level data from observational studies. OAC therapy was associated with a significant reduction in thromboembolic events (summary relative risk [sRR], 0.51 [95% CI, 0.30–0.86], heterogeneity $I^2=2\%$; $P=0.39$, n=5 studies) and all-cause mortality (sRR, 0.52 [95% CI, 0.38–0.71], heterogeneity $I^2=0\%$; $P=0.44$, n=3 studies). OAC therapy was not associated with an increased risk of recurrent ICrH (sRR, 1.44 [95% CI, 0.38–5.46], heterogeneity $I^2=70\%$, $P=0.02$, n=5 studies). Nonvitamin K antagonist OACs were more effective at reducing the risk of thromboembolic events (sRR, 0.65 [95% CI, 0.44–0.97], heterogeneity $I^2=72\%$, $P=0.03$, n=3 studies) and were associated with a lower risk of recurrent ICrH (sRR, 0.52 [95% CI, 0.40–0.67], heterogeneity $I^2=0\%$, $P=0.43$, n=3 studies) than warfarin.

CONCLUSIONS: In nontraumatic ICrH survivors with atrial fibrillation, OAC therapy is associated with a reduced risk of thromboembolic events and all-cause mortality without significantly increasing risk of recurrent ICrH. This finding is primarily based on observational data, and further larger randomized controlled trials are needed to corroborate or refute these findings.

GRAPHIC ABSTRACT: A [graphic abstract](#) is available for this article.

Key Words: anticoagulant ■ atrial fibrillation ■ intracranial hemorrhage ■ ischemic stroke ■ systematic review

Long-term oral anticoagulation (OAC) is the main treatment for ischemic stroke prevention in patients with atrial fibrillation (AF) and at least 1 additional stroke risk factor,¹ but all OAC therapy is associated with

an increased risk of bleeding. Intracranial hemorrhage (ICrH) is a potential complication of OAC² and is associated with significant mortality and morbidity.^{3,4} ICrH survivors are at risk of sustaining further hemorrhage or

Correspondence to: Deirdre A. Lane, PhD, Liverpool Centre for Cardiovascular Science, University of Liverpool, 6 West Derby St, Liverpool, L7 8TX, United Kingdom. Email deirdre.lane@liverpool.ac.uk

This manuscript was sent to Kazunori Toyoda, Guest Editor, for review by expert referees, editorial decision, and final disposition.

Supplemental Material is available at <https://www.ahajournals.org/doi/suppl/10.1161/STROKEAHA.122.038752>.

For Sources of Funding and Disclosures, see page 3045.

© 2022 The Authors. *Stroke* is published on behalf of the American Heart Association, Inc., by Wolters Kluwer Health, Inc. This is an open access article under the terms of the [Creative Commons Attribution](#) License, which permits use, distribution, and reproduction in any medium, provided that the original work is properly cited. *Stroke* is available at www.ahajournals.org/journal/str

Nonstandard Abbreviations and Acronyms

AF	atrial fibrillation
ICH	intracerebral hemorrhage
ICrH	intracranial hemorrhage
NOAC	nonvitamin-K antagonist oral anticoagulant
OAC	oral anticoagulant
RCT	randomized controlled trial
RR	relative risk
VKA	vitamin-K antagonist

an ischemic stroke, particularly if AF is present.^{5,6} As a result, the clinical dilemma about what, if any, stroke prevention therapy should be offered to ICrH survivors with AF persists. There are few published data from randomized controlled trials (RCTs),^{7,8} and most of the available evidence is from observational studies.

This review aims to systematically assess the effectiveness and safety of OAC in patients with AF who have sustained a nontraumatic ICrH.

METHODS

Data Availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Eligibility Criteria

The review was conducted in accordance with the Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) guidelines and was registered with the PROSPERO database of systematic reviews (<https://www.clinicaltrials.gov>; Unique identifier: CRD42020223266).

Participants

To be eligible for inclusion, articles had to report on adults (aged ≥ 18 years) with AF who had survived a nontraumatic spontaneous ICrH of any size and any type (lobar, brain stem, deep, cerebellar, subdural, epidural or subarachnoid location; see [Table S1](#) for definitions of terms) or had cerebral microbleeds.

Intervention

The intervention of interest was long-term OAC and/or antiplatelets for stroke prevention in AF. Short-term and/or non-oral anticoagulation therapy or OAC for other reasons were excluded.

Comparators

Any form of oral, long-term anticoagulation therapy and/or antiplatelet therapy, or no comparator (no therapy) were considered.

Outcomes

The primary outcomes were thromboembolic events and recurrent ICrH. Thromboembolic events were chosen as an outcome to reflect the range of definitions used in the included studies (eg, ischemic stroke and/or systemic embolism, and thromboembolic events). Secondary outcomes were major bleeding, all-cause mortality, cardiovascular mortality, fatal hemorrhage or stroke, incidence of clinically significant nonmajor bleeding or thromboembolic events (other than ischemic stroke).

Search Strategy

The following electronic databases were searched: CENTRAL (29/06/20, 07/12/20, and 25/10/21), MEDLINE (03/07/20, 25/10/20, and 25/10/21), EMBASE and CINAHL (30/06/20, 25/09/20, and 25/10/21). Search terms and index terms associated with AF, intracerebral hemorrhage (ICH), major bleeding, and anticoagulant medications were included ([Table S2](#)). Only full-text articles were included. Searches were not limited by language but were restricted to the year 2000 onward.

Study Selection

Two researchers (E.I. and L.A.R.) independently assessed the suitability of articles for inclusion against the eligibility criteria. Any disagreements were resolved through examination of the original data and discussion, with recourse to a third reviewer (D.A.L.) where necessary.

Data Extraction

Two researchers (E.I. and L.A.R.) extracted relevant data from the articles using a standardized tabulated data extraction form. One author provided additional unpublished data.⁹

Risk of Bias Assessment

Observational studies were assessed using the Risk of Bias Assessment Tool for Nonrandomized Studies (RoBANS)¹⁰ ([Figure S1](#)) and RCTs were assessed using the Cochrane Collaboration's tool for assessing risk of bias¹¹ ([Figure S2](#)) independently by 2 researchers (E.I. and L.A.R.).

Data Synthesis

Included studies were assessed for clinical and statistical heterogeneity. Meta-analyses were performed if studies reported similar designs, had the same outcomes, comparable interventions and comparators, and pooling the results was appropriate. Studies that could not be included in meta-analyses are reported narratively.

Statistical Analysis

A random effects model was used in all meta-analyses. Event data for control and intervention groups was compared using risk ratios and associated 95% CIs. Sensitivity analyses were performed according to outcome and follow-up period, where appropriate.

Statistical heterogeneity was evaluated using the I^2 statistic. An I^2 value of 0% to 40% indicated low heterogeneity, 30% to

60% moderate heterogeneity, 50% to 90% substantial heterogeneity, and $\geq 75\%$ considerable heterogeneity.

RESULTS

The searches identified 4429 citations, and 4 titles were identified through hand-searching. After removal of duplicates, 3053 titles and 211 abstracts were assessed for eligibility. Reasons for exclusion at the abstract and full-text stages are provided in Figure 1. A total of 20 articles were included in the review.

Characteristics of the Included Studies

The systematic review included articles published between 2015 and 2021, comprising a total of 50 470 participants (mean age ranging from 67.9 years¹² to 83.6 years¹³; 24%⁹ to 71.3%¹² female, ICrH sustained on OAC ranging from 0% to 100%). Eight articles^{13–20} reported on prospective observational cohorts, 6^{9,12,21–24} on retrospective cohorts, 2^{25,26} on nationwide cohorts, 2^{27,28} meta-analyzed individual-level data from observational studies, 1 reported on the pilot phase of an RCT,⁸ and 1 reported on a Phase 2 trial⁷ (Table S3). Nine studies^{8,12–15,23–26} included patients with an index ICrH (intracerebral, subdural, subarachnoid, or epidural hemorrhages) and 11 studies^{7,9,16–22,27,29} included patients with an index ICH. Eight studies could not be included in meta-analyses, either due to differences in reported outcomes^{21,27,28} or because raw event data were not available.^{14,15,17,18,24}

The included articles reported on a mixture of OAC-naïve patients and patients who had their index event while on OAC and/or antiplatelets. The intervention ranged from vitamin-K antagonist (VKA) only,^{12,15,22,25,27,28} nonvitamin K antagonist oral anticoagulants (NOAC) only,^{7,16,21,23,26} a mixture of VKA and NOAC,^{9,13,14,17,20,24} or OAC and/or antiplatelets.^{8,18,19} The most commonly reported outcomes were ischemic stroke and recurrent ICrH. There were variations in how the outcome of ischemic stroke was defined, including cerebral infarct, ischemic stroke, thromboembolic events, major vascular events, and the combined outcome of ischemic event/systemic embolism (Table).

Primary Outcomes

Thromboembolic Events

Seventeen articles (n=35 441) reported on the primary outcome of ischemic stroke/systemic embolism or alternative definitions of stroke. Of these, 9 studies reported on ischemic stroke alone,^{7–9,16,19,23,25–27} 5 on ischemic stroke/systemic embolism combined,^{13–15,18,20} 1 on thromboembolic events,¹² 1 on ischemic stroke combined with transient ischemic attack (TIA),²⁴ and 1 on cerebral infarction.²² One article included OAC-naïve

participants,²³ 8 included participants who sustained an ICrH on OAC^{7,15,19,20,22,24,26,27} and 8 included a combination of both OAC-naïve and current OAC users.^{8,9,12–14,16,18,25} Follow-up ranged from a median of 17 days⁹ to a median of 48.6 months.²⁷

Oral Anticoagulation Versus No Therapy

Five studies^{9,12,13,20,22} (n=1187 participants) compared the effect of OAC with no therapy on the risk of thromboembolic events (defined in the included articles as cerebral infarction, ischemic stroke, ischemic stroke/systemic embolism) and were entered into a meta-analysis (Figure 2A), which revealed a significant reduction in thromboembolic events with OAC compared with no therapy (relative risk [RR], 0.51 [95% CI, 0.30–0.86]; $P=0.01$, $I^2=2\%$).

Three studies could not be entered into a meta-analysis,^{15,24,27} either because raw event data were not available^{15,24} or because the study compared the effect of OAC therapy solely in patients with lobar and nonlobar ICH (with no control group).²⁷ Nielsen et al¹⁵ reported that OAC therapy was associated with a nonsignificant reduction in the rate of ischemic stroke and systemic embolism compared with no therapy (event rate 3.3 versus 8.9 per 100 person-years, adjusted HR, 0.49 [95% CI, 0.24–1.02]). Biffi et al²⁷ reported that restarting VKA was associated with a reduced risk of sustaining an ischemic stroke in both lobar (HR, 0.48 [95% CI, 0.25–0.75]; $P=0.003$) and nonlobar (HR, 0.39 [95% CI, 0.21–0.74]; $P=0.004$) ICH patients. Newman et al²⁴ reported no statistical significance between the OAC and no therapy groups for the outcome of stroke/TIA (adjusted HR, 0.87 [95% CI, 0.62–121]).

Oral Anticoagulation and/or Antiplatelets Versus No Therapy

Three observational studies^{18,19,25} compared OAC and/or antiplatelets with no therapy and reported on the outcome of ischemic events. The study by Pennlert et al¹⁸ was not included in the meta-analysis due to unavailability of raw event data. The pooled relative risk for the other 2 studies (n=13 063)^{19,25} was RR, 0.93 [95% CI, 0.43–2.04]; $P=0.87$, $I^2=77\%$ (Figure 2B).

Oral Anticoagulation Versus Antiplatelets or No Therapy

Three observational studies^{14,18,25} (n=17 287) and 2 RCTs^{7,8} (n=304) compared OAC versus antiplatelet or no therapy. Three studies were included in a meta-analysis^{7,8,25} (n=13 221), which found no significant difference in the risk of thromboembolic events between OAC and antiplatelet or no therapy (RR, 0.58 [95% CI, 0.23–1.46]; $P=0.25$, $I^2=74\%$; Figure 2C).

Two studies could not be entered into the meta-analysis due to lack of raw event data. Pennlert et al¹⁸ reported that the cumulative incidence of thromboembolic events 3-years post-index ICH was 6.3% in patients assigned to OAC versus 18.8% in the antiplatelet group and 13.8% in the no therapy group. Nielsen et al¹⁴ reported that the




Figure 1. Flow-diagram depicting the selection of included studies.

incidence rate of ischemic stroke/systemic embolism, per 100 person-years, was 5.3 (95% CI, 3.3–8.5) in the OAC group, 10.3 (95% CI, 7.3–14.4) in the antiplatelet group, and 10.4 (95% CI, 8.2–13.1) in the no therapy group.

NOAC Versus Warfarin

Three studies ($n=8711$)^{16,23,26} compared NOAC with warfarin and reported a significant reduction in the risk of thromboembolic events with NOAC compared with

Table. Summary of Event Data by Intervention and Comparator for Articles Included in the Meta-Analyses

Author, y, country	Intervention (n)	Comparison (n)	Length of follow-up	Types of ICrH included and ICrH diagnosis technique(s)	Number of events: ischemic stroke	Number of events: intracranial/intracerebral hemorrhage	Number of events: all-cause mortality
NOAC vs warfarin							
Lee et al, ²³ 2020, Korea	NOAC (n=1115)*	Warfarin (n=2434)	Median, year 0.6 (IQR, 0.2–1.7)	Intracranial hemorrhage. Diagnosis based on clinical presentation, hospitalization, CT, and/or MRI scan.	NOAC: 45/1115* Warfarin: 191/2434 HR, 0.729 (95% CI, 0.522–1.017)	NOAC: 19/1115* Warfarin: 92/2434 HR, 0.628 (95% CI, 0.379–1.039)	NOAC: 78/1115* Warfarin: 260/2434 HR 0.907 (95% CI, 0.699–1.719)
Tsai et al, ²⁶ 2020, Taiwan	NOAC (n=3493)	Warfarin (n=1047)	Not reported	Intracranial hemorrhage. Diagnosis method not reported (data from registry).	NOAC: 226/3493 Warfarin: 78/1047 aHR, 0.879 (95% CI, 0.678–1.141)	NOAC: 83/3493 Warfarin: 50/1047 aHR, 0.556 (95% CI, 0.389–0.796)	NOAC: 682/3493 Warfarin: 421/1047 aHR, 0.517 (95% CI, 0.457–0.585)
Nielsen et al, ¹⁶ 2019, Denmark	NOAC (n=348)	Warfarin (n=274)	1 and 3 y	Intracerebral hemorrhage. Diagnosis method not reported (data from registry).	NOAC: 15/348 Warfarin: 21/274 Weighted risk difference, 3.78% (95% CI, –0.15% to 7.71%)*	NOAC: 18/348 Warfarin: 19/274 Weighted risk difference, 1.93% (95% CI, –2.02% to 5.87%)	Not reported
OAC vs no therapy							
Sadighi et al, ²⁰ 2020, United States	Warfarin or NOAC (n=38)	No therapy (n=55)	Mean, mo OAC: 22.7 (22.4) No OAC: 28.8 (22.0)	Spontaneous nontraumatic intracerebral hemorrhage. Initial method of diagnosis not reported.	OAC: 7/38† No therapy: 10/55 RR, 0.9 (95% CI, 0.3–2.7)	OAC: 5/38 No therapy: 3/55 RR, 2.9 (95% CI, 0.3–30.8)	OAC: 10/38 No therapy: 20/55 RR, 0.8 (95% CI, 0.3–1.9)
Sakamoto et al, ⁹ 2019, Japan	VKA or NOAC (n=29)	No therapy (n=4)	To hospital discharge. Median hospital stay 17 d (IQR, 11–26)	Lobar or nonlobar intracerebral hemorrhage. Brain imaging (CT/MRI) available for all participants.	OAC: 2/29 No therapy: 0/4 HR not reported	OAC: 0/29 No therapy: 0/4 HR not reported	Not reported
Perreault et al, ¹³ 2019, Canada	VKA or NOAC (n=max. 125)	No therapy (n=max. 249)	1 y	Intracranial hemorrhage. Initial method of diagnosis not reported.	OAC: 1/125† No therapy: 7/247 HR not reported	OAC: 4/123 No therapy: 23/249 HR not reported	OAC: 20/125 No therapy: 80/246 HR not reported
Park et al, ¹² 2016, Korea	Warfarin (n=254)	No therapy (n=174)	Mean, mo 39.5±31.9	Cerebral (47%), subdural (36.4%), subarachnoid (11%), cerebellar (3%), epidural (0.7%), intraventricular (0.7%) hemorrhages. Diagnosed by clinical presentation, CT, and/or MRI scan.	Warfarin: 7/254 No therapy: 14/174 RR, 0.19 (95% CI, 0.08–0.47)	Warfarin: 13/254§ No therapy: 0/174 RR not reported	Warfarin: 13/254 No therapy: 22/174 RR not reported
Kura-matsu et al, ²² 2015, Germany	VKA (n=108)	No therapy (n=153)	1 y	Deep (45.1%)*, lobar (37.1%), cerebellar (10.3%), brain stem (4.3%), intraventricular (3.2%) hemorrhages. Diagnosed using brain imaging (CT/MRI).	VKA: 4/108¶ No therapy: 16/153 HR not reported	VKA: 4/108 No therapy: 5/153 HR not reported	Not reported

(Continued)

Author, y, country	Intervention (n)	Comparison (n)	Length of follow-up	Types of ICrH included and ICrH diagnosis technique(s)	Number of events: ischemic stroke	Number of events: intracranial/intracerebral hemorrhage	Number of events: all-cause mortality
OAC and/or antiplatelet therapy vs no therapy							
Chao et al, ²⁵ 2016, Taiwan	Warfarin (n=1154) or antiplatelet (n=3552)	No therapy (n=8211)	Mean, y, 3.3±3.6	Intracerebral (68.6%), subarachnoid (12.3%), epidural (2.5%), subdural (12.6%), nonspecified (4%) hemorrhages. Diagnosis method not reported (registry data).	Warfarin: 130/1154 aHR, 0.66 (95% CI, 0.55–0.79) Antiplatelet: 581/3552 aHR, 0.90 (95% CI, 0.81–1.01) No therapy: 954/8211 aHR, 1	Warfarin: 241/1154 aHR, 1.60 (95% CI, 1.38–1.86) Antiplatelet: 628/3552 aHR, 1.35 (95% CI, 1.21–1.51) No therapy: 730/8211 aHR, 1	Not reported
Poli et al, ¹⁹ 2018, Italy	Warfarin or NOAC (n=55) or antiplatelet (n=29)	No therapy (n=62)	Median, mo 18.0	Lobar (cortex or cerebellar) and nonlobar (basal ganglia, thalamus, or brain stem) intracerebral hemorrhages. Diagnosed by CT or MRI scan.	OAC: 2/55† wHR, 0.09 (95% CI, 0.02–0.40) Antiplatelet: 8/29 wHR, 1.07 (95% CI, 0.40–2.83) No therapy: 13/62 wHR, 1	Not reported	OAC: 5/55 wHR, 0.23 (95% CI, 0.08–0.68) Antiplatelet: 7/29 wHR, 0.84 (95% CI, 0.31–2.25) No therapy: 12/62 wHR, 1
OAC vs antiplatelet therapy or no therapy							
Chao et al, ²⁵ 2016, Taiwan	Warfarin (n=1154) or antiplatelet (n=3552)	No therapy (n=8211)	Mean, y, 3.3±3.6	Intracerebral (68.6%), subarachnoid (12.3%), epidural (2.5%), subdural (12.6%), nonspecified (4%) hemorrhages. Diagnosis method not reported (registry data).	Warfarin: 130/1154 aHR, 0.66 (95% CI, 0.55–0.79) Antiplatelet: 581/3552 aHR, 0.90 (95% CI, 0.81–1.01) No therapy: 954/8211 aHR, 1	Warfarin: 241/1154 aHR, 1.60 (95% CI, 1.38–1.86) Antiplatelet: 628/3552 aHR, 1.35 (95% CI, 1.21–1.51) No therapy: 730/8211 aHR, 1	Not reported
APACHE-AF Investigators, ⁷ 2021, the Netherlands	Apixaban (n=50)	Antiplatelet or no therapy (n=51)	Median 1.9 y (IQR, 1.0–3.1)	Lobar, nonlobar, brain stem, cerebellar, and intraventricular intracerebral hemorrhages. Diagnosis confirmed by imaging (not defined).	Apixaban: 6/50 Antiplatelets or no therapy: 6/51 aHR, 0.96 (95% CI, 0.31–2.97)	Apixaban: 4/50# Antiplatelets or no therapy: 1/51 aHR, 4.08 (95% CI, 0.45–36.91)	Apixaban: 9/50 Antiplatelets or no therapy: 11/51 HR not available
SoSTART Collaboration, ⁸ 2021, United Kingdom	NOAC or VKA (n=101)	Antiplatelet or no therapy (n=102)	Median 1.2 y (IQR, 0.97–1.95)	Lobar and nonlobar spontaneous intracerebral, nonaneurysmal subarachnoid, intraventricular or subdural hemorrhage. Diagnosis confirmed by CT or MRI scan.	OAC: 3/101 Antiplatelets or no therapy: 19/102 HR not available	OAC: 8/101 Antiplatelets or no therapy: 4/102 aHR, 2.42 (95% CI, 0.72–8.09)	OAC: 15/101 Antiplatelets or no therapy: 11/102 HR not available

AF indicates atrial fibrillation; aHR, adjusted hazard ratio; CT, computerized tomography; HR, hazard ratio; ICD-10, International Classification of Diseases and Health Related Problems, Tenth Revision; ICrH, intracranial hemorrhage; IQR, interquartile range; MRI, magnetic resonance imaging; N/A, not available; NOAC, nonvitamin K antagonist oral anticoagulant; OAC, oral anticoagulant; RR, relative risk; VKA, vitamin K antagonist; and wHR, weighted hazard ratio.

*Participant numbers and event rates given are before propensity matching.

†Outcome reported is ischemic stroke/systemic embolism.

#Participant numbers reported for participants with AF only.

§Outcome reported is recurrent CNS bleeding.

||Event rate data reported is after propensity matching.

¶Outcome reported is cerebral infarction.

warfarin (RR, 0.65 [95% CI, 0.44–0.97]; $P=0.03$, $I^2=72\%$) but there was considerable heterogeneity (Figure 2D).

Recurrent ICrH

Fifteen studies^{7–9,12–16,20,22–27} ($n=32\,579$ patients with AF and ICrH) reported on the outcome of recurrent ICrH. Eight studies^{8,13–15,23–26} included patients with an index ICrH (intracerebral, subdural, subarachnoid, or epidural hemorrhages) and 7 studies^{7,9,14,16,20,22,27} included patients with an index ICH. One study included OAC-naïve participants,²³ 7 studies included participants who sustained an ICrH on OAC,^{7,15,20,22,24,26,27} and 7 studies included a combination of OAC-naïve patients and patients who sustained an ICrH on OAC.^{8,9,12–14,16,25} Follow-up ranged from a median of 17 days⁹ to a median 48.6 months.²⁷

Oral Anticoagulation Versus No Therapy

Five observational studies ($n=1187$)^{9,12,13,20,22} compared OAC versus no therapy on the risk of sustaining a recurrent ICrH and the pooled estimate revealed no statistically significant difference (RR, 1.44 [95% CI, 0.38–5.46]; $P=0.59$, $I^2=70\%$; Figure 3A).

Three observational studies could not be entered into a meta-analysis.^{15,24,27} Biffi et al²⁷ reported that VKA resumption was associated with a nonsignificant increase in the risk of sustaining an ICH (lobar HR, 1.21 [95% CI, 0.86–1.70]; $P=0.27$; nonlobar HR, 1.10 [95% CI, 0.94–1.28]; $P=0.23$). Nielsen et al¹⁵ reported that warfarin treatment was associated with a nonsignificant increase in the risk of recurrent ICrH (adjusted HR, 1.31, [95% CI, 0.68–2.50]). Newman et al²⁴ reported an association between OAC therapy (VKA or NOAC) post-ICrH and a reduction in the risk of recurrent ICrH (incidence 3.29 versus 5.80 events per 100 patient years, adjusted HR=0.62 [95% CI, 0.41–0.95]).

Oral Anticoagulation Versus Antiplatelets or No Therapy

Two RCTs⁷⁸ ($n=304$) and 2 observational studies^{14,25} ($n=14\,669$) compared OAC versus antiplatelet therapy or no therapy. Three studies^{78,25} ($n=13\,221$) were entered into a meta-analysis; OAC was associated with a higher risk of recurrent ICrH versus antiplatelet or no therapy (RR, 1.82 [95% CI, 1.61–2.05]; $P<0.01$, $I^2=0\%$; Figure 3B).

Nielsen et al.¹⁴ reported that the incidence rate of recurrent ICrH per 100 person-years, was 8.0 (95% CI, 5.4–11.8) in the OAC group, 5.3 (95% CI, 3.3–8.4) in the antiplatelet group, and 8.6 (95% CI, 6.6–11.2) in the no therapy group.

NOAC Versus Warfarin

Three studies ($n=8711$)^{16,23,26} compared the effect of NOAC versus warfarin on recurrent ICrH with the pooled relative risk demonstrating that NOAC significantly reduced risk of recurrent ICrH compared with warfarin (RR, 0.52 [95% CI, 0.40–0.67]; $P<0.00001$, $I^2=0\%$; Figure 3C).

Secondary Outcomes

All-Cause Mortality

Five observational studies ($n=11\,456$)^{12,13,20,23,26} and 2 RCTs⁷⁸ ($n=304$) reported on all-cause mortality. Three studies included patients who had sustained their index ICrH on OAC,^{7,20,26} 1 study included OAC-naïve patients,²³ and another 3 included a mixture of OAC-naïve patients and patients who sustained their ICrH on OAC.^{8,12,13} Two studies included patients who sustained an index ICH^{7,20} and 5 articles included patients who sustained an index ICrH.^{8,12,13,23,26} The longest follow-up was median 39.9 months.¹²

Oral Anticoagulation Versus No Therapy

Three studies ($n=891$)^{12,13,20} examined the impact of OAC versus no therapy on all-cause mortality and reported a significant reduction in death associated with OAC (RR, 0.52 [95% CI, 0.38–0.71]; $P<0.01$, $I^2=0\%$; Figure 4A).

Oral Anticoagulation Versus Antiplatelets or No Therapy

Two RCTs ($n=304$) examined the impact of OAC versus antiplatelets or no therapy on all-cause mortality and were entered into a meta-analysis, which was not statistically significant (RR, 1.09 [95% CI, 0.64–1.87]; $P=0.74$, $I^2=0\%$; Figure 4B).

NOAC Versus Warfarin

Two studies ($n=8089$)^{23,26} examined the impact of NOAC versus warfarin on the risk of all-cause mortality and the pooled estimate demonstrated that NOACs were significantly associated with a reduced risk of all-cause mortality (RR, 0.55 [95% CI, 0.41–0.74]; $P<0.00001$, $I^2=81\%$; Figure 4C).

Subgroup and Sensitivity Analyses

Subgroup and sensitivity analyses are provided in Table S4. OAC therapy significantly reduced the risk of thromboembolic events at 1 year follow-up (RR, 0.34 [95% CI, 0.13–0.87], $I^2=0\%$) but not at >1 year (RR, 0.59 [95% CI, 0.20–1.72], $I^2=66\%$). The risk of sustaining a recurrent ICrH did not differ by follow-up time. Examining only the RCT data⁷⁸ ($n=304$), no difference was found in the risk of ischemic stroke (RR, 0.41 [95% CI, 0.06–2.64], $I^2=82\%$) or recurrent ICH (RR, 2.37 [95% CI, 0.85–6.62], $I^2=0\%$) when comparing OAC with antiplatelet or no therapy.

Risk of Bias Assessment

The overall risk of bias assessment for all included studies is presented in Figures S1 and S2. The categories addressing participant selection, incomplete outcome data, and selective outcome reporting among observational studies were assessed as having the lowest risk of bias.

DISCUSSION

This systematic review included 20 studies ($n=50\,470$, 304 were enrolled in RCTs) and updates previously




Figure 2. Forest plot depicting the risk of thromboembolic events in patients postintracranial hemorrhages with atrial fibrillation receiving oral anticoagulant (OAC) versus no therapy, OAC and/or antiplatelets versus no therapy, OAC versus antiplatelets/no therapy, or nonvitamin K antagonist oral anticoagulant (NOAC) versus warfarin.

published reviews.^{29–31} Our main findings are that OAC significantly reduced the risk of thromboembolic events and all-cause mortality in patients with AF and ICrH, without significantly increasing the risk of recurrent ICrH. Second, NOACs were associated with a lower risk of thromboembolic events and recurrent ICrH than warfarin.

Oral Anticoagulation for Stroke Prevention

This PRISMA-compliant systematic review found that OAC therapy significantly reduced the risk of an ischemic stroke in patients with AF and a history of ICrH when compared with no therapy (RR, 0.51 [95% CI, 0.30–0.86]; $P=0.01$). Previous meta-analyses that investigated the effect of restarting OAC post-ICrH reported that OAC generally and VKA specifically were associated with a reduction in thromboembolic events.^{29,30,32} The

current review also found that NOAC therapy is more effective at preventing thromboembolic events than warfarin (RR, 0.65 [95% CI, 0.44–0.97]). Although trials comparing NOAC and warfarin largely excluded patients with ICrH, and findings in this review are based on observational data, there is evidence to suggest that NOACs are more effective at preventing thromboembolic events than warfarin.³³ However, 2 recently completed RCTs found that restarting OAC was not associated with a significant decrease in the risk of thromboembolic events in patients with AF and a history of ICH or ICrH.^{7,8}

Oral Anticoagulation and All-Cause Mortality

The significant reduction in all-cause mortality among patients with AF who received OAC therapy following ICrH compared with those who received no therapy




Figure 3. Forest plot depicting the risk of repeat intracranial hemorrhage in patients post-intracerebral hemorrhage with atrial fibrillation receiving oral anticoagulant (OAC) versus no therapy, OAC versus antiplatelets/no therapy, or nonvitamin-K antagonist oral anticoagulant (NOAC) versus warfarin.

supports the use of OAC in the post-ICrH population with AF. This finding is confirmed by a previous meta-analysis that examined OAC resumption in patients who sustained an OAC-associated ICrH.³² The main limitation with assessing the impact of OAC therapy following ICrH is confounding by indication, as studies have shown differences between those who did and those who did not receive OAC post-ICrH.^{34–36} Two studies also reported that patients exposed to OAC at the time of their ICrH were less likely to restart OAC post-ICrH.^{12,13} It is possible that OAC is associated with improved survival in AF patients post-ICrH as OAC is more likely to be prescribed to those who are more likely to survive. However, OAC use post-ICrH has been shown to be associated with improved functional outcomes among patients with poor functional status (modified Rankin Scale score >3) at hospital discharge.^{27,28}

Oral Anticoagulation and the Risk of Recurrent ICrH

The current review found that OAC therapy was not associated with a statistically significantly increased

risk of recurrent ICrH (RR, 1.44 [95% CI, 0.38–5.46]; $P=0.59$). Of the 10 studies examining the association between OAC and/or antiplatelet therapy and the risk of recurrent ICrH, 2 studies^{12,25} reported a significant increase in the risk of an ICrH (defined in 1 article as CNS bleeding), 2 studies^{13,24} reported a reduction in the risk of repeat ICrH, and 6^{7,8,14,20,22,27} studies reported no significant difference in the risk of recurrent ICrH. There was heterogeneity in the type of OAC therapy used and the baseline characteristics of the patients who were commenced on OAC therapy. Furthermore, there was considerable heterogeneity in the participant cohorts and follow-up periods reported, and an unclear risk of bias regarding measurement of participants' exposure to OAC therapy in all but one of the included articles.

From the patient's perspective, the key attribute of OAC therapy is stroke prevention, although risk of bleeding is the second most important attribute when choosing OAC.³⁷ However, patients report variability in the number of acceptable bleeds associated with OAC therapy and considerable differences in the percentage of patients who were not willing to consider OAC therapy.³⁸ The study by Chao et al²⁵ reported that




Figure 4. Forest plot depicting the risk of all-cause mortality in patients postintracranial hemorrhage with atrial fibrillation receiving oral anticoagulation (OAC) versus no therapy, OAC versus antiplatelets/no therapy, or nonvitamin K antagonist oral anticoagulant (NOAC) versus warfarin.

patients who survive an ICrH are at increased risk of repeat ICrH regardless of whether they receive OAC therapy post-ICrH or not, and that OAC therapy with VKA post-ICrH ought to be reserved for patients with $\text{CHA}_2\text{DS}_2\text{-VASc} \geq 6$. NOACs may alleviate some concerns about OAC-related ICrH because apixaban and dabigatran have been shown to be associated with reduced risk of major bleeding when compared with warfarin,³⁹ and NOACs have reversal agents which may prevent exacerbating the ICrH. However, results from APACHE-AF⁷ show that there were more recurrent ICH in the apixaban group than in the antiplatelet or no therapy group (8% versus 2%), although this difference was not significant (adjusted hazard ratio, 4.08 [95% CI, 0.45–36.91]). Furthermore, SoSTART⁸ reported that OAC could not be considered noninferior to no therapy due to the increased risk of mortality and recurrent ICrH.

Strengths and Limitations of This Review

Several bibliographic databases were searched to ensure that all contemporary relevant literature was captured, and 2 authors independently selected the included studies and extracted the data. Sensitivity analyses and subgroup analyses were undertaken.

The primary limitation of this review is that most are observational cohort studies. The included studies were heterogeneous, both clinically and methodologically, with most reporting on both intracerebral and other types of ICrH combined. This is a limitation since intracerebral (or parenchymal) hemorrhage is associated with a higher intrinsic risk of thrombotic events than subarachnoid hemorrhage. In addition, not all studies could be included in the meta-analyses due to unavailable data or differences in reported outcomes. Finally, it was difficult to accurately assess the measurement of participants' exposure to OAC therapy, since several studies were retrospective in design and utilized patient data from large databases. Therefore, the results of this systematic review should be interpreted with caution. Several RCTs addressing the efficacy and safety of OAC for stroke prevention in patients with AF who have survived an ICH or ICrH are ongoing.^{40–45} The findings reported by these RCTs will be critical to confirm or refute the findings of this review.

CONCLUSIONS

OAC use after ICrH in patients with AF significantly reduces the risk of thromboembolic events and all-cause mortality, without significantly increasing the risk of recurrent ICrH. NOACs are preferable to warfarin as they

are associated with preventing thromboembolic events with a lower risk of recurrent ICrH. Nevertheless, the available evidence is mostly observational, with considerable clinical and methodological heterogeneity, including differences in intervention, comparators, outcomes, and follow-up time. Thus, further evidence from ongoing RCTs is urgently needed to corroborate these findings.

ARTICLE INFORMATION

Received January 18, 2022; final revision received May 5, 2022; accepted May 25, 2022.

Affiliations

Liverpool Centre for Cardiovascular Science (E.I., L.A.R., G.Y.H.L., R.R.L., D.A.L.) and Department of Cardiovascular and Metabolic Medicine, Institute of Life Course and Medical Sciences (E.I., L.A.R., G.Y.H.L., D.A.L.), University of Liverpool, United Kingdom. Liverpool Heart and Chest Hospital, United Kingdom (G.Y.H.L., D.A.L.). Department of Clinical Medicine, Aalborg University, Denmark (G.Y.H.L., D.A.L.). School of Nursing and Allied Health, Faculty of Health, Liverpool John Moores University, United Kingdom (R.R.L.). Stroke Research Centre, University College London, Queen Square Institute of Neurology, United Kingdom (D.J.W.).

Acknowledgments

The authors would like to thank Dr Yuki Sakamoto and colleagues for their provision of supplementary data to accompany the article included in the review.

Sources of Funding

This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 754517.

Disclosures

E. Ivany is a Research Associate on the PRESTIGE-AF study, which has received funding from the European Union's Horizon 2020 research and innovation programme under Grant Agreement No. 754517. Dr Lane has received investigator-initiated educational grants from Bristol-Myers Squibb (BMS), has been a speaker for Bayer, Boehringer Ingelheim, and BMS/Pfizer and has consulted for BMS, and Boehringer Ingelheim. Dr Lane is a co-investigator on the PRESTIGE-AF study, which has received funding from the European Union's Horizon 2020 research and innovation programme under Grant Agreement No. 754517. Dr Lip is a consultant and speaker for BMS/Pfizer, Boehringer Ingelheim, and Daiichi-Sankyo. No fees are received personally. Dr Lip is a co-investigator on the PRESTIGE-AF study, which has received funding from the European Union's Horizon 2020 research and innovation programme under Grant Agreement No. 754517. Dr Lip was part of the steering committee for the SoSTART trial. Dr Werring received honoraria for consulting or lecturing from Alexion, Alnylam, Bayer, Novo Nordisk, and Portola. The other authors report no conflicts.

Supplemental Material

Tables S1–S4
Figures S1–S2
MOOSE and PRISMA Checklists

REFERENCES

- Lip GYH, Banerjee A, Boriani G, Chiang CE, Fargo R, Freedman B, Lane DA, Ruff CT, Turakhia M, Werring D, et al. Antithrombotic therapy for atrial fibrillation: CHEST guideline and expert panel report. *Chest*. 2018;154:1121–1201. doi: 10.1016/j.chest.2018.07.040
- Hart RG, Boop BS, Anderson DC. Oral anticoagulants and intracranial hemorrhage. Facts and hypotheses. *Stroke*. 1995;26:1471–1477. doi: 10.1161/01.str.26.8.1471
- Poon MTC, Fonville AF, Salman RA-S. Long-term prognosis after intracerebral hemorrhage: Systematic review and meta-analysis. *J Neurol Neurosurg Psychiatry*. 2014;85:660–667. doi: 10.1136/jnnp-2013-306476
- Delcourt C, Zheng D, Chen X, Hackett M, Arima H, Hata J, Heeley E, Al-Shahi Salman R, Woodward M, Huang Y, et al; INTERACT Investigators. Associations with health-related quality of life after intracerebral hemorrhage: pooled analysis of INTERACT studies. *J Neurol Neurosurg Psychiatry*. 2017;88:70–75. doi: 10.1136/jnnp-2016-314414
- Murthy SB, Diaz I, Wu X, Merkler AE, Iadecola C, Safford MM, Sheth KN, Navi BB, Kamel H. Risk of arterial ischemic events after intracerebral hemorrhage. *Stroke*. 2020;51:137–142. doi: 10.1161/STROKEAHA.119.026207
- Vermeer SE, Algra A, Franke CL, Koudstaal PJ, Rinkel GJ. Long-term prognosis after recovery from primary intracerebral hemorrhage. *Neurology*. 2002;59:205–209. doi: 10.1212/wnl.59.2.205
- Schreuder FHBM, van Nieuwenhuizen KM, Hofmeijer J, Vermeer SE, Kerkhoff H, Zock E, Luijckx GJ, Messchendorp GP, van Tuijl J, Bienfait HP, et al; APACHE-AF Trial Investigators. Apixaban versus no anticoagulation after anticoagulation-associated intracerebral hemorrhage in patients with atrial fibrillation in the Netherlands (APACHE-AF): a randomised, open-label, phase 2 trial. *Lancet Neurol*. 2021;20:907–916. doi: 10.1016/S1473-4422(21)00298-2
- Collaboration S. Effects of oral anticoagulation for atrial fibrillation after spontaneous intracranial haemorrhage in the UK: a randomised, open-label, assessor-masked, pilot-phase, non-inferiority trial. *Lancet Neurol*. 2021;20:842–853. doi: 10.1016/S1473-4422(21)00264-7
- Sakamoto Y, Nito C, Nishiyama Y, Suda S, Matsumoto N, Aoki J, Shimoyama T, Kanamaru T, Suzuki K, Nishimura T, et al. Safety of anticoagulant therapy including direct oral anticoagulants in patients with acute spontaneous intracerebral hemorrhage. *Circ J*. 2019;83:441–446. doi: 10.1253/circj.CJ-18-0938
- Kim SY, Park JE, Lee YJ, Seo HJ, Sheen SS, Hahn S, Jang BH, Son HJ. Testing a tool for assessing the risk of bias for nonrandomized studies showed moderate reliability and promising validity. *J Clin Epidemiol*. 2013;66:408–414. doi: 10.1016/j.jclinepi.2012.09.016
- Higgins JP, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD, Savovic J, Schulz KF, Weeks L, Sterne JA; Cochrane Bias Methods Group; Cochrane Statistical Methods Group. The cochrane collaboration's tool for assessing risk of bias in randomised trials. *BMJ*. 2011;343:d5928. doi: 10.1136/bmj.d5928
- Park YA, Uhm JS, Pak HN, Lee MH, Joung B. Anticoagulation therapy in atrial fibrillation after intracranial hemorrhage. *Heart Rhythm*. 2016;13:1794–1802. doi: 10.1016/j.hrthm.2016.05.016
- Perreault S, Côté R, White-Guay B, Dorais M, Oussaid E, Schnitzer ME. Anticoagulants in older patients with nonvalvular atrial fibrillation after intracranial hemorrhage. *J Stroke*. 2019;21:195–206. doi: 10.5853/jos.2018.02243
- Nielsen PB, Larsen TB, Skjøth F, Gorst-Rasmussen A, Rasmussen LH, Lip GY. Restarting anticoagulant treatment after intracranial hemorrhage in patients with atrial fibrillation and the impact on recurrent stroke, mortality, and bleeding: a nationwide cohort study. *Circulation*. 2015;132:517–525. doi: 10.1161/CIRCULATIONAHA.115.015735
- Nielsen PB, Larsen TB, Skjøth F, Lip GY. Outcomes associated with resuming warfarin treatment after hemorrhagic stroke or traumatic intracranial hemorrhage in patients with atrial fibrillation. *JAMA Intern Med*. 2017;177:563–570. doi: 10.1001/jamainternmed.2016.9369
- Nielsen PB, Skjøth F, Søgaard M, Kjældgaard JN, Lip GYH, Larsen TB. Non-Vitamin K antagonist oral anticoagulants versus warfarin in atrial fibrillation patients with intracerebral hemorrhage. *Stroke*. 2019;50:939–946. doi: 10.1161/STROKEAHA.118.023797
- Pennert J, Asplund K, Carlberg B, Wiklund PG, Wisten A, Åsberg S, Eriksson M. Antithrombotic treatment following intracerebral hemorrhage in patients with and without atrial fibrillation. *Stroke*. 2015;46:2094–2099. doi: 10.1161/STROKEAHA.115.009087
- Pennert J, Overholser R, Asplund K, Carlberg B, Van Rompaye B, Wiklund PG, Eriksson M. Optimal timing of anticoagulant treatment after intracerebral hemorrhage in patients with atrial fibrillation. *Stroke*. 2017;48:314–320. doi: 10.1161/STROKEAHA.116.014643
- Poli L, Grassi M, Zedde M, Marcheselli S, Silvestrelli G, Sessa M, Zini A, Paciaroni M, Azzini C, Gamba M, et al; Multicenter Study on Cerebral Hemorrhage in Italy (MUCH-Italy) Investigators. Anticoagulants resumption after warfarin-related intracerebral haemorrhage: The Multicenter Study on Cerebral Hemorrhage in Italy (MUCH-Italy). *Thromb Haemost*. 2018;118:572–580. doi: 10.1055/s-0038-1627454
- Sadighi A, Wasko L, DiCristina H, Wagner T, Wright K, Capone K, Monczewski M, Kester M, Bourdages G, Griessenauer C, et al. Long-term outcome of resuming anticoagulation after anticoagulation-associated intracerebral hemorrhage. *eNeurologicalSci*. 2020;18:100222. doi: 10.1016/j.ensci.2020.100222
- Kato Y, Hayashi T, Suzuki K, Maruyama H, Kikkawa Y, Kurita H, Takao M. Resumption of direct oral anticoagulants in patients with acute spontaneous

- intracerebral hemorrhage. *J Stroke Cerebrovasc Dis*. 2019;28:104292. doi: 10.1016/j.jstrokecerebrovasdis.2019.07.008
22. Kuramatsu JB, Gerner ST, Schellinger PD, Glahn J, Endres M, Sobesky J, Flechsenhar J, Neugebauer H, Jüttler E, Grau A, et al. Anticoagulant reversal, blood pressure levels, and anticoagulant resumption in patients with anticoagulation-related intracerebral hemorrhage. *JAMA*. 2015;313:824–836. doi: 10.1001/jama.2015.0846
 23. Lee SR, Choi EK, Kwon S, Jung JH, Han KD, Cha MJ, Oh S, Lip GYH. Oral anticoagulation in asian patients with atrial fibrillation and a history of intracranial hemorrhage. *Stroke*. 2020;51:416–423. doi: 10.1161/STROKEAHA.119.028030
 24. Newman TV, Chen N, He M, Saba S, Hernandez I. Effectiveness and safety of restarting oral anticoagulation in patients with atrial fibrillation after an intracranial hemorrhage: Analysis of medicare part d claims data from 2010–2016. *Am J Cardiovasc Drugs*. 2020;20:471–479. doi: 10.1007/s40256-019-00388-8
 25. Chao TF, Liu CJ, Liao JN, Wang KL, Lin YJ, Chang SL, Lo LW, Hu YF, Tuan TC, Chung FP, et al. Use of oral anticoagulants for stroke prevention in patients with atrial fibrillation who have a history of intracranial hemorrhage. *Circulation*. 2016;133:1540–1547. doi: 10.1161/CIRCULATIONAHA.115.019794
 26. Tsai CT, Liao JN, Chiang CE, Lin YJ, Chang SL, Lo LW, Hu YF, Tuan TC, Chung FP, Chao TF, et al. Association of ischemic stroke, major bleeding, and other adverse events with warfarin use vs non-vitamin K antagonist oral anticoagulant use in patients with atrial fibrillation with a history of intracranial hemorrhage. *JAMA Netw Open*. 2020;3:e206424. doi: 10.1001/jamanetworkopen.2020.6424
 27. Biffi A, Kuramatsu JB, Leasure A, Kamel H, Kourkoulis C, Schwab K, Ayres AM, Elm J, Gurol ME, Greenberg SM, et al. Oral anticoagulation and functional outcome after intracerebral hemorrhage. *Ann Neurol*. 2017;82:755–765. doi: 10.1002/ana.25079
 28. Murphy MP, Kuramatsu JB, Leasure A, Falcone GJ, Kamel H, Sansing LH, Kourkoulis C, Schwab K, Elm JJ, Gurol ME, et al. Cardioembolic stroke risk and recovery after anticoagulation-related intracerebral hemorrhage. *Stroke*. 2018;49:2652–2658. doi: 10.1161/STROKEAHA.118.021799
 29. Murthy SB, Gupta A, Merkler AE, Navi BB, Mandava P, Iadecola C, Sheth KN, Hanley DF, Ziai WC, Kamel H. Restarting anticoagulant therapy after intracranial hemorrhage: a systematic review and meta-analysis. *Stroke*. 2017;48:1594–1600. doi: 10.1161/STROKEAHA.116.016327
 30. Korompoki E, Filippidis FT, Nielsen PB, Del Giudice A, Lip GYH, Kuramatsu JB, Huttner HB, Fang J, Schulman S, Martí-Fàbregas J, et al. Long-term antithrombotic treatment in intracranial hemorrhage survivors with atrial fibrillation. *Neurology*. 2017;89:687–696. doi: 10.1212/WNL.0000000000004235
 31. Zhou Z, Yu J, Carcel C, Delcourt C, Shan J, Lindley RI, Neal B, Anderson CS, Hackett ML. Resuming anticoagulants after anticoagulation-associated intracranial haemorrhage: Systematic review and meta-analysis. *BMJ Open*. 2018;8:e019672. doi: 10.1136/bmjopen-2017-019672
 32. Chai-Adisaksopha C, Iorio A, Hillis C, Siegal D, Witt DM, Schulman S, Crowther M. Warfarin resumption following anticoagulant-associated intracranial hemorrhage: a systematic review and meta-analysis. *Thromb Res*. 2017;160:97–104. doi: 10.1016/j.thromres.2017.11.001
 33. Ruff CT, Giugliano RP, Braunwald E, Hoffman EB, Deenadayalu N, Ezekowitz MD, Camm AJ, Weitz JJ, Lewis BS, Parkhomenko A, et al. Comparison of the efficacy and safety of new oral anticoagulants with warfarin in patients with atrial fibrillation: a meta-analysis of randomised trials. *Lancet*. 2014;383:955–962. doi: 10.1016/S0140-6736(13)62343-0
 34. Hsu JC, Freeman JV. Underuse of vitamin K antagonist and direct oral anticoagulants for stroke prevention in patients with atrial fibrillation: a contemporary review. *Clin Pharmacol Ther*. 2018;104:301–310. doi: 10.1002/cpt.1024
 35. O'Brien EC, Holmes DN, Ansell JE, Allen LA, Hylek E, Kowey PR, Gersh BJ, Fonarow GC, Koller CR, Ezekowitz MD, et al. Physician practices regarding contraindications to oral anticoagulation in atrial fibrillation: findings from the Outcomes Registry for Better Informed Treatment of Atrial Fibrillation (ORBIT-AF) registry. *Am Heart J*. 2014;167:601–609.e1. doi: 10.1016/j.ahj.2013.12.014
 36. Wu J, Alsaeed ES, Barrett J, Hall M, Cowan C, Gale CP. Prescription of oral anticoagulants and antiplatelets for stroke prophylaxis in atrial fibrillation: nationwide time series ecological analysis. *Europace*. 2020;22:1311–1319. doi: 10.1093/europace/eaab126
 37. Lane DA, Meyerhoff J, Rohner U, Lip GYH. Atrial fibrillation patient preferences for oral anticoagulation and stroke knowledge: Results of a conjoint analysis. *Clin Cardiol*. 2018;41:855–861. doi: 10.1002/clc.22971
 38. Wilke T, Bauer S, Mueller S, Kohlmann T, Bauersachs R. Patient preferences for oral anticoagulation therapy in atrial fibrillation: a systematic literature review. *Patient*. 2017;10:17–37. doi: 10.1007/s40271-016-0185-9
 39. Adeboyeje G, Sylwestrzak G, Barron JJ, White J, Rosenberg A, Abarca J, Crawford G, Redberg R. Major bleeding risk during anticoagulation with warfarin, dabigatran, apixaban, or rivaroxaban in patients with nonvalvular atrial fibrillation. *J Manag Care Spec Pharm*. 2017;23:968–978. doi: 10.18553/jmcp.2017.23.9.968
 40. ClinicalTrials.gov. NCT02998905. NOACs for stroke prevention in patients with atrial fibrillation and previous ICH (NASPAF-ICH). 2016.
 41. ClinicalTrials.gov. NCT03186729. Study of antithrombotic treatment after intracerebral haemorrhage (STATICH). 2017.
 42. ClinicalTrials.gov. NCT03907046. Anticoagulation in ICH survivors for stroke prevention and recovery (ASPIRE). 2019.
 43. ClinicalTrials.gov. NCT03996772. Prevention of stroke in intracerebral haemorrhage survivors with atrial fibrillation (PRESTIGE-AF). 2019.
 44. ClinicalTrials.gov. NCT03950076. Edoxaban for intracranial hemorrhage survivors with atrial fibrillation (ENRICH-AF). 2019.
 45. ClinicalTrials.gov. NCT04891861. Restart TICrH alpha pilot protocol, restarting doacs after traumatic intracranial hemorrhage. 2021.