



LJMU Research Online

Bidula, S, Sexton, DW and Schelenz, S

Serum opsonin ficolin-A enhances host-fungal interactions and modulates cytokine expression from human monocyte-derived macrophages and neutrophils following *Aspergillus fumigatus* challenge.

<http://researchonline.ljmu.ac.uk/id/eprint/2011/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Bidula, S, Sexton, DW and Schelenz, S (2015) Serum opsonin ficolin-A enhances host-fungal interactions and modulates cytokine expression from human monocyte-derived macrophages and neutrophils following *Aspergillus fumigatus* challenge. *Medical Microbiology Immunology*. ISSN

LJMU has developed [LJMU Research Online](http://researchonline.ljmu.ac.uk/) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

1 **Serum opsonin ficolin-A enhances host-fungal interactions and modulates**
2 **cytokine expression from human monocyte-derived macrophages and**
3 **neutrophils following *Aspergillus fumigatus* challenge**

4 Stefan Bidula ^{1,4}, Darren W. Sexton ^{1,2} and Silke Schelenz ^{1,3}, †

5 ¹ Biomedical Research Centre, Norwich Medical School, University of East Anglia, Norwich
6 Research Park, Norwich NR4 7TJ, UK

7 ² School of Pharmacy and Biomolecular Science, Liverpool John Moores University, Byrom
8 Street, Liverpool, L3 3AF, UK

9 ³ Department of Microbiology, Royal Brompton Hospital, Sydney Street, London, SW3 6NP,
10 UK

11 ⁴ Aberdeen Fungal Group, School of Medical Sciences, Institute of Medical Sciences,
12 University of Aberdeen, Aberdeen, AB25 2ZD UK.

13

14

15

16

17

18

19

20

21 †Corresponding author. Royal Brompton Hospital, Department of Microbiology, Sydney
22 Street, London, SW3 6NP. Tel: +44 (0)20 7352 8121. E-mail:sschelenz@doctors.org.uk

23 **Abstract**

24 Invasive aspergillosis is a devastating invasive fungal disease associated with a high mortality
25 rate in the immunocompromised; such as leukaemia patients, transplant patients and those
26 with HIV/AIDS. The rodent serum orthologue of human L-ficolin, ficolin-A, can bind to and
27 opsonize *A. fumigatus*, the pathogen that causes invasive aspergillosis, and may participate in
28 fungal defence. Using human monocyte-derived macrophages and neutrophils isolated from
29 healthy donors, we investigated conidial association and fungal viability by flow cytometry
30 and microscopy. Additionally, cytokine production was measured via cytometric bead arrays.
31 Ficolin-A opsonization was observed to significantly enhance association of conidia, while
32 also inhibiting hyphal growth and contributing to increased fungal killing following
33 incubation with monocyte-derived macrophages and neutrophils. Additionally, ficolin-A
34 opsonization was capable of manifesting a decrease in IL-8, IL-1 β , IL-6, IL-10 and TNF- α
35 production from MDM and IL-1 β , IL-6 and TNF- α from neutrophils 24 h post-infection. In
36 conclusion, rodent ficolin-A is functionally comparable to human L-ficolin and is capable of
37 modulating the innate immune response to *A. fumigatus*, down-regulating cytokine
38 production and could play an important role in airway immunity.

39

40 Key words; aspergillosis; macrophage; neutrophil; cytokines; innate immunity

41

42

43

44

45 **Introduction**

46 *Aspergillus fumigatus* (*A. fumigatus*) is the most common invasive mold pathogen and is the
47 primary causative species of the devastating disease, invasive pulmonary aspergillosis (IA) in
48 immunocompromised hosts. IA is an ever increasing challenge in the developed world and is
49 associated with a mortality rate of around 30-40% if treated and 100% if left untreated [1].

50 Those most at risk of disease include leukaemia sufferers, solid and haematopoietic transplant
51 patients, neutropenic and diabetic patients or those undergoing corticosteroid therapies [2-4].

52 *A. fumigatus* infects via the propagation of its conidia (spores) in to the air, which are then
53 inhaled by the host. Due to the small size of the conidia, those that evade mucocilliary
54 clearance can invade the depths of the alveolar space where they can germinate into
55 filamental (hyphal) structures. This invasive stage can lead to thrombosis, tissue necrosis and
56 dissemination to other organs such as the skin and brain, leading to death [5,6].

57 Defence is initiated by the innate immune system and predominantly comprises of type II
58 epithelial cells, alveolar macrophages and neutrophils. Macrophages are traditionally
59 acknowledged to be essential in the phagocytosis and removal of conidia, whereas
60 neutrophils are necessary for protection against the invasive hyphal stage whereby they
61 produce fungistatic neutrophil extracellular traps (NETs) and degranulate, releasing
62 antimicrobial compounds [7-10]. Serum opsonins such as ficolins, proteins similar to the
63 collectins mannose-binding lectin (MBL) and the surfactant proteins (SP)-A and -D, can
64 assist and enhance the functions of these host cells.

65 Ficolins are novel opsonins composed of an N-terminal collagen-like domain and a C-
66 terminal fibrinogen-like (FBG) domain with lectin activity predominantly for the acetylated
67 carbohydrate, *N*-acetylglucosamine (GlcNAc). However, ficolins do not bind exclusively to
68 acetylated carbohydrates and can recognise pathogen specific structures such as (1,3)- β -D-

69 glucan, lipotechoic acid and LPS, in addition to interacting with acute phase proteins such as
70 C-reactive protein (CRP) and pentraxin-3 (PTX3) [11]. Humans possess three types of
71 ficolin; the membrane-bound M-ficolin and the serum ficolins L-ficolin and H-ficolin. Only
72 orthologues of human L-ficolin and M-ficolin can be found in rodents and are termed ficolin-
73 A and ficolin-B, respectively.

74 Others have demonstrated that ficolin-A can recognise *A. fumigatus* conidia and we have
75 recently demonstrated that ficolin-A is capable of recognizing *A. fumigatus* conidia,
76 increasing the quantity of conidia associating with the A549 type II epithelial cell line and
77 magnifying the production of pro-inflammatory IL-8 [12,13]. However, the protective roles
78 of ficolins and whether opsonization can enhance phagocytosis by professional phagocytes
79 during *Aspergillus* defence is poorly characterised. Therefore, we utilized ficolin-A and
80 investigated if ficolin-A opsonization of *A. fumigatus* could enhance conidial associations,
81 contribute to killing or modulate inflammatory cytokine production following incubation with
82 human macrophages and neutrophils.

83

84

85

86

87

88

89

90 **Materials and Methods**

91 *Fungal pathogens*

92 A clinical isolate of *A. fumigatus* was used as described previously [14]. Resting conidia
93 were obtained after *A. fumigatus* was subcultured on Sabouraud dextrose agar (Oxoid,
94 Basingstoke, UK) at 37 °C for 7 days, and conidia were harvested using sterile physiological
95 saline (Oxoid, Basingstoke, UK). Resting live conidia were used immediately or fixed in 4%
96 PBS–formaldehyde for 10 min at room temperature (RT), washed, and resuspended in PBS.
97 Fixed *A. fumigatus* conidia were stored at 4 °C for up to 1 month until further use. To
98 fluorescently label *A. fumigatus* conidia, up to 10⁹ particles were incubated for 24 h at 4°C
99 with 0.5 mg mL⁻¹ fluorescein isothiocyanate (FITC; Sigma) in FITC labelling buffer (500
100 mM NaHCO₃, 500 mM Na₂CO₃, 0.9% (w/v) NaCl), washed repeatedly with PBS and
101 centrifuged at 775 g for 4 min until the supernatant was clear in colour.

102

103 *Ethical approval*

104 Ethical approval for blood donation by healthy participants was obtained from the Faculty of
105 Health Research Ethics Committee (Ref. Mechanisms of airway diseases – 2008042). Blood
106 was acquired through venepuncture of healthy participants who gave informed consent at the
107 time of collection. All donors were not on medication at the time of collection. All
108 procedures performed in studies involving human participants were in accordance with the
109 ethical standards of the institutional and/or national research committee and with the 1964
110 Helsinki declaration and its later amendments or comparable ethical standards.

111

112 *Cells and reagents*

113 All experiments were conducted using human monocyte-derived macrophages (MDM) or
114 peripheral blood neutrophils. MDM and neutrophils were isolated from healthy donor blood
115 via a 68% percoll gradient modified from Walsh et al (1999) [15]. Monocytes were selected
116 for by adherence to tissue culture plastic ware for 1h and differentiated in RPMI-1640
117 supplemented with 10% autologous human serum and 50 I.U ml⁻¹ penicillin and 50 µg ml⁻¹
118 streptomycin over 5-9 days. Granulocytes were removed from a separate Percoll layer and
119 were stained with Kimura to determine the percentage of neutrophils in the population.
120 Neutrophils were briefly maintained in RPMI-1640 supplemented with 10% heat-inactivated
121 foetal calf serum and 50 I.U mL⁻¹ penicillin and 50 µg ml⁻¹ streptomycin (All purchased from
122 Invitrogen, Paisley, UK). Granulocyte preparations containing greater than 90% neutrophils
123 and exhibiting >98% viability (as determined by trypan blue staining) were placed in culture.
124 Recombinant ficolin-A was produced from CHO-DXB11 cells containing ficolin-A cDNA
125 and purified by affinity chromatography on *N*-acetylglucosamine (GlcNAc)-Sepharose
126 columns as previously described [16]. Ficolin-A protein was eluted with 50 mM Tris-HCl
127 (pH 7.5) containing 150 mM NaCl and 300 mM GlcNAc (Sigma-Aldrich, United Kingdom).
128 Finally, GlcNAc was removed by dialysis and ficolin-A presence confirmed by SDS-PAGE.
129 This was kindly provided by the laboratory of Prof. Russell Wallis (University of Leicester,
130 UK). Experiments were all performed in serum-free conditions and data are representative of
131 a minimum of three independent experiments.

132

133 *Association assays*

134 MDM or human neutrophils were seeded in 24-well plates (Thermo Scientific,
135 Loughborough, UK) in supplemented RPMI-1640 at 37 °C in a 5% CO₂ atmosphere. FITC-
136 labelled fixed *A. fumigatus* conidia (FL1-A) were opsonized with 5 µg mL⁻¹ ficolin-A for 1 h
137 at 37°C. Ficolin-A-opsonized *A. fumigatus* conidia (5 x 10⁵) were incubated for 2 h with

138 adherent MDM or neutrophils in suspension (ratio of conidia to cells of 5:1) at pH 5.7 or pH
139 7.4 at 37 °C in a 5% CO₂ atmosphere. Following incubation, non-adherent conidia and cells
140 were removed and the adherent cells were washed with warm supplemented RPMI 1640.
141 Adherent cells were subsequently removed by the use of trypsin-EDTA and gentle trituration
142 prior to washing. Cells were fixed in 4% PBS–formaldehyde for 10 min at RT and analyzed
143 by flow cytometry using a BD Accuri C6 flow cytometer with BD CFlow software. The
144 percentage of MDMs and neutrophils associated with *A. fumigatus* conidia was determined
145 by gating on the cell populations based upon their size and granularity (forward scatter
146 channel/side scatter channel [FSC/SSC]). Additionally, the cells staining positive for conidia-
147 derived FITC fluorescence were gated to quantify conidial association with cells (FL1-A
148 [Exλ 488 nm, Emλ 530/33 nm]).

149

150 *Fungal viability assays*

151 Fungal viability was measured using a LIVE/DEAD[®] viability kit (Invitrogen, Paisley, UK).
152 Assays were conducted as outlined in the protocol. In brief, live resting conidia were
153 opsonised with 5 µg mL⁻¹ prior to incubation with MDM or neutrophils for 6 h. Following
154 cell challenge, conidia were stained with 15 µM FUN-1 cell stain and incubated in the
155 absence of light for 30 min at 30 °C. FUN-1 stain is metabolised inside the fungus from
156 yellow-green to orange-red fluorescence if the fungi are metabolically active and possess
157 intact membranes. Dead fungi display diffuse bright green staining. Following incubation,
158 supernatants were aspirated and the cells were washed with warm supplemented RPMI-1640.
159 Adherent cells were subsequently removed by the use of trypsin/EDTA and gentle trituration.
160 Cells and supernatants were fixed in 4% PBS/formaldehyde for 10 min at RT. Fungal
161 viability was quantitated by flow cytometry using a BD Accuri C6 flow cytometer with BD
162 CFlow[®] Software and measuring FL1-A (Exλ 488 nm, Emλ 530/33 nm). Microscopic

163 analysis was conducted after 24 h and growth was observed using an Axiovert 40 CFL
164 microscope (Zeiss) at 10x objective.

165

166 *Cytokine determination*

167 Cytokine production from supernatants was quantitated using a BD cytometric bead array
168 (CBA) Human Inflammatory Cytokines kit (BD Biosciences, Oxford, UK). Assays were
169 conducted as outlined in the protocol. In brief, live un-opsonized or ficolin-A opsonized
170 conidia (ratio of conidia to cells of 5:1) were added to MDM or neutrophils for 24 h prior to
171 collection of the supernatant and storage at -80°C. Capture beads for the measurement of IL-
172 8, IL-1 β , IL-6, IL-10 and TNF- α were mixed together prior to their addition to the
173 supernatant sample and standard tubes. Following the addition of capture beads to the
174 samples, Human Inflammatory Cytokine PE Detection Reagent was added to all tubes and
175 incubated for 3 h in the absence of light. Following incubation, samples were washed in wash
176 buffer for 5 min at 200 g prior to aspiration of the supernatant, re-suspension in wash buffer
177 and flow cytometry (Ex λ 488 nm, Em λ 585/40nm) and (Ex λ 633 nm, Em λ 780/30 nm) on a
178 BD Accuri C6 flow cytometer with BD CFlow[®] Software, collecting 1500 events as outlined
179 in the protocol.

180

181 *Statistical analysis*

182 Results were expressed as mean \pm SD. Descriptive and two-tailed Students *t*-test analyses
183 were performed using GraphPad prism software (version 5). One-way ANOVA's were
184 performed using SigmaStat software (version 3.5). A value of $p < 0.05$ was considered
185 statistically significant.

186 **Results**

187 *Ficolin-A opsonization enhances association of A. fumigatus with human monocyte-derived*
188 *macrophages and contributes to enhancing fungal killing.*

189 Macrophages are the resident lung phagocytes responsible for the removal of *A. fumigatus*
190 conidia. Therefore, we investigated whether ficolin-A opsonization of *A. fumigatus* could
191 enhance association with macrophages and contribute to fungal killing. Experiments were
192 conducted both at pH 5.7 conditions as well as pH 7.4, as we had previously observed acidic
193 conditions to be optimal for ficolin binding [12].

194 MDM were gated based on size and granularity (FSC-A/SSC-A; Fig. 1A) and association
195 was determined by the detection of FITC positive cells (Fig. 1B and C). We observed that the
196 percentage of the MDM population staining positive for conidial association was not altered
197 (Fig. 1D). However, we observed an increase in fluorescence within the gated cell population
198 suggesting that the number of FITC-labelled conidia associated with the MDM was
199 significantly enhanced following ficolin-A opsonization in acidic conditions (based upon the
200 median fluorescence intensity) (Fig. 1E; $p=0.00728$).

201 Next, we investigated the importance of ficolin-A opsonization to potentiate the ability of
202 MDM to kill *A. fumigatus* conidia. Both the free and cell-associated *A. fumigatus* were gated
203 based on size and granularity (Fig. 1F). Fungal viability assays demonstrated a significant
204 reduction in fungal viability (demonstrated by an increase in green fluorescence) following
205 opsonization by ficolin-A in acidic conditions (pH 5.7), as observed by flow cytometry (Fig.
206 1G; $p=2.2 \times 10^{-5}$).

207

208 *Ficolin-A opsonization enhances association of A. fumigatus with human neutrophils and*
209 *contributes to reducing fungal viability*

210 Neutropenic patients are at significant risk for invasive aspergillosis, highlighting the
211 importance of neutrophils in *Aspergillus* defence. In particular they are integral in the
212 destruction of the invasive hyphal stage. Therefore, we also investigated whether ficolin-A
213 opsonization of *A. fumigatus* could enhance phagocytosis and contribute to the reduction of
214 fungal viability by human neutrophils that had been isolated from healthy donor blood.

215

216 Neutrophils were gated based on size and granularity (FSC-A/SSC-A; Fig. 2A) and
217 association was determined by the detection of FITC positive cells (Fig. 2B and C). As for
218 the macrophages, flow cytometry analysis indicated no difference in the percentage of the
219 neutrophil population associated with conidia (Fig. 2D). Again however, a significant
220 increase in the fluorescence indicated that the number of FITC-labelled conidia associated with
221 the neutrophils (based upon the median fluorescence intensity) was observed following
222 ficolin-A opsonization in acidic conditions (Fig. 2E; $p=0.03215$).

223

224 As before, we were interested in investigating fungal viability following neutrophil challenge
225 with un-opsonized or ficolin-A opsonized conidia. Light microscopy indicated that in the
226 absence of ficolin-A in pH 5.7 or pH 7.4 conditions, hyphal growth was very dense (Fig. 2F,
227 G). Conversely, following challenge with ficolin-A opsonized conidia in pH 5.7 conditions,
228 hyphal growth was significantly reduced and branching much less dense (Fig. 2H). A
229 minimal reduction in hyphal growth was also observed following ficolin-A opsonization in
230 pH 7.4 conditions (Fig. 2I).

231

232 Additionally, the free and cell-associated *A. fumigatus* were gated based on size and
233 granularity (Fig. 2J), fungal viability assays demonstrated a significant decrease in fungal
234 viability following opsonization by ficolin-A in pH 5.7 conditions, as observed by flow
235 cytometry (Fig. 2K; $p=0.00262$).

236

237 *Ficolin-A opsonization modulates the secretion of inflammatory cytokines from MDM and*
238 *neutrophils in response to A.fumigatus*

239 Previous observations demonstrated the ability of ficolin-A opsonization to elicit an increase
240 in IL-8 production from the A549 cell line following live conidial challenge [12]. Whether
241 ficolin-A could modulate cytokine production from other cell types involved in *Aspergillus*
242 defence was poorly understood. Therefore, we utilised cytometric bead arrays and measured
243 the concentration of IL-8, IL-1 β , IL-6, IL-10 and TNF- α produced by MDM and neutrophils
244 following challenge with un-opsonized or ficolin-A opsonized live *A. fumigatus* conidia.

245

246 Indeed, challenge of MDM with ficolin-A opsonized conidia led to a significant reduction in
247 production of all of the cytokines tested (Fig. 3A-E). Ficolin-A itself in the absence of
248 conidia was capable of stimulating an increase in IL-6, IL-10 and TNF- α production (Fig.
249 3C-E). Additionally, following challenge of neutrophils with ficolin-A opsonized conidia we
250 observed a significant reduction in the production of IL-1 β , IL-6 and TNF- α (Fig. 4B, C, E).
251 As with the MDM, unbound ficolin-A was capable of stimulating a significant increase in the
252 production of IL-8, IL-1 β , and TNF- α from neutrophils (Fig. 4A, B, E).

253

254

255

256 **Discussion**

257 Our study focused on whether opsonization of *A. fumigatus* by ficolin-A could significantly
258 enhance *Aspergillus*-phagocyte interactions and modulate cytokine production. This led to a
259 few new observations. Firstly, ficolin-A opsonization enhanced the quantity of *A. fumigatus*
260 conidia associated with MDM and neutrophils. Next, ficolin-A opsonization significantly
261 inhibited hyphal growth and contributed to the reduction of fungal viability in the presence of
262 phagocytes. Finally, ficolin-A opsonization manifested a reduction in inflammatory cytokine
263 production from MDM and neutrophils. This led us to postulate that ficolin-A could play an
264 important role in airway immunity via efficient recognition and removal of *A. fumigatus* and
265 modulation of cytokine production.

266 We recently demonstrated that in an acidic pH representative of infection/inflammation, the
267 affinity of ficolin-A binding to *A. fumigatus* conidia was greatly increased and this increase in
268 recognition subsequently led to enhanced conidia-epithelium interactions [12]. However, the
269 interactions of ficolin-A opsonized *A. fumigatus* with professional phagocytes were unknown.

270 Following inhalation of *A. fumigatus*, macrophages are essential at ameliorating the early
271 stages of infection whereby conidia are phagocytosed and destroyed in acidic
272 phagolysosomes [17,18]. Conversely, neutrophils are traditionally recognised to be recruited
273 to the site of infection during the later stages (in response to IL-8) where they assist the
274 inhibition of fungal invasion by degranulation and the production of fungistatic NETs
275 following adherence to the hyphal cell wall, albeit recently they have also been implicated in
276 the early phagocytosis of conidia [10,19-21].

277 Initial observations indicated that ficolin-A was capable of enhancing conidial associations
278 with the murine macrophage cell line Raw 264.7 (data not shown) which prompted the use of
279 primary cells. Additionally, ficolin-A was also capable of enhancing the association of

280 conidia by both human MDM and neutrophils but only in acidic conditions. We have recently
281 described that the binding of ficolin-A to *A. fumigatus* occurs with greatest affinity in acidic
282 conditions, therefore functional enhancement in these conditions comes as no surprise [12].
283 Potentiation of cell-microbe interactions is not uncommon amongst serum opsonins, as the
284 human orthologue of ficolin-A, L-ficolin, in addition to the functionally and structurally
285 similar molecules MBL, SP-A and SP-D, have all been observed to enhance phagocytosis of
286 pathogenic microorganisms by macrophages and polymorphonuclear cells [22,23,12,24-27].

287 Although phagocytosis is important, conidial killing is even more crucial to sterilize infected
288 tissues. Therefore we utilised fungal viability assays and investigated the contribution of
289 ficolin-A to killing. In our study, we demonstrated that ficolin-A opsonization enhanced the
290 ability of macrophages and neutrophils to significantly inhibit hyphal growth and reduce
291 fungal viability. The most striking observation was how significantly ficolin-A opsonization
292 reduced hyphal density following incubation with neutrophils. Conversely, the presence of
293 ficolin-A almost completely inhibited germination when incubated with MDM.

294 Recent *in vitro* and patient evidence has highlighted that L-ficolin can also enhance
295 opsonophagocytic killing by human MDM in addition to neutrophils and is present in the
296 fungal infected lung [28]. Moreover, L-ficolin can interact with the acute phase protein,
297 PTX3 to enhance recognition and complement deposition [29], indicating a role in antifungal
298 immunity. However, to date, no *in vivo* studies have been conducted confirming the
299 importance of ficolin-A in opsonophagocytosis and defence against *A. fumigatus*. However,
300 there is a body of evidence describing the integral role of the related opsonins, SP-D and
301 MBL, in airway immunity *in vivo* [25,27,30]. Deficiencies in human L-ficolin have also been
302 observed to exacerbate susceptibility to recurrent respiratory infections, suggesting that its
303 rodent orthologue, ficolin-A, could also potentially play a significant role in airway immunity

304 in a transgenic mouse model [31]. This is something we are currently investigating in our
305 laboratory.

306 As indicated earlier, we have previously observed that ficolin-A opsonization leads to an
307 increase in IL-8 secretion from A549 cells, a cytokine that is crucial for the recruitment of
308 neutrophils during *Aspergillus* infection. Cytokines are key regulators of inflammation and
309 play an essential role in the defence against fungal challenge. Conidial challenge can induce
310 the production of a wide array of cytokines and chemokines including IL-1 β , IL-2, IL-5, IL-6,
311 IL-8, IL-10, IL-13, IL-17A, IL-22, IFN- γ , TNF- α , GM-CSF and MCP-1 [20,12,32-37]. In
312 this study we highlighted that MDM challenged with ficolin-A opsonized conidia produced
313 significantly less IL-8, IL-1 β , IL-6, IL-10 and TNF- α , whereas only IL-1 β , IL-6 and TNF- α
314 production was lower from neutrophils. Ficolins have been described to have the potential to
315 be both pro- or anti-inflammatory, dependent upon the cell type involved [38,12,39,40]. We
316 have previously observed that L-ficolin opsonization can modulate an anti-inflammatory
317 response from MDM and neutrophils, while additionally, others have reported that ficolin-A
318 can dampen LPS-induced inflammatory responses on mast cells [39,28]. In addition, we
319 again show that unbound ficolin-A can increase cytokine production. The mechanisms by
320 which these effects are achieved are currently unknown but preliminary studies in our lab
321 have indicated the potential importance of the MAPK signalling cascades and TLRs.

322 In conclusion, we demonstrate that ficolin-A plays an important role in potentiating the
323 functions of macrophages and neutrophils against *A. fumigatus* challenge *in vitro* and is
324 functionally comparable to human L-ficolin. Additionally, we highlight that ficolin-A may be
325 important in the down-regulation of cytokine production post-infection but relevant ficolin-A
326 *in vivo* studies are lacking.

327

328 **Acknowledgements**

329 We are grateful to Dr. Orla Jupp (University of East Anglia), for processing blood and
330 providing us with monocytes and neutrophils. We would also like to thank Prof. Russell
331 Wallis and Dr. Umakhanth Venkatraman Girija for supplying us with recombinant rat ficolin-
332 A. This work was supported by a Faculty of Health, University of East Anglia, UK PhD
333 studentship (FMH 04.4.66 C4).

334 **Conflict of interest**

335 The authors have no conflicts of interest.

336

337 **References**

- 338 1. Segal BH (2009) Aspergillosis. *N Engl J Med* 360 (18):1870-1884.
339 doi:10.1056/NEJMra0808853
- 340 2. Denning DW, Pleuvry A, Cole DC (2013) Global burden of allergic bronchopulmonary
341 aspergillosis with asthma and its complication chronic pulmonary aspergillosis in adults. *Med*
342 *Mycol* 51 (4):361-370. doi:10.3109/13693786.2012.738312
- 343 3. Steinbach WJ, Marr KA, Anaissie EJ, Azie N, Quan SP, Meier-Kriesche HU, Apewokin S,
344 Horn DL (2012) Clinical epidemiology of 960 patients with invasive aspergillosis from the
345 PATH Alliance registry. *J Infect* 65 (5):453-464. doi:10.1016/j.jinf.2012.08.003
- 346 4. Taccone FS, Van den Abeele AM, Bulpa P, Misset B, Meersseman W, Cardoso T, Paiva
347 JA, Blasco-Navalpotro M, De Laere E, Dimopoulos G, Rello J, Vogelaers D, Blot SI, Asp
348 ICUSI (2015) Epidemiology of invasive aspergillosis in critically ill patients: clinical
349 presentation, underlying conditions, and outcomes. *Critical care (London, England)* 19 (1):7.
350 doi:10.1186/s13054-014-0722-7
- 351 5. Martin TR, Frevert CW (2005) Innate immunity in the lungs. *Proc Am Thorac Soc* 2:403-
352 411

- 353 6. Schelenz S, Goldsmith DJ (2003) *Aspergillus* endophthalmitis: an unusual complication of
354 disseminated infection in renal transplant patients. *J Infect* 47 (4):336-343
- 355 7. Luther K, Torosantucci A, Brakhage AA, Heesemann J, Ebel F (2007) Phagocytosis of
356 *Aspergillus fumigatus* conidia by murine macrophages involves recognition by the dectin-1
357 beta-glucan receptor and Toll-like receptor 2. *Cell Microbiol* 9 (2):368-381.
358 doi:10.1111/j.1462-5822.2006.00796.x
- 359 8. Aratani Y, Kura F, Watanabe H, Akagawa H, Takano Y, Suzuki K, Dinauer MC, Maeda
360 N, Koyama H (2002) Relative contributions of myeloperoxidase and NADPH-oxidase to the
361 early host defense against pulmonary infections with *Candida albicans* and *Aspergillus*
362 *fumigatus*. *Med Mycol* 40 (6):557-563
- 363 9. Segal BH, Han W, Bushey JJ, Joo M, Bhatti Z, Feminella J, Dennis CG, Vethanayagam
364 RR, Yull FE, Capitano M, Wallace PK, Minderman H, Christman JW, Sporn MB, Chan J,
365 Vinh DC, Holland SM, Romani LR, Gaffen SL, Freeman ML, Blackwell TS (2010) NADPH
366 oxidase limits innate immune responses in the lungs in mice. *PLoS ONE* 5 (3):e9631.
367 doi:10.1371/journal.pone.0009631
- 368 10. Bruns S, Kniemeyer O, Hasenberg M, Aimanianda V, Nietzsche S, Thywissen A, Jeron
369 A, Latge JP, Brakhage AA, Gunzer M (2010) Production of extracellular traps against
370 *Aspergillus fumigatus* in vitro and in infected lung tissue is dependent on invading
371 neutrophils and influenced by hydrophobin RodA. *PLoS Pathog* 6 (4):e1000873.
372 doi:10.1371/journal.ppat.1000873
- 373 11. Matsushita M (2013) Ficolins in complement activation. *Mol Immunol* 55 (1):22-26.
374 doi:10.1016/j.molimm.2012.08.017
- 375 12. Bidula S, Kenawy H, Ali Y, Sexton D, Schwaeble W, Schelenz S (2013) Role of ficolin-
376 A and lectin complement pathway in the innate defense against pathogenic *Aspergillus*
377 species. *Infect Immun* 81:1730-1740

- 378 13. Hummelshøj T, Ma YJ, Munthe-Fog L, Bjarnsholt T, Moser C, Skjødt K, Romani L,
379 Fujita T, Endo Y, Garred P (2012) The interaction pattern of murine serum ficolin-A with
380 microorganisms. *PLoS ONE* **7** (5). doi:10.1371/journal.pone.0038196
- 381 14. Bidula S, Sexton D, Schelenz S (2013) Serum ficolins enhance phagocytosis and killing
382 of *Aspergillus fumigatus* leading to modulation of the pro-inflammatory cytokine response.
383 *Immunology* **140** ((Suppl. 1)):121
- 384 15. Walsh GM, Sexton DW, Blaylock MG, Convery CM (1999) Resting and cytokine-
385 stimulated human small airway epithelial cells recognize and engulf apoptotic eosinophils.
386 *Blood* **94** (8):2827-2835
- 387 16. Girija UV, Dodds AW, Roscher S, Reid KBM, Wallis R (2007) Localization and
388 characterisation of the mannose-binding lectin (MBL)-associated-serine-protease-2 binding
389 site in rat ficolin-A; equivalent binding sites within the collagenous domains of MBLs and
390 ficolins. *J Immunol* **179**:455-462
- 391 17. Romani L (2004) Immunity to fungal infections. *Nat Rev Immunol* **4** (1):1-23.
392 doi:10.1038/nri1255
- 393 18. Ibrahim-Granet O, Philippe B, Boleti H, Boisvieux-Ulrich E, Grenet D, Stern M, Latge JP
394 (2003) Phagocytosis and intracellular fate of *Aspergillus fumigatus* conidia in alveolar
395 macrophages. *Infect Immun* **71** (2):891-903
- 396 19. Bianchi M, Hakkim A, Brinkmann V, Siler U, Seger RA, Zychlinsky A, Reichenbach J
397 (2009) Restoration of NET formation by gene therapy in CGD controls aspergillosis. *Blood*
398 **114** (13):2619-2622. doi:10.1182/blood-2009-05-221606
- 399 20. Schelenz S, Smith DA, Bancroft GJ (1999) Cytokine and chemokine responses following
400 pulmonary challenge with *Aspergillus fumigatus*: obligatory role of TNF-alpha and GM-CSF
401 in neutrophil recruitment. *Med Mycol* **37** (3):183-194

- 402 21. Mircescu MM, Lipuma L, van Rooijen N, Pamer EG, Hohl TM (2009) Essential role for
403 neutrophils but not alveolar macrophages at early time points following *Aspergillus*
404 *fumigatus* infection. *J Infect Dis* 200 (4):647-656. doi:10.1086/600380
- 405 22. Matsushita M, Endo Y, Taira S, Sato Y, Fujita T, Ichikawa N, Nakata M, Mizuochi T
406 (1996) A novel human serum lectin with collagen- and fibrinogen-like domains that functions
407 as an opsonin. *J Biol Chem* 271:2448-2454
- 408 23. Fujieda M, Aoyagi Y, Matsubara K, Takeuchi Y, Fujimaki W, Matsushita M, Bohnsack
409 JF, Takahashi S (2012) L-ficolin and capsular polysaccharide-specific IgG in cord serum
410 contribute synergistically to opsonophagocytic killing of serotype III and V group B
411 streptococci. *Infect Immun* 80 (6):2053-2060
- 412 24. Allen MJ, Harbeck R, Smith B, Voelker DR, Mason RJ (1999) Binding of rat and human
413 surfactant proteins A and D to *Aspergillus fumigatus* conidia. *Infect Immun* 67 (9):4563-4569
- 414 25. Madan T, Kishore U, Singh M, Strong P, Hussain EM, Reid KB, Sarma PU (2001)
415 Protective role of lung surfactant protein D in a murine model of invasive pulmonary
416 aspergillosis. *Infect Immun* 69 (4):2728-2731. doi:10.1128/iai.69.4.2728-2731.2001
- 417 26. Madan T, Eggleton P, Kishore U, Strong P, Aggrawal SS, Sarma PU, Reid KB (1997)
418 Binding of pulmonary surfactant proteins A and D to *Aspergillus fumigatus* conidia enhances
419 phagocytosis and killing by human neutrophils and alveolar macrophages. *Infect Immun* 65
420 (8):3171-3179
- 421 27. Kaur S, Gupta VK, Thiel S, Sarma PU, Madan T (2007) Protective role of mannan-
422 binding lectin in a murine model of invasive pulmonary aspergillosis. *Clin Exp Immunol* 148
423 (2):382-389. doi:10.1111/j.1365-2249.2007.03351.x
- 424 28. Bidula S, Sexton DW, Abdolrasouli A, Shah A, Reed A, Armstrong-James D, Schelenz S
425 (2015) Serum opsonin, L-ficolin, is detected in human lungs of transplant patients following

- 426 fungal infections and modulates inflammation and killing of *A. fumigatus*. *J Infect Dis*.
427 doi:10.1093/infdis/jiv027
- 428 29. Ma YJ, Doni A, Hummelshøj T, Honore C, Bastone A, Mantovani A, Thielens NM,
429 Garred P (2009) Synergy between ficolin-2 and pentraxin 3 boosts innate immune
430 recognition and complement deposition. *J Biol Chem* **284** (41):28263-28275
- 431 30. Lambourne J, Agranoff D, Herbrecht R, Troke PF, Buchbinder A, Willis F, Letscher-Bru
432 V, Agrawal S, Doffman S, Johnson E, White PL, Barnes RA, Griffin G, Lindsay JA,
433 Harrison TS (2009) Association of mannose-binding lectin deficiency with acute invasive
434 aspergillosis in immunocompromised patients. *Clin Infect Dis* 49 (10):1486-1491.
435 doi:10.1086/644619
- 436 31. Cedzynski M, Atkinson A, Swierzko A, MacDonald S, Szala A, Zeman K, Buczylko K,
437 Bak-Romaniszyn L, Wiszniewska M, Matsushita M, Szemraj J, Banasik M, Turner M,
438 Kilpatrick D (2009) L-ficolin (ficolin-2) insufficiency is associated with combined allergic
439 and infectious respiratory diseases in children. *Mol Immunol* **47**:415-419
- 440 32. Zhang Z, Liu R, Noordhoek J, Kauffman H (2005) Interaction of airway epithelial cells
441 (A549) with spores and mycelium of *Aspergillus fumigatus* *J Infect* **51**:375-382
- 442 33. Balloy V, Sallenave J, Wu Y, Touqui L, Latgé J, Si-Tahar M, Chignard M (2008)
443 *Aspergillus fumigatus* -induced Interleukin-8 synthesis by respiratory epithelial cells is
444 controlled by the phosphatidylinositol 3-kinase, p38 MAPK, and ERK 1/2 pathways and not
445 by the toll-like receptor-MyD88 pathway. *JBiol Chem* **283**:30513-30521
- 446 34. Koth L, Rodriguez M, Bernstein X, Chan S, Huang X, Charo I, Rollins B, Erle D (2004)
447 *Aspergillus* antigen induces robust Th2 cytokine production, inflammation, airway
448 hyperreactivity and fibrosis in the absence of MCP-1 or CCR2. *Resp Res* **5**:12
- 449 35. Graziutti ML, Rex JH, Cowart RE, Anaissie EJ, Ford A, Savary CA (1997) *Aspergillus*
450 *fumigatus* conidia induce a Th1-type cytokine response. *J Infect Dis* 176 (6):1579-1583

- 451 36. Werner JL, Gessner MA, Lilly LM, Nelson MP, Metz AE, Horn D, Dunaway CW,
452 Deshane J, Chaplin DD, Weaver CT, Brown GD, Steele C (2011) Neutrophils produce
453 interleukin 17A (IL-17A) in a dectin-1- and IL-23-dependent manner during invasive fungal
454 infection. *Infect Immun* 79 (10):3966-3977. doi:10.1128/iai.05493-11
- 455 37. Gessner MA, Werner JL, Lilly LM, Nelson MP, Metz AE, Dunaway CW, Chan YR,
456 Ouyang W, Brown GD, Weaver CT, Steele C (2012) Dectin-1-dependent interleukin-22
457 contributes to early innate lung defense against *Aspergillus fumigatus*. *Infect Immun* 80
458 (1):410-417. doi:10.1128/iai.05939-11
- 459 38. Ma YJ, Doni A, Romani L, Jurgensen HJ, Behrendt N, Mantovani A, Garred P (2013)
460 Ficolin-1-PTX3 complex formation promotes clearance of altered self-cells and modulates
461 IL-8 production. *J Immunol* 191 (3):1324-1333. doi:10.4049/jimmunol.1300382
- 462 39. Ma YJ, Kang HJ, Kim JY, Garred P, Lee MS, Lee BL (2013) Mouse mannose-binding
463 lectin-A and ficolin-A inhibit lipopolysaccharide-mediated pro-inflammatory responses on
464 mast cells. *BMB Rep* 46 (7):376-381
- 465 40. Luo F, Sun X, Wang Y, Wang Q, Wu Y, Pan Q, Fang C, Zhang XL (2013) Ficolin-2
466 Defends against Virulent Mycobacteria Tuberculosis Infection In Vivo, and Its Insufficiency
467 Is Associated with Infection in Humans. *PLoS ONE* 8 (9):e73859.
468 doi:10.1371/journal.pone.0073859

469

470

471

472

473

474

475 **Figure legends**

476 **Fig. 1.** Association and killing of ficolin-A opsonized *A. fumigatus* conidia with MDM.
477 FITC-labelled *A. fumigatus* conidia (5×10^5) were opsonized with $5 \mu\text{g mL}^{-1}$ ficolin-A prior
478 to incubation with MDM (conidia:MDM ratio of 5:1) in pH 5.7 and pH 7.4 conditions for 2 h.
479 To measure killing, live unlabelled conidia were incubated with MDM for 24 h prior to
480 LIVE/DEAD staining. **A** Gating strategy used to distinguish macrophages. **B** MDM in the
481 absence of FITC-labelled conidia. **C** MDM in the presence of FITC-labelled conidia. **D** The
482 percentage of MDM associated with conidia in pH 5.7 and pH 7.4 conditions in the presence
483 or absence of ficolin-A. **E** The relative number of associated FITC-labelled conidia (based
484 upon the median fluorescence intensity; FL1-A) either un-opsonized (-ficolin-A) or following
485 opsonization by ficolin-A (+ficolin-A). **F** Gating strategy used to measure fungal viability. **G**
486 Fungal death after incubation with MDMs based upon the median fluorescence intensity of
487 FL1-A. Heat killed conidia were used as a positive control for fungal killing. Results are
488 representative of the average of all data points gained from three independent experiments.
489 Error bars represent SD and significance was determined via two-tailed Students *t*-test. An
490 asterisk indicates a significant difference; $p < 0.05$.

491

492 **Fig. 2.** Association and killing of ficolin-A opsonized *A. fumigatus* conidia with neutrophils.
493 FITC-labelled *A. fumigatus* conidia (5×10^5) were opsonized with $5 \mu\text{g mL}^{-1}$ ficolin-A prior
494 to incubation with neutrophils (conidia:neutrophil ratio of 5:1) in pH 5.7 and pH 7.4
495 conditions for 2 h. To measure killing, live unlabelled conidia were incubated with
496 neutrophils for 24 h prior to LIVE/DEAD staining. **A** Gating strategy used to distinguish
497 neutrophils. **B** Neutrophils in the absence of FITC-labelled conidia. **C** Neutrophils in the
498 presence of FITC-labelled conidia. **D** The percentage of neutrophils associated with conidia
499 in pH 5.7 and pH 7.4 conditions in the presence or absence of ficolin-A. **E** The relative

500 number of associated FITC-labelled conidia (based upon the median fluorescence intensity;
501 FL1-A) either un-opsonized (-ficolin-A) or following opsonization by ficolin-A (+ficolin-A).
502 **F** Hyphal germination following incubation of un-opsonized conidia in pH 5.7 or. **G** in pH
503 7.4. **H** Hyphal germination following incubation of ficolin-A opsonized conidia in pH 5.7 or.
504 **I** in pH 7.4. **J** Gating strategy used to measure fungal viability. **K** Fungal death after
505 incubation with neutrophils based upon the median fluorescence intensity of FL1-A. Heta
506 killed conidia were used as a positive control for fungal killing. Results are representative of
507 the average of all data points gained from three independent experiments. Error bars represent
508 SD and significance was determined via two-tailed Students *t*-test. An asterisk indicates a
509 significant difference; $p < 0.05$.

510

511 **Fig. 3.** Inflammatory cytokine release from MDM following challenge with un-opsonized or
512 ficolin-A opsonized conidia. Supernatants were collected after 8 h and 24 h time points
513 during challenge of MDM with live *A. fumigatus* conidia (5×10^5) either un-opsonized or
514 ficolin-A opsonized ($5 \mu\text{g mL}^{-1}$) prior to the conduction of cytometric bead arrays
515 (conidia:MDM ratio of 5:1). **A** Quantity of IL-8 production. **B** Quantity of IL-1 β production.
516 **C** Quantity of IL-6 production. **D** Quantity of IL-10 production. **E** Quantity of TNF- α
517 production. Following *A. fumigatus* challenge. **MDM** is representative of MDM alone. +
518 **ficolin-A** represents MDM in the presence of ficolin-A alone. + **AF** and +**AF +ficolin-A** are
519 representative of un-opsonized *A. fumigatus* or ficolin-A opsonized conidia, respectively.
520 Results are representative of the average of all the data points gained from three independent
521 experiments. Error bars represent the SD. Significance was determined via one-way ANOVA
522 and pair-wise comparisons were conducted using the Student-Newman-Keuls method. An
523 asterisk indicates a significant difference: $p < 0.05$.

524

525 **Fig. 4.** Inflammatory cytokine release from neutrophils following challenge with un-
526 opsonized or ficolin-A opsonized conidia. Supernatants were collected after 8 h and 24 h time
527 points during challenge of neutrophils with live *A. fumigatus* conidia (5×10^5) either un-
528 opsonized or ficolin-A opsonized ($5 \mu\text{g mL}^{-1}$) prior to the conduction of cytometric bead
529 arrays (conidia:neutrophil ratio of 5:1). **A** Quantity of IL-8 production. **B** Quantity of IL-1 β
530 production. **C** Quantity of IL-6 production. **D** Quantity of IL-10 production. **E** Quantity of
531 TNF- α production. Following *A. fumigatus* challenge. **Neutrophils** is representative of
532 neutrophils alone. + **ficolin-A** represents neutrophils in the presence of ficolin-A alone. + **AF**
533 and +**AF +ficolin-A** are representative of un-opsonized *A. fumigatus* or ficolin-A opsonized
534 conidia, respectively. Results are representative of the average of all the data points gained
535 from three independent experiments. Error bars represent the SD. Significance was
536 determined via one-way ANOVA and pair-wise comparisons were conducted using the
537 Student-Newman-Keuls method. An asterisk indicates a significant difference: $p < 0.05$.
538
539