



LJMU Research Online

Bidula, S, Sexton, DW, Abdolrasouli, A, Shah, A, Reed, A, Armstrong-James, D and Schelenz, S

The serum opsonin L-ficolin is detected in lungs of human transplant recipients following fungal infections and modulates inflammation and killing of *Aspergillus fumigatus*.

<https://researchonline.ljmu.ac.uk/id/eprint/2012/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Bidula, S, Sexton, DW, Abdolrasouli, A, Shah, A, Reed, A, Armstrong-James, D and Schelenz, S (2015) The serum opsonin L-ficolin is detected in lungs of human transplant recipients following fungal infections and modulates inflammation and killing of *Aspergillus fumigatus*. *Journal of*

LJMU has developed [LJMU Research Online](#) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

1 **Serum opsonin, L-ficolin, is detected in human lungs of transplant patients**
2 **following fungal infections and modulates inflammation and killing of *A.***
3 ***fumigatus***

4 Running title: L-ficolin and immunity to *A. fumigatus*

5 Stefan Bidula,¹ Darren W. Sexton,¹ Alireza Abdolrasouli,² Anand Shah,² Anna Reed,³ Darius
6 Armstrong-James,² Silke Schelenz,^{1†}

7 ¹Biomedical Research Centre, Norwich Medical School, University of East Anglia, Norwich
8 Research Park, Norwich NR4 7TJ, UK

9 ²Section of Infectious Diseases and Immunity, Imperial College London, Flowers Building,
10 Armstrong Road, London SW7 2AZ, UK

11 ³Department of Lung Transplantation, Harefield Hospital, Hill End Road, Middlesex, UB9
12 6JH, UK

13 Abstract word count: 200

14 Manuscript word count: 3498

15

16

17

18

19 †Corresponding author. Mailing address: Biomedical Research Centre, Norwich Medical
20 School, University of East Anglia, Norwich NR4 7TJ, United Kingdom. Phone:44 1603
21 591699. Fax: 44 1603 591750. E-mail:sschelenz@doctors.org.uk

22 **FOOTNOTES**

23 This work was supported by the Medical Research Council [MR/K002708/1 to A.A, A.S,
24 A.R and D.A.J and a Faculty of Health, University of East Anglia, UK PhD studentship FMH
25 04.4.66 C4 to S.B, D.S and S.S].

26 Potential conflicts of interest. All authors have reported no conflicts. All authors have
27 submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest.

28 Presented in part: British Society for Immunology Annual Congress, 2013, Liverpool, UK
29 and British Society for Medical Mycology Annual Meeting, 2014, Manchester, UK.

30 Correspondence:

31 Corresponding author. Mailing address: Biomedical Research Centre, Norwich Medical
32 School, University of East Anglia, Norwich NR4 7TJ, United Kingdom. Phone:44 1603
33 591699. Fax: 44 1603 591750. E-mail:sschelenz@doctors.org.uk

34

35 ¹ Biomedical Research Centre, Norwich Medical School, University of East Anglia, Norwich
36 Research Park, Norwich NR4 7TJ, UK

37 ²Section of Infectious Diseases and Immunity, Imperial College London, Flowers Building,
38 Armstrong Road, London SW7 2AZ, UK

39 ³Department of Lung Transplantation, Harefield Hospital, Hill End Road, Middlesex, UB9
40 6JH, UK

41

42

43

44

45 **ABSTRACT**

46 **Background.** Invasive aspergillosis (IA) is a life threatening systemic fungal infection in the
47 immunocompromised caused by *Aspergillus fumigatus*. The human serum opsonin, L-ficolin
48 has been observed to recognise *A. fumigatus* and could participate in fungal defence.

49 **Methods.** Using lung epithelial cells, primary human monocyte-derived macrophages
50 (MDM) and neutrophils from healthy donors, we assessed phagocytosis and killing of L-
51 ficolin opsonized *A. fumigatus* live conidia by flow cytometry and microscopy. Additionally,
52 cytokines were measured by cytometric bead array and L-ficolin was measured in
53 bronchoalveolar lavage (BAL) fluid from lung transplant recipients by ELISA.

54 **Results.** L-ficolin opsonization increased conidial uptake and enhanced killing of *A.*
55 *fumigatus* by MDM and neutrophils. Opsonization was also shown to manifest an increase in
56 IL-8 release from A549 lung epithelial cells but decrease IL-1 β , IL-6, IL-8, IL-10 and TNF-
57 α release from MDM and neutrophils 24 h post-infection. The concentration of L-ficolin was
58 significantly higher in BAL of patients with fungal infection than in control subjects
59 ($p=0.00087$) and ROC curve analysis highlighted the diagnostic potential of L-ficolin for
60 lung infection (AUC=0.842; $p<0.0001$).

61 **Conclusions.** L-ficolin modulates the immune response to *A. fumigatus*. Additionally, for the
62 first time, L-ficolin has been demonstrated to be present in human lungs.

63 **Keywords.** L-ficolin, *Aspergillus fumigatus*, macrophage, neutrophil, epithelial, phagocytosis,
64 cytokines, lung transplant

65

66

67 **Introduction**

68 *Aspergillus fumigatus* (*A. fumigatus*) is a major worldwide prevalent pathogenic mold and the
69 primary cause of invasive pulmonary aspergillosis (IA) in immunocompromised hosts. [1]. In
70 those at risk such as leukaemia, solid-organ and haematopoietic transplant patients or those
71 with neutropenia, IA is associated with a mortality rate of up to 30% if treated and 100% in
72 untreated patients [2-6]. Infection is initiated following the inhalation of small hydrophobic
73 conidia from the environment which have the propensity to germinate into filamental
74 (hyphal) structures. These invade local tissues causing thrombosis, necrosis and
75 dissemination of the fungus to other organs such as the skin and brain, ultimately leading to
76 death [7-9].

77 Alveolar macrophages, neutrophils, complement and pattern recognition proteins; such as the
78 ficolins and collectins, all work synergistically to remove *Aspergillus*. The process of
79 phagocytosis by macrophages is an integral aspect in innate host defence against *A. fumigatus*
80 conidia [10, 11]. Neutrophils have also been observed to be important in the early stages of
81 conidial removal, but are essential in the destruction of the large hyphal structures following
82 degranulation and the production of neutrophil extracellular traps (NETs) [12-14].

83 Ficolins are a family of proteins composed of an N-terminal collagen-like domain and a C-
84 terminal fibrinogen-like domain with lectin activity (highly specific for *N*-acetylglucosamine
85 (GlcNAc)). Human serum L-ficolin has the potential to enhance phagocytosis via direct
86 binding to pathogens [15] but the protective roles of ficolins in *Aspergillus* defence are still
87 poorly characterised.

88 We have recently demonstrated that L-ficolin is able to enhance the binding of *Aspergillus*
89 conidia to the lung epithelium, but little is known about the functional consequences
90 following ficolin opsonization [16]. We therefore utilised L-ficolin to investigate its roles in

91 phagocytosis and killing of *A. fumigatus* by phagocytes in addition to its role in modulating
92 cytokine production. In this study we have also shown for the first time that L-ficolin is
93 present in BAL from lung transplant patients suffering from fungal pneumonia compared to
94 uninfected controls. Additionally, we highlight the potential of L-ficolin as a tool for the
95 diagnosis of fungal infections following lung transplants.

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110 **Materials and Methods**

111 *Patients and Ethical approval*

112 Evidence of fungal infection was based on clinical European Organization for Research and
113 Treatment of Cancer/Mycoses Study Group (EORTC/MSG) criteria [17]. BAL sampling of
114 lung transplant patients from the Royal Brompton and Harefield NHS Trust was performed
115 under Biomedical Research Unit ethics approval (RBH/AS1).

116 Ethical approval for blood donation by healthy participants was obtained from the Faculty of
117 Health Research Ethics Committee (Ref. Mechanisms of airway diseases – 2008042). Blood
118 was acquired through venepuncture of healthy participants who gave informed consent at the
119 time of collection. All donors were not on medication at the time of collection.

120 Informed consent was obtained from patients and human experimentation guidelines of the
121 United States Department of Health and Human Services were adhered to in the conduct of
122 clinical research.

123 *Cells and reagents*

124 All experiments were conducted using the A549 adenocarcinomic human alveolar basal
125 epithelial cell line, human monocyte-derived macrophages (MDM) or peripheral blood
126 neutrophils. MDM and neutrophils were isolated from healthy donor blood via a 68% percoll
127 gradient modified from Walsh et al (1999) [18]. Monocytes were selected for by adherence to
128 tissue culture plastic ware for 1h and differentiated in RPMI-1640 supplemented with 10%
129 autologous serum and 50 I.U mL⁻¹ penicillin and 50 µg mL⁻¹ streptomycin over 5-9 days.
130 A549 cells and neutrophils were briefly maintained in RPMI-1640 supplemented with 10%
131 heat-inactivated foetal calf serum and 50 I.U mL⁻¹ penicillin and 50 µg mL⁻¹ streptomycin.
132 Polymorphonuclear preparations containing greater than 90% neutrophils and exhibiting

133 >98% viability (as determined by trypan blue staining) were placed in culture. Experiments
134 were all performed in serum-free conditions. Recombinant L-ficolin was purchased from
135 R&D Systems. FITC was purchased from Sigma-Aldrich. A clinical *A. fumigatus* strain
136 isolated from a respiratory specimen was used in all experiments and maintained/harvested as
137 previously described [16].

138 ***Detection of infection and L-ficolin in bronchoalveolar lavage***

139 BAL fluid was collected from lung transplant recipients at Royal Brompton and Harefield
140 NHS Trust by instilling 200 mL sterile saline into distal airway segments under flexible
141 bronchoscopy. BAL return was centrifuged at 1500 rpm for 10 minutes. *Aspergillus* antigens,
142 indicative of invasive aspergillosis, were detected via the lateral-flow device as previously
143 described [19] and/or via detection of galactomannan (GM) using a Platelia™ *Aspergillus*
144 antigen kit (Bio-Rad). For BAL samples, an index of < 0.5 was considered negative, an index
145 of ≥ 0.5 was considered positive for GM [20]. Samples were tested for a panel of respiratory
146 viruses (multiplex PCR) and bacteria by culture (B57, UK standard for microbiology
147 investigations) [21]. High resolution computed tomography (HRCT) chest imaging was
148 reviewed for evidence of findings consistent with fungal infection [21]. The presence of L-
149 ficolin in the BAL fluid of lung transplant patients was detected using a ficolin-2 human
150 ELISA kit (Hycult). Patients were categorised for possible, probable and proven invasive
151 fungal infection according to revised EORTC/MSG criteria [17].

152 ***Phagocytosis assays***

153 FITC-labelled live *A. fumigatus* conidia were opsonized with $5 \mu\text{g ml}^{-1}$ L-ficolin as
154 previously described [16]. MDM or human neutrophils were seeded in 24-well plates (Nunc)
155 prior to challenge with ficolin-opsonized FITC-labelled *A. fumigatus* conidia (5×10^5 ;
156 conidia:cell ratio of 5:1) for 2 h at 37°C. Adherent cells were subsequently removed by the

157 use of trypsin/EDTA, gentle trituration and scraping. Neutrophils in suspension were pelleted
158 at 300 g for 5 mins. Cells were fixed in 4% PBS/formaldehyde for 10 min at RT before re-
159 suspension in PBS. Phagocytosis was analysed by flow cytometry (Exλ 488 nm, Emλ 533/30
160 nm) on a BD Accuri C6 flow cytometer with BD CFlow[®] Software (BD Biosciences)
161 collecting 5000 events. To yield quantitative counts, positively phagocytic cells were
162 identified and expressed as a percentage of all phagocytes present and the relative abundance
163 of conidia contained within the positively ingesting phagocytes was determined by the
164 relative fluorescence intensity (FL1-A) of the positively phagocytic cells.

165 ***Ficolin binding, visualising fungal growth and fungal killing assays***

166 L-ficolin binding assays were conducted as previously described [16] and data was collected
167 by flow cytometry as described above.

168 MDM or human neutrophils were seeded in 24-well plates prior to challenge with ficolin-
169 opsonized live *A. fumigatus* conidia (5×10^5 ; conidia:cell ratio of 5:1) for 24 h at 37°C as
170 above. Growth was observed using an Axiovert 40 CFL microscope (Zeiss) at 10x objective
171 for neutrophils and 20x objective for MDM.

172 Fungal killing was measured using a LIVE/DEAD[®] Fungal Viability Kit (Invitrogen). In
173 brief, fungi were stained with 15 μM FUN-1 prior to the measurement of green fluorescence
174 (FL1-A, Exλ 488 nm, Emλ 585/40nm), an increase in which, represents a reduction in fungal
175 viability. Fluorescence was quantified by flow cytometry as above.

176 ***Cytokine determination***

177 Cytokine protein concentrations from the supernatants of *A. fumigatus* challenged A549,
178 MDM and neutrophils were determined using a BD cytometric bead array (CBA) Human
179 Inflammatory Cytokines kit (BD Biosciences). Data was gained by flow cytometry (Exλ 488

180 nm, Em λ 585/40nm) and (Ex λ 633 nm, Em λ 780/30 nm) on a BD Accuri C6 flow cytometer
181 with BD CFlow[®] Software, collecting 1800 events as outlined in the protocol.

182 ***Statistical analysis***

183 Results were expressed as mean \pm SD. Descriptive and 2-tailed Students *t*-test analyses were
184 performed using GraphPad prism software (version 5). One-way ANOVA's were performed
185 using SigmaStat software (version 3.5). A value of $p < 0.05$ was considered statistically
186 significant. Receiver operating characteristics (ROC) curve analysis was conducted using
187 MedCalc (version 13.1.1).

188

189 **Results**

190 ***L-ficolin opsonization enhances phagocytosis and killing of *A. fumigatus* by human*** 191 ***monocyte-derived macrophages***

192 We, and others, have previously acknowledged that L-ficolin is capable of binding to
193 *A.fumigatus* [16,22]. Here we verify that L-ficolin can recognize *A.fumigatus* live conidia
194 ($p=2.7 \times 10^{-5}$; Figure 1A) and we demonstrate enhanced binding in acidic pH (5.7)
195 ($p=0.00089$; Figure 1B).

196

197 We have shown that the phagocytosis of conidia by the airway epithelial cell line, A549, is
198 enhanced following L-ficolin opsonization [16]. Another integral cell type involved in the
199 early defence against *Aspergillus* conidia are macrophages.

200 Initially, the ability of L-ficolin to enhance phagocytosis was investigated using FITC-
201 labelled *A. fumigatus* conidia opsonized with L-ficolin prior to incubation with adherent
202 human MDM for 2 hours. MDM were gated (Figure 2A) and the percentage of FITC negative

203 and positive MDM were used to identify phagocytic cells (Figure 2B and Figure 2C). The
204 proportion of phagocytic MDM was unaffected in physiological (pH 7.4) or acidic conditions
205 (pH 5.7) (Figure 2D), however, the number of FITC labelled L-ficolin opsonized conidia
206 ingested per MDM (based upon the median fluorescence intensity of phagocytic MDM) was
207 significantly enhanced in inflammatory (pH 5.7) conditions but not at pH 7.4 (Figure 2E and
207 Figure 2F) ($p=6.6 \times 10^{-5}$).

208 Additionally, light microscopy demonstrated that MDM inhibited conidial germination
209 following opsonization by L-ficolin in inflammatory conditions (Figure 2G-J). L-ficolin in
210 the absence of phagocytes had no effect on *A. fumigatus* growth (data not shown). Moreover,
211 following gating (Figure 2K), fungal viability assays demonstrated a significant increase in
212 fungal killing following opsonization by L-ficolin in these conditions; as quantitated by flow
213 cytometry ($p=0.00249$) (Figure 2L and Figure 2M). When ingested *A. fumigatus* and free
214 *A. fumigatus* populations were gated separately, death-associated green-yellow fluorescence
215 emitted by *A. fumigatus* within MDM was observed to be significantly greater compared to the
216 un-associated fungi, highlighting potent intracellular killing (Supplementary Figure 1).

218

219 ***L-ficolin opsonization enhances phagocytosis and killing of A. fumigatus by human*** 220 ***neutrophils***

221 Neutropenia poses a significant risk factor for developing aspergillosis which led us to
222 investigate the importance of neutrophils in the recognition and removal of *A. fumigatus*
223 conidia following ficolin opsonization.

224 The association of L-ficolin opsonized conidia with human neutrophils was investigated as
225 per MDM-protocols. In this case, neutrophils were gated (Figure 3A) and the percentage of

226 FITC negative and positive neutrophils were used to identify phagocytic cells (Figure 3B and
227 Figure 3C). Again, the percentage of cells phagocytosing was unaffected in physiological (pH
228 7.4) or acidic conditions (pH 5.7) (Figure 3D). However, as for the macrophages, flow
229 cytometric analysis indicated a significant increase in the number of conidia phagocytosed
230 per neutrophil following L-ficolin opsonization, but only in pH 5.7 conditions (p=0.01056)
231 (Figure 3E and Figure 3F).

232 Light microscopy demonstrated that in the absence of L-ficolin opsonization in pH 5.7 and
233 pH 7.4 or L-ficolin opsonization in pH 7.4 conditions, hyphal growth appeared very dense
234 (Figure 3G, 3H and 3J). Following opsonization by L-ficolin at pH 5.7, hyphal growth
235 appeared significantly less dense and clumping was observed (Figure 3I). Following gating
236 (Figure 3K), the viability assays demonstrated a significant decrease in fungal viability
237 following opsonization by L-ficolin in these conditions (p=0.04324) (Figure 3L and Figure
238 3M). As for MDM, death-associated green-yellow fluorescence emitted by *A.fumigatus* within
239 neutrophils was observed to be significantly greater compared to the unassociated fungi
240 (Supplementary Figure 2A). Conversely, the fluorescence of the free *A.fumigatus* in the
241 presence of neutrophils was significantly greater when compared to *A.fumigatus* in the absence
242 of neutrophils, suggesting augmentation of extracellular killing mechanisms (Supplementary
243 Figure 2B)

244

245 ***L-ficolin opsonization modulates the secretion of inflammatory cytokines in response to A.***
246 ***fumigatus***

247 We utilised cytometric bead arrays to investigate the concentration of IL-8, IL-1 β , IL-6, IL-
248 10 and TNF- α secreted from A549 type II alveolar cells, MDM and human neutrophils
249 following challenge by L-ficolin-opsonized *A. fumigatus* conidia.

250 From the cytokine panel tested, IL-8 was the only cytokine significantly modulated in A549
251 cells in response to L-ficolin (Supplementary Figure 3). L-ficolin opsonization induced a
252 significant increase in the secretion of pro-inflammatory IL-8 compared to challenge with un-
253 opsonized conidia after 8 h and 24 h (Supplementary Figure 3). L-ficolin in the absence of
254 conidia induced a significant spike in IL-8 secretion at 8 h which was maintained up to 24 h
255 (Supplementary Figure 3).

256 L-ficolin opsonization also modulated cytokine secretion from MDM. Following MDM
257 challenge with conidia opsonized by L-ficolin an anti-inflammatory effect was observed. The
258 secretion of IL-8, IL-1 β , IL-6, IL-10 and TNF- α from MDM cells 24 h post-infection were
259 decreased (Figure 4A-E). Again L-ficolin alone appeared capable of significantly increasing
260 the cytokine concentrations of all tested (Figure 4A-E).

261 Additionally, L-ficolin opsonization led to significantly decreased secretion of IL-8, IL-1 β ,
262 IL-6 and TNF- α from neutrophils, compared to un-opsonized conidia (Figure 5A-E). We
263 observed that IL-10 was only secreted at baseline levels regardless of any challenges (Figure
264 5D). L-ficolin was also observed to have the ability to increase the secretion of IL-8, IL-1 β
265 and TNF- α in the absence of *A. fumigatus* (Figure 5A, B and E).

266

267 ***L-ficolin is present in the bronchoalveolar lavage fluid of lung transplant recipients with*** 268 ***fungal pneumonia***

269 Based upon our recent observations [16], it was important to investigate whether L-ficolin
270 was detectable in the lungs of patients with invasive fungal infections, particularly as L-
271 ficolins have never formally been described to be present in lungs.

272 Here, we utilised an L-ficolin-specific ELISA to detect the presence of L-ficolin in the BAL
273 samples of lung transplant recipients. In patients who were diagnosed with probable or
274 proven invasive pulmonary fungal infection based on EORTC/MSG criteria and/or positive

275 fungal biomarkers (GM/lateral-flow), L-ficolin was detected at significantly higher
276 concentrations ($p= 0.00087$; Figure 6A) compared to uninfected control patients. L-ficolin
277 was only detected once in the BAL samples that tested negative for fungal growth or fungal
278 radiology features, albeit at a very low concentration (Figure 6A). An ROC curve analysis
279 was conducted to investigate whether the detection of L-ficolin could be used as a potential
280 biomarker/diagnostic tool for fungal infection in the lung. The area under the curve (AUC)
281 was calculated to be 0.842 which suggested there was an 84.2% chance that fungal infected
282 transplant patients would have L-ficolin present in their BAL fluid ($p<0.0001$; Figure 6B).

283

284 **Discussion**

285 Our study focused on the functional consequences of L-ficolin opsonization of *A. fumigatus*;
286 in particular, its effect on *Aspergillus*-phagocyte interactions. In order to translate our in vitro
287 findings to clinical infections, we also investigated whether L-ficolin is present in human
288 lungs during fungal pneumonia. As a result a number of new observations have been made.
289 Firstly, L-ficolin opsonization led to enhanced uptake of *A. fumigatus* conidia by MDM and
290 neutrophils under inflammatory conditions. Secondly, this opsonization led to enhanced
291 inhibition of hyphal formation and an increase in *A. fumigatus* killing by MDM and
292 neutrophils. Thirdly, opsonization of *A. fumigatus* conidia by L-ficolin, evoked an anti-
293 inflammatory cytokine response from MDM and neutrophils. Finally, for the first time we
294 provide evidence that L-ficolin is present in the BAL fluid of lung transplant recipients
295 diagnosed with fungal infections, which could potentially be used as a diagnostic tool for
296 fungal infection in a clinical setting.

297 Initially, we showed that L-ficolin bound to *A. fumigatus* at low pH (5.7) which was similar
298 to ficolin-A [16]. The ability of such pattern recognition molecules to function at decreased

299 pH is important in the defence against microorganisms, with pH at the local site of infection
300 being observed to drop as low as pH 5.5 during inflammation [23].

301 Another key participant during infection induced inflammation is the macrophage, which is
302 the most prominent phagocyte in the lung in the early stages of *A. fumigatus* infection [24].
303 We have demonstrated here that L-ficolin enhances conidial uptake by primary MDM from
304 healthy donors. Opsonophagocytosis was enhanced at inflammatory pH, which is also
305 optimal for ficolin binding.

306 The other essential phagocyte in the defence against *Aspergillus* is the neutrophil which is
307 known to prevent fungal growth, although the mechanism has not been fully elucidated [25].
308 We have shown here that L-ficolin enhances neutrophil function by increasing conidial
309 uptake following opsonization.

310 Our observations are adding to the knowledge of previous reports on L-ficolin enhancing the
311 opsonophagocytosis of not only bacteria such as *Salmonella typhimurium* and *Streptococcus*
312 *agalactiae*, but also of fungi [26, 27]. It is likely that ficolins work together with other pattern
313 recognition molecules (SP-A, SP-D and mannose-binding lectin (MBL)) and receptors
314 (dectin-1 and Toll-like receptor 2), which have also been observed to bind *A. fumigatus*
315 conidia and enhance phagocytic uptake [11, 16, 28-32]. Although binding and phagocytosis
316 is important, ultimately, killing of the fungi is crucial in order to sterilize infected tissues.

317 Macrophages are usually able to kill conidia in their acidic phagolysosomes [33], but if
318 conidia escape this process and germinate into hyphae, they become too large a structure to
319 be phagocytosed. Neutrophils are then recruited to the site of infection (in response to IL-8)
320 where they assist the inhibition of fungal invasion by degranulation and the production of
321 fungistatic NETs following adherence to the hyphal cell wall [14, 24, 34].

322 In our study we observed that L-ficolin opsonization potentiated the ability of macrophages
323 and neutrophils to significantly enhance fungal killing. Macrophages appeared more capable
324 of inhibiting germination of conidia in comparison to neutrophils. Gating separately on the
325 neutrophil/MDM populations containing *A. fumigatus* or the free *A. fumigatus* further
326 illuminated the roles of these cells in killing. These observations were in keeping with
327 previous reports that macrophages are involved in early conidial phagocytosis and killing
328 whereas neutrophils are recruited for help at a later stage whereby extracellular killing
329 mechanisms are integral [35].

330 This represents the first observation of the ability of ficolins to enhance killing of *A.*
331 *fumigatus* by phagocytes which is supporting observations of others who have reported this
332 as a characteristic of the related surfactant proteins [30]. The importance of surfactant
333 proteins was further highlighted by their protective role against *A. fumigatus* in an *in vivo*
334 model [31]. Additionally, MBL has been observed to be a key component in systemic
335 *Aspergillus* infections, further emphasizing that humoral pattern recognition molecules play
336 an important role in the defence against fungi [32, 36]. However, we are currently
337 investigating the role of L-ficolin in the *in vivo* defence against aspergillosis.

338 As indicated earlier, we have previously observed that ficolin-A opsonization leads to an
339 increase in IL-8 secretion from A549 cells, a cytokine that is crucial for the recruitment of
340 neutrophils during *Aspergillus* infection. It is known that in response to *A. fumigatus*, a
341 plethora of cytokines are secreted from various host cells, including; IL-2, IL-5, IL-6, IL-8,
342 IL-13, IL-17A, IL-22, IFN- γ , TNF- α , GM-CSF and MCP-1 [24, 37-42].

343 In the present study, we found that L-ficolin opsonized conidia were also capable of inducing
344 an increase in IL-8 as previously observed for ficolin-A opsonized conidia [16]. In contrast,
345 opsonization of *A. fumigatus* by L-ficolin led to a significant decrease in IL-8, IL-1 β , IL-6,

346 IL-10 and TNF- α production from MDM and neutrophils. In support of our observations, it
347 was recently observed that ficolin-A could act in an anti-inflammatory manner by binding to
348 lipopolysaccharide (LPS) and inhibiting LPS-mediated pro-inflammatory responses on
349 murine mast cells [43]. Additionally, the pattern recognition proteins SP-A and -D modulate
350 an anti-inflammatory cytokine profile in response to viruses, LPS-induced cytokine and nitric
351 oxide production, and allergens [44-46].

352 Our work represents the first observations that unbound ficolins may have the potential to
353 increase cytokine secretion. The mechanisms of this interaction are still not fully understood
354 but it most likely depends on the orientation of ficolin binding. Interestingly, both SP-A and -
355 D have been observed to function in both an anti- and pro-inflammatory manner, dependent
356 upon the interaction of their globular heads with SIRP α or their collagenous tails with
357 calreticulin/CD91, respectively [47]. Some data suggests that L-ficolin binds to calreticulin
358 but there has been no demonstrable binding to SIRP α [48]. Another caveat to be aware of is
359 that in its native state, L-ficolin normally exists as quiescent polymers but the recombinant
360 form used in this study is in a depolymerised state and may not be completely representative
361 of normal *in vivo* function. This is an area of research that is currently ongoing in our
362 laboratory.

363 The most important clinical observation of our study was the detection of the serum L-ficolin
364 in BAL of patients lungs diagnosed with invasive *A.fumigatus* infection. Moreover, L-ficolin
365 could also be detected in the lungs of recipient's infected with *A.flavus*, *Penicillium*
366 spp., *Acremonium* spp., *Scedoporium apiospermum* and at very low concentration in one
367 incidence of *S.aureus* infection. This ficolin has, until now, not been reported to be present in
368 the lung. We postulate that L-ficolin, which is normally produced by the liver, enters the
369 alveolar space during infection from the blood stream similarly to the related acute phase

370 protein, MBL (a serum collectin), which has also been found in the BAL fluid from infected

371 lungs [49]. Although the current sample size is small (39 patients), ROC analysis has
372 indicated that the presence of L-ficolin in the lungs of transplant patients could be linked with
373 fungal infection, but this diagnostic potential will need to be further investigated in larger
374 clinical trials.

375 In conclusion, L-ficolin is present in fungal infected lungs of transplant patients and has
376 immunomodulatory properties that highlight an important role in the innate defence against
377 *Aspergillus* through enhancing opsonophagocytosis by macrophages and neutrophils,
378 increasing fungal killing and manifesting an anti-inflammatory cytokine profile post-
379 infection. Future research will be concerned with understanding the signalling pathways
380 involved in immune defence and utilizing ficolin-deficient transgenic animal models to
381 elucidate the function of ficolins in the defence against *Aspergillus in vivo*.

382

383 **Funding**

384 This work was supported by the Medical Research Council [MR/K002708/1 to A.A, A.S,
385 A.R and D.A.J and a Faculty of Health, University of East Anglia, UK PhD studentship FMH
386 04.4.66 C4 to S.B, D.S and S.S].

387

388 **Acknowledgements**

389 The authors would like to acknowledge Dr. Orla Jupp (University of East Anglia) for
390 providing us with isolated human monocytes and neutrophils in addition to Professor
391 Christopher Thornton (University of Exeter) for providing lateral-flow devices.

392

393 **References**

- 394 1. Morgan J, Wannemuehler KA, Marr KA, et al. Incidence of invasive aspergillosis
395 following haematopoietic stem cell and solid organ transplantation: interim results of a
396 prospective multicenter surveillance program. *Med Mycol* **2005**; 43:S49-S58.
- 397 2. Denning DW. Invasive aspergillosis. *Clin Infect Dis* **1998**; 26:781-803.
- 398 3. Marr KA, Carter RA, Boeckh M, Martin P, Corey L. Invasive aspergillosis in allogeneic
399 stem cell transplant recipients; changes in epidemiology and risk factors. *Blood* **2002**;
400 100:4358-66.
- 401 4. Wiederhold NP, Lewis RE, Kontoyiannis DP. Invasive aspergillosis in patients with
402 hematologic malignancies. *Pharmaco* **2003**; 23:1592-610.
- 403 5. Pagano L, Caira M, Candoni A, et al. Invasive aspergillosis in patients with acute myeloid
404 leukaemia: a SEIFEM-2008 registry study. *Haematol* **2010**; 95:644-50.
- 405 6. Segal BH. Aspergillosis. *N Engl J Med* **2009**; 360:1870-84.
- 406 7. Martin T, Frevert C. Innate immunity in the lungs. *Proc Am Thorac Soc* **2005**; 2:403-11.
- 407 8. Falvey D, Streifel A. Ten-year air sample analysis of *Aspergillus* prevalence in a
408 University hospital. *J Hosp Infect* **2007**; 67:35-41.
- 409 9. Schelenz S, Goldsmith DJ. *Aspergillus* endophthalmitis: an unusual complication of
410 disseminated infection in renal transplant patients. *J Infect* **2003**; 47:336-43.
- 411 10. Ibrahim-Granet O, Jouvion G, Hohl TM, et al. In vivo bioluminescence imaging and
412 histopathologic analysis reveal distinct roles for resident and recruited immune effector
413 cells in defense against invasive aspergillosis. *BMC Microbiol* **2010**; 10:105.
- 414 11. Luther K, Torosantucci A, Brakhage AA, Heesemann J, Ebel F. Phagocytosis of
415 *Aspergillus fumigatus* conidia by murine macrophages involves recognition by the dectin-1
416 beta-glucan receptor and Toll-like receptor 2. *Cell Microbiol* **2007**; 9:368-81.

- 417 12. Aratani Y, Kura F, Watanabe H, et al. Relative contributions of myeloperoxidase and
418 NADPH-oxidase to the early host defense against pulmonary infections with *Candida*
419 *albicans* and *Aspergillus fumigatus*. *Med Mycol* **2002**; 40:557-63.
- 420 13. Segal BH, Han W, Bushey JJ, et al. NADPH oxidase limits innate immune responses in
421 the lungs in mice. *PLoS ONE* **2010**; 5:e9631.
- 422 14. Bruns S, Kniemeyer O, Hasenberg M, et al. Production of extracellular traps against
423 *Aspergillus fumigatus* in vitro and in infected lung tissue is dependent on invading
424 neutrophils and influenced by hydrophobin RodA. *PLoS Pathog* **2010**; 6:e1000873.
- 425 15. Taira S, Kodama N, Matsushita M, Fujita T. Opsonic function and concentration of
426 human serum ficolin/P35. *J Med Sci* **2000**; **46**:13-23.
- 427 16. Bidula S, Kenawy H, Ali Y, Sexton D, Schwaeble W, Schelenz S. Role of ficolin-A and
428 lectin complement pathway in the innate defense against pathogenic *Aspergillus* species.
429 *Infect Immun* **2013**; **81**:1730-40.
- 430 17. De Pauw B, Walsh TJ, Donnelly JP, et al. Revised definitions of invasive fungal disease
431 from the European Organization for Research and Treatment of Cancer/Invasive Fungal
432 Infections Cooperative Group and the National Institute of Allergy and Infectious Diseases
433 Mycoses Study Group (EORTC/MSG) Consensus Group. *Clin Infect Dis* **2008**; 46:1813-21.
- 434 18. Walsh GM, Sexton DW, Blaylock MG, Convery CM. Resting and cytokine-stimulated
435 human small airway epithelial cells recognize and engulf apoptotic eosinophils. *Blood* **1999**;
436 94:2827-35.
- 437 19. Thornton CR. Development of an immunochromatographic lateral-flow device for rapid
438 serodiagnosis of invasive aspergillosis. *Clin Vaccine Immunol* **2008**; 15:1095-105.
- 439 20. D'Haese J, Theunissen K, Vermeulen E, et al. Detection of galactomannan in
440 bronchoalveolar lavage fluid samples of patients at risk for invasive pulmonary aspergillosis:
441 analytical and clinical validity. *J Clin Microbiol* **2012**; 50:1258-63.

- 442 21. Investigation of Bronchoalveolar Lavage, Sputum and Associated Specimens. UK
443 Standards for Microbiology Investigations. **2014**; B57:<http://www.hpa.org.uk/SMI/pdf>. Date
444 accessed 21/12/14.
- 445 22. Ma YJ, Doni A, Hummelshøj T, et al. Synergy between ficolin-2 and pentraxin 3 boosts
446 innate immune recognition and complement deposition. *J Biol Chem* **2009**; **284**:28263-75.
- 447 23. Martinez D, Vermeulen M, Trevani A, et al. Extracellular acidosis induces neutrophil
448 activation by a mechanism dependent on activation of phosphatidylinositol 3-kinase/Akt and
449 ERK pathways. *J Immunol* **2006**; **176**:1163-71.
- 450 24. Schelenz S, Smith DA, Bancroft GJ. Cytokine and chemokine responses following
451 pulmonary challenge with *Aspergillus fumigatus*: obligatory role of TNF-alpha and GM-CSF
452 in neutrophil recruitment. *Med Mycol* **1999**; **37**:183-94.
- 453 25. Mircescu MM, Lipuma L, van Rooijen N, Pamer EG, Hohl TM. Essential role for
454 neutrophils but not alveolar macrophages at early time points following *Aspergillus*
455 *fumigatus* infection. *J Infect Dis* **2009**; **200**:647-56.
- 456 26. Matsushita M, Endo Y, Taira S, et al. A novel human serum lectin with collagen- and
457 fibrinogen-like domains that functions as an opsonin. *J Biol Chem* **1996**; **271**:2448-54.
- 458 27. Fujieda M, Aoyagi Y, Matsubara K, et al. L-ficolin and capsular polysaccharide-specific
459 IgG in cord serum contribute synergistically to opsonophagocytic killing of serotype III and
460 V group B streptococci. *Infect Immun* **2012**; **80**:2053-60.
- 461 28. Aïmanianda V, Bayry J, Bozza S, et al. Surface hydrophobin prevents immune
462 recognition of airborne fungal spores. *Nature* **2009**; **460**:1117-21.
- 463 29. Allen MJ, Harbeck R, Smith B, Voelker DR, Mason RJ. Binding of rat and human
464 surfactant proteins A and D to *Aspergillus fumigatus* conidia. *Infect Immun* **1999**; **67**:4563-9.

- 465 30. Madan T, Eggleton P, Kishore U, et al. Binding of pulmonary surfactant proteins A and D
466 to *Aspergillus fumigatus* conidia enhances phagocytosis and killing by human neutrophils
467 and alveolar macrophages. *Infect Immun* **1997**; 65:3171-9.
- 468 31. Madan T, Kishore U, Singh M, et al. Protective role of lung surfactant protein D in a
469 murine model of invasive pulmonary aspergillosis. *Infect Immun* **2001**; 69:2728-31.
- 470 32. Kaur S, Gupta VK, Thiel S, Sarma PU, Madan T. Protective role of mannan-binding
471 lectin in a murine model of invasive pulmonary aspergillosis. *Clin Exp Immunol* **2007**;
472 148:382-9.
- 473 33. Ibrahim-Granet O, Philippe B, Boleti H, et al. Phagocytosis and intracellular fate of
474 *Aspergillus fumigatus* conidia in alveolar macrophages. *Infect Immun* **2003**; 71:891-903.
- 475 34. Bianchi M, Hakkim A, Brinkmann V, et al. Restoration of NET formation by gene
476 therapy in CGD controls aspergillosis. *Blood* **2009**; 114:2619-22.
- 477 35. Schaffner A. Host-parasite relation in invasive aspergillosis. *Nihon Ishinkin Gakkai*
478 *Zasshi* **2002**; 43:161.
- 479 36. Lambourne J, Agranoff D, Herbrecht R, et al. Association of mannose-binding lectin
480 deficiency with acute invasive aspergillosis in immunocompromised patients. *Clin Infect Dis*
481 **2009**; 49:1486-91.
- 482 37. Zhang Z, Liu R, Noordhoek J, Kauffman H. Interaction of airway epithelial cells (A549)
483 with spores and mycelium of *Aspergillus fumigatus* *J Infect* **2005**; **51**:375-82.
- 484 38. Balloy V, Sallenave J, Wu Y, et al. *Aspergillus fumigatus* -induced Interleukin-8
485 synthesis by respiratory epithelial cells is controlled by the phosphatidylinositol 3-kinase, p38
486 MAPK, and ERK 1/2 pathways and not by the toll-like receptor-MyD88 pathway. *JBiol*
487 *Chem* **2008**; **283**:30513-21.

488 39. Koth L, Rodriguez M, Bernstein X, et al. *Aspergillus* antigen induces robust Th2 cytokine
489 production, inflammation, airway hyperreactivity and fibrosis in the absence of MCP-1 or
490 CCR2. *Resp Res* **2004**; **5**:12.

491 40. Graziutti ML, Rex JH, Cowart RE, Anaissie EJ, Ford A, Savary CA. *Aspergillus*
492 *fumigatus* conidia induce a Th1-type cytokine response. *J Infect Dis* **1997**; **176**:1579-83.

493 41. Werner JL, Gessner MA, Lilly LM, et al. Neutrophils produce interleukin 17A (IL-17A)
494 in a dectin-1- and IL-23-dependent manner during invasive fungal infection. *Infect Immun*
495 **2011**; **79**:3966-77.

496 42. Gessner MA, Werner JL, Lilly LM, et al. Dectin-1-dependent interleukin-22 contributes
497 to early innate lung defense against *Aspergillus fumigatus*. *Infect Immun* **2012**; **80**:410-7.

498 43. Ma YJ, Kang HJ, Kim JY, Garred P, Lee MS, Lee BL. Mouse mannose-binding lectin-A
499 and ficolin-A inhibit lipopolysaccharide-mediated pro-inflammatory responses on mast cells.
500 *BMB Rep* **2013**; **46**:376-81.

501 44. Harrod KS, Trapnell BC, Otake K, Korfhagen TR, Whitsett JA. SP-A enhances viral
502 clearance and inhibits inflammation after pulmonary adenoviral infection. *Am J Physiol*
503 **1999**; **277**:L580-8.

504 45. Borron P, McIntosh JC, Korfhagen TR, Whitsett JA, Taylor J, Wright JR. Surfactant-
505 associated protein A inhibits LPS-induced cytokine and nitric oxide production in vivo. *Am J*
506 *Physiol Lung Cell Mol Physiol* **2000**; **278**:L840-7.

507 46. Schleh C, Rothen-Rutishauser BM, Blank F, et al. Surfactant protein D modulates
508 allergen particle uptake and inflammatory response in a human epithelial airway model. *Resp*
509 *Res* **2012**; **13**:8.

510 47. Gardai SJ, Xiao YQ, Dickinson M, et al. By binding SIRPalpha or calreticulin/CD91,
511 lung collectins act as dual function surveillance molecules to suppress or enhance
512 inflammation. *Cell* **2003**; **115**:13-23.

513 48. Kuraya M, Ming Z, Liu X, Matsushita M, Fujita T. Specific binding of L-ficolin and H-
514 ficolin binding to apoptotic cells leads to complement activation. *Immunobiology* **2005**;
515 **209**:689-97.

516 49. Summerfield JA. The role of mannose-binding protein in host defence. *Biochem Soc*
517 *Trans* **1993**; 21:473-7.

518

519

520

521

522

523

524

525

526

527

528 **Figure legends**

529 **Figure 1. L-ficolin binding to live *A. fumigatus* conidia.** Live *A. fumigatus* conidia (5×10^5)
530 were opsonized with $5 \mu\text{g ml}^{-1}$ L-ficolin in the presence or absence of Ca^{2+} and in a range of
531 pH's from 3.7-10.7 prior to staining and flow cytometric analysis. **(A)** Binding of L-ficolin to
532 *A. fumigatus* in the presence or absence of Ca^{2+} . **AF** represents *A. fumigatus* alone. **+ Abs**
533 represents the antibody background fluorescence. **BSA** was used as a negative control for
534 binding. **(B)** Binding of L-ficolin to *A. fumigatus* in pH 3.7-pH 10.7 conditions. Results are
535 representative of the average of all data points gained from three independent experiments.
536 Error bars represent SD and significance was determined via two-tailed Students *t*-test. An
537 asterisk indicates a significant difference: $p < 0.05$. MFI, median fluorescence intensity; AF, *A.*
538 *fumigatus*, Abs, antibodies.

539

540 **Figure 2. Phagocytosis and fungal viability following incubation of L-ficolin-opsonized**
541 ***A. fumigatus* conidia with monocyte-derived macrophages.** FITC-labelled or live freshly
542 harvested *A. fumigatus* conidia (5×10^5) were opsonized with $5 \mu\text{g ml}^{-1}$ L-ficolin prior to
543 incubation with MDM (conidia:MDM ratio of 5:1) in pH 5.7 and pH 7.4 conditions for 2 h
544 or 24 h for phagocytosis and viability assays, respectively. **(A)** Gate P1 on MDM used to
545 produce figures **B-F**. Some points were removed for clarity. **(B)** Representative flow data
546 depicting % MDM phagocytosing in the absence of FITC-labelled *A. fumigatus* conidia (Q1).
547 **(C)** or in the presence of FITC-labelled *A. fumigatus* conidia (Q1). **(D)** The percentage of
548 MDM phagocytosing conidia in pH 5.7 or pH 7.4 conditions in the presence or absence of L-
549 ficolin. **(E)** Representative histogram depicting the uptake of conidia in pH 5.7 conditions in
550 the presence or absence of L-ficolin. **(F)** The relative number of phagocytosed FITC-labelled
551 conidia (based upon the median fluorescence intensity; FL1-A) either un-opsonized (–L-

552 ficolin) or following opsonization by L-ficolin (+L-ficolin). **(G)** Hyphal germination
553 following incubation of un-opsonized conidia in pH 5.7. The black arrows point to
554 macrophages containing conidia which makes macrophages appear dark. The white arrow
555 heads are used to trace single hyphae from MDM. Many hyphae are visible, some of which
556 are blurry as they are growing in three-dimensions and are out of the focal plane. Or. **(H)** in
557 pH 7.4. **(I)** Hyphal germination following incubation of L-ficolin opsonized conidia in pH
558 5.7. Hyphae are present although growth is much less dense. Or. **(J)** in pH 7.4. **(K)** Gate P2
559 on MDM and *A. fumigatus* used to produce figures **L** and **M**. **(L)** Representative histogram
560 depicting the killing of conidia in pH 5.7 conditions in the presence or absence of L-ficolin.
561 Increased FL1-A depicts enhanced killing. **(M)** Viability of un-opsonized or L-ficolin
562 opsonized conidia after incubation with MDMs. Results are representative of the average of
563 all data points gained from three independent experiments. Error bars represent SD and
564 significance was determined via two-tailed Students *t*-test. An asterisk indicates a significant
565 difference: $p < 0.05$. FITC, fluorescein isothiocyanate; MDM, monocyte-derived macrophage;
566 MFI, median fluorescence intensity; AF, *A. fumigatus*; SSC-A, side scatter; FSC-A, forward
567 scatter; FL1-A, fluorescence.

568

569 **Figure 3. Phagocytosis and fungal viability following incubation of ficolin-opsonized *A.***
570 ***fumigatus* conidia with human neutrophils.** FITC-labelled or live freshly harvested *A.*
571 *fumigatus* conidia (5×10^5) were opsonized with $5 \mu\text{g ml}^{-1}$ L-ficolin prior to incubation with
572 neutrophils (conidia:neutrophil ratio of 5:1) in pH 5.7 and pH 7.4 conditions for 2 h or 24 h
573 for phagocytosis and viability assays, respectively. **(A)** Gate P1 on neutrophils used to
574 produce figures **B-F** **(B)** Representative flow data depicting % neutrophils phagocytosing in
575 the absence of FITC-labelled *A. fumigatus* conidia (Q1). **(C)** or in the presence of FITC-
576 labelled *A. fumigatus* conidia (Q1). **(D)** The percentage of neutrophils phagocytosing conidia

577 in pH 5.7 or pH 7.4 conditions in the presence or absence of L-ficolin. (E) Representative
578 histogram depicting the uptake of conidia in pH 5.7 conditions in the presence or absence of
579 L-ficolin. (F) The relative number of phagocytosed FITC-labelled conidia (based upon the
580 median fluorescence intensity; FL1-A) either un-opsonized (-L-ficolin) or following
581 opsonization by L-ficolin (+L-ficolin). (G) Hyphal germination following incubation of un-
582 opsonized conidia in pH 5.7 or. (H) in pH 7.4. (I) Hyphal germination following incubation
583 of L-ficolin opsonized conidia in pH 5.7 or. (J) in pH 7.4. (K) Gate P2 on neutrophils and *A.*
584 *fumigatus* used to produce figures L and M. (L) Representative histogram depicting the
585 killing of conidia in pH 5.7 conditions in the presence or absence of L-ficolin. Increased FL1-
586 A depicts enhanced killing. (M) Viability of un-opsonized conidia or L-ficolin opsonized
587 conidia after incubation with neutrophils. Results are representative of the average of all the
588 data points gained from three independent experiments. Error bars represent the SD and
589 significance was determined via two-tailed Students *t*-test. An asterisk indicates a significant
590 difference: $p < 0.05$. FITC, fluorescein isothiocyanate; SSC-A, side scatter; FSC-A, forward
591 scatter; FL1-A, fluorescence.

592

593 **Figure 4. Inflammatory cytokine release from monocyte-derived macrophages following**
594 **challenge by un-opsonized or L-ficolin-opsonized conidia.** Supernatants were collected
595 after 8h and 24 h time points during challenge with live *A. fumigatus* conidia (5×10^5) either
596 un-opsonized or L-ficolin opsonized ($5 \mu\text{g ml}^{-1}$) prior to the conduction of cytometric bead
597 arrays. (A) The concentration of IL-8 secreted. (B) The concentration of IL-1 β secreted. (C)
598 The concentration of IL-6 secreted. (D) The concentration of IL-10 secreted. (E) The
599 concentration of TNF- α secreted. Following *A. fumigatus* challenge. **MDM** is representative
600 of MDM alone. **+L-ficolin** represents MDM in the presence of L-ficolin alone. **+AF** and
601 **+AF+L-ficolin** are representative of un-opsonized *A. fumigatus* or L-ficolin opsonized

602 conidia, respectively. Results are representative of the average of all the data points gained
603 from three independent experiments. Error bars represent the SD. Significance was
604 determined via one-way ANOVA and pair-wise comparisons were conducted using the
605 Student-Newman-Keuls method. An asterisk indicates a significant difference: $p < 0.05$.
606 MDM, monocyte-derived macrophage; IL, interleukin.

607

608 **Figure 5. Inflammatory cytokine release from neutrophils following challenge by un-**
609 **opsonized or L-ficolin opsonized conidia.** Supernatants were collected after 8h and 24 h
610 time points during challenge with live *A. fumigatus* conidia (5×10^5) either un-opsonized or
611 L-ficolin opsonized ($5 \mu\text{g ml}^{-1}$) prior to the conduction of cytometric bead arrays. **(A)** The
612 concentration of IL-8 secreted. **(B)** The concentration of IL-1 β secreted. **(C)** The
613 concentration of IL-6 secreted. **(D)** The concentration of IL-10 secreted. **(E)** The
614 concentration of TNF- α secreted. Following *A. fumigatus* challenge. **Neutrophils** is
615 representative of neutrophils alone. **+L-ficolin** represents neutrophils in the presence of L-
616 ficolin alone. **+AF** and **+AF+L-ficolin** are representative of un-opsonized *A. fumigatus* or L-
617 ficolin opsonized conidia, respectively. Results are representative of the average of all the
618 data points gained from three independent experiments. Error bars represent the SD.
619 Significance was determined via one-way ANOVA and pair-wise comparisons were
620 conducted using the Student-Newman-Keuls method. An asterisk indicates a significant
621 difference: $p < 0.05$. IL, interleukin.

622

623 **Figure 6. L-ficolin is found in the bronchoalveolar lavage fluid of lung transplant**
624 **recipients.** BAL fluid was collected following bronchoscopies from lung transplant
625 recipients. **(A)** BAL samples were considered **positive** or **negative** for invasive fungal

626 infection dependent upon patients classification according to EORTC/MSG criteria. All
627 samples were tested for fungal infection via *Aspergillus* antigen detection, radiology and
628 culture. **(B)** ROC curve analysis for L-ficolin detection in fungal-infected transplant patients
629 compared to non-infected transplant patients. Results are representative of the data points
630 gained from three independent experiments (19 positive and 20 negative patients). Bars
631 represent the median and significance was determined via two-tailed Students *t*-test ($p=$
632 0.00087). Abbreviation: BAL, bronchoalveolar lavage.

633

634 Supplementary Figure 1. Intracellular and extracellular killing of L-ficolin opsonized
635 conidia by MDM.

635 Live freshly harvested *A. fumigatus* conidia (5×10^5) were opsonized with $5 \mu\text{g ml}^{-1}$ L-
636 ficolin prior to incubation with MDM (conidia:MDM ratio of 5:1) in pH 5.7 and
637 pH 7.4 conditions for 24 h. **(A)** The death-associated green-yellow fluorescence emitted by
638 intracellular L-ficolin opsonized or un-opsonized *A. fumigatus*, after incubation with MDM.
639 **(B)** The death-associated green-yellow fluorescence emitted by extracellular L-ficolin
640 opsonized or un-opsonized *A. fumigatus*, after incubation with MDM. Results are
641 representative of the average of all the data points gained from three independent
642 experiments. Error bars represent the SD and significance was determined via two-tailed
Students *t*-test. An asterisks indicated difference: $p < 0.05$.

643

644 Supplementary Figure 2. Intracellular and extracellular killing of L-ficolin opsonized
646 conidia by neutrophils.

645 Live freshly harvested *A. fumigatus* conidia (5×10^5) were 647 opsonized with $5 \mu\text{g ml}^{-1}$
646 L-ficolin prior to incubation with neutrophils (conidia:neutrophil ratio of 5:1) in pH 5.7 and

647 pH 7.4 conditions for 24 h. (A) The death-associated green-yellow fluorescence emitted by
648 intracellular L-ficolin opsonized or un-opsonized *A. fumigatus*, after incubation with
649 neutrophils. (B) The death-associated green-yellow fluorescence emitted by
650 extracellular L-ficolin opsonized or un-opsonized *A. fumigatus*, after incubation with
651 neutrophils. Results are representative of the average of all the data points gained from three
652 independent experiments. Error bars represent the SD and significance was determined via
653 two-tailed Students t-test. An asterisks indicated difference: $p < 0.05$.

654

655 Supplementary Figure 3. IL-8 production from A549 cells following challenge by
656 unopsonized or L-ficolin opsonized conidia.

657 Supernatants were collected after 8h and 24 h 657 time points during challenge with live
658 *A.fumigatus* conidia (5×10^5) either un-opsonized or 658 L-ficolin opsonized ($5 \mu\text{g ml}^{-1}$
659 prior to the conduction of cytometric bead array for the measurement of IL-8. **A549** is
660 representative of A549 cells alone. **+L-ficolin** represents A549 cells in the presence of L-
661 ficolin alone. **+AF** and **+AF+L-ficolin** are representative of un-opsonized *A.fumigatus* or
662 L-ficolin opsonized conidia, respectively. Results are representative of the average of all the
663 data points gained from three independent experiments. Error bars represent the SD.
664 Significance was determined via one-way ANOVA and pair-wise comparisons were
665 conducted using the Student-Newman-Keuls method. An asterisk indicated a significant
666 difference: $p < 0.05$.