



## LJMU Research Online

**Basar, N, Othman, SNAM, Nahar, L and Sarker, SD**

**The ethnomedicinal, phytochemical and pharmacological properties of Phaleria macrocarpa (Scheff). Boerl**

<http://researchonline.ljmu.ac.uk/id/eprint/222/>

### Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Basar, N, Othman, SNAM, Nahar, L and Sarker, SD (2014) The ethnomedicinal, phytochemical and pharmacological properties of Phaleria macrocarpa (Scheff). Boerl. TANG – Humanitas Traditional Medicine, 4 (4). e22-1-e22-12. ISSN 2233-8985**

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)

<http://researchonline.ljmu.ac.uk/>

Review

## The ethnomedicinal, phytochemical and pharmacological properties of *Phaleria macrocarpa* (Scheff.) Boerl.

Siti Nur Atiqah Md Othman<sup>1</sup>, Satyajit Dey Sarker<sup>2</sup>, Lutfun Nahar<sup>2</sup>, Norazah Basar<sup>1,2,\*</sup>

<sup>1</sup>Department of Chemistry, Faculty of Science, Universiti Teknologi Malaysia, 81310 Johor Bahru, Johor, Malaysia; <sup>2</sup>Medicinal Chemistry and Natural Products Research Group, School of Pharmacy and Biomolecular Sciences, Faculty of Science, Liverpool John Moores University, James Parsons Building, Byrom Street, Liverpool L3 3AF, England, UK

### ABSTRACT

*Phaleria macrocarpa* (Scheff.) Boerl. is a dense evergreen tree of the family Thymelaeaceae. This plant is popular with the name of Mahkota dewa, which is literally translated as God's Crown. All parts of this plant including fruits, seeds, stem, and leaves have well known therapeutic properties and have been extensively used in traditional medicine for the treatment of various diseases such as cancer, diabetes mellitus, allergies, kidney disorders, blood diseases, stroke, and acne with satisfactory results. Scientific findings on bioactivities of *P. macrocarpa* also demonstrated different pharmacological properties of various parts of this plant including cytotoxic, antidiabetic, antioxidant, anti-inflammatory, antibacterial, and antihypertensive activities. Phytochemicals studies of *P. macrocarpa* revealed the presence of several classes of compounds such as benzophenones, terpenoids, xanthenes, lignans, acids, and sugars. This review aims to provide a critical overview on botanical description, traditional usage, phytochemicals, and pharmacological activities of *P. macrocarpa*.

**Keywords** *Phaleria macrocarpa*, Thymelaeaceae, phytochemicals, pharmacological activities

### INTRODUCTION

Nature plays an important role in providing the basic needs of human in the production of food-stuffs, shelters, clothing, means of transportation, fertilizers, flavours and fragrances, and, not the least, medicines for the treatment of various diseases (Cragg and Newmann, 2013; Gordaliza, 2007). Mineral, animal and plant products were utilized as the main sources of drugs, and the use of natural products with therapeutic properties is as ancient as human civilization (De Pasquale, 1984; Rates, 2001; Samuelsson, 2004). Most of the plant compounds that have been found to be medicinally useful and interesting tend to be secondary metabolites including alkaloids, phenolics, acetogenins and terpenoids. Secondary metabolites represent features that can be expressed in terms of ecological, taxonomic and biochemical differentiation and diversity. The wide chemical diversity of secondary metabolites throughout the plant kingdom represents an extremely rich biogenic resource for the discovery of novel drugs and for developing innovative remedies (Gurib-Fakim, 2006). To date, natural products and their derivatives represent more than 50% of all the drugs in clinical use in the world. Higher plants contribute no less than 25% of the total. During the last 40 years, at least a dozen potent drugs were reported from flowering plants (Gurib-Fakim, 2006).

The Thymelaeaceae family is a cosmopolitan family of flowering plants, which is established by Hanus-Fajerska et al.

(2012). This family consists of 45 genera and 700 - 800 species, and is widely distributed in both hemispheres (Herber, 2002; 2003). Nine genera and 89 species of the Thymelaeaceae plants are endemic to China (Zheng et al., 1999). In the other large genera of the Thymelaeaceae are *Gnidia* with approximate number of species 160, *Pimelea* (110), *Daphne* (95), *Wikstroemia* (70), *Daphnopsis* (65), *Struthiola* (35), *Lachnaea* (30), *Thymelaea* (30), *Phaleria* (30), and *Gonystylus* (25) (Kubitzki and Bayer, 2003). The species of this family include mostly shrubs or small trees, rarely herbs, evergreen or deciduous. Most species are toxic but some have medicinal properties. The phloem contains strong fibers, which make the bark of many species beneficial in manufacturing of high quality paper especially bank notes. The stems have characteristics of supple and are difficult to break, and used as a substitute for string (Zheng et al., 1999). One of the plants within the Thymelaeaceae is *Phaleria macrocarpa* (Scheff.) Boerl. which was first described by Scheffer as *Drimyspermum macrocarpum* based on fruiting specimens collected by Teysmann near Doré, in western New Guinea (Angiosperm Phylogeny Group, 2003). The other botanical name of this plant is *Phaleria papuana* Warb var. *Wichanii* (Val) Back (Hou, 1960). This plant is popular with the name of 'Mahkota dewa', which is literally translated as God's Crown. It is locally known as 'Simalakama' in Sumatra (Malay) and Depok (West Java) and 'Makutadewa', 'Makuto rajo', 'Makuto ratu' or 'Makuto mewa' in Java (Harmanto, 2005).

\*Correspondence: Norazah Basar

E-mail: norazahb@utm.my

Received April 14, 2014; Accepted October 31, 2014; Published November 30, 2014

doi: <http://dx.doi.org/10.5667/tang.2014.0018>

© 2014 by Association of Humanitas Medicine

This is an open access article under the CC BY-NC license.

(<http://creativecommons.org/licenses/by-nc/3.0/>)

### Botanical descriptions of *P. macrocarpa* (Scheff.) Boerl.

*P. macrocarpa* is a shrub or small tree that grows throughout the year. This plant usually reaches the height of 5 m but sometimes its height could also reach up to 18 m (Harmanto, 2003; Stevens, 1974; Winarto, 2003). It grows in areas of 10 - 1,200 m above the sea level and the most productive age of this



**Fig. 1.** The (A) leaves, (B) fruit and (C) pit of *P. macrocarpa*

plant is in between 10 – 20 years (Saufi, 2007). The plant of *P. macrocarpa* has features of many-branched crown with 1-metre long straight root exuding sap, a brownish green bark and white wood. The leaves are green, sharp edge and tapering from 10 – 15 cm in length and 3 – 5 cm in wide (Fig. 1A). Its flowers appear in white colour with trumpet-like shape and produce pleasant smell. The fruits have an eclipse shape; occur in various sizes with diameter ranging from 3 – 5 cm. Its fruits have smooth surface and changing their colour from green when young into red or maroon when ripening (Fig. 1B). The pit is round, white and poisonous (Fig. 1C) (Altaf et al., 2013; Hendra et al., 2011; Saufi, 2007)

#### **Traditional usage of various parts of *P. macrocarpa* (Scheff.) Boerl.**

*P. macrocarpa* is frequently used as a therapeutic healing alternative in health system of the Indonesians and lower course of Malaysia (Ali et al., 2012). All parts of this plant including fruits, seeds, stems and leaves have well known therapeutic properties and have been extensively used in traditional medicine (Tjandrawinata et al., 2010; Winarto, 2003). Specifically, the fruits of *P. macrocarpa* are used to treat flu, rheumatism, heart diseases and cancer; the leaves are used to treat dysentery, allergy, tumour and impotency while the stems are beneficial in the treatment of bone cancer. The eggshells of seeds are used to counter breast cancer, cervix cancer, lung disease, liver, and heart diseases. This plant especially the seed part cannot be consumed directly due to its high toxicity which can cause swelling, numbness and unconsciousness. However, the seeds can be used as an external medicine for the treatment of skin conditions and for ornamental cultivation purposes, which act as a traditional biopesticide (De Padua et al., 1999; Harmanto, 2003).

#### **Phytochemical studies**

Several research groups especially from Indonesia and China had extensively carried out studies to find chemical constituents from *P. macrocarpa*. The studies resulted in the isolation of several classes of compounds such as benzophenones, terpenoids, xanthenes, lignans, acids and sugars. Chemical investigation on the fruits, leaves and bark of *P. macrocarpa* afforded eight benzophenone derivatives, identified as phalerin (1) (Altaf et al., 2013; Oshimi et al., 2008), isophalerin A (2), isophalerin B (3) (Susilawati, 2012), Mahkoside A (4) (Zhang et al., 2006), Mahkoside B (5) (Zhang et al., 2012), 6,4'-dihydroxy-4-methoxybenzophenone-2-O- $\alpha$ -D-glucopyranoside (6) (Tambunan and Simanjutak, 2006), 6,4'-dihydroxy-4-methoxybenzophenone-2-O- $\beta$ -D-glucopyranoside (7) (Susilawati, 2012; Winarno and Katrin, 2009) and 2,6,4'-trihydroxy-4-methoxybenzophenone (8) (Simanjutak, 2008; Susilawati et al., 2011). Several triterpenoids derivatives known as icaraside C3 (9) (Oshimi et al., 2008), phalerielide (10) (Susilawati, 2012),  $\beta$ -sitosterol (11), stigmaterol (12) and cyloargetanol (13) (Simanjutak, 2008) were successfully isolated from the fruits of *P. macrocarpa*. Phytochemical studies on the fruits by Kurnia et al. (2008) reported a new 29-norcucurbitacin derivative named as desacetyl-fevicordin A

(14), together with fevicordin A (15), fevicordin A glucoside (16) and fevicordin D glucoside (17).

In addition, studies on chemical constituents from the fruits of *P. macrocarpa* revealed the isolation of an ester compound, ethyl stearate (18) (Zhang, 2006) and acid derivatives including palmitic acid (19) (Simanjutak, 2008; Zhang et al., 2006), oleic acid (20), linoleic acid (21), linolenic acid (22), dodecanoic acid (23) (Simanjutak, 2008), naphthoic acid (24) (Susilawati, 2012) and gallic acid (25) (Fariet et al., 2007). A novel lignan named as macronone (26) (Susilawati et al., 2012) and a known lignan, syringaresinol (27) (Lisdawati et al., 2007) were obtained from the ethyl acetate extract of the bark and mesocarp of *P. macrocarpa*, respectively. The investigation on the chemical constituents of this plant yielded a xanthone and flavonoid compound identified as mangiferin (28) (Kim et al., 2010; Oshimi et al., 2008; Zhang et al., 2006) and kaempferol-3-O- $\beta$ -D-glucoside (29) (Zhang et al., 2006). Moreover, two sugar molecules known as glucose (30) and sucrose (31) (Simanjutak, 2008; Zhang et al., 2006) were isolated from the aqueous extract of *P. macrocarpa* fruits. Furthermore, the quantitative analysis on various parts of *P. macrocarpa* fruits revealed the presence of five major flavonoids named as kaempferol (32), myricetin (33), quercetin (34), naringin (35), and rutin (36). Qualitative analysis of the flavonoids was carried out by reversed-phased high performance liquid chromatography (RP-HPLC) using an analytical column C18 60Å 4 $\mu$ m, 3.9  $\times$  150 mm, Waters, NANPA, MA (USA). The flavonoids were detected at 365 nm of UV-Vis photodiode array (DAD) detector (Hendra et al., 2011).

#### **Pharmacological activities**










Empirically, Indonesian people have as often utilized the fruits bark and leave of *P. macrocarpa* for the treatment of various diseases such as cancer, diabetes mellitus, allergies, kidney disorders, blood diseases, stroke and acne with satisfactory results. Therefore, many scientific evaluations on bioactivities of *P. macrocarpa* have been conducted in order to prove the traditional claims on the medicinal values of this plant.

#### **Anticancer activity**















Cytotoxic activities of the methanolic extract from different parts of *P. macrocarpa* fruits were evaluated against the human colon adenocarcinoma cell line (HT-29), human breast adenocarcinoma cell line (MCF-7), human cervical cell line (HeLa) and normal human hepatocytes cell line (Chang liver cell). The viability of cells was measured using the MTT assay. The fruits were divided into pericarp, mesocarp and seed. Results obtained indicated that all parts had potential cytotoxic activity against the MCF-7 and HeLa cancer cell lines with IC<sub>50</sub> values ranging from 25.5 – 40.8  $\mu$ g/ml. The results also showed that the seeds exhibited potential cytotoxic effect against HT-29 with an IC<sub>50</sub> value of 29.5  $\mu$ g/ml while the pericarp and mesocarp exhibited mild cytotoxicity with IC<sub>50</sub> values between 63.8 – 70.1  $\mu$ g/ml (Hendra et al., 2011).

An in vitro study on the cytotoxic effect of fruit extract was carried out against the human uterine cervical carcinoma cell line (HeLa) (Rahmawati et al., 2006). Various concentrations of fruit extract were used to determine the inhibition activity against the HeLa cell line after 24, 72 and 120 h of incubation. The results showed good inhibitory activity against the HeLa cell after 72 h of incubation with an IC<sub>50</sub> value of 5.09 ppm (Pertamawati, 2007). Another investigation on the anticancer activity of the ethanol extract of the fruit pulps of *P. macrocarpa* against mouse mammary tumour was induced by transplantation. This study concluded that the ethanol extract did not inhibit the mouse mammary tumour growth at doses of








**Table 1.** Phytochemicals of *P. macrocarpa* according to its part and types of extract used

No	Name of compound	Structure	Plant part	Types of extract	References
(1)	Phalerin		Fruits	Chloroform	Oshimi et al., 2008
			Leaves	Methanol	Altaf et al., 2013
(2)	Isophalerin A		Fruits	Ethyl acetate	Susilawati, 2012
(3)	Isophalerin B		Leaves	Ethyl acetate	Susilawati, 2012
(4)	Mahkoside A		Nutshell	Ethyl acetate	Zhang et al., 2006
(5)	Mahkoside B		Nutshell	Ethyl acetate	Zhang et al., 2012
(6)	6,4'-Dihydroxy-4-methoxybenzophenone-2-O- $\alpha$ -D-glucopyranoside		Fruits	<i>n</i> -butanol	Tambunan and Simanjutak, 2006
			Fruits	Aqueous	Simanjutak, 2008
(7)	6,4'-Dihydroxy-4-methoxybenzophenone-2-O- $\beta$ -D-glucopyranoside		Fruits	Ethyl acetate	Susilawati, 2012
			Bark	Ethyl acetate	Winarno and Katrin, 2009
(8)	2,6,4'-Trihydroxy-4-methoxybenzophenone		Fruits	Methanol	Simanjutak, 2008
			Leaves	Ethyl acetate	Susilawati, 2011
(9)	Icariside C <sub>3</sub>		Fruits	Chloroform	Oshimi et al., 2008
(10)	Phalerielide		Fruits	Methanol	Susilawati, 2012

**Table 1.** Phytochemicals of *P. macrocarpa* according to its part and types of extract used (continued)

No	Name of compound	Structure	Plant part	Types of extract	References
(11)	$\beta$ -Sitosterol		Fruits	Ethyl acetate	Simanjutak, 2008
(12)	Stigmasterol		Fruits	Ethyl acetate	Simanjutak, 2008
(13)	Cycloartanol		Fruits	Ethyl acetate	Simanjutak, 2008
(14)	Desacetyl-fevicordin A		Seed	Ethyl acetate	Kurnia et al., 2008
(15)	Fevicordin A		Seed	Ethyl acetate	Kurnia et al., 2008
(16)	Fevicordin A glucoside		Seed	Ethyl acetate	Kurnia et al., 2008
(17)	Fevicordin D glucoside		Seed	Ethyl acetate	Kurnia et al., 2008
(18)	Ethyl stearate		Fruits	Methanol	Simanjutak, 2008
(19)	Palmitic acid		Fruits	Chloroform <i>n</i> -Hexane	Oshimi et al., 2008 Simanjutak, 2008
(20)	Oleic acid		Fruits	<i>n</i> -Hexane	Simanjutak, 2008
(21)	Linoleic acid		Fruits	<i>n</i> -Hexane	Simanjutak, 2008
(22)	Linolenic acid		Fruits	<i>n</i> -Hexane	Simanjutak, 2008
(23)	Dodecanoic acid		Fruits	<i>n</i> -Hexane	Simanjutak, 2008
(24)	Naphtoic acid		Fruits	Ethyl acetate	Susilawati, 2012

**Table 1.** Phytochemicals of *P. macrocarpa* according to its part and types of extract used (continued)

No	Name of compound	Structure	Plant part	Types of extract	References
(25)	Gallic acid		Fruits	Ethyl acetate	Faried et al., 2007
(26)	Macronone		Bark	Ethyl acetate	Susilawati, 2012
(27)	Syringaresinol		Fruits	Ethyl acetate	Lisdawati et al., 2007
(28)	Mangiferin		Fruits	Methanol	Oshimi et al., 2008
			Fruits	Ethyl acetate	Zhang et al., 2006
			Leaves	Aqueous	Kim et al., 2010
(29)	Kaempferol-3-O-β-D-glucoside		Fruits	Ethyl acetate	Zhang et al., 2006
(30)	Glucose		Fruits	Aqueous	Simanjutak, 2008
(31)	Sucrose		Fruits	Chloroform	Zhang et al., 2006
			Fruits	Aqueous	Simanjutak, 2008

20, 40 and 80 fold human doses given orally after tumour transplantation for 30 days. However, the results demonstrated the significant increase of apoptosis at the dose of 80 times human dose (Rahmawati et al., 2006).

Previous cytotoxic study on the fruits of *P. macrocarpa* reported the non-toxic effect of the ethanol extract from the seeds and fruits flesh towards human mononuclear peripheral normal cell but slightly toxic to the vero cell line. The extract also did not increase p53 and decrease bcl-2 gene expression that suggested the mechanism of dying cell was caused by necrosis and not by apoptosis (Altaf R et al., 2013). Bioactivity study of irradiated *P. macrocarpa* leaves showed that the n-hexane, ethyl acetate and methanol extracts exhibited strong cytotoxic effect against the mouse leukaemia L1210 cell line with  $IC_{50}$  values of 12.4, 10.3 and 24.1  $\mu\text{g/ml}$ , respectively (Katrin et al., 2011). In addition, cytotoxic activity of the n-hexane, chloroform, ethyl acetate and methanol extracts from the leaves of *P. macrocarpa* plant were investigated against the human hepatoma cell lines (HepG2). The ethyl acetate and methanol extracts were found to have mild cytotoxic effect ( $IC_{50}$  32.5 and 40  $\mu\text{g/ml}$ , respectively) (Yosie et al., 2011).

The investigations on cytotoxic effects of isolated compounds were conducted against several cancer cell lines. Study on cytotoxic activity of phalerin (1) from the methanolic

extract of *P. macrocarpa* leaves was investigated against the myeloma cell line (NS-1). Phalerin (1) was non-toxic towards NS-1 cell line ( $IC_{50}$  83  $\mu\text{g/ml}$ ) (Altaf et al., 2013). Two benzophenone glucosides, mahkoside A (4) and mahkoside B (5) were found to have low cytotoxic effect towards several human cancer cell lines including the prostate cancer cell line (PC-3), stomach cancer cell line (MGC-803) and esophageal cancer cell lines (EC109 and EC9706), with  $IC_{50}$  values exceeding 100  $\mu\text{M/L}$  (Zhang et al., 2012). However, the inhibitory activity of another benzophenone glucoside named as 6,4'-dihydroxy-4-methoxybenzophenone-2-O-β-D-glucopyranoside (7) displayed that this compound had strong cytotoxic properties against the mouse leukaemia cell line (L1210) ( $IC_{50}$  5.1  $\mu\text{g/ml}$ ) (Winarno and Katrin, 2009). Additionally, fevicordin A (15) demonstrated strong anticancer properties against the murine leukaemia cell line (P388) and cervix cancer cell line (HeLa) ( $IC_{50}$  0.01 and 1.16  $\mu\text{g/ml}$ , respectively). This compound also exhibited moderate anticancer activity against another cervix cancer cell line (CasKi) and oesophagus cancer cell line (TE-8) with  $IC_{50}$  values of 12 and 14.6  $\mu\text{g/ml}$ , respectively (Diantini et al., 2012).

Faried et al. (2007) studied the anticancer properties of gallic acid (25), isolated from the fruits of *P. macrocarpa*. The cell proliferation activity was performed using the MTT assay against the human esophageal cancer cell line (TE-2), gastric

**Table 2.** Summary of pharmacological activities of chemical constituent from *P. macrocarpa*

Activity	Compound	Assay	Organism / cells used	Results / Observation	Reference
Cytotoxic activity	(1)	MTT assay	Myeloma cancer cell line (NS-1)	Non-toxic towards NS-1 cell line (IC <sub>50</sub> 83 µg/ml)	Altaf et al., 2013
	(4)	Sulforhodamine B (SRB assay)	Esophageal cancer cell lines (EC109 and EC9706), stomach cancer cell line (MGC-803), prostate cancer cell line (PC-3)	Low cytotoxic effect against PC-3, MGC-803, EC109 and EC9706 cell lines (all IC <sub>50</sub> >100µM/l)	Zhang et al., 2012
	(5)				
	(7)	MTT assay	Leukemia cancer cell line (L1210)	Strong cytotoxic effect against L1210 cell line (IC <sub>50</sub> 5.1 µg/ml)	Winarno and Katrin, 2009
	(15)	MTT assay	Murine leukemia cancer cell line (P388), cervix cancer cell lines (HeLa and CaSki), oesophagus cancer cell line (TE-8)	-Strong anticancer properties against P388 (IC <sub>50</sub> 0.01 µg/ml) and HeLa cell lines (IC <sub>50</sub> 1.16 µg/ml) Moderate anticancer activity against CasKi (IC <sub>50</sub> 12 µg/ml) and TE-8 (IC <sub>50</sub> 14.6 µg/ml)	Diantini et al., 2012
	(25)	MTT assay	Esophageal cancer cell line (TE-2), gastric cancer cell line (MKN-28), colon cancer cell lines (HT-29 and Colo201), breast cancer cell line (MCF-7), CaSki cell line, mouse colon cancer cell line (colon 26)	Have significant anticancer properties against TE-2, MKN-28, HT-29, Colo201, MCF-7, CaSki and colon 26 cancer cell lines	Faried et al., 2007
Antioxidant activity	(8)	DPPH assay	-	Strong antioxidant activity (IC <sub>50</sub> 10.57 µg/ml)	Susilawati et al., 2011
	(26)	DPPH assay	-	Weak antioxidant activity (IC <sub>50</sub> 240.14 µg/ml)	Susilawati et al., 2012
Anti-inflammatory activity	(1)	Lipoxygenase assay (LOX) Hyaluronidase assay (HYA) Xanthine Oxidase assay (XO)	-	-Displayed mild anti-inflammatory properties in XO and LOX assay with percentage of inhibition 34.83% and 23.47%, respectively Does not exhibited as inflammatory effect on HYA assay	Fariza et al., 2012
Vasorelaxant activity	(1)	Vasodilator assay	Noradrenaline-induced contraction of isolated rat aorta	No vasorelaxant effect	Oshimi et al., 2008
	(9)			Slow vasorelaxant activity	
	(28)			No vasorelaxant effect	
Toxicity activity	(14)	-	Brine shrimp ( <i>A. Salina</i> )	All compounds showed variable toxicity with LD <sub>50</sub> ranging from 3 ppm – 12 ppm	Kurnia et al., 2008
	(15)				
	(16)				
	(17)				

cancer cell line (MKN-28), colon cancer cell lines (HT-29 and Colo201), breast cancer cell line (MCF-7), cervix cancer cell line (CaSki), mouse colon cancer cell line (colon 26) with one normal human esophageal cell line (CHEK-1). Interestingly, gallic acid (25) showed significant anticancer properties towards all tested cancer cell lines and induced apoptosis in TE-2 cell line. Apart from the cytotoxic screening activities, the study on molecular mechanism of the extract from this plant on human breast cancer cell line (MDA-MB-231) was performed using a bioactivity-guided DLBS1425. In a previous study, DLBS1425 was found to confer antiproliferative and proapoptosis effects via eicosanoid pathway (Tjandrawinata et al., 2010). DLBS1425 was shown as a potent anticancer agent that targets genes involved in cell survival and apoptosis in the MDA-MB-231 cell line (Tandrasasmita et al., 2010).

#### Antidiabetic activities

The investigation on the  $\alpha$ -glucosidase inhibitory activity and hypoglycemic effect by oral administration of fruit extracts from *P. macrocarpa* in rats were evaluated (Sugiwati et al., 2006). The highest  $\alpha$ -glucosidase inhibitory activity was displayed by the n-butanol extract of young and ripe fruits followed by the ethyl acetate and methanol extracts. The boiled water and n-butanol extracts also displayed hypoglycemic effect by reducing the blood glucose concentration of rats with

effective dose ranging between  $1.0 \times 10^{-3} - 6.5 \times 10^{-3}$  mg/g. The results suggested that this plant could be suitable as a traditional antidiabetic drug (Sugiwati et al., 2006). The hypoglycemic activity of this plant was further investigated by evaluating the effect of the fruits powder on blood glucose levels in glucose loading healthy volunteers. The effective dose of mesocarp fruit powder to decrease the blood glucose level in healthy volunteers was significantly found at the concentration of 500 mg (Meiyanti et al., 2006).

Another study revealed the potential of the ethyl acetate extract of fruits to improve insulin sensitivity in hyperglycemic rats by decreasing the blood glucose level. The highest activity of insulin activity was 18.3%, 9 min after the rats being injected with glucose and exogenous insulin (Muhtadi et al., 2008). The *in vitro* mechanism study demonstrated the antidiabetic properties of the ethanol extract of the fruits as  $\alpha$ -glucosidase inhibitor and as insulin secretagog in clonal glucose responsive insulin secreting cell (BRIN-BD11). In the same study, the qualitative analysis of phytochemicals from the ethanol extract revealed the presence of flavonoid, alkaloid, tannin and steroid. Among these classes of compounds, flavonoid was suggested to be responsible for the antidiabetic properties of the extract in that study (Suparto et al., 2008). In addition, the methanol extract from the fruits pericarp was found to have potential anti-hyperglycemic effect by lowering blood glucose at 56.3% and 58.3% after 12 days of treatment.

**Table 3.** Summary of pharmacological activities of extracts from *P. macrocarpa*

Activity	Extract	Plant part	Assay / Method	Organism / cells used	Results / observation	Reference
Cytotoxic activity	Methanol	Pericarp, mesocarp and seed		Colon cancer cell line (HT-29), breast cancer cell line (MCF-7), cervix cancer cell line (HeLa)	-All parts show potential cytotoxic activity against the MCF-7 and HeLa cancer cell lines with IC <sub>50</sub> values ranging from 25.5 – 40.8 µg/ml.  - The seed part exhibited as potential cytotoxic effect against HT-29 with IC <sub>50</sub> 29.5 µg/ml while the pericarp and mesocarp exhibited as mild cytotoxicity with IC <sub>50</sub> between 63.8 – 70.1 µg/ml.	Hendra et al., 2011
	Ethanol	Fruit	Various concentrations of extract used to determine the inhibition activity after 24, 72 and 120 hours of incubation.	Cervix cancer cell line (HeLa)	Showed the good inhibition activity of HeLa cell occur after 72 hours of incubation with IC <sub>50</sub> 5.09 ppm	Pertamawati, 2007
	Ethanol	Fruit pulp	Silver colloidal stain (AgNOR)	C3H mouse mammary tumour induced by transplantation	-Did not inhibit the C3H mouse mammary tumour growth at dose 20, 40 and 80 fold human dose given orally after tumour transplantation for 30 days.  -The results demonstrated the significant increase of apoptosis at dose of 80 time human dose.	Rahmawati et al., 2006
	Ethanol	Fruit and seed	MTT assay and immunocytochemistry	Human mononuclear perifer normal cell line and vero cell line	-Non-toxic effect of ethanol extract from the seed and fruits flesh towards human mononuclear perifer normal cell but slightly toxic to vero cell line.  -The ethanol extract also did not increase <i>p53</i> and decrease <i>bcl-2</i> gene expression that suggested the mechanism of dying cell was caused by necrosis and not by apoptosis.	Altaf et al., 2013
	<i>n</i> -Hexane, ethyl acetate and methanol	Irradiated leaves	MTT assay	Mouse leukemia cell line (L1210),	The <i>n</i> -hexane, ethyl acetate and methanol extracts exhibited strong cytotoxic effect against mouse leukemia L1210 cell line with IC <sub>50</sub> 12.4, 10.3 and 24.1 µg/ml, respectively.	Katrin et al., 2011
	<i>n</i> -Hexane, chloroform, ethyl acetate and methanol	Leaves	MTT assay	Human hepatoma cell lines (HepG2)	- The <i>n</i> -hexane and chloroform did not show cytotoxic effect -Ethyl acetate and methanol extracts were found to have significant mild cytotoxic effect with IC <sub>50</sub> 32.5 and 40 µg/ml, respectively.	Yosie et al., 2011
	-	-	Bioactivity-guided DLBS1425	Breast cancer cell line (MDA-MB-231)	Results suggested that DLBS1425 exhibited as potent anticancer agent that targets genes involved in cell survival and apoptosis in MDA-MB-231 cell line.	Tandrasmita et al., 2010 and Tjandrawinata et al., 2010
Antidiabetic activity	<i>n</i> -Butanol, ethyl acetate, methanol and boiled water	Fruit	$\alpha$ -glucosidase inhibitory activity and hypoglycemic effect by oral administration in rats	Male rats strain Wistar, about 6 months old, weight 250 – 350 g	-The highest $\alpha$ -glucosidase inhibitory activity was shown by the <i>n</i> -butanol extract of young and ripe fruits followed by ethyl acetate and methanol extract.  -The boiled water and <i>n</i> -butanol extracts were also found to have hypoglycemic effect by reducing the blood glucose concentration of rats with effective dose ranging between $1.0 \times 10^{-3}$ – $6.5 \times 10^{-3}$ mg/g.	Sugiwati et al., 2006

**Table 3.** Summary of pharmacological activities of extracts from *P. macrocarpa* (continued)

Activity	Extract	Plant part	Assay / Method	Organism / cells used	Results / observation	Reference
Antidiabetic activity	-	Fruit powder	Hypoglycemic activity on blood glucose levels in glucose loading healthy volunteers	Healthy volunteers	The effective dose of mesocarp fruit powder to decrease the blood glucose level in healthy volunteers was significantly found at concentration of 500 mg.	Meiyanti et al., 2006
	Ethyl acetate	Fruit	Insulin activity	Male white mice, 3 - 4 weeks old, weight 20-30 g	-The study revealed the potential of ethyl acetate extract to improve insulin sensitivity in hyperglycemic rats by decreasing the blood glucose level. -The highest activity of insulin activity was 18.3%, 9 min after the rats being injected with glucose and exogenous insulin.	Muhtadi et al., 2008
	Ethanol	Fruit	Insulin secretion assay	Insulin secreting cell (BRIN-BD11)	The study has demonstrated the antidiabetic properties of ethanol extract as $\alpha$ -glucosidase inhibitor and as insulin secretagog in clonal glucose responsive insulin secreting cell (BRIN-BD11).	Suparto et al., 2008
	Methanol	Fruits pericarp	Anti-hyperglycemic activity	Normal and diabetic rats	The methanol extract was found to have potential anti-hyperglycemic effect by lowering blood glucose at 56.3% and 58.3% after 12 days of treatment.	Ali et al., 2012
	Methanol and water	Fruits	Protective effect of extracts on renal histological changes of alloxan-induced diabetes	Male Sprague Dawly rats, 6-8 weeks old, weight 160 - 200 g	Both extracts have potential to restore glomerular hypertrophy and improved glomerulosclerosis in alloxan-induced diabetes.	Sulistyo-ningrum et al., 2013
	Fraction: Methanol Subfractions: Chloroform, ethyl acetate, <i>n</i> -butanol and aqueous	Fruits	Bioassay-guided antidiabetic activity	-	-The study identified the sub-fraction that which contain flavonoid in abundance. -The sub-fraction was proved to have potent antidiabetic activity by inhibiting rat intestinal glucose transportation and absorption.	Atangwho et al., 2012
Anti-inflammatory activity	Pericarp, mesocarp and seed	Fruits	Nitric oxide (NO) synthesis induced by LPS/IFN- $\gamma$ assay	Macrophage RAW 264.7 cell lines	Extracts from pericarp and mesocarp showed notable anti-inflammatory effect with percentage of inhibition 63.4% and 69.5%, respectively.	Hendra et al., 2011
Antioxidant activity	Methanol and ethanol	Different parts of young and old fruits	DPPH assay	-	Most of the parts from young and old fruits showed potent antioxidant activity with scavenging inhibition ranging between 38.4 - 48.1%.	Soeksmanto et al., 2007
	Ethyl acetate, <i>n</i> -butanol and water				The highest antioxidant activity was shown by <i>n</i> -butanol extracts of young fruits with IC <sub>50</sub> 41.07 ppm.	
	<i>n</i> -Hexane, chloroform, ethyl acetate and methanol	Leaves	DPPH assay	-	-The <i>n</i> -hexane and chloroform extracts were exhibited as moderate antioxidant properties with percentage of inhibition 59% and 69%, respectively. - Ethyl acetate and methanol extracts were exhibited as strong antioxidant activity with percentage of inhibition 76% and 79%, respectively.	Yosie et al., 2011
	Methanol	Pericarp, mesocarp and seed	Ferric thiocyanate (FTC), thiobarbituric acid (TBA), radical scavenging method (DPPH), ferric-reducing antioxidant power (FRAP) and nitric oxide scavenging method (NO)	-	The results revealed the strong antioxidant properties of pericarp and mesocarp on FRAP assays with percentage inhibition of 92.35% and 78.78%, respectively.	Hendra et al., 2011

**Table 3.** Summary of pharmacological activities of extracts from *P. macrocarpa* (continued)

Activity	Extract	Plant part	Assay / Method	Organism / cells used	Results / observation	Reference
Antibacterial activity	Methanol	Pericarp, mesocarp and seed	Disc diffusion method	<i>B. cereus</i> , <i>B. subtilis</i> , <i>E. aerogenes</i> , <i>E. coli</i> , <i>K. pneumoniae</i> , <i>M. luteus</i> , <i>P. aeruginosa</i> and <i>S. aureus</i>	All parts including pericarp, mesocarp and seed were exhibited weak to moderate antibacterial activity against all pathogenic tested bacteria with inhibition zone ranging from 0.93 cm – 2.33 cm.	Hendra et al., 2011
	<i>n</i> -Hexane, chloroform, ethyl acetate and methanol	Leaves	Mueller Hinton agar well diffusion method	<i>E. coli</i> , <i>K. pneumoniae</i> , <i>P. aeruginosa</i> , <i>S. ubellis</i> , <i>S. aureus</i> and <i>B. cereus</i>	The highest activity was shown by ethyl acetate and methanol extracts against <i>B. cereus</i> and <i>S. aureus</i> with inhibition zone diameter ranging between 15 mm – 27 mm.	Yosie et al., 2011
Antifungal activity	Methanol	Pericarp, mesocarp and seed	Agar well diffusion assay	<i>A.niger</i> , <i>F.oxysporum</i> , <i>G. lucidum</i> and <i>M. indicus</i>	The results showed that only seed extract was displayed antifungal properties against <i>A. niger</i> at concentration of 0.3 mg/well.	Hendra et al., 2011
Vasorelaxant activity	Chloroform and methanol	Fruits	Vasodilator assay	Noradrenaline-induced contraction of isolated rat aorta	The results demonstrated the moderate vasorelaxant activity of chloroform extract.	Oshimi et al., 2008
Antihypertensive activity	Petroleum ether, ethyl acetate and methanol	Fruits and leaves	Inhibitory activity	Angiotensin converting enzyme (ACE)	All extracts from the fruits and leaves were found to have the highest inhibitory activity against ACE with IC <sub>50</sub> ranging between 102 – 189 µg/ml.	Rinayanti et al., 2013

This finding led to phytochemical screening, revealing the presence of flavonoids, terpenoids, and tannins in the methanol extract. These classes of compounds were suggested to be the major contributors to its antidiabetic properties (Ali et al., 2012).

Sulistiyoningrum et al. (2013) discovered the protective effect of the methanol and water extracts of *P. macrocarpa* on renal histological changes of alloxan-induced diabetes. The results concluded that both extracts were able to restore glomerular hypertrophy and improved glomerulosclerosis in alloxan-induced diabetes. Bioassay-guided antidiabetic study on the extract from the fruits successfully identified the active sub-fraction that had flavonoids in abundance. The sub-fraction was proved to have potent antidiabetic activity by inhibiting rat intestinal glucose transportation and absorption. (Atangwho et al., 2012).

#### Anti-inflammatory activities

Different parts of the fruits of *P. macrocarpa* were screened for their anti-inflammatory activity using the nitric oxide (NO) synthesis in macrophage RAW 264.7 cell lines induced by the LPS/IFN- $\gamma$  assay. Extracts from the pericarp and mesocarp showed notable anti-inflammatory effect with percentage of inhibition of 63.4% and 69.5%, respectively (Hendra et al., 2011). Study on anti-inflammatory activity was performed on the major compound from the fruits identified as phalerin (1). This compound showed low inflammatory effect since it decreased the inflammation twice lower than the standard, Naproxen at dose of 22.5 mg/kg body weight (Mariani et al., 2010). Anti-inflammatory activity of phalerin (1) was also determined by using the lipoxygenase (LOX), hyaluronidase (HYA) and xanthine oxidase (XO) assays. The results showed that phalerin (1) had mild anti-inflammatory properties in the XO and LOX assays with percentage of inhibition 34.8% and 23.5%, respectively. Meanwhile, phalerin (1) did not exhibit any inflammatory effect in the HYA assay (Fariza et al., 2012).

#### Antioxidant activities

Antioxidant activity of the methanol and ethanol extracts from different parts of young and old fruits of *P. macrocarpa* was

evaluated using the free-radical-scavenging method (DPPH). Most of the parts from young and old fruits showed potent antioxidant activity with scavenging inhibition ranging between 38.4 - 48.1%. Further fractionation of the active extracts was carried out to give ethyl acetate, *n*-butanol and water extracts. The highest antioxidant activity was observed in the *n*-butanol extract of the young fruits with IC<sub>50</sub> of 41.07 ppm (Soeksmanto et al., 2007). The same DPPH method was also performed to determine the antioxidant activity of different polarity of extracts from the leaves of *P. macrocarpa*. Ethyl acetate and methanol extracts exhibited strong antioxidant activity with 76% and 79% inhibitions, respectively. Meanwhile, the *n*-hexane and chloroform extracts displayed moderate antioxidant properties with 59% and 69% inhibitions, respectively (Yosie et al., 2011).

Various *in vitro* model systems such as ferric thiocyanate, thiobarbituric acid, DPPH, ferric-reducing antioxidant power (FRAP) and nitric oxide (NO) scavenging method were used to characterize the antioxidant properties of different parts of the fruits. The results revealed the strong antioxidant properties of the pericarp and mesocarp in the FRAP assays with percentage inhibition of 92.35% and 78.78%, respectively (Hendra et al., 2011). In addition, antioxidant study of 2,6,4'-trihydroxy-4-methoxybenzophenone (8) and macronone (26) performed using the DPPH method. Interestingly, compound (8) displayed strong antioxidant properties (IC<sub>50</sub> 10.57 µg/ml) while macronone (26) had weak activity (IC<sub>50</sub> 240.14 µg/ml) (Susilawati et al., 2011; 2012).

#### Antimicrobial activities

Antibacterial activity of various parts of *P. macrocarpa* fruits was studied using the disc diffusion method against eight bacterial strains, i.e., *Bacillus cereus*, *Bacillus subtilis*, *Enterobacter aerogenes*, *Escherichia coli*, *Klebsiella pneumoniae*, *Micrococcus luteus*, *Pseudomonas aeruginosa* and *Staphylococcus aureus*. All parts including the pericarp, mesocarp and seeds exhibited weak to moderate antibacterial activity against all pathogenic bacteria strains with inhibition zones ranging from 9.3 – 23.3 mm (Hendra et al., 2011). In the same study, the antifungal activity was evaluated using the agar well diffusion assay against *Aspergillus niger*, *Fusarium*

*oxysporum*, *Ganoderma lucidum* and *Mucor indicus*. The results showed that only seed extract was active against *A. niger* at a concentration of 0.3 mg/well (Hendra et al., 2011).

Different polarities of extracts from the leaves of *P. macrocarpa* including the n-hexane, chloroform, ethyl acetate and methanol extracts were evaluated for their antibacterial activity against *E. coli*, *K. pneumonia*, *P. aeruginosa*, *Streptococcus ubellis*, *Streptococcus aureus* and *B. cereus*. Mueller Hinton agar well diffusion method was used to determine the susceptibility of bacteria tests. The highest activity was shown by ethyl acetate and methanol extracts against *B. cereus* and *S. aureus* with inhibition zone diameter ranging between 1527 mm (Yosie et al., 2011).

### Toxicity

The general toxicity of 29-norcucurbitacin derivatives; desacetyl-fevicordin A (14), fevicordin A (15), fevicordin A glucoside (16) and fevicordin D glucoside (17), isolated from this plant was evaluated by the brine shrimp (*Artemia salina*) lethality assay. All compounds showed variable general toxicity with LD<sub>50</sub> values ranging from 3 – 12 ppm (Kurnia et al., 2008).

### Vasorelaxant activity

The vasorelaxant activity of the extracts and compounds (1, 9, 28) isolated from the fruits of *P. macrocarpa* was evaluated against noradrenaline-induced contraction of isolated rat aorta. The results demonstrated the moderate vasorelaxant activity of the chloroform extract, while icariside C3 (9) showed a slow vasorelaxant activity. Phalerin (1) and mangiferin (28) did not show any vasorelaxant effect (Oshimi et al., 2008).

### Antihypertensive activity

Study on the antihypertensive activity of nine medicinal plants from Indonesia was conducted against angiotensin converting enzyme. The nine tested plants were *Scurulla artopurpurea*, *catharanthus roseus*, *Swietenia mahogany*, *Persea Americana*, *Oxalis corniculata*, *P. macrocarpa*, *Gynura procumbens*, *Melia azedarach* and *Hisbiscus rosinensis*. Interestingly, all extracts from the fruits and leaves of *P. macrocarpa* displayed the highest level of inhibitory activity against acetylcholine esterase with IC<sub>50</sub> ranging between 102 – 189 µg/ml (Rinayanti et al., 2013).

### CONCLUSION

In this review, we have reviewed the relevant literature to assemble the ethnomedicinal, phytochemical and pharmacological properties of *P. macrocarpa* (Scheff.) Boerl. This plant is used as folk remedies in both traditional as well as modern system of medicine to treat various diseases and illnesses. Various types of compounds with diverse chemical structures present in this plant are responsible for varied pharmacological and medicinal properties. Reported data show that the plant possesses promising anticancer, antidiabetic, anti-inflammatory, antioxidant, antimicrobial, antihypertensive, toxicity and vasorelaxant activities. However, in view of the wide range of medicinal uses of *P. macrocarpa*, it is necessary to conduct further clinical and pharmacological studies at molecular level to investigate the potential of this plant, because most of the activity reported is based only on their *in vitro* assays. Similarly, additional studies have to be carried out in order to establish the potential of the extracts of *P. macrocarpa* in the development of new therapeutic drugs and to provide the basis for future research on the application of medicinal plants.

### ACKNOWLEDGEMENTS

The authors acknowledge Ministry of Higher Education (MOHE), Malaysia for the financial support under Research University Grant (RUG) vote number Q.J130000.2626.08J14 and the Faculty of Science, Universiti Teknologi Malaysia for providing the necessary support for this study.

### CONFLICT OF INTEREST

The authors declare no conflict of interest.

### REFERENCES

Ali RB, Atangwho IJ, Kuar N, Mohamed EAH, Mohamed AJ, Asmawi MZ and Mahmud R. Hypoglycemic and anti-hyperglycemic study of *Phaleria macrocarpa* fruits pericarp. J Med Plants Res. 2012;6:1982-1990.

Altaf R, Asmawi MZ, Dewa A, Sadikun A and Umar MI. Phytochemistry and medicinal properties of *Phaleria macrocarpa* (Scheff.) Boerl. extracts. Pharmacogn Rev. 2013;7:73-80.

Angiosperm Phylogeny Group. An update of the angiosperm phylogeny group classification for the orders and families of flowering plants: APG II. Botanical J Linnean Society. 2003;141:399-436.

Atangwho IJ, Ali RB, Kuar N, Asmawi MZ Mahmud R and Ahmad M. Bioassay-guided antidiabetic study of *Phaleria macrocarpa* fruit extract. Molecules. 2012;17:4986-5002.

Cragg GM, Newmann DJ. Natural products: A continuing source of novel drug leads. Biochim Biophys Acta. 2013;1830:3670-3695.

De Padua LS, Bunyapraphatsara N, Lemmens RHMS. Plant resources of South East Asia, Medical and poisonous plants 1 (PROSEA). (Leiden, Netherlands: Backhuys Publishers), pp. 36-38, 1999.

De Pasquale A. Pharmacognosy: the oldest modern science. J Ethnopharmacol. 1984;11:1-16.

Diantini A, Subarnas A, Supriyatna, Levita J, Abdulah R, Achmad TH, Faried A, Faried LS, Julaeha E, Kurnia D, Wardhani SR and Koyama H. Cytotoxicity of fevicordin-A from *Phaleria macrocarpa* (Scheff.) Boerl. on P388, HeLa, Caski, TE-2, TE-8 and Prepuce's Fibroblast cells. J Med Res. 2012;1:1-5.

Faried A, Kurnia D, Faried LS, Usman N, Miyazaki T, Kato H, and Kuwano H. Anticancer effects of gallic acid isolated from Indonesia herbal medicine, *Phaleria macrocarpa* (Scheff.) Boerl, on human cancer cell lines. Int J Oncol. 2007;30:605-613.

Fariza IN, Fadzureena J, Zunoliza A, Chuah AL, Pin KY and Adawiyah I. Anti-inflammatory activity of the major compound from methanol extract of *Phaleria macrocarpa* leaves. J Appl Sci. 2012;12:1195-1198.

Gordaliza M. Natural products as leads to anticancer drugs.

- Clin Transl Oncol. 2007;9:767-776.
- Gurib-Fakim A. Medicinal plants: Traditions of yesterday and drugs of tomorrow. Mol Aspects Med. 2006;27:1-93.
- Hanus-Fajerska E, Wiszniewska A, Czaicki P. Effectiveness of Daphne L. (Thymelaeaceae) In vitro propagation, rooting of microshoots and acclimatization of Plants. Acta Agrobotanica. 2012;65:21-28.
- Harmanto N. Conquering disease in unison with mahkota dewa, *Phaleria macrocarpa*. 1<sup>st</sup> ed. (North Jakarta, Indonesia: Mahkotadewa Indonesia), p.14, 2003.
- Hendra R, Ahmad S, Sukari A, Shukor MY and Oskoueian E. Antioxidant, anti-inflammatory and cytotoxicity of *Phaleria macrocarpa* (Boerl.)Scheff fruit. BMC Complement Altern Med. 2011;11:1-10.
- Hendra R, Ahmad S, Sukari A, Shukor MY and Oskoueian E. Flavanoid analyses and antimicrobial activity of various parts of *Phaleria macrocarpa* (Scheff.) Boerl fruit. Int J Mol Sci. 2011;12:3422-3431.
- Herber BE. Pollen morphology of the Thymelaeaceae in relation to its taxonomy. Plant Sys Evol. 2002;232:107-121.
- Herber BE. Flowering plants dicotyledons. In The families and genera of vascular plants. Vol. 5. Bayer C, Kubitzki K ed. (New York, USA: Springer Berlin Heidelberg), pp. 373-396, 2003.
- Hou D. Thymelaeaceae. In Flora malesiana series I. Vol. 6. Van Steenis, C.G.G.J. ed. (Groningen, Netherlands: Wolter-Noordhoff Publishing), pp. 1-15, 1960.
- Katrin E, Selvie, Winarno H. Chromatogram profiles and cytotoxic activity of irradiated mahkota dewa (*Phaleria macrocarpa* (Scheff.) Boerl.) leaves. Atom Indones. 2011;37:17-23.
- Kim WJ, Veriansyah B, Lee YW, Kim J, Kim JD. Extraction of mangiferin from mahkota dewa (*Phaleria macrocarpa*) using subcritical water. J Ind Eng Chem. 2010;16:425-430.
- Kubitzki K and Bayer C. The families and genera of vascular plants: Flowering plants dicotyledons. Springer Berlin Heidelberg. 2003;5:383.
- Kurnia D, Akiyama K and Hayashi H. 29-Norcucurbitacin derivatives isolated from the Indonesian medicinal plant, *Phaleria macrocarpa* (Scheff.) Boerl. Biosci Biotechnol Biochem. 2008;72:618-620.
- Lisdawati V, Wiryowidagdo S, Kardono L and Broto S. Isolasi dan elusidasi struktur senyawa lignin dan asam lemak dari ekstrak daging buah *Phaleria macrocarpa*. Bul Penel Kesehatan. 2007;35:115-124.
- Mariani R, Wirasutisna K, R, Nawawi A and Adnyana IK. Anti-inflammatory activity of dominant compound of mahkota dewa fruit *Phaleria macrocarpa* (Scheff.) Boerl. Indonesian J Pharm. 2010;21:129-133.
- Meiyanti, Dewoto, HR and Suyatna FD. Hypoglycemic effect of mahkota dewa mesocarp fruit (*Phaleria macrocarpa* (Scheff.) Boerl.) on glucose blood level in glucose loading healthy volunteers. Universa Medicina. 2006;25:114-120.
- Muhtadi A, Susilawati Y and Zakaria AD. The activity of isolate from ethyl acetate fraction of mahkota dewa (*Phaleria macrocarpa* (Scheff.) Boerl.) fruits on insulin sensitivity in hyperglycaemic mouse. Proceeding Int Seminar on Chemistry, Jatinangor, Indonesia. 2008;527-530.  
<http://chemistry.unpad.ac.id/isc-proceeding/2008/Pdf/PP/0527-0530%20PP053%20Ahmad%20Muhtadi.pdf>
- Oshimi S, Zaima K, Matsuno Y, Hirasawa Y, Izuka T, Studiawan H, Indrayanto G, Zaini NC and Morita H. Studies on the constituents from the fruits of *Phaleria macrocarpa*. J Nat Med. 2008;62:207-210.
- Pertamawati. Pengaruh sitotoksik ekstrak buah mahkota dewa (*Phaleria macrocarpa* (Scheff.) Boerl.) terhadap sel kanker lestari HeLa. J Sains dan Teknologi Indones. 2007;9:39-43.
- Rahmawati E, Dewoto HR and Wuyung PE. Anticancer activity study of ethanol extract of Mahkota dewa fruit pulp (*Phaleria macrocarpa* (Scheff.) Boerl.) in C3H mouse mammary tumor induced by transplantation. Med J Indones. 2006;15:217-222.
- Rates SMK. Plants as source of drugs. Toxicon. 2001;39:603-613.
- Rinayanti A, Radji M, Mun'im A and Suyatna FD. Screening angiotensin converting enzyme (ACE) inhibitor activity of antihypertensive medicinal plants from Indonesia. International Journal of Pharmacy Teaching and Practices. 2013;4:527-532.
- Samuelsson G. A textbook of pharmacognosy. In Drugs of natural origin. (Stockholm, Sweden: Swedish Pharmaceutical Press), pp.473-575, 2004.
- Saufi A. Lignans in *Phaleria macrocarpa* (Scheff.) Boerl. and in *Linum flavum* var. *compactum* L. (Maastricht, Germany: Shaker Verlag GmbH), pp.13-14, 2007.
- Simanjutak P. Identifikasi senyawa kimia dalam buah Mahkota dewa (*Phaleria macrocarpa*), Thymelaeaceae. J Ilmu Kefarmasian Indones. 2008;6:23-28.
- Soeksmanto A, Hapsari Y, Simanjutak P. Antioxidant content of parts of mahkota dewa, *Phaleria macrocarpa* [Scheff] Boerl. (Thymelaeaceae). Biodivertas. 2007;8:92-95.
- Stevens PF. Three new species of *Phaleria* (Thymelaeaceae) from Papuasia. J Arnold Arboretum. 1974;55:264-268.
- Sugiwati S, Kardono LBS and Bintang M.  $\alpha$ -Glucosidase inhibitory activity and hypoglycemic effect of *Phaleria macrocarpa* fruit pericarp extracts by oral administration to rats. J Appl Sci. 2006;6:2312-2316.
- Sulistiyoningrum E, Setiawati and Ismaulidiya FR. *Phaleria macrocarpa* (Scheff.) Boerl. improved renal histological changes in alloxan-induced diabetic rats. Int J Med Plants Alt Med. 2013;1:87-92.
- Suparto IH, Arfianti N, Septiawati T, Triwahyuni W and Iskandriati D. Ethanol extract of mahkota dewa (*Phaleria macrocarpa* (Scheff.) Boerl.) fruit with in-vitro antidiabetic activities. Proceeding. International Seminar on Chemistry,

Jatinangor, Indonesia. 2008;285-288.

<http://www.chemistry.unpad.ac.id/isc-proceeding/2008/Pdf/OP/0285-0288%20OP072%20-%20Irma%20Suparto.pdf>

Susilawati. Isolasi metabolit sekunder dari buah, kulit batang dan daun Mahkota Dewa (*Phaleria macrocarpa* (Scheff.) Boerl.) serta uji antioksidannya. (Yogyakarta, Indonesia: PhD dissertation of Universitas Gadjadara Mada), pp. 250-267, 2012. [http://repository.ugm.ac.id/digitasi/download.php?file=3169\\_RD-201301037-susillawati.pdf](http://repository.ugm.ac.id/digitasi/download.php?file=3169_RD-201301037-susillawati.pdf)

Susilawati, Matsjeh S, Pranowo HD and Anwar C. Antioxidant activity of 2,6,4'-trihydroxy-4-methoxybenzophenone from ethyl acetate extract of leaves of mahkota dewa (*Phaleria macrocarpa* (Scheff.) Boerl.). *Indo J Chem.* 2011;11:180-185.

Susilawati, Matsjeh S, Pranowo HD and Anwar C. Macrone, a novel diepoxy lignan from bark of mahkota dewa (*Phaleria macrocarpa* (Scheff.) Boerl.) and its antioxidant activity. *Indo J Chem.* 2012;12:62-69.

Tambunan RM and Simanjutak P. Determination of chemical structure of antioxidant compound benzophenone glycoside from n-butanol extract of the fruits of mahkota dewa (*Phaleria macrocarpa* (Scheff.) Boerl.). *Majalah Farmasi Indones.* 2006;17:184-189.

Tandrasasmita OM, Lee JS, Baek SH and Tjandrawinata R. Induction of cellular apoptosis in human breast cancer by DLBS1425, a *Phaleria macrocarpa* compound extract, via downregulation of P13-kinase/AKT pathway. *Cancer Biol Ther.* 2010;10:814-823.

Tjandrawinata RR, Arifin PF, Tandrasasmita OM, Rahmi D and Aripin A. DLBS1425, a *Phaleria macrocarpa* (Scheff.) Boerl. extract confers antiproliferative and proapoptosis effects via eicosanoid pathway. *J Exp Ther Oncol.* 2010;8:187-201.

Wang Y, Gilbert MG, Mathew B, Brickell CD, and Nevling LI. Flora of China In Thymelaeaceae. Vol. 13. Wu ZY, Raven PH, Hong DY. eds. (Beijing, China; Science Press), pp. 213-250, 2007.

Winarno H and Katrin WE. Benzophenone glucoside isolated from the ethyl acetate of the bark of mahkota dewa (*Phaleria macrocarpa* (Scheff.) Boerl.) and its inhibitory activity on leukemia L1210 cell line. *Indo J Chem.* 2009;9:142-145.

Yosie A, Effendy MAW, Sifzizul TMT and Habsah M. Antibacterial, radical-scavenging activities and cytotoxicity properties of *Phaleria macrocarpa* (Scheff.) Boerl leaves in HEPG2 cell lines. *Int J Pharm Sci Res.* 2011;2:1700-1706.

Zhang YB, Xu XJ, Liu HM. Chemical constituents from mahkota dewa. *J Asian Nat Prod Res.* 2006;8:119-123.

Zhang SY, Zhang QH, Zhao W, Zhang X, Zhang Q, Bi YF and Zhang YB. Isolation, characterization and cytotoxic activity of benzophenone glucopyranoside from mahkota dewa (*Phaleria macrocarpa* (Scheff.) Boerl.). *Bioorg Med Chem Lett.* 2012;22:6862-6866.