



LJMU Research Online

Kinderman, P, Tai, S, Pontin, E, Schwannauer, M, Jarman, I and Lisboa, P

Causal and mediating factors for anxiety, depression and well-being

<http://researchonline.ljmu.ac.uk/id/eprint/2265/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Kinderman, P, Tai, S, Pontin, E, Schwannauer, M, Jarman, I and Lisboa, P (2015) Causal and mediating factors for anxiety, depression and well-being. BRITISH JOURNAL OF PSYCHIATRY, 206 (6). pp. 456-460. ISSN 0007-1250

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

Causal and mediating factors for anxiety, depression and well-being

Peter Kinderman, Sara Tai, Eleanor Pontin, Matthias Schwannauer, Ian Jarman and Paulo Lisboa

Background

The relationship between well-being and mental ill health is complex; people may experience very low levels of well-being even in the absence of overt mental health problems.

Aims

This study tested the hypothesis that anxiety, depression and well-being have different causal determinants and psychological mediating mechanisms.

Method

The influence of causal and mediating factors on anxiety, depression and well-being were investigated in a cross-sectional online questionnaire survey hosted on a UK national broadcasting website.

Results

Multivariate conditional independence analysis of data from 27 397 participants revealed different association pathways for the two constructs. Anxiety and depression were associated with negative life events mediated by rumination; low levels of subjective well-being were associated with material deprivation and social isolation, mediated by adaptive coping style.

Conclusions

Our findings support the 'two continua' model of the relationship between psychological well-being and mental health problems, with implications for both treatment and prevention.

Declaration of interest

None.

The relationships between mental health, well-being and mental illness are complex, and people may experience very low levels of well-being even in the absence of overt mental health problems.¹ Despite general agreement that these three concepts are important targets for health and social care, there is no consensus about the relationship between them. This conceptual distinction is best exemplified in the 'two continua' model,² which suggests that mental illness and mental well-being are related yet distinct phenomena, rather than ends of a single measurement continuum. One important way to disambiguate 'psychological well-being', 'mental health' and 'mental health problems' is to explore their distinct causal determinants. We report here on an analysis of a unique database of 27 397 participants, using the novel but powerful techniques of conditional independence graphical analysis,^{3,4} testing whether the same patterns of these causal factors and psychological mediators are implicated in the development of mental health problems and mental well-being. Although conditional independence mapping has been used to analyse a range of data in both social and physical sciences, this is the first time that the technique has been used on multivariate data such as these in such a substantial data-set. Conditional independence mapping has similarities with covariance or structural equation modelling, but makes fewer *a priori* assumptions about the expected relationships, allowing all possible permutations of association to be tested simultaneously in a single multivariate model. Identified significant associations can then be explored in more detail to explore the contingent relationships between selected variables.

Method

Participants were self-selected respondents to an online battery of questionnaires hosted on the British Broadcasting Corporation (BBC) website (the Stress Test). In total 32 827 respondents (age 18–85 years) completed the test battery; their mean age was 40.5 years (s.d. = 14.3), 61% were women, 93% were 'White British'

or 'White other' and 88% were from the UK. To determine whether the sample was representative of the UK population, results were compared with national data for England and Wales.⁵

Procedure

The Stress Test was promoted via multimedia formats (television, radio and online) and launched on BBC Radio 4's *All in the Mind*, a flagship documentary focusing on issues of the human mind. The test's URL (www.bbc.co.uk/labuk/experiments/stress/) was publicised on subsequent episodes, related television broadcasts and made available on BBC web pages and social networking sites. Visitors to the Stress Test's homepage accessed the test by signing in using BBC online membership. The test was 'live' for a 4-week period. The test had 12 sections, which took approximately 20 min to complete in total. It was designed by authors P.K. and S.T. and developed in collaboration with BBC Lab UK. Questionnaire items were completed in a fixed order and answers were selected from a drop-down menu. Some tasks were constrained within time limits. On completion, an overview of scores was displayed on a results home page and URL links for comprehensive and tailored feedback based on test scores were presented. Once participants had completed the test they were not permitted to retake it.

Measures

Measures were selected on the basis of theoretical principles and empirical research to provide indicators of latent constructs representative of the components of the biopsychosocial model of mental health and well-being.⁶ Where measures were modified or created for the purposes of this study, other analyses have demonstrated their *post hoc* construct validity.⁷ Demographic data collected included age, gender, ethnic group, occupation, gross annual or weekly household earnings, highest level of formal schooling, occupational status, parents' income, relationship status and number of children. Measured variables to represent

Author:
Title modified
to avoid
anticipating
results
(AQ1)

Author:
Details of the
sample moved
to 'results'
section
(AQ2)

the biological component of the theoretical model were a yes/no response to indicate participants' reports of familial mental health diagnoses by a psychiatrist or general practitioner,⁸ and performance on two cognitive tests to detect response to negative feedback and negative and positive stimuli. These were the 'delayed match to sample' (DMS) and the 'affective go/no go' (AGN) tasks adapted from the Cambridge Neuropsychological Test Automated Battery (CANTAB).^{9–11} These tests were chosen to measure neurological functioning previously associated with anxiety and depression,^{9,10} and therefore supplement the more subjective participant report of familial history of mental health problems. The social component of the model was represented by a novel 11-item questionnaire designed to indicate social relationships with friends and family, and participation in social activities.¹² These items were a combination of Likert scale and yes/no responses. Indicators of the circumstantial component included recent and historical life events. Recent life events were measured using the List of Threatening Experiences Questionnaire (LTE-Q).¹³ Participants also indicated on a five-point Likert scale whether they believed they had historically been physically, sexually or emotionally abused, or bullied at school.¹⁴ The two key psychological processes (response style and attributional style) were measured using an adapted Response Style Questionnaire (RSQ).¹⁵ In the modified version for the Stress Test, participants indicated on a Likert scale their response to stressful situations from a list of coping strategies pertaining to rumination, problem-solving/adaptive or dangerous activities. Rumination was characterised by endorsement of statements such as 'I think about my shortcomings, failings, faults and mistakes'; adaptive coping was characterised by endorsement of statements such as 'I make a plan to overcome a problem'; and dangerous activities encompassed strategies such as 'go shopping with no regard for the debts I may run up' or 'drink alcohol excessively'. Attributional style was measured using an adapted and modified version of the Internal, Personal and Situational Attributions Questionnaire (IPSAQ),¹⁶ to determine the degree to which individuals generate internal, personal or situational causes for an event or situation. Finally, mental health problems were assessed using the well-validated Goldberg Anxiety and Depression (GAD) scales,¹⁷ a paired nine-item self-report symptom inventory designed to assess anxiety and depression. The GAD scale was chosen for its brevity and because it has been found to be a valid and acceptable method of detecting psychological disturbance in the general population.¹⁷ Well-being was measured using the BBC Subjective Well-Being (BBC-SWB) scale,¹⁸ a 24-item questionnaire designed to measure people's subjective experiences across the wide breadth of domains commonly included in the definitions of well-being: psychological well-being, physical health and satisfaction with relationships. The BBC-SWB scale has been found to be a reliable and valid measure for the online assessment of well-being in the general population with good psychometric properties.¹⁹

Statistical analysis

The data-set was subjected to conditional independence graphical analysis,^{3,4} to test the conditional independence between paired variables. Since mutual covariance is common in multivariate analyses – and especially common in the social sciences – the algorithm used in this analysis ensures that each specific test for significance of association is conditional on all associations with other variables connected in the model. The resulting multivariate association structure is represented in a conditional independence map. Any pair of variables that do not carry significant information about each other are disconnected in the map. The remaining associations are then directed according to the relative

strength of conditional probabilities, and subjected also to additional constraints so that the final map is consistent and acyclic, so removing any closed loops. Specific associations between variables may be investigated further using covariance tables. Missing data were deleted listwise to produce complete data for 27 397 participants (retention of 83.5% of the original sample, $n = 32\,827$). The neurocognitive data accounted for a large proportion of missing data owing to invalid recording, probably because of technical error. Missing case analysis revealed no significant difference between those with and without missing data on demographic variables and a selection of measured variables.

The software used in this study, CiMap version 1.0,³ employs a computationally efficient algorithm,²⁸ with adjustments for repeated tests. It starts from a fully connected undirected map and deletes conditionally independent associations in multiple statistical tests. Conditional mutual information is computed using the equation

$$\tilde{I}(i, j, A) = \sum_{x_i, x_j, x_A} p_{ijA}(x_i, x_j, x_A) \log_2 \frac{p_{ijA}(x_i, x_j | x_A)}{p_{i|A}(x_i | x_A) p_{j|A}(x_j | x_A)}$$

where p is the joint mass function that is typically approximated with the point estimate n_{ijA}/N .

Larger values of \tilde{I} imply stronger dependencies, and significance tests assumed a χ^2 distribution for independent or conditional independent variables.²⁰

Conditional independence graphical analysis conducts simultaneous tests of association on categorical data. Variables with more than 15 categories (attribution style, dangerous activities response style, rumination response style, well-being score and GAD score) were transformed into deciles. Frequency of childhood abuse was transformed into six categories, owing to the distribution of responses. Family history of mental health problems was transformed to two simple 'yes/no' categories. Other variables (highest level of formal schooling achieved, occupational status, total gross annual or weekly household income, parents' income while growing up, relationship status, number of children, relationships with friends, relationships with family, social activities, frequency of social contacts, total number of life events, adaptive response style and dangerous activities response style) were used in their raw form, with each integer treated as a separate category. In the case of total gross annual income, 12% of participants recorded either 'don't know' or declined to answer the question. As these responses are meaningful (and were associated with occupational status as students or unemployed), these categories were included in the analysis rather than excluded as missing. A summary of the data processing applied to each variable is given as an online supplement (Table DS1).

Results

The majority of respondents held an undergraduate, professional qualification or postgraduate degree (73%), were in full-time employment (52%), earned £20 000 to £29 999 per annum (16%) and were in a relationship (35%), with no children (54%). Compared with national data for England and Wales,⁵ more UK respondents were White, had slightly higher earnings and were better educated than the general population, although comparable on other demographic features. Demographic data for the full sample are presented in Table 1.

The results of the analysis are summarised in the conditional independence map (Fig. 1), which shows variables that are

Author:
Insertion OK?
Please specify
platform
(AQ3)

Author:
Other variables
not defined;
can you insert,
or refer to
explanation
elsewhere?
(AQ4)

Author:
Table 2 too
large for print
version, has
therefore been
changed to an
online supplement. Other
tables renumbered
(AQ5)

Author:
73% in Table 1,
or did that
value include
part-time
work?
(AQ6)

Author:
Sentence not
very clear;
minority in re-
lationship (cf.
Table 1)? Please
confirm %s
(AQ7)

AUTHOR'S PROOF

Table 1 Demographic details of whole sample and sample after removal of missing data

	Whole sample (<i>n</i> = 32 827) % (<i>n</i>)	Missing data removed (<i>n</i> = 27 397) % (<i>n</i>)
Ethnic group		
White (British, Irish, other)	85.7 (28 105)	92.8 (25 434)
Black minority ethnic	5.6 (1850)	5.8 (1612)
Rather not say or missing	8.7 (2872)	1.3 (351)
Highest level of schooling achieved		
Did not complete schooling	2.3 (747)	2.2 (601)
In education until age 18 years	24.4 (8032)	24.7 (6766)
Degree or professional qualification	73.2 (24 048)	73.1 (20 030)
Occupational status		
In education	11.8 (3885)	11.4 (3109)
In employment	73.1 (23 991)	73.7 (20 195)
Other	15.1 (4951)	14.9 (4093)
Total gross annual household income		
Up to £39 999 (\$65 000) per annum	51.6 (16 936)	51.9 (14 206)
Above £39 999 (\$65 000) per annum	34.2 (11 225)	36.0 (9851)
Don't know/prefer not to say	14.2 (4666)	12.1 (3340)
Estimated parents' income while growing up		
Lower than 50% population	51.2 (16 824)	50.8 (13 913)
Higher than 50% population	48.8 (16 003)	49.2 (13 484)
Relationship status		
In a relationship	74.3 (24 389)	73.2 (20 062)
Single	25.7 (8438)	26.8 (7335)
Number of children		
None	54.4 (17 900)	53.7 (14 717)
One or more	45.6 (14 927)	46.3 (12 680)

Author:
Should 85.7
be 85.6?
(AQ10)

Author:
Should 5.8
be 5.9?
(AQ11)

Author:
Should 12.1
be 12.2?
(AQ12)

Author:
Should 48.8
be 48.7?
(AQ13)

Author:
Should 54.4
be 54.5?
(AQ14)

significantly associated ($P < 0.05$), having tested for all possible associations. Each connecting line in Fig. 1 therefore represents the associated pairwise mutual information between the respective variables. Higher mutual information corresponds to a stronger pairwise association, so for this map the strongest pairwise association is 0.35 and the weakest 0.01, but all are significant at the 95% confidence level. Inspecting the map in more detail reveals that the variable 'mental health problems' was directly and significantly associated with two other variables: 'rumination' (0.35) and 'number of negative life events' (0.05). This means that, given all the possible associations of 'mental health problems' with every other variable in the data, it was only these two variables that conveyed statistically significant conditional mutual information. In contrast, 'well-being' was directly and significantly associated with eight other variables: adaptive coping (0.1), frequency of social contacts (0.05), social activities (0.19), relationships with friends (0.11), relationships with family (0.05), parental income (0.02), household income (0.03) and highest level of education (0.02). Surprisingly, well-being and mental health had no directly associated variables in common. All the variables included in the analysis (listed in online Table DS1) were represented on the map, with the exception of 'attribution style', which had no significant association with any other variable.

Examination of the map of conditional associations reveals a cluster of social variables associated with well-being, accompanied by adaptive coping, previously identified as a mediating variable.⁷ In contrast, a cluster of variables including negative life events, abuse and a familial history of mental health problems was associated with mental health problems (anxiety and depression), accompanied by rumination, again previously identified as a mediating variable.⁷ This analysis does not suggest that well-being and mental health problems are unrelated, merely that conditional

mutual information concerning their association is best represented through a pattern of indirect associations as presented in Fig. 1 and involving several different pathways. The analysis does, however, strongly suggest that the two concepts are unlikely to be two poles of a single continuum. These associations were explored in greater detail using simple contingency tables based on median splits. Table 2 presents the data from the BBC-SWS and the GAD, with a clear inverse relationship between the two variables.

Different pathways for mental health problems and well-being

Contingency tables also permit more detailed exploration of three-way associations revealed by the conditional independence map. These analyses reveal clearly differentiated mediated pathways leading from causal factors to well-being and mental health problems. The contingent relationship between life events, rumination and mental health problems is presented in Table 3, with simple binomial categorisation to permit straightforward interpretation. For the whole sample the relationship between life events and mental health problems was relatively weak, with only a slightly disproportionate number of people falling into the cell representing both higher levels of mental health problems and more life events. The mediating effects of rumination are clear, however, in the two succeeding parts of Table 3. For participants reporting low levels of rumination, prevalence of mental health problems was low, even in the presence of more frequent life events (17%). In contrast, for participants reporting high levels of rumination, more frequent life events were associated with a markedly higher rate of mental health problems (56%).

Similar contingency tables explored the three-way associations between well-being, social activities and adaptive coping (Table 4). For the whole sample the relationship between social activities and well-being was clear, with higher levels of social activities associated with higher levels of subjective well-being. Again, however, adaptive coping acted as a mediating protective factor, as high levels of adaptive coping were associated with disproportionately fewer people reporting low levels of subjective well-being in the absence of social activity.

Discussion

Our findings provide clear quantitative support to the hypothesis that subjective well-being and mental health problems have distinct causal pathways,^{1,2} with different causal factors and psychological mediators, despite the high correlation between these concepts. These findings are strongly supportive of previous reports that a family history of mental health difficulties, social deprivation and traumatic or abusive life experiences are associated with higher levels of anxiety and depression, mediated by adaptive coping and rumination.⁷ The mediating role of psychological processes was strongly evidenced in the relationship between life events and mental health problems, in that there was a striking distinction between the rates of mental health problems following frequent life events for people with low, as opposed to high, levels of rumination. Low levels of subjective well-being were associated with social isolation and low levels of adaptive coping. On the other hand, mental health problems were associated with negative life events and rumination.

Because this study was conducted on a cross-sectional rather than a longitudinal basis, we were able to identify patterns of association rather than direct causal relationships. For instance, rumination is often considered a symptom of depression rather than a cause or a mediator. Our findings require further

AUTHOR'S PROOF

Table 4 Contingent relationship between well-being and social activity, stratified by levels of adaptive coping, based on median splits of each variable

	Low level of social activity <i>n</i> (%)	High level of social activity <i>n</i> (%)
All participants (<i>n</i> = 27 397)		
Well-being		
Low	8995 (32.8)	4434 (16.2)
High	4259 (15.5)	9709 (35.4)
Low level of adaptive coping (<i>n</i> = 10 914)		
Well-being		
Low	4875 (44.7)	1930 (17.7)
High	1396 (12.8)	2713 (24.9)
High level of adaptive coping (<i>n</i> = 16 483)		
Well-being		
Low	4120 (25.0)	2504 (15.2)
High	2863 (17.4)	6996 (42.4)

relationships, can become depressed. These findings imply that if bad things happen to them (they experience negative life events), and if they ruminate about those events, they are much more likely to become anxious or depressed. Clearly, this analysis can be extended to a wide range of other causes and mediators not included in this study. These findings can also partially explain how people who do not show signs of specific mental health problems such as anxiety and depression can have low levels of well-being or fail to flourish. If socially isolated and relatively lacking in adaptive coping responses (such as problem-solving and emotional self-regulation), people are more likely to experience low levels of well-being, but are unlikely to experience mental health problems if otherwise not exposed to the causal mechanisms described above.

Implications of the study

Practically, these findings suggest that interventions aimed at maximising well-being and interventions aimed at preventing or treating mental health problems should be complementary but different. We found that people who ruminated were three times more likely to develop anxiety or depression following negative life events, and people who lacked adaptive coping skills were three times more likely to have low subjective well-being in the face of social deprivation, but that these two pathways were independent of each other. These findings have important implications for current policies in health and social care. Specifically, they strongly suggest that conceptually distinct interventions are needed; services aimed at treatment and prevention of mental health problems may be very different from those needed to promote and improve subjective well-being. All such clinical decisions must be dictated by individual case formulations, but it seems reasonable to suggest that interventions and social policy aimed at improving social circumstances, addressing financial equity and social justice, building local communities, supporting families, and facilitating relationships on the societal level, and developing individuals' personal capacity to cope adaptively with stressful life circumstances may be best suited to developing subjective well-being. Local authority social services may have an immediate responsibility here, but genuine progress may necessitate wider societal and political action. Interventions aimed at preventing abuse, mitigating the consequences of traumatic personal life events, and evidence-based interventions targeting psychological processes such as rumination (including the now ubiquitous cognitive-behavioural therapy) may be best suited to helping people recover from mental health problems.

Peter Kinderman, PhD, Institute of Psychology, Health and Society, University of Liverpool; **Sara Tai**, ClinPsyD, School of Psychological Science, Manchester; **Eleanor Pontin**, DClPsych, Institute of Psychology, Health and Society, University of Liverpool; **Matthias Schwannauer**, PhD, School of Health in Social Science, Edinburgh; **Ian Jarman**, PhD, **Paulo Lisboa**, PhD, School of Computing and Mathematical Sciences, Liverpool John Moores University, Liverpool, UK

Correspondence: Professor Peter Kinderman, Institute of Psychology, Health and Society, University of Liverpool, Waterhouse Building, Liverpool L69 3GL, UK. Email: p.kinderman@liverpool.ac.uk

First received 24 Feb 2014, final revision 30 Sep 2014, accepted 6 Oct 2014

Acknowledgements

We thank the staff of BBC Lab UK for their support for the development and conduct of the study, and J. Rees for critical discussions and reading of the manuscript.

References

- Keyes CLM. Mental health and/or mental illness? Investigating axioms of the complete state model of health. *J Consult Clin Psychol* 2005; **73**: 539–48.
- Westerhof GJ, Keyes CLM. Mental illness and mental health: the two continua model across the lifespan. *J Adult Dev* 2010; **17**: 110–9.
- Bacciu D, Etschells TA, Lisboa PJG, Whittaker J. Efficient identification of independence networks using mutual information. *Comput Stat* 2013; **28**: 621–46.
- Spirtes P, Glymour C, Scheines R. *Causation, Prediction and Search* (2 ed). MIT Press, 2000.
- Office for National Statistics. *Labour Force Survey: Household Datasets, Spring Quarter*. ONS, 2003.
- Kinderman P. A psychological model of mental disorder. *Harv Rev Psychiatry* 2005; **13**: 206–17.
- Kinderman P, Schwannauer M, Pontin E, Tai S. Psychological processes mediate the impact of familial risk, social circumstances and life events on mental health. *PLoS One* 2013; **8**: e76564.
- Milne BJ, Caspi A, Crump R, Poulton R, Rutter M, Sears MR, et al. The validity of the family history screen for assessing family history of mental disorders. *Am J Med Genet B* 2008; **50**: 41–9.
- Elliott R, Sahakian BJ, Herrod JJ, Robbins TW, Paykel ES. Abnormal response to negative feedback in unipolar depression: evidence for a diagnosis specific impairment. *J Neurol Neurosurg Psychiatry* 1977; **63**: 74–82.
- Murphy FC, Sahakian BJ, Rubinsztein JS, Michael A, Rogers RD, Robbins TW, et al. Emotional bias and inhibitory control processes in mania and depression. *Psychol Med* 1999; **29**: 1307–21.
- Chamberlain SR, Sahakian BJ. The neuropsychology of mood disorders. *Curr Psychiatry Rep* 2006; **8**: 458–63.
- National Development Team for Social Inclusion. *Reviewing Measures of Social Exclusion. National Social Inclusion Programme, 2003*.
- Brugha TS, Cragg D. The list of threatening experiences: the reliability and validity of a brief life events questionnaire. *Acta Psychiatr Scand* 1990; **82**: 77–81.
- Bernstein DP, Fink L. Initial reliability and validity of a new retrospective measure of child abuse and neglect. *Am J Psychiatry* 1994; **151**: 1132–6.
- Nolen-Hoeksema S, Wisco BE, Lyubomirsky S. Rethinking rumination. *Perspec Psychol Sci* 2008; **3**: 400–24.
- Kinderman P, Bentall RP. A new measure of causal locus: the internal, personal and situational attributions questionnaire. *Pers Individ Dif* 1996; **20**: 261–4.
- Goldberg D, Bridges K, Duncan P, Grayson D. Detecting anxiety and depression in general medical settings. *BMJ* 1989; **297**: 897–9.
- Kinderman P, Schwannauer M, Pontin E, Tai S. The development and validation of a general measure of well-being: the BBC well-being scale. *Qual Life Res* 2011; **20**: 1035–42.
- Pontin E, Schwannauer M, Tai S, Kinderman P. A UK validation of a general measure of subjective well-being: the modified BBC subjective well-being scale (BBC-SWB). *Health Qual Life Outcomes* 2013; **11**: 150–9.
- Goebel B, Dawy Z, Hagenauer J, Mueller JC. An approximation to the distribution of finite sample size mutual information estimates. *IEEE Intl Conf Comm South Korea* 2005; **2**: 1102–6.
- Weich S, Brugha T, King M, McManus S, Bebbington P, Jenkins R, et al. Mental well-being and mental illness: findings from the Adult Psychiatric Morbidity Survey for England 2007. *Br J Psychiatry* 2011; **199**: 23–8.

Author:
Could not
access through
URL provided;
is reference
acceptable as
now listed?
If not, please
provide new
URL
(AQ9)

Data supplement

Table DS1 Data-processing of each variable for conditional graphical analysis		
Composite variable from the Stress Test	Categories	No. of categories
Familial mental health history	Yes No	2
Relationships with friends	I have no friends I have few friends I have some close friends that I see occasionally I have few close friends that I see often I have many close friends that I see often	5
Relationships with family	I never see my family I have limited contact with members of my family I have reasonable contact with members of my family I am very close to members of my family	4
Social activities	I don't really do anything active or sociable I have only a limited social life I have a reasonably active social life I have an interesting and enjoyable social life	4
Frequency of social contacts	(None of the below) See parent weekly See sibling weekly See other relative weekly Attend an evening class Given up time for charity or local groups Involved in club/organisation/religious group Participated in sports/physical activity Go to the cinema	9
Severity of childhood trauma	Scores on 5-point Likert scale ranging from; 1 ('never') to 5 ('over many years') on: In the past I believe I was physically abused In the past I believe I was sexually abused In the past I believe I was emotionally abused In the past I believe I was bullied at school	6
Number of life events	Total number of life events (0–12) on the Brugha life events scale ¹³	13
Highest level of education	Did not complete GCSE/CSE/O-levels or equivalent Completed GCSE/CSE/O-levels or equivalent Completed post-16 vocational course A-levels or equivalent Undergraduate degree or professional qualification Postgraduate degree	6
Occupational status	Still at school At university In full-time employment In part-time employment Self-employed Unemployed Retired In full or part-time voluntary employment	8
Household income	Up to £9999 per annum £10 000 to £19 999 per annum £20 000 to £29 999 per annum £30 000 to £39 999 per annum £40 000 to £49 999 per annum £50 000 to £74 999 per annum £75 000 or more per annum Don't know Prefer not to say	9
Parents' income while growing up	Much lower than others (lowest 25% of population) Slightly lower than others (lowest 50% of population) Slightly higher than others (upper 50% of population) Much higher than others (top 25% of population)	4
Relationship status	In a relationship (but not married, in a civil partnership or living together) Living together (but not married or in a civil partnership) Married (first marriage) Civil partnership Divorced Divorced and remarried Separated (but legally still married) Widowed Widowed and remarried Single (never married)	10

(continued)

AUTHOR'S PROOF

Table DS1 Data-processing of each variable for conditional graphical analysis (<i>continued</i>)		
Composite variable from the Stress Test	Categories	No. of categories
Number of children	None	7
	One	
	Two	
	Three	
	Four	
	Five	
	Six or more	
Attributional style internal score	Scores on continuous variable	10
Rumination response style	Scores on continuous variable	10
Adaptive coping response style	Scores ranging from 4 to 16 entered as integers	13
Dangerous activities response style	Scores ranging from 3 to 12 entered as integers	10
Mental health problems: total Goldberg		
Anxiety and Depression scale score	Scores on continuous variable	10
Well-being	Scores on continuous variable	10