



LJMU Research Online

Fisher, N, Lattimore, P and Malinowski, P

Attention with a mindful attitude attenuates subjective appetitive reactions and food intake following food-cue exposure.

<http://researchonline.ljmu.ac.uk/id/eprint/2561/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Fisher, N, Lattimore, P and Malinowski, P (2015) Attention with a mindful attitude attenuates subjective appetitive reactions and food intake following food-cue exposure. *Appetite*, 99. pp. 10-16. ISSN 0195-6663

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.


For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

Accepted Manuscript

Attention with a mindful attitude attenuates subjective appetitive reactions and food intake following food-cue exposure

Naomi Fisher, PhD, Paul Lattimore, PhD, Peter Malinowski, PhD



PII: S0195-6663(15)30118-5

DOI: [10.1016/j.appet.2015.12.009](https://doi.org/10.1016/j.appet.2015.12.009)

Reference: APPET 2793

To appear in: *Appetite*

Received Date: 30 April 2015

Revised Date: 8 December 2015

Accepted Date: 11 December 2015

Please cite this article as: Fisher N., Lattimore P. & Malinowski P., Attention with a mindful attitude attenuates subjective appetitive reactions and food intake following food-cue exposure, *Appetite* (2016), doi: 10.1016/j.appet.2015.12.009.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Title: Attention with a mindful attitude attenuates subjective appetitive reactions and food intake following food-cue exposure

Article type: Full length Paper

Key words: Mindfulness; Hedonic reactions; Hunger; Food cue exposure; Food intake.

Abbreviations: MAI = Mindful attention induction; FCE = Food cue exposure

Highlights:

- Mindful attention can attenuate tendencies to eat in response to hedonic properties of food
- Effects of attention with and without a mindful attitude were compared
- Subjective reactions to the hedonic properties of energy-dense foods and food intake were examined
- Following attention with a mindful attitude fullness increased and hunger did not whereas without a mindful attitude hunger increased and fullness did not
- Significantly fewer cookies were eaten ten minutes post-exposure following the mindful attention induction.

Title page

Attention with a mindful attitude attenuates subjective appetitive reactions and food intake following food-cue exposure

Authors: Naomi Fisher, PhD^a; Paul Lattimore, PhD^a; Peter Malinowski, PhD^a.

^aSchool of Natural Sciences and Psychology, Liverpool John Moores University, Tom Reilly Building, Byrom Street, Liverpool, L3 3AF.

Corresponding author: Naomi Fisher, PhD

School of Natural Sciences and Psychology,
Liverpool John Moores University,
Tom Reilly Building, Byrom Street,
Liverpool,
L3 3AF.

Email: N.R.Fisher@ljmu.ac.uk
Telephone: 0151 904 6472

1 Abstract:

2 **Background:** Excessive energy intake that contributes to overweight and obesity is arguably
3 driven by pleasure associated with the rewarding properties of energy-dense palatable foods.

4 It is important to address influences of external food cues in food-abundant societies where
5 people make over 200 food related decisions each day. This study experimentally examines
6 protective effects of a mindful attention induction on appetitive measures, state craving and
7 food intake following exposure to energy-dense foods.

8 **Method:** Forty females were randomly allocated to a standard food-cue exposure condition
9 in which attention is brought to the hedonic properties of food or food-cue exposure
10 following a mindful attention induction. Appetitive reactions were measured pre, post and ten
11 minutes after post-cue exposure, after which a plate of cookies was used as a surreptitious
12 means of measuring food intake.

13 **Results:** Self-reported hunger remained unchanged and fullness significantly increased for
14 the mindful attention group post-cue exposure whereas hunger significantly increased for the
15 standard attention group and fullness remained unchanged. There was no significant between-
16 group difference in state craving post-cue exposure and ten minutes later. Significantly more
17 cookies were eaten by the standard attention group ten minutes post-cue exposure although
18 no significant between-group differences in appetitive and craving measures were reported at
19 that time.

20 **Conclusion:**

21 Our results point to a promising brief intervention strategy and highlights the importance of
22 distinguishing mindful attention from attention. Results also demonstrate that mindful
23 attention can influence food intake even when craving and hunger are experienced.

24 **Key words:** Mindfulness; Hedonic reactions; Hunger; Food cue exposure; Food intake.

25

26

Introduction

27 At present one in four adults can be described as obese and it is predicted that, if
28 current trends continue, nine in ten adults will be overweight or obese by 2050 (Department
29 of Health, 2013). The causes of obesity reflect complex interactions between genetic,
30 behavioural, environmental and psychosocial factors (Butland et al., 2007; Jebb, 1997). In
31 food-abundant environments where people make an estimated 200 food related decisions
32 each day research indicates that eating predominately occurs to prevent hunger (Lowe, Van
33 Steenburgh, Ochner, & Coletta, 2009; Wansink & Sobal, 2007). That is, eating happens
34 before significant energy depletion and associated physiologic signals that form part of the
35 homeostatic system are experienced (Lowe et al., 2009). It is recognised that much of this
36 excessive energy intake that contributes to overweight and obesity, is driven by pleasure or
37 the rewarding properties of readily available energy-dense palatable foods (Appelhans, 2009).
38 Food consumption, in the absence of physical signals or energy deficit, is driven by hedonic
39 hunger and reactions to hedonic properties of foods (e.g. sight, smell) rather than homeostatic
40 mechanisms (Lowe & Butryn, 2007). Hedonic hunger, the motivation to consume food for
41 pleasure, is often associated with increased susceptibility to environmental food cues
42 presenting a barrier to behaviour change and weight management (Lowe & Butryn, 2007;
43 O'Neil, Theim, Boeka, Johnson, & Miller-Kovach, 2012). In experimental settings this is
44 demonstrated by evidence that exposure to high-calorie food-cues increases appetitive
45 responses such as hunger and desire to eat cued and non-cued foods (Ferriday & Brunstrom,

46 2008, 2010; Jansen, Nederkoorn, Van Baak, Kierse, & Guerrieri, 2009). The food-cue
47 exposure paradigm, a reliable method for examining the effect of exposure to food, has also
48 been shown to effect subsequent food intake of similar or identical cued foods (e.g., Jansen et
49 al., 2009; Ferriday & Brunstrom, 2010).

50 Characteristics of the individual (e.g. emotional needs; Evers, Stok, & de Ridder,
51 2010), food or the food environment may exert influences that individuals may not wholly be
52 aware of (e.g. Herman & Polivy, 2005; Marchiori & Papies, 2014). Unrecognised somatic
53 and mental phenomena can trigger automatic reward-motivated behaviours, including eating
54 (Caldwell, Baime, & Wolever, 2012). The role of automatic habitual tendencies associated
55 with hedonic hunger are an obstacle to dietary educational approaches (Rothman, Sheeran, &
56 Wood, 2009). Alternative and complementary approaches are required to understand and
57 address automatic reward motivated behaviours associated with excessive food intake. In this
58 respect the concept of mindfulness has received considerable attention (Mantzios & Wilson,
59 2015). Mindfulness, as defined by Kabat-Zinn (2003) encompasses receptive attention to
60 whatever arises in the present moment with an open, curious non-judgmental attitude.
61 Compared to normal functioning a mindful state is one of enhanced receptive awareness and
62 attention to present reality (Brown & Ryan, 2003). Mindfulness techniques have been shown
63 to moderate eating behaviours influenced at a perceptual or preconscious level (Kahn &
64 Wansink, 2004; Wansink, 2010).

65 Mindfulness training interventions have been shown to increase discrimination
66 between externally cued hunger and hunger associated with emotions (Baer, Fischer, & Huss,
67 2006), and to attenuate hedonic hunger reducing automatic relations between cravings and
68 food intake (Alberts, Mulkens, Smeets, & Thewissen, 2010). Increasing awareness and
69 attention to internal cues and cued responses can serve a “de-automatisation” function (Bargh,
70 1997; Baumeister, Heatherton, & Tice, 1994; Lattimore & Maxwell, 2004), improve health
71 outcomes, enable weight regulation (Dalen et al., 2010), and facilitate successful self-
72 regulation (Papies, Barsalou, & Custers, 2012).

73 In experimental settings mindfulness techniques that increase attention with a mindful
74 attitude (e.g. non-reactive, non-judgemental) can influence both psychological and
75 behavioural outcomes (e.g. Arch & Craske, 2006; Erisman & Roemer, 2010; Verplanken &
76 Fisher, 2013). Specifically, the ability to mindfully observe thoughts and emotions has been
77 shown to reduce craving (Lacaille et al., 2014), chocolate consumption (Jenkins & Tapper,
78 2014) and approach responses to appetitive foods (Papies et al., 2012). Under everyday living
79 conditions the use of a brief mindfulness exercise (see Papies et al., 2012) changed
80 participants’ levels of hunger so it no longer influenced the attractiveness of unhealthy foods
81 and eating choices (Papies, Pronk, Keesman, & Barsalou, 2014). In addition to these
82 mindfulness inductions, the effects of mindfulness practices that are part of standard
83 mindfulness-based intervention programmes have been investigated. For instance, following
84 a guided ‘body scan meditation’ (14 minutes), one of the first exercises taught in

85 Mindfulness-Based Stress Reduction (MBSR; Kabat-Zinn, 1982), has been shown to make
86 the translation of hunger into unhealthy snacking behaviour less likely compared to listening
87 to an audiobook (Marchiori & Papies, 2014). However, the body-scan does not directly relate
88 to or address automatic eating nor does it encourage a mindful attitude to thoughts and
89 emotions around eating (Mantzios & Wilson, 2015).

90 Although research indicates that mindfulness techniques show promise in altering
91 habitual or automatic eating behaviour there is considerable variation in the content and
92 structure of techniques used (e.g. Alberts et al., 2010; Forman et al., 2007; Jenkins & Tapper,
93 2014; Moffitt, Brinkworth, Noakes, & Mohr, 2012). As a consequence caution is required
94 when interpreting these findings and attributing beneficial effects to mindfulness per se, or to
95 common practices in mindfulness-based interventions (Grossman & Van Dam, 2011). The
96 main aim of the current study was to test the influence of mindful attention on eating
97 behaviour. The “mindful attention induction” (MAI) was developed based on a systematic
98 review of existing inductions and incorporates key elements of mindful breath awareness
99 practice (Malinowski, 2013) that is a core technique of multicomponent MBSR programmes
100 (Brown, Ryan, & Cresswell, 2007; Kabat-Zinn, 1990). The objective of the mindful breath
101 awareness practice is to foster a state of present moment awareness involving a non-reactive
102 and non-judgemental attitude. The development of the MAI was motivated by a need to
103 qualify the use of mindfulness within research, clearly stating how it has been operationalised
104 or manipulated each context (Davidson, 2010). In doing so this study begins to address the
105 considerable variation in the use of mindfulness techniques in eating related research. By
106 combining our MAI approach with an established food-cue exposure methodology this study
107 examines how brief mindful attention practice may alter habitual or automatic reactivity to
108 food cues that typically leads to overeating.

109 In the current study participants were randomly allocated to either an attention
110 (control) or brief mindfulness attention induction (MAI). This was followed by a standard
111 food-cue exposure task (Jansen et al., 2009) and thus participants were either subjected to a
112 standard food-cue exposure (Standard-FCE) or to a food-cue exposure following a mindful
113 attention induction (Mindful-FCE). The Mindful-FCE fostered a decentred non-reactive
114 observational stance to phenomena, thus inviting attention with a mindful attitude. By
115 contrast, the Standard-FCE brought attention to food properties without prior guidance on the
116 processing of cues or the automatic quality of reactions to cues, thus representing attention
117 without a mindful attitude. Based on evidence suggesting that mindful attention can influence
118 both psychological and behavioural outcomes we expected that compared to the Standard-
119 FCE participants Mindful-FCE participants would experience lower increases in hunger,
120 feeling like eating, desire to eat and craving, but an increase in fullness immediately post
121 food-cue-exposure. These effects were expected to be short-lived therefore appetitive
122 measures were repeated 10 minute post-cue exposure, directly before measuring food intake.
123 It was predicted that Standard-FCE would result in greater food intake compared to Mindful-
124 FCE. Aspects of state mindfulness were measured to see if they would be influenced by the
125 MAI. Liking and desire to eat the cued foods, mood and awareness of the experimental
126 hypotheses were measured to examine alternative influences on measures of appetite and
127 food intake.

128

129

Methods

130 Design

131 A mixed factorial design was employed. Experience of Standard-FCE or Mindful-
132 FCE served as the between subjects factor and time of assessment the within subjects factor
133 (pre-exposure vs. post-exposure vs. end-of-delay). Outcomes were assessed using visual
134 analogue scales for hunger, fullness, feeling like eating and desire to eat, and self-report
135 Likert scales for craving. Food intake was measured as number of items consumed. After
136 participants had completed the pre-exposure assessments the experimenter (NF) opened an
137 envelope for each participant containing their group assignment. These envelopes had been
138 prepared using a random allocation algorithm by a third party blind to the nature of
139 conditions. Participants had an equal chance of assignment to either condition. (See Figure 1
140 for a visual presentation of the experimental design).

141 Participants

142 Females (18-50yrs) from a university research participants panel, and university staff,
143 were invited to take part in a “Food and Attention” study. Ethical approval was obtained from
144 the University’s Research Ethics Committee. A brief screening telephone interview ensured
145 participants met inclusion criteria: 1) regularly eating between meals and 2) liking crisps and
146 chocolate. Exclusion criteria were 1) Body Mass Index (BMI) < 18.5 or > 39.5; 2) currently
147 pregnant; 3) presence of food allergy; 4) diabetes diagnosis; 5) having sought medical help in
148 past six months for eating disorder and/or mental health problems; 6) current use of anti-
149 depressant and/or weight-loss medication; 7) any previous formal or informal meditation
150 experience (including yoga and self-help books or audio recordings); and 8) actively trying to
151 reduce weight (independently or on weight loss programme). Eighty-seven women expressed
152 an interest in taking part. Of the sixty-three eligible participants invited to take part forty-one

153 (M/SD: Age 30yrs, ± 7.7 ; BMI $25.4\text{kg/m}^2 \pm 0.7$) completed the online survey and the
154 subsequent experiment. Of the 24 participants who did not meet eligibility criteria six were
155 actively trying to lose weight, five had a BMI > 39.5 , five were on anti-depressant and/or
156 weight-loss medication, four had previous experience of mindfulness training, two were
157 unable to attend, one was pregnant and another had a diabetes diagnosis. All of the forty-one
158 participants reported liking and eating chocolate and crisps, and 72.5 % ate between meals
159 almost every day. Data from one participant were excluded from analyses as she indicated
160 that due to personal circumstances she had been unable to provide reliable responses, leaving
161 a total of 40 participants (20 in each group).

162 Measures

163 *Pre-exposure control measures.* Dispositional Mindfulness was assessed with the 39-
164 item Five-Facet Mindfulness Questionnaire (FFMQ; Baer, Smith, Hopkins, Krietemeyer, &
165 Toney, 2006) using a five-point Likert scale ranging from “*Never or very rarely true*” to
166 “*Very often or always true*”. The current study reports the total score as an overall measure of
167 dispositional mindfulness with higher scores indicating greater dispositional mindfulness
168 (Baer, Smith, et al., 2006). Internal consistency was satisfactory for the total score ($\alpha = 0.88$).
169 Eating attitudes relating to hedonic eating behaviour were assessed with the 18-item Three
170 Factor Eating Questionnaire (TFEQ-R18V2) with subscales measuring uncontrolled eating
171 (UE), emotional eating (EE), and cognitive restraint (CR; Cappelleri et al., 2009). The four-
172 point Likert scale ranged from “*Definitely true*” to “*Definitely false*” with responses
173 transformed to a 0-100 scale in line with common practice for the TFEQ. Higher scores
174 indicate greater uncontrolled and emotional eating and greater cognitive restraint. Internal
175 consistency was satisfactory for UE, EE and CR ($\alpha = 0.83, 0.93, 0.74$ respectively). The
176 FFMQ, TFEQ, age and BMI recorded by self-report and anthropomorphic measures were
177 included to ensure groups did not differ in these characteristics.

178 *Appetitive ratings & Food intake.* Four Visual Analogue Scales (VAS) were used to
179 assess hunger, fullness, feeling like eating and desire to eat pre-, post-exposure and ten
180 minutes after post-exposure (end-of –delay). Each VAS were preceded by the phrase “*Right*
181 *now, I feel...*” followed by a 0-100mm line used to indicate the responses: hungry (not at all
182 /very hungry), full (not at all /very full), feel like eating (not at all/very much), and desire to
183 eat food (absolutely no desire/very strong). Participants could not refer to previous appetitive
184 VAS ratings. Seven additional VAS, assessing aspects of sociability and self-pride, were
185 included to reduce the likelihood that participants would guess the experimental hypotheses.
186 VAS scales in appetitive research have shown good test-retest reliability and sensitivity to
187 subtle changes in appetite (Stubbs et al., 2000). Twelve Maryland chocolate chip cookies
188 were presented as a surreptitious ad libitum eating opportunity 10 minutes after post-cue
189 exposure. The number of cookies consumed served as a measure of food intake.

190 *State craving.* State craving was measured using the 15 item state Food Cravings
191 Questionnaire (Nijs, Franken, & Muris, 2007). The FCQ provides a total score and five
192 subscales: 1) An intense desire to eat; 2) Anticipation of relief from negative states and
193 feelings as a result of eating; 3) Craving as a physiological state; 4) Obsessive preoccupation
194 with food or lack of control over eating; 5) Anticipation of positive reinforcement that may
195 result from eating. A five-point Likert scale was used that ranged from “*Strongly Agree*” to
196 “*Strongly Disagree*”. Higher scores indicate greater state craving. The total and subscales had
197 good internal consistency with Cronbach coefficients ranging between $\alpha = 0.77$ and 0.97.

198 *Liking or desire for cued food and current mood.* VAS (0-100mm line) were also
199 administered to assess reactions to cued foods in terms of liking (not at all/really like this
200 food) and desire to eat the food (absolutely no desire/a very strong desire to eat this food).
201 Current mood was assessed using VAS in terms of happiness (not at all/very happy) and

202 relaxation (not at all/ very relaxed). These measures were included in order to rule out
203 alternative explanations for any between group differences.

204 *State mindfulness.* Aspects of state mindfulness were measured using five VAS items
205 adapted from validated mindfulness scales. Items (M1-M5) measured the extent to which
206 participants noticed internal and external phenomena: M1) “*I feel myself getting carried away*
207 *by my thoughts rather than just noticing them*”; M2) “*I pay attention to my thoughts and*
208 *feelings*”; M3) “*I am aware of my thoughts, feelings and bodily sensations*”; M4). “*Food*
209 *affects my thoughts and feelings*”; and M5) “*I notice how food affects my thoughts and*
210 *feelings*”. Participants responded using a 0-100mm line (never/all the time). Higher scores
211 indicate greater perceived levels of aspects of mindfulness.

212 *Mindful attention induction.* The MAI script was developed following systematic
213 analyses of the constituent components of published experimental mindfulness inductions and
214 review of current literature (the detailed analysis is in preparation for publication). The MAI
215 script included the identified constituent components: descriptions and practice of
216 mindfulness using breath as an object of focus; bringing attention with a quality of non-
217 reactive and non-judgemental to the observation of self; and used of rhetorical devices. As
218 such the MAI largely followed Kabat-Zinn’s (2002) sitting mindful breath awareness
219 meditation used in Verplanken and Fisher (2013). In brief, participants in the Mindful-FCE
220 read a description of mindfulness and then the experimenter read a guided breath awareness
221 meditation in which they were directed to notice arising thoughts, emotions and physical
222 sensations without reaction or judgement.

223 *Control attention condition.* In the control condition the presence of the experimenter
224 and effects of being given information in written and oral forms were matched as closely as
225 possible to the format of the MAI. The mindfulness scripts were substituted with a script

226 describing an exploration of the Venezuelan Rain forest as used in a previous study
227 (Lattimore & Mead, 2015). This Venezuelan Rain forest text was read in the same tone and
228 for the same duration as the mindful attention induction.

229 *Food-cue exposure task.* Four high-calorie foods (Cheese and Onion Pringles, Tesco's
230 Rocky Road Clusters, Green and Blacks organic Milk Chocolate, and Mini Twix's) were
231 used as exposure stimuli. The exposure activity was described and "modelled" by the
232 experimenter in a timed procedure based on Jansen et al. (2009). Two pieces of each food
233 item were presented in separate opaque sealed Tupperware. A stopwatch was used to time
234 exposure and a bowl of water and napkin provided to clean fingers between each food item.
235 Participants were instructed to hold and smell each item intensely, touch them against their
236 lips, rotate them between fingers and look intensively at each one. They were told not to eat
237 or taste the food. Participants took a sip of water between food cue exposures.

238 Procedure

239 The FFMQ and TFEQ were administered via Bristol Online Survey two weeks prior
240 to the experimental session. To control for readiness to eat participants were asked not to eat
241 or have any caffeinated drinks two hours prior to the experimental session. Participants were
242 tested individually in the laboratory. An overview of participant flow through the procedure
243 and assessments at different stages is provided in Figure 1. On arrival, participants were
244 informed they would be taking part in a "food task" that was being piloted for a different
245 study. After giving informed consent they completed appetite, state mindfulness and mood
246 VAS before random allocation to either the Standard-FCE or the Mindful-FCE group. The
247 Mindful-FCE group completed the MAI, whereas the Standard-FCE listened to information
248 read in the same tone and duration as the MAI. Participants then completed the food-cue
249 exposure task lasting 10 minutes, followed by post-exposure assessment of appetite, state

250 mindfulness and mood VAS, and of food craving (FCQ). Following post-exposure
251 assessments participants in the Mindful-FCE condition were instructed to practice the
252 mindful breathing meditation taught during the MAI as they were waiting for the next part of
253 the experiment, whereas participants in the Standard-FCE were simply told to reflect on their
254 experience up to that point. During this 10-minute delay period all participants remained in
255 the presence of the cue exposure foods left on a table in their product packaging (unopened).
256 Subsequently, the experimenter returned to inform the participants that the study was almost
257 over and took them to another room to complete end-of-delay appetite, state mindfulness and
258 mood VAS, and measure of food craving (FCQ). Additionally participants completed VAS
259 ratings of their liking and desire to eat cued foods. To maintain the cover story of piloting a
260 food tasting task participants were prompted to provide feedback about the cue-exposure task
261 as an open-ended question. Participants were then given a plate of 12 cookies and a glass of
262 water from which they could have as much as they wanted as ‘a token of appreciation and as
263 they had not eaten for two hours and may have to return to work or drive somewhere’.

264 Participants were left unobserved with the cookies for five minutes. Finally, weight and
265 height were measured in a separate room. Participants, having been told the study was
266 completed were asked to suggest what they thought the experiment had been examining.
267 None had disputed the cover story. Suggestions about what the study was measuring
268 included: attention/ concentration/ distraction, the attractiveness of sensory properties of
269 food, and piloting of the food cue procedure. Importantly, no participant suggested that it was
270 about food intake.

271 Data analysis strategy

272 Assumptions required for parametric testing were examined prior to any inferential
273 analysis. Parametric test assumptions were met for all analyses with the exception of food
274 intake. Box plots and normality tests indicated that the distributions for the number of

275 cookies eaten were non-normal with multiple extreme scores in the Standard-FCE condition.
276 A Mann-Whitney test was used, due to the normality violation, to test differences in food
277 intake between the Mindful-FCE and Standard-FCE groups. Analysis of variance (ANOVA)
278 was used to test hypotheses. Where appropriate, Bonferroni corrected t-tests were used to
279 probe significant main effects and interactions. Summary statistics are presented as means
280 (M) and standard deviation of means (SD) unless otherwise stated.

281

282

Results:

283 A series of one way ANOVAs were carried out to examine whether the two groups
284 differed in terms of BMI, age, dispositional mindfulness, uncontrolled eating, emotional
285 eating, cognitive restraint, and time since last eating. No significant differences were found
286 (see Table 1). Two multivariate ANOVAs revealed no significant between-group differences
287 (all $p > .05$) on 1) pre-exposure appetite (hunger, fullness, feeling like eating and desire to eat
288 food) and 2) current mood ratings (happy, relaxed).

289 *Appetitive ratings*

290 To test the hypotheses concerning the overall effects of Mindful-FCE vs Standard-
291 FCE on appetite VAS (hunger, fullness, feeling like eating and desire to eat food) separate 2
292 (Group: Mindful-FCE vs. Standard-FCE) x 3 (Time: pre vs. post vs. end-of-delay) mixed
293 factorial ANOVAs were conducted (See Table 2). There were no significant main effects of
294 Group on any of the appetite ratings. A significant main effect of Time for hunger showed an
295 increase from pre-cue exposure to post-cue exposure to end-of-delay. Planned comparisons
296 indicated significant differences in hunger pre- to end-of-delay, and post-cue to end-of-delay

297 ($ps < .05$). There were no significant main effects of Time for the remaining appetitive
298 measures.

299 The Time-by-Group interaction for hunger approached significance ($p = .076$). Based
300 on the hypothesis that compared to the Standard-FCE participants Mindful-FCE participants
301 would experience lower increases in hunger, the Time-by-Group interaction for hunger was
302 investigated. Bonferroni adjusted pairwise comparisons revealed significant increases in
303 hunger pre to post-cue exposure ($p = .05$) and post-cue to end-of delay ($p < .01$) for the
304 Standard-FCE group but no significant increases pre to post-cue ($p = 1.0$) and post-cue to end-
305 of delay ($p = .06$) for the Mindful-FCE group. A significant Time-by-Group interaction was
306 found for fullness. Bonferroni adjusted pairwise comparisons revealed significant increases in
307 fullness pre to post-cue exposure ($p = .03$) and a significant decrease post-cue to end-of delay
308 ($p = .03$) for the Mindful-FCE group but no significant changes between pre, post or end of
309 delay for the Standard-FCE group ($ps > .05$). There were no other significant interaction
310 effects.

311 *State craving*

312 To examine the effects of the Mindful-FCE vs Standard-FCE on total scores and
313 subscales of state craving separate 2 (Group: Mindful-FCE vs. Standard-FCE) x 2 (Time: pre
314 vs. post) mixed factorial ANOVAs were conducted. There were no significant main effects or
315 interactions for total scores or subscales of the state FCQ (see Table 3).

316 *Food intake*

317 A Mann-Whitney test on the number of cookies eaten confirmed the hypothesis that
318 the MAI would affect food intake. Significantly fewer cookies were eaten by the Mindful-

319 FCE group (Range between 0-3; mean = 0.7; *Mdn* = 0) than by the Standard-FCE group
320 (Range between 0-7, *M* = 2.2; *Mdn* = 2), $U = 69.0$, $z = -3.7$, $p < .001$, $r = -.58$.

321

322 *State mindfulness*

323 To examine how Mindful-FCE vs Standard-FCE influenced aspects of state
324 mindfulness throughout the experimental session separate 2 (Group: Mindful-FCE vs.
325 Standard-FCE) by 3 (Time: pre vs. post vs. end-of-delay) mixed factorial ANOVAs were
326 conducted. There were no main effects or interactions for responses to the mindfulness items
327 M1, M2 and M3. There was a main effect of Time for item M4: “*Food affects my thoughts*
328 *and feelings*” ($F(2,76) = 3.53$, $p < .05$, $\eta_p^2 = .16$). Bonferroni contrasts ($p = .016$) indicated
329 that food affected participants thoughts and feelings significantly more end-of-delay ($M =$
330 69.9 ± 20.7) compared to pre-cue ($M = 58.6 \pm 21.5$). There was no significant difference
331 post-cue ($M = 69.9 \pm 24.3$) vs end-of-delay ($p < .016$). There was also a main effect of Time
332 for item M5: “*I notice how food affects my thoughts and feelings*”, ($F(2,76) = 5.55$, $p < .01$,
333 $\eta_p^2 = .12$). Bonferroni contrasts ($p = .016$) revealed significant differences between pre-cue (M
334 $= 56.9 \pm 25.95$) and end-of-delay ($M = 70.4 \pm 23.12$; $p < .01$) but no significant differences
335 for the other contrasts ($ps > .016$). There were no significant interaction effects for items M4
336 and M5.

337 *Ruling out alternative explanations*

338 To rule out that the observed differences in food intake were merely based on liking
339 or the desire to eat at the moment when the cookies were offered, two separate multivariate
340 between subjects ANOVAs (Mindful-FCE vs. Standard-FCE) were conducted on liking and
341 desire to eat VAS for the four cued foods at end-of-delay. There were no significant
342 multivariate effects for liking (Pillai’s Trace = .71, $F = .65$, $df = (4, 34)$, $p > .05$) or desire to
343 eat cued foods (Pillai’s Trace = .08, $F = .77$, $df = (4, 34)$, $p > .05$). To examine whether

344 changes in mood might contribute to any of the observed mindfulness-specific effects
345 separate 2 (Group: Mindful-FCE vs. Standard-FCE) x 3 (Time: pre vs. post vs. end-of-delay)
346 mixed factorial ANOVAs were conducted for “happy” and “relaxed”. There was a significant
347 main effect of Time on relaxed ratings ($F(2,76)= 6.75, p<.01, \eta_p^2 =.15$). Bonferroni contrasts
348 (adjusted alpha level: $p=.016$) indicated that participants were significantly more relaxed
349 post-cue ($M = 75.23, SD = 17.08$) compared to pre-cue ($M = 65.30, SD = 14.16$), and end-of-
350 delay ($M = 74.18, SD = 18.69$) compared to pre-cue. There was no significant difference
351 post-cue vs end-of-delay ($ps >.016$). There were no significant main effects of Time or Group
352 on happy ratings and, importantly, no time-by-condition interactions for both mood ratings.

353

354

Discussion

355 The present study examined the effects of a brief mindful attention induction on
356 appetitive reactions immediately following exposure to energy-dense food cues, ten
357 minutes post exposure and subsequent food intake. The outcomes partially support our
358 hypotheses. Firstly, the hypothesis that the mindful attention induction would attenuate
359 appetitive reactions to cued foods was confirmed for hunger and fullness. There was no
360 change in hunger and an increase in fullness pre to post exposure following attention with
361 a mindful attitude. In contrast fullness remained the same and hunger increased in the
362 standard attention group. However, desire to eat and feeling like eating were unaffected.
363 Regarding the longevity of effects, the differences in hunger and fullness between groups
364 were not seen ten minutes post exposure. Contrary to expectations there were no between
365 group differences in state craving post-cue exposure or after delay. The hypothesis
366 regarding food intake was wholly supported as the Standard-FCE resulted in significantly
367 more intake compared to the Mindful-FCE. Potential alternative explanations for the
368 observed pattern of differences in appetite measures and intake, such as differences in

369 liking or desire to eat cued food or in mood, were ruled out as no between-group
370 differences were found.

371 The effects on subjective hunger and fullness in this sample were short-lived.
372 Participants in both conditions reported comparable levels of hunger, fullness, feeling like
373 eating and craving after the delay period, when given an eating opportunity. Although the two
374 groups did not differ on any of these measures participants in the Standard-FCE group, who
375 brought attention without a mindful attitude to the qualities of foods, ate significantly more
376 than those in the Mindful-FCE group. The current findings support the assertion that mindful
377 attention can disrupt relations between internal experiences and observable behaviours (e.g.
378 Bargh, 1994; Verplanken & Fisher, 2013). These findings differ from Marchiori and Papies
379 (2014) in which a mindfulness exercise (body scan) was shown not to reduce portion size
380 effects but did reduce effects of hunger on unhealthy food choice. However the current study
381 did not compare preferences for healthy/unhealthy food or small/large portions.

382 The current pattern of results are consistent with mindfulness-based intervention
383 studies evidencing a modulation of the translation of motivational states into eating behaviour
384 (Alberts et al., 2010; Papies et al., 2014) and reduced external eating (Daubenmier et al.,
385 2011). As such this controlled experimental examination of the effect of mindful attention on
386 reactions to exposure to foods may offer insights into underlying mechanisms of effects
387 suggested in previous research (e.g. Daubenmier et al., 2011; Hooper, Sandoz, Ashton,
388 Clarke, & McHugh, 2012). In comparing mindful attention with attention to the properties of
389 cued foods the between-group differences in food intake and post-exposure hunger and
390 fullness can reasonably be attributed to distinctive qualities of mindful attention. However,
391 inferences and suggestions made about the differences in hunger are offered with caution as
392 the interaction did not meet conventional significance levels. One suggestion is that the
393 elicited non-reactive, non-judgemental attitude allowed participants to bring attention to the

394 sensory properties of cued foods and their reactions reducing the likelihood of reacting
395 automatically. A further explanation for the lack of translation from subjective experience to
396 behaviour following mindful attention is that mindful attention practices influence how
397 thoughts are processed rather than changing the content (Brown et al., 2007). This emphasis
398 on the *how* thoughts are perceived rather than changing the content of thoughts is consistent
399 with findings that the number of cravings experienced is less relevant to the control of eating
400 behaviour than how the cravings are perceived (Hooper et al., 2012). This raises questions as
401 to the importance or value of focusing on the content or number of food related cravings or
402 thoughts when attempting to address the relations between experienced thoughts and
403 behavioural outcomes. There was no difference between the groups in terms of state craving
404 post-cue or prior to the measure of food intake. However craving measures have been shown
405 to be influenced by a mindful attention exercise (noticing, accepting non-judgementally) used
406 in the presence of smoking cues (Westbrook et al., 2013). The current lack of difference may
407 reflect unmeasured differences in state craving pre food-cue exposure rather than the MAI
408 not having an effect on state craving. State craving was not measured pre-cue exposure to
409 avoid participants becoming aware of the experimental hypotheses.

410 An important conclusion from this study is that a brief mindful attention induction can
411 lead to demonstrable beneficial effects without involving traditional meditation practices,
412 suggesting that such an approach might increase accessibility for people not able, willing or
413 ready to engage more formally with meditation (Mantzios & Wilson, 2015). Further research
414 is required to ascertain the acceptability and practicality of applying mindful attention in this
415 way and if it has the longer term efficacy required to manage weight including the absence of
416 rebound effects (Hooper et al., 2012).

417 Confidence in attributing effects to the mindful attention induction is increased by the
418 randomised and controlled design and lack of between group differences on measures of

419 dispositional mindfulness, trait eating patterns (uncontrolled or emotional eating or cognitive
420 restraint) and feelings of relaxedness. Additionally, advertising the study as an examination
421 of food and attention, reduced bias associated with recruiting participants willing to
422 participate in mindfulness meditation experiments. The intention was to minimise the
423 potential for enthusiasm for such practices to create a placebo effect (discussed further in
424 Mantzios & Wilson, 2015). The lack of between-group differences in measured aspects of
425 mindfulness using single item measures may in part reflect a particularly pertinent limitation
426 of self-report measures when considering the accuracy of mindfulness measures: the ability to
427 accurately measure 'mindfulness' is reliant on participants' 'mindfulness' (Grossman, 2011).

428 Limitations of the study include the laboratory setting, the sample size which limits
429 the statistical power of the analyses, and the representativeness of the sample which limits
430 generalisability. For these reasons, effect sizes are provided to give further information about
431 which findings may be important to pursue in future studies in and out of experimental
432 settings. Further studies are required to ascertain if more enduring effects are only attainable
433 through regular meditation training and if the effects demonstrated in studies such as these
434 differ from that of long term practice. The fact that short-lived effects can be obtained,
435 nevertheless suggests that mindfulness is a powerful and interesting state of consciousness
436 worth further exploration (Verplanken & Fisher, 2013). The current findings indicate that
437 attention with a mindful attitude may promote better eating behaviours in the short-term, and
438 adds to the evidence base justifying the examination of components of mindfulness-based
439 interventions within the context of obesity prevention and management.

440

441

442

443 References

- 444 Alberts, H. J. E. M., Mulkens, S., Smeets, M., & Thewissen, R. (2010). Coping with food
445 cravings. Investigating the potential of a mindfulness-based intervention. *Appetite*, *55*,
446 160-163.
- 447 Appelhans, B. M. (2009). Neurobehavioral Inhibition of Reward driven Feeding:
448 Implications for Dieting and Obesity. *Obesity Research*, *17*(4), 640-647.
- 449 Arch, J. J., & Craske, M. G. (2006). Mechanisms of Mindfulness: Emotion Regulation
450 Following a Focused Breathing Induction *Behaviour research and therapy*, *44*, 1849-
451 1858.
- 452 Baer, R. A., Fischer, S., & Huss, D. B. (2006). Mindfulness and acceptance in the treatment
453 of disordered eating. *Journal of Rational-Emotive & Cognitive - Behaviour Therapy*,
454 *23*(4), 281-300.
- 455 Baer, R. A., Smith, G. T., Hopkins, J., Krietemeyer, J., & Toney, L. (2006). Using self-report
456 assessment methods to explore facets of mindfulness. *Assessment*, *13*, 27-45.
- 457 Bargh, J. A. (1994). *The Four Horsemen of Automaticity: Awareness, Intention, Efficiency,*
458 *and Control in Social Cognition* (Vol. 1). Hillsdale, NJ: Erlbaum.
- 459 Bargh, J. A. (1997). *Automaticity in social psychology*. New York: Guilford.
- 460 Baumeister, R. F., Heatherton, T. F., & Tice, D. M. (1994). *Losing control: How and why*
461 *people fail at self-regulation*. San Diego Academic.
- 462 Brown, K. W., & Ryan, R. M. (2003). The Benefits of Being Present: Mindfulness and it's
463 Role in Psychological Well-being. *Journal of Personality and Social Psychology*, *84*,
464 822-848
- 465 Brown, K. W., Ryan, R. M., & Cresswell, J. D. (2007). Mindfulness: Theoretical Foundations
466 and Evidence for it's Salutory effects *Psychological Inquiry*, *18*(4), 211-237.


- 467 Butland, B., Jebb, S., Kopelman, P., McPherson, K., Thomas, S., Mardell, J., & Parry, V.
468 (2007). *Foresight. Tackling obesity: future choices*. Retrieved from UK:
- 469 Caldwell, K. L., Baime, M. J., & Wolever, R. Q. (2012). Mindfulness based approaches to
470 obesity and weight loss maintenance. *Journal of Mental Health Counseling, 34*(3),
471 269-282.
- 472 Cappelleri, J. C., Bushmakin, A. G., Gerber, R. A., Leidy, N. K., Sexton, C. C., Lowe, M. R.,
473 & Karlsson, J. (2009). Psychometric analysis of the Three-Factor Eating
474 Questionnaire-R21: results from a large diverse samples of obese and non-obese
475 participants. *International Journal of Obesity, 33*, 611-620.
- 476 Dalen, J., Smith, B. W., Shelley, B. M., Sloan, A. L., Leahigh, L., & Begay, D. (2010). Pilot
477 study: Mindful eating and living (MEAL): Weight, eating behavior, and psychological
478 outcomes associated with a mindfulness-based intervention for people with obesity.
479 *Complementary Therapies in Medicine, 18*(6), 260-264.
- 480 Daubenmier, J., Kristeller, J. L., Hecht, F. M., Maninger, N., Kuwata, M., Jhaveri, K., . . .
481 Epel, E. (2011). Mindfulness intervention for stress eating to reduce cortisol and
482 abdominal fat among overweight and obese women: an exploratory randomized
483 controlled study. *Journal of obesity, 1-13*.
- 484 Davidson, R. J. (2010). Empirical Explorations of Mindfulness: Conceptual and
485 Methodological Conundrums. *Emotion, 10*(1), 8-11.
- 486 Department of Health. (2013). Obesity information. Retrieved from
487 [<http://www.dh.gov.uk/en/Publichealth/Healthimprovement/Obesity/index.htm>]
- 488 Erisman, S. M., & Roemer, L. (2010). A Preliminary Investigation of the Effects of
489 Experimentally Induced Mindfulness on Emotional Responding to Film Clips.
490 *Emotion, 10*(1), 72-82.

- 491 Evers, C. F., Stok, F., & de Ridder, D. T. D. (2010). Feeding your feelings: Emotion
492 regulation strategies and emotional eating. *Personality and Social Psychology Bulletin*,
493 36, 792-804.
- 494 Ferriday, D., & Brunstrom, J. M. (2008). How does food-cue exposure lead to larger meal
495 sizes? *British Journal of Nutrition*, 100(6), 1325-1332.
- 496 Ferriday, D., & Brunstrom, J. M. (2010). 'I just can't help myself': effects of food-cue
497 exposure in overweight and lean individuals. *International Journal of Obesity*, 35(1),
498 142-149.
- 499 Forman, E. M., Hoffman, K. L., McGrath, K. B., Herbert, J. D., Brandsma, L. L., & Lowe, M.
500 R. (2007). A comparison of acceptance-and control-based strategies for coping with
501 food cravings: An analog study. *Behaviour research and therapy*, 45(10), 2372-2386.
- 502 Grossman, P. (2011). Defining mindfulness by how poorly I think I pay attention during
503 everyday awareness and other intractable problems for psychology's (re) invention of
504 mindfulness: comment on Brown et al.(2011). *Psychological assessment*, 23(4), 1034-
505 1040. doi:DOI: 10.1037/a0022713
- 506 Grossman, P., & Van Dam, N. T. (2011). Mindfulness, by another name...: Trials and
507 tribulations of Sati in Western psychology and Science. *Contemporary Buddhism*, 12,
508 219-239.
- 509 Herman, C. P., & Polivy, J. (2005). Normative influences on food intake. *Physiology &*
510 *behavior*, 86(5), 762-772.
- 511 Hooper, N., Sandoz, E. K., Ashton, J., Clarke, A., & McHugh, L. (2012). Comparing thought
512 suppression and acceptance as coping techniques for food cravings. *Eating Behaviors*,
513 13(1), 62-64.

- 514 Jansen, A., Nederkoorn, C., Van Baak, L., Kierse, C., & Guerrieri, R. (2009). High-restrained
515 eaters only overeat when they are also impulsive. *Behaviour research and therapy*, *47*,
516 105-110.
- 517 Jebb, S. (1997). Aetiology of obesity. *British Medical Bulletin*, *53*(2), 264-285.
- 518 Jenkins, K. T., & Tapper, K. (2014). Resisting chocolate temptation using a brief mindfulness
519 strategy. *British journal of health psychology*.
- 520 Kabat-Zinn, J. (1982). An outpatient program in behavioral medicine for chronic pain
521 patients based on the practice of mindfulness meditation: Theoretical considerations
522 and preliminary results. *General Hospital Psychiatry*, *4*, 33-47.
- 523 Kabat-Zinn, J. (1990). *Full catastrophe living: using the wisdom of your body and mind to*
524 *face stress, pain and illness*. New York: Delacorte.
- 525 Kabat-Zinn, J. (2002). 20 Minute Sitting Meditation. *On Guided Mindfulness Meditation:*
526 *Series Two*. USA: UMass Center for Mindfulness.
- 527 Kabat-Zinn, J. (2003). Mindfulness-based interventions in context: Past, present, and future.
528 *Clinical Psychology-Science and Practice*, *10*(2), 144-156. Retrieved from <Go to
529 ISI>://000182986300002
- 530 Kahn, B. E., & Wansink, B. (2004). The influence of assortment structure on perceived
531 Variety and consumption quantities. *Journal of Consumer Research*, *30*, 519-533.
- 532 Lacaille, J., Ly, J., Zacchia, N., Bourkas, S., Glaser, E., & Knäuper, B. (2014). The effects of
533 three mindfulness skills on chocolate cravings. *Appetite*, *76*, 101-112.
- 534 Lattimore, P., & Maxwell, L. (2004). Cognitive load, stress and disinhibited eating. *Eating*
535 *Behaviors*, *5*, 315-324.
- 536 Lowe, M. R., & Butryn, M. L. (2007). Hedonic hunger: A new dimension of appetite?
537 *Physiology and Behavior*, *91*, 432-439.

- 538 Lowe, M. R., Van Steenburgh, J., Ochner, C., & Coletta, M. (2009). Neural correlates of
539 individual differences related to appetite. *Physiology and Behavior*, *91*, 432-439.
- 540 Malinowski, P. (2013). Neural mechanisms of attentional control in mindfulness meditation.
541 *Frontiers in Neuroscience*, *7*(8), 1-11.
- 542 Mantzios, M., & Wilson, J. C. (2015). Mindfulness, Eating Behaviours, and Obesity: A
543 Review and Reflection on Current Findings. *Current Obesity Reports*(1-6).
- 544 Marchiori, D., & Papies, E. K. (2014). A brief mindfulness intervention reduces unhealthy
545 eating when hungry, but not the portion size effect. *Appetite*, *75*, 40-45.
- 546 Moffitt, R., Brinkworth, G., Noakes, M., & Mohr, P. (2012). A comparison of cognitive
547 restructuring and cognitive defusion as strategies for resisting a craved food.
548 *Psychology & Health*, *27*(sup2), 74-90.
- 549 Nijs, I. M. T., Franken, I. H. A., & Muris, P. (2007). The modified Trait and State Food-
550 Cravings Questionnaires: Development and validation of a general index of food
551 craving. *Appetite*, *49*(1), 38-46.
- 552 O'Neil, P. M., Theim, K. R., Boeka, A., Johnson, G., & Miller-Kovach, K. (2012). Changes
553 in weight control behaviors and hedonic hunger during a 12-week commercial weight
554 loss program. *Eating Behaviors*. doi:10.1016/j.eatbeh.2012.06.002
- 555 Papies, E. K., Barsalou, L. W., & Custers, R. (2012). Mindful attention prevents mindless
556 impulses. *Social Psychological and Personality Science*, *3* (3), 291-299.
- 557 Papies, E. K., Pronk, T. M., Keesman, M., & Barsalou, L. W. (2014). The Benefits of Simply
558 Observing: Mindful Attention Modulates the Link Between Motivation and Behavior.
559 *Journal of Personality and Social Psychology*. doi:10.1037/a0038032
- 560 Rothman, A. J., Sheeran, P., & Wood, W. (2009). Reflexive and automatic processes in the
561 initiation and maintenance of dietary change. *Annals Behav Med*, *38*(Suppl 1), S4-S17.

- 562 Stubbs, R. J., Hughes, D. A., Johnstone, A. M., Rowley, E., Reid, C., Elia, M., & Blundell, J.
563 E. (2000). The use of visual analogue scales to assess motivation to eat in human
564 subjects: a review of their reliability and validity with an evaluation of new hand-held
565 computerized systems for temporal tracking of appetite ratings. *British Journal of*
566 *Nutrition*, 84(04), 405-415.
- 567 Verplanken, B., & Fisher, N. (2013). Habitual Worrying and Benefits of Mindfulness.
568 *Mindfulness*. doi:10.1007/s12671-013-0211-0
- 569 Wansink, B. (2010). From mindless eating to mindlessly eating better. *Physiology and*
570 *Behaviour*, 100, 454-463.
- 571 Wansink, B., & Sobal, J. (2007). Mindless Eating: The 200 Daily Food Decisions We
572 Overlook. *Environment and Behavior*, 39(1), 106-123.
- 573 Westbrook, C., Creswell, J. D., Abibnia, G., Julson, E., Kober, H., & Tindle, H. A. (2013).
574 Mindful attention reduces neural and self-reported cue-induced craving in smokers.
575 *Social cognitive and affective neuroscience*, 8(1), 73-84.
- 576



577

578

579 Figure1. Participant flow and assessment points from pre-exposure to post-exposure to end-of-delay and food intake task.

580 Table 1. Baseline measures of individual differences and ANOVA summary values.

	<i>Range</i>	Mindful attention (N = 20)		Attention Control (N = 20)		<i>F</i> (1,38)
		<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	
Age	(21- 46)	30.65	9.15	29.50	6.12	.22
BMI (kg/m ²)	(20-39)	25.40	3.72	25.40	4.84	.00
TFEQ-UE	(11-78)	49.07	19.03	43.33	16.25	.37
TFEQ-EE	(0-72)	49.44	32.14	44.16	24.74	2.18
TFEQ-CR	(0-100)	38.05	16.55	46.11	17.94	1.05
FFMQ Total	(95-172)	128.00	17.73	125.55	17.25	.34
Last Ate	(1-15)	3.96	3.29	3.44	3.36	.50

581 Note: TFEQ UE = uncontrolled eating; TFEQ EE = emotional eating; TFEQ CR=
582 cognitive restraint; FFMQ Total= mindfulness; last ate = hours and minutes since last
583 ate.

584 Table 2. Appetite ratings (*M/SD*) and ANOVA summary values pre-vs post-cue exposure vs end of delay.

Variable	Mindful attention (N=20)						Attention control (N=20)						F(2,76)		
	Pre		Post		End-of - Delay		Pre		Post		End-of - Delay		T	C	T x C
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>			
Hungry	54.8	30.1	47.0	26.6	63.1	29.5	44.9	25.2	57.9	22.6	64.4	23.5	5.39*	.01	2.67
													[.28]	[.00]	[.12]
Full	28.7	21.9	48.0	25.1	33.6	24.8	36.4	22.5	35.4	20.1	34.1	21.8	2.91	.08	3.16*
													[.07]	[.00]	[.08]
Feel like eating	70.5	17.8	63.0	24.3	67.5	27.6	57.3	25.0	63.8	20.2	67.1	23.0	.68	.52	2.29
													[.05]	[.01]	[.08]
Desire to eat food	64.7	23.9	64.6	23.5	68.1	25.2	57.2	25.3	66.9	19.3	68.5	24.3	1.99	.07	.96
													[.08]	[.00]	[.04]

585 Note: * $p < .05$. T = Time main effect; G= Group main effect; T x G= Time-by-Group interaction. Partial Eta squared effect sizes in [].

586 Table 3. General state craving total and subscales post-cue exposure (T2) and end-of -

Variable	Mindful attention (N=20)				Attention control (N=20)				F(1,38)		
	Post		End-of - delay		Post		End-of- delay				
	M	SD	M	SD	M	SD	M	SD	T	G	T x G
GSC-total	41.7	13.0	42.4	16.4	42.6	9.8	41.2	11.4	.15 [.00]	.00 [.00]	1.61 [.04]
GSC-IDE	7.3	3.8	7.3	4.1	6.8	2.4	6.7	2.9	0.2 [.00]	.30 [.01]	0.00 [.00]
GSC-ARFN	9.0	2.7	9.0	3.6	8.8	2.6	8.7	2.6	.01 [.00]	.10 [.00]	.01 [.00]
GSC-CPS	8.0	2.6	7.9	3.3	8.5	2.8	8.0	2.9	2.18 [.05]	.09 [.00]	6.33 [.02]
GSC-OPF	9.4	3.4	9.6	3.7	10.5	2.5	10.0	3.3	.37 [.01]	.57 [.02]	2.62 [.07]
GSC-APR	8.0	3.0	8.7	3.4	8.2	2.3	7.9	2.4	.84 [.02]	.15 [.00]	3.73 [.10]

587 delay (T3).

588

589 Note: IDE= Intense desire to eat; ARFN= Anticipation of relief from negative states and feelings;

590 CPS = Craving as a physiological state; OPF= Obsessive preoccupation with food or lack of

591 control over eating; APR= Anticipation of positive reinforcement that may result from eating.

592 Range = the minimum and maximum scores for each subscale and total.

593