



LJMU Research Online

Moreira, A, Bradley, PS, Carling, C, Arruda, AF, Spigolon, L, Franciscon, C and Aoki, M

Effect of a Congested Match Schedule on Immune-Endocrine responses, Technical Performance, and Session-RPE in Elite Youth Soccer Players

<http://researchonline.ljmu.ac.uk/id/eprint/3758/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Moreira, A, Bradley, PS, Carling, C, Arruda, AF, Spigolon, L, Franciscon, C and Aoki, M (2016) Effect of a Congested Match Schedule on Immune-Endocrine responses, Technical Performance, and Session-RPE in Elite Youth Soccer Players. Journal of Sports Sciences. 34 (24). pp. 2255-2261.

LJMU has developed [LJMU Research Online](#) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

**Effect of a Congested Match Schedule on Immune-Endocrine responses,
Technical Performance, and Session-RPE in Elite Youth Soccer Players**

Running head: Effect of Match Congestion in Youth Soccer

Alexandre Moreira¹, Paul S. Bradley², Chris Carling³, Ademir F.S. Arruda¹,
Leandro M.P. Spigolon⁴, Clovis Franciscon⁴, Marcelo S. Aoki⁵,

¹Department of Sport, School of Physical Education and Sport, University of Sao Paulo, Sao Paulo, Brazil.

²RISES, Liverpool John Moores University, UK.

³ Institute of Coaching and Performance, University of Central Lancashire, Preston, UK.

⁴Desportivo Brazil, Porto Feliz, Brazil.

⁵ School of Arts, Sciences and Humanities, University of Sao Paulo, Sao Paulo, Brazil

Corresponding author

Dr Alexandre Moreira

School of Physical Education and Sport, University of Sao Paulo, Sao Paulo, Brazil.

Av. Prof. Mello Moraes, 65, Cidade Universitária, São Paulo, Brazil. Zip Code 05508-030. E-mail: alemoreira@usp.br

Telephone: +55-11-30918789

Fax: +55-11-38135091

**Effect of a Congested Match Schedule on Immune-Endocrine responses,
Technical Performance, and Session-RPE in Elite Youth Soccer Players**

Running head: Effect of Match Congestion in Youth Soccer Players

Key-words: football; testosterone; cortisol; saliva; Immunoglobulin A.

ABSTRACT

This study investigated the effects of a congested match schedule (7 matches played in 7 days) on steroid hormone concentrations, mucosal immunity, session rating of perceived exertion (S-RPE) and technical performance in 16 elite youth soccer players (14.8 ± 0.4 yr; 170.6 ± 9.4 cm; 64.9 ± 7 kg). No change was observed for salivary cortisol concentration across multiple matches ($p=0.33$; Effect size [ES]=0.13-0.48). In contrast, there was a decrease in salivary testosterone and IgA concentrations (SIgA) from the 1st compared to the last time-point ($p=0.01$ and 0.001 , ES=0.42 and 0.67, respectively). The SIgA concentration varied across time points ($p<0.001$) with the highest value observed at the 3rd time-point (rest-day) (3rd versus all time-point; ES=0.47-0.73). No changes were observed for S-RPE across matches ($p> 0.05$). A higher number of tackles and interceptions were observed during the 4th match vs. 1st and 7th matches ($p<0.001$; ES=2.25 and 1.90, respectively). The present data demonstrate that accumulated fatigue related to participation in a congested match schedule might induce a decrease in testosterone concentration in youth players' and negatively affect their mucosal immunity and capacity to perform certain technical actions.

Key-words: football; competition; testosterone; cortisol; saliva; Immunoglobulin A.

INTRODUCTION

There is a growing body of literature on the effect of congested match schedules on physical and physiological performance in elite senior soccer players (Dellal et al., 2013; Dupont et al., 2010; Odetoyinbo, Wooster, & Lane, 2009). However, official youth tournaments in which elite young soccer players participate can also involve congested schedules (Arruda et al., 2015; Buchheit, Horobeau, Mendez-Villanueva, Simpson, & Bourdon, 2011; Rowsell, Coutts, Reaburn, & Hill-Haas, 2011). While these studies have shown that running activity can be affected due to match congestion, relatively little is known about the potential effect of playing successive matches in a short period on technical performance metrics in youth populations.

This knowledge is important as it have been reported that physical fatigue represented by a reduction in running activity can affect skill-related performance in soccer match-play (Rampinini, Impellizzeri, Castagna, Coutts, & Wisløff, 2009). Indeed, Stone and Oliver (2009) suggested that resistance to fatigue plays a role in the effectiveness of players' ability to maintain optimal performance and perform precise movements during match play. Rampinini et al. (2009) reported a decline in technical and physical performance when comparing first and second halves of play in elite senior players. In contrast, Carling and Dupont (2011) demonstrated that neither physical nor skill-related performance was affected across three consecutive games within a period of ≤ 7 days in an elite senior soccer team. Research is warranted to examine physical and technical performance metrics in elite youth soccer populations.

Similarly, the effects of a very congested match schedule on neuro-immune-endocrine responses of youth players are still relatively unclear. Findings from previous studies suggest that successive matches and periods of increased

training load may affect psycho-physiological responses in youth players. Rowsell et al. (2009) demonstrated increased concentrations of muscle damage markers (creatine kinase and lactate dehydrogenase) in male junior players completing four competitive matches in 4 days with ~24 h between matches. Freitas et al. (2014) provided evidence that intensive periods of training may lead to an increase in salivary cortisol concentration but not stress tolerance in elite youth soccer players. In addition, Mortatti et al. (2012) reported that a decrement in mucosal immunity, as measured by salivary IgA concentration, can lead to a greater incidence of upper respiratory tract illness (URTI) in elite Under 19 soccer players monitored in a series of 7 matches in 20 days.

Given the importance of extending current knowledge in elite soccer regarding the impact of successive matches played in a short time interval using a more holistic and integrated approach (Carling et al., 2015), the aim of this study was to analyse the effects of a congested match schedule on steroid hormones concentrations (salivary testosterone and cortisol), mucosal immunity, technical performance parameters, and session rating of perceived exertion (S-RPE) in elite Under-15 players during a major national tournament. It was hypothesized that successive matches played in a short time interval would negatively affect technical performance, and simultaneously lead to negative changes in neuro-immune-endocrine parameters.

METHODS

Participants

Eighteen elite Under-15 Brazilian soccer players initially volunteered to participate in this study. Of these, 16 outfield players participated in matches and

were thus considered for further analysis (age 14.8 ± 0.4 yr; 1.5 ± 0.7 yr from peak height velocity; height 170.6 ± 9.4 cm and body mass 64.9 ± 7 kg). Players typically participated in ~15 h of training per week (5-8 soccer training sessions, 3-4 strength training sessions, 1-2 conditioning sessions) and a single match. Informed consent was obtained from each player and their parents or guardians and the study was approved by the appropriate institutional ethics committee. All players underwent a thorough medical assessment to verify their health status prior to participation and were free from illness or injury at the time of this study.

Procedures

Player performances in a reference team were monitored during an annual national tournament in which seven matches were played in seven days during the second week of May. The first two matches were played on the first day of the competition (two halves of 20 min interspersed by 10 min rest). The 1st match of the day was played at 9:00 am and the 2nd at 3:00 pm. The 3rd and 4th matches were played on the second day and the 5th and 6th matches of the tournament were played on the sixth day of the week, after a 3 day lay-off. On the final day of the tournament, there was only one match, played at 9:00 am. The mean (\pm SD) temperature and humidity during the tournament were, 20°C (± 4) and 77% (± 5), respectively. The reference team won matches 1-4, drew matches 5 and 6 and lost match 7 (Table 1). No post-match recovery strategies were implemented during this study period. Saliva samples were collected on mornings in a fasted state (no food 6-7 hr before sampling) on each match day. Players were woken up approximately 15 min prior to sampling. Saliva sampling was also undertaken on day 3 (without match) at the same time of day (fasted state). Session rating of

perceived exertion (S-RPE) was used to quantify the perceived intensity of each match (Foster, 1998). All players were familiar with these procedures.

*****INSERT TABLE 1 NEAR HERE*****

Video Analysis and Technical Performance

Video recordings were collected using two cameras (Panasonic, 60Hz frequency acquisition). One was located 15 m above and to one side of the long axis of the pitch, and the other was placed 5 m to one side of the pitch to facilitate the player's identification. SportsCode GameBreaker Plus 10 software (Sportstec, Sydney, Australia) was used to code match performance. The inter- and intra-reliability was found to be excellent when analyzing two trials for each match using two experienced analysts. The Kappa values for the analysed variables ranged between 0.90-0.98 (intra-observer) to 0.82-0.91 (inter-observers). The technical events were chosen to fall in line with those used in previous research literature (Rampinini et al., 2009; Waldron & Worsfold, 2010) and included attacking and defending variables. The frequencies of the following technical actions were recorded for the reference team as a whole and each respective opponent:

- Involvements with the ball: all situations where the player was in contact with the ball.
- Goal attempts: number of attempts to score a goal.
- Total passes: number of short and long foot passes performed by a player.

- Successful (completed) passes: number of short and/or long passes performed by a player and successfully received by another player of his own team.
- Total headers: number of times where a player touched the ball with his head.
- Tackles and interceptions: number of situations where a player contested the ball with an opponent player irrespective of whether these situations involved or not clear physical contact between players.
- Total involvements: sum of all assessed technical actions.

Attacking variables included: involvements with the ball, goal attempts, passes (total passes and successful [completed] passes); defending variables included: tackles and interceptions made. Headers were included but not classified according to whether these were attacking or defending actions. Descriptive data for the assessed team as a whole (total number of team technical actions) and its opponents are presented.

Saliva Collection and Analysis

All players remained in a seated position, with eyes open, head tilted slightly forward, and making minimal orofacial movement during saliva collection. Unstimulated saliva was collected in a preweighed sterile 15 mL centrifuge tube over 5 min and stored at -80°C until assayed. The tubes were re-weighed before analysis, so that saliva volume could be estimated with saliva density assumed to be $1.00\text{ g}\cdot\text{mL}^{-1}$. Salivary flow rate was determined by dividing the volume of saliva collected by the duration of the sampling period. After thawing and centrifugation (10,000 g for 10 min at 4°C), the samples were tested for cortisol,

testosterone and salivary IgA (SIgA) concentrations in duplicate using enzyme-linked immunosorbent assays (ELISA, Salimetrics™) in accordance with standard procedures (Freitas et al., 2014; Moreira et al., 2013; Mortatti et al., 2012). The SIgA secretion rate (SIgA_{rate} - micrograms per minute) was calculated by multiplying the absolute SIgA_{abs} by salivary flow rate (milliliters per minute), in accordance with the procedure adopted by Mortatti et al. (2012). The mean intra-assay coefficient of variation for cortisol, testosterone, and SIgA concentrations was 3.9, 4.5, and 5.6%, respectively.

Match Intensity

In order to quantify match intensity, the S-RPE was assessed after each match (Foster, 1998). Each player rated the match intensity using the CR-10 sliding scale 30 min post-match. This method has been shown to be a valid method for monitoring load in young players (Freitas et al., 2014; Mortatti et al., 2012).

Statistical analyses

Data were analyzed using statistical software (version 12.0, StatSoft). Data normality was assessed by the Shapiro-Wilk's test and homoscedasticity by Levene's test. The technical performance data were not normally distributed and therefore the Friedman test was used to detect differences between matches. A Wilcoxon test with a Bonferroni adjustment was used to verify differences. The Friedman test was also used to compare S-RPE across the 7 matches. ANOVAs with repeated measures were used to compare cortisol, testosterone, and SIgA concentrations (absolute concentration and SIgA rate) in a fasted state across matches and the day without match (3rd day of the study). Fisher's LDS post hoc test was used when a significance was observed. Statistical significance was set at

$p \leq 0.05$. Effect size (ES) values of 0.20–0.49, 0.50–0.79, and ≥ 0.8 were considered to represent small, medium, and large differences between values respectively (Cohen, 1988).

RESULTS

Figure 1 illustrates resting and fasted salivary measures across the investigation period. No change was observed for salivary cortisol concentration across multiple matches ($F = 1.17$; $p=0.33$; $ES= 0.13-0.48$ [trivial to small]; Figure 1A). However, there was a significant change in testosterone concentration across the investigated period ($F = 2.97$; $p=0.02$). Post hoc tests showed a decrease on the 3rd compared to the 1st collection time-point ($p= 0.007$; $ES = 0.47$ [small]); an increase in testosterone concentration from the 3rd to the 4th collection time-point ($p = 0.003$; $ES=0.36$ [small]; Figure 1B); and a decrease in concentration when comparing the first with the last time-point ($p = 0.01$; $ES = 0.42$ [small]; Figure 1B). A significant change in SIgA concentration was also observed across time-points ($F = 4.21$; $p=0.004$). A decrease in SIgA concentration was reported between the first and last time-point ($p < 0.001$; $ES = 0.67$ [medium]; Figure 1C). The SIgA concentration at the 3rd time-point (without matches) was different (higher value) compared to all other time-points except the first time-point. In addition, SIgA concentration was higher on the day without matches in comparison to all other time-points ($p < 0.001$ for all time-point comparisons; $ES = 0.47$ [small], 0.48 [small], 0.73 [medium], for 3rd vs 2nd, 3rd vs. 4th and 3rd vs 5th, respectively). No difference was evident for SIgA rate ($F = 0.64$; $p = 0.63$; $ES=0.03-0.37$ [trivial-small]; Figure 1D) or saliva flow rate in a fasted state ($F = 0.85$; $p = 0.50$; $ES=0.03-0.10$ [trivial]). No change was observed for S-RPE across the experimental period ($\chi^2 = 10.83$; $p=0.09$). The mean and SD for S-RPE values

across the matches were: 7.7 ± 0.4 , 7.7 ± 0.6 , 7.1 ± 1.2 , 8.2 ± 0.7 , 7.2 ± 1.3 , 8.0 ± 0.8 , 7.7 ± 0.8 , for matches 1, 2, 3, 4, 5, 6, and 7, respectively.

*****INSERT FIGURE 1 NEAR HERE*****

Technical performance data are presented in figure 2 and table 2. Figure 2 shows the defending variables which were the only technical parameters that changed over the 7 played matches. A significant change in total tackles and interceptions was observed ($\chi^2 = 17.24$; $p=0.008$). A higher value for tackles and interceptions during the 4th match (second match played on the second day of the competition) was observed compared to the 1st (13.1 ± 4.8 vs 4.2 ± 3.1 ; ES = 2.25 [large]) and 7th matches (vs 5.6 ± 2.8 ; ES = 1.90 [large]). Moreover, successful tackles and interceptions also changed during the congested match schedule ($\chi^2 = 12.47$; $p=0.05$). Successful tackles and interceptions were decreased when comparing values for the 4th vs. the 7th match (5.0 ± 2.7 vs 1.0 ± 1.5 ; ES = 1.90 [large]) and between the 5th (4.0 ± 2.2) and the 7th matches (ES = 0.41 [trivial]). No change was observed for any other technical performance parameters ($p>0.05$). Table 2 shows the total teams occurrence of technical performance parameters over the 7 assessed matches. A higher number of total tackles and interceptions for the 3rd and 4th matches compared to all other matches was observed. The opponent team was the same (same players in the team) for the 3rd and last match (please see Table 1). During the former this team performed 163 tackles and interceptions against the reference team (vs 108), while 75 tackles and interceptions (vs 71) were performed during the final match.

*****INSERT FIGURE 2 NEAR HERE*****

*****INSERT TABLE 2 NEAR HERE*****

DISCUSSION

The aim of this study was to analyze the effects of a congested fixture schedule on steroid hormone concentrations, mucosal immunity, S-RPE and technical match performance parameters in Under-15 players during a major national tournament. No changes in salivary cortisol concentration were observed over the congested match schedule; however, there was a significant decrease in the salivary concentration of testosterone (small ES) and SIgA (moderate ES) from the beginning to the end of the competition. Despite changes in resting testosterone and SIgA, there was no observed change in the session-RPE score across matches. Regarding technical performance data, defensive performance changed across the competition but attacking performance was unaffected.

The observed resting salivary cortisol response in the present study is in agreement with Mortatti et al. (2012). The authors investigated the subsequent impact of a competition and training period (7 matches over a 20-day period) on resting hormonal and immune parameters in young players from a reference team. The authors reported no alteration in salivary cortisol concentration, despite a progressive increment in internal training load. Given the present results and those reported by Mortatti et al. (2012) it could be suggested that both a short-congested period (7 matches in 7 days with short match halves duration) and a longer (7 matches in 20 days with long match halves duration) did not affect significantly the HPA axis responses of youth soccer players, at least, considering their resting cortisol values. The absence of a significant change in resting cortisol during the present match schedule in youth players might suggest that this congested period of matches does not represent a higher physiological stimulus than the habitual

training schedule of these players. In addition, the absence of changes in resting salivary cortisol maybe, at least in part, attributed to the high-fitness level of the assessed players and to their habitual high daily and weekly-volume training schedule.

Nevertheless, the cortisol response to the congested match schedule should be viewed with caution. In the present study only resting salivary concentration was examined. Some authors have proposed that the examination of exercise induced hormonal responses rather than resting level of hormones may offer a better scenario when regarding hormonal changes, for example during training or even during states of non-functional overreaching and syndrome of overtraining (Hough et al., 2013; Meeusen et al., 2004; 2010). However, it is reasonable to suggest that the practical usefulness of using this approach to assess hormonal changes during a fixture congested match study may be limited.

In contrast to cortisol responses, a significant change in testosterone concentration was observed notably on the 3rd day of collection (morning of the day without match) compared to the 1st and the 4th collection day. A decrease in testosterone concentration has previously been reported after different types of exercise, lasting for hours to days (Peñailillo, Maya, Niño, Torres, & Zbinden-Foncea, 2015; Urhausen and Kindermann, 1987). Peñailillo et al. (2015) for example, compared changes in testosterone concentration from pre- to post-match (5-10 min after match), and demonstrated that salivary testosterone decreased by ~30% after a soccer match in elite soccer players. The authors also reported that players who showed smaller decreases in testosterone level had covered more distance and illustrated less pronounced alterations to their immune function. While the mechanisms underlying such decrements are beyond the scope of the present investigation, previous observations have nevertheless suggested potential

mechanisms: 1) a disturbance in gonadotrophin-releasing hormone (GnRH) and testicular luteinising hormone (LH) pulsatile release; 2) a desensitizing effect on LH receptors; 3) an increased metabolic rate of testosterone and 4) a catecholamine-induced increase in the testicular vascular resistance (Urhausen, Gabriel, & Kindermann, 1995).

The current results suggest that the successive matches played during the first 2 days of the competition (4 matches in 2 days) potentially affected resting testosterone concentration. Nevertheless, after a recovery period (the time between the 3rd and 4th collection days) the players' testosterone concentration returned to initial levels (higher values at the 4th time-point compared to the 3rd). The accumulated fatigue potentially generated by the congested competitive schedule seems to lead to a decrement in testosterone as a decrease was also observed on the 5th collection day.

Previous research found that testosterone concentration was affected by tiredness and the magnitude of fatigue (Maso, Lac, Filaire, Michaux, & Robert, 2004). The authors suggested that a low testosterone level could be used a marker of fatigue, and might reflect the residual effect of a previous accumulated training load and an insufficient recovery period. Thus, the lower salivary testosterone induced by the congested competition may reflect accumulation of fatigue linked to insufficient recovery intervals. In addition, Papadopoulos et al. (2014) observed a decrease in testosterone concentration during a week leading to competition as compared to a control (noncompetition) week, in competitive swimmers (11-15 yr old). Based on previous findings reported by Maso et al. (2004) and Papadopoulos et al. (2014), it seems plausible that the observed testosterone response in the present study is associated to the accumulative fatigue generated during the congested fixture competition. This finding is novel and adds important

information to the literature regarding the effect of a short-term congested match schedule on salivary testosterone concentration in youth players, and the use of this measure to monitor fatigue in youth players.

Results showed that mucosal immunity was also affected over the congested competitive period. A decrease in SIgA was observed when comparing the first and the final saliva collection time-point (fasted state). It is noteworthy that the mean SIgA value at the 3rd time-point was significantly higher than all other saliva collection time-points. An insufficient recovery period between successive training sessions can cause suppression of the immune system (Bishop & Gleeson, 2009). Hence it is reasonable to suggest that the increased stress related to the assessed congested fixture schedule on youth players may have compromised players' mucosal immunity. This result suggests that the additional stress of participating in a congested match schedule competition may lead to changes in mucosal immune function in youth players. A significant change in SIgA concentration was reported by Nazem et al. (2011) in young female handball players. The authors reported a decrement in SIgA concentration following matches considered by the players as 'hard' (perceived difficulty of the matches was rated by players). Collectively, the results from the present study and those by Nazem et al. (2011), suggest that additional factors, such as match intensity, competitive anxiety, perceived difficulty of the matches, pressure to perform well, and successive matches, could impose a high level of stress on young team sports players, leading therefore to alterations in their mucosal immunity.

During the present congested match schedule period, S-RPE values were unchanged. However, it is noteworthy that despite the unchanged S-RPE values over the congested match schedule, the matches were played in a high intensity

zone (≥ 7 ; Lovell, Sirotic, Impellizzeri, & Coutts, 2013; Moreira et al., 2015) demonstrating the high demands placed on the players. This result suggests that despite the likely accumulated fatigue represented by the results from testosterone and SIgA responses, the players might have employed pacing strategies in order to regulate their efforts to avoid fatigue and consequently their perceived effort was unaltered (Bradley & Noakes, 2013). Whilst, the absence of data on players' running activity is a limitation of the present study for discussing whether the assessed soccer players employed some form of pacing strategy to avoid fatigue, this possibility should be mentioned, although it is evidently speculative. Indeed, it is worth mentioning that the presence of a pacing strategy in young team players has previously been reported (Sampson, Fullagar, & Gabbett, 2014), and that schoolchildren at a more advanced stage of cognitive development (Under 14) exhibited the well-known U-shaped pacing strategy during the first 15% and last 20% of a 4-min run (Micklewright et al., 2012).

Contrary to the hypothesis, a congested match schedule did not affect most of the technical performance parameters assessed in the present study. However, it is worth noting that there was a change in defensive actions, which might be associated to the fixture congestion. This assumption may be reinforced by the fact that these actions were also affected for the opponent teams in the competition. Analysis of the descriptive data of the opponents showed a higher number of tackles and interceptions for the 3rd and 4th matches (163 and 130 tackles and interceptions for the opponent teams, respectively), whilst only 75 tackles and interceptions were observed for the final match. It is worth noting that the opponent team was the same for the 3rd and last match (Table 1). Additionally, it should be highlighted that there was a meaningful decrement (3rd vs last match)

regarding defensive parameters for the assessed team and the opponent team which in turn affected their total involvements.

One potential explanation for the decrement in defensive actions may be the accumulated fatigue associated to the congested competition schedule. Folgado, Duarte, Marques, and Sampaio (2015) speculated that the lower percentage of dyadic synchronization of players observed during their study may be explained by accumulated mental fatigue that impairs a players' capacity to be synchronized with teammates during periods of low-intensity. It is noteworthy that a meaningful reduction in the frequency of defensive actions of the reference players was observed for the last two matches (vs 4th match) of the competition. Notably, the 6th match ended tied and the assessed team lost the final match of the competition. In addition, as a decrease in the salivary testosterone was also observed, it could be inferred that a fatigue state would have actually impaired the capacity of the player to perform, notably in executing defensive actions.

While this is a novel and interesting finding of the effect of a congested match schedule on technical performance in youth players, further empirical studies should be conducted to corroborate the assumption that this reduction in the defensive performance might in fact be attributed to the fatigue state. For example, an investigation into the effect of consecutive running simulations on performance in technical skill tests would be pertinent (Carling et al, 2015). Additionally, including psychometric tools (questionnaires) in order to monitor both the perceived fatigue and perceived effort would add important information regarding the effect of a successive soccer physical demand on players' skills.

CONCLUSION

The data demonstrate that the congested period impacted upon salivary immune-endocrine, notably on testosterone and SIgA responses. Moreover, a congested period led to a change in defensive but not in attacking performance metrics. The present results suggest that accumulated fatigue induced by a congested schedule resulted in a decrease in testosterone concentration in youth players and negatively affected their mucosal immunity and capacity to perform certain technical actions. Changes in testosterone have been linked to training motivation, performance, training outcomes, and effort during competition; and declines in SIgA have been associated with an increased risk of upper respiratory tract infections. Hence coaches and support staff should adopt strategies to monitor these parameters during training and competition, in order to evaluate players' ability to cope with the demands of athletic training and competition. However, it is important to highlight that the present findings are representative of a very unique congested match schedule for youth players and thus the results should not be generalized to other situations (i.e. normal training week and other kinds of congested schedules competition or senior players).

REFERENCES

- Arruda, A. F., Carling, C., Zanetti, V., Aoki, M. S., Coutts, A. J., & Moreira, A. (2015). Effects of a very congested match schedule on body load impacts, accelerations, and running measures in youth soccer players. *International Journal of Sports Physiology and Performance*, *10*, 248–252.
- Bishop, N., & Gleeson, M. (2009). Acute and chronic effects of exercise on markers of mucosal immunity. *Frontiers in Bioscience*, *14*, 4444–4456.

- Bradley, P. S., & Noakes, T. D. (2013). Match running performance fluctuations in elite soccer: Indicative of fatigue, pacing or situational influences? *Journal of Sports Sciences, 31*, 1627–1638.
- Buchheit, M., Horobeanu, C., Mendez-Villanueva, A., Simpson, B. M., & Bourdon, P. C. (2011). Effects of age and spa treatment on match running performance over two consecutive games in highly trained young soccer players. *Journal of Sports Sciences, 29*, 591–598.
- Carling, C., & Dupont, G. (2011). Are declines in physical performance associated with a reduction in skill-related performance during professional soccer match-play? *Journal of Sports Sciences, 29*, 63–71.
- Carling, C., Gregson, W., McCall, A., Moreira, A., Wong, D. P., & Bradley, P. S. (2015). Match running performance during fixture congestion in elite soccer: research issues and future directions. *Sports Medicine, 45*, 605–613.
- Cohen J. (1988). *Statistical power analysis for the behavioral sciences*. Hillsdale: Lawrence Erlbaum.
- Dellal, A., Lago-Peñas, C., Rey, E., Chamari, K., & Orhant, E. (2013). The effects of a congested fixture period on physical performance, technical activity and injury rate during matches in a professional soccer team. *British Journal of Sports Medicine, 49*, 390-394.
- Dupont, G., Nedelec, M., McCall, A., McCormack, D., Berthoin, S., & Wisløff, U. (2010). Effect of 2 soccer matches in a week on physical performance and injury rate. *The American Journal of Sports Medicine, 38*, 1752–1758.
- Folgado, H., Duarte, R., Marques, P., & Sampaio, J. (2015). The effects of congested fixtures period on tactical and physical performance in elite football. *Journal of Sports Sciences, 33*, 1238–1247.
- Foster, C. (1998). Monitoring training in athletes with reference to overtraining

syndrome. *Medicine and Science in Sports and Exercise*, 40, 1164–1168.

Freitas, C. G., Aoki, M. S., Francison, C. A., Arruda, A. F. S., Carling, C., & Moreira, A. (2014). Psychophysiological responses to overloading and tapering phases in elite young soccer players. *Pediatric Exercise Science*, 26, 195–202.

Hough, J., Corney, R., Kouris, A., & Gleeson, M. (2013). Salivary cortisol and testosterone responses to high-intensity cycling before and after an 11-day intensified training period. *Journal of Sports Sciences*, 31, 1614–1623.

Lovell, T. W. J., Sirotic, A. C., Impellizzeri, F. M., & Coutts, A. J. (2013). Factors affecting perception of effort (session rating of perceived exertion) during rugby league training. *International Journal of Sports Physiology and Performance*, 8, 62–69.

Maso, F., Lac, G., Filaire, E., Michaux, O., & Robert, A. (2004). Salivary testosterone and cortisol in rugby players: correlation with psychological overtraining items. *British Journal of Sports Medicine*, 38, 260–263.

Meeusen, R., Piacentini, M. F., Busschaert, B., Buyse, L., De Schutter G, & Stray-Gundersen, J. (2004). Hormonal responses in athletes: The use of a two bout exercise protocol to detect subtle differences in (over)training status. *European Journal of Applied Physiology*, 91, 140–146.

Meeusen, R., Nederhof, E., Buyse, L., Roelands, B., de Schutter, G., & Piacentini, M. F. (2010). Diagnosing overtraining in athletes using the two-bout exercise protocol. *British Journal of Sports Medicine*, 44, 642–648.

Micklewright, D., Angus, C., Suddaby, J., St Clair Gibson, A. S. C., Sandercock, G., & Chinnasamy, C. (2012). Pacing strategy in schoolchildren differs with age and cognitive development. *Medicine & Science in Sports & Exercise*, 44, 362–369.

Moreira, A., Coutts, A. J., Moreira, A., Bilsborough, J. C., Sullivan, C. J.,

Cienciosi, M., ... Coutts, A. J. (2015). Training periodization of professional australian football players during an entire australian football league season. *International Journal of Sports Physiology and Performance*, *10*, 566–571.

Moreira, A., Mortatti, A., Aoki, M., Arruda, A., Freitas, C., & Carling, C. (2013). Role of free testosterone in interpreting physical performance in elite young brazilian soccer players. *Pediatric Exercise Science*, *25*, 186–197.

Mortatti, A., Moreira, A., Aoki, M. S., Crewther, B., Castagna, C., Arruda, A., & Filho, J. (2012). Effect of competition on salivary cortisol, immunoglobulin A, and upper respiratory tract infections in elite young soccer players. *The Journal of Strength and Conditional Reserach*, *26*, 1396–1401.

Nazem, G., Sharifi, G. R., Taghian, F., Jourkesh, M., Ostoic, S., Calleja-Gonzalez, J., & Keikhai, B. (2011). The effects of successive official competitions on salivary cortisol and immunogolobulin responses in women handballers. *Serbian Journal of Sports Sciences*, *5*, 67–73.

Odetoyinbo, K., Wooster, B., & Lane, A. (2009). The effect of a succession of matches on the activity profiles of professional soccer players. In T. Reilly & F. Korkusuz (Eds.), *Science and Football* (pp. 182–185). London, United Kingdom: Routledge.

Papadopoulos, E., Muir, C., Russell, C., Timmons, B.W., Falk, B., & Klentrou, P. Markers of biological stress and mucosal immunity during a week leading to competition in adolescent swimmers. *Journal of Immunology Research*. Epub 2014, June 12. doi: 10.1155/2014/234565.

Peñailillo, L., Maya, L., Niño, G., Torres, H., & Zbinden-Foncea, H. (2015). Salivary hormones and IgA in relation to physical performance in football. *Journal of Sports Sciences*, *33*, 2080-2087.

Rampinini, E., Impellizzeri, F. M., Castagna, C., Coutts, A. J., & Wisløff, U.

(2009). Technical performance during soccer matches of the Italian Serie A league: effect of fatigue and competitive level. *Journal of Science and Medicine in Sport*, 12, 227–233.

Rowell, G. J., Coutts, A. J., Reaburn, P., & Hill-Haas, S. (2009). Effects of cold-water immersion on physical performance between successive matches in high-performance junior male soccer players. *Journal of Sports Sciences*, 27, 565–573.

Rowell, G. J., Coutts, A. J., Reaburn, P., & Hill-Haas, S. (2011). Effect of post-match cold-water immersion on subsequent match running performance in junior soccer players during tournament play. *Journal of Sports Sciences*, 29, 1–6.

Sampson, J.A., Fullagar, H.H., & Gabbett, T. (2014). Knowledge of bout duration influences pacing strategies during small-sided games. *Journal of Sports Sciences*, 33, 85-98.

Stone, K. J., & Oliver, J. L. (2009). The effect of 45 minutes of soccer-specific exercise on the performance of soccer skills. *International Journal of Sports Physiology and Performance*, 4, 163–175.

Urhausen, A., Gabriel, H., & Kindermann, W. (1995). Blood hormones as markers of training stress and overtraining. *Sports Medicine*, 20, 251–276.

Urhausen, A., Kindermann, W. (1987). Behavior of testosterone, sex hormone binding globulin (SHBG), and cortisol before and after a triathlon competition. *International Journal of Sports Medicine*, 8, 305-308.

Waldron, M., & Worsfold, P. (2010). Differences in the Game Specific Skills of Elite and Sub-Elite Youth Football Players: Implications for Talent Identification. *International Journal of Performance Analysis in Sport*, 9–24.

Table 1. Competition schedule and results

Match	Opponent	Result	Assessed team match outcome	Day of the investigation; Period
First	SE Palmeiras	1 – 0	Won	1 st ; Morning
Second	Santos FC	3 - 2	Won	1 st ; Afternoon
Third	SC Internacional	1 – 0	Won	2 nd ; Morning
Fourth	Audax	1 – 0	Won	2 nd ; Afternoon
Fifth	Bahia EC	1 – 1	Draw	6 th ; Morning
Sixth	Botafogo RJ	0 – 0	Draw	6 th ; Afternoon
Seventh	SC Internacional	0 - 3	Lost	7 th ; Morning

Table 2. Total teams (assessed versus opponent teams) occurrence of technical performance parameters over the 7 assessed matches

Assessed team score vs opponent team score							
Variable	1st match	2nd match	3rd match	4th match	5th match	6th match	7th match
Goal attempts	6 vs 7	7 vs 7	4 vs 4	10 vs 3	10 vs 12	4 vs 4	3 vs 13
Total passes	166 vs 142	149 vs 168	174 vs 183	119 vs 213	158 vs 137	123 vs 186	135 vs 139
Successful passes	128 vs 107	112 vs 118	128 vs 112	77 vs 158	118 vs 87	76 vs 137	95 vs 93
Unsuccessful passes	38 vs 35	37 vs 50	46 vs 71	42 vs 55	40 vs 50	47 vs 49	40 vs 46
Total headers	8 vs 16	14 vs 30	29 vs 18	13 vs 17	16 vs 15	5 vs 20	25 vs 29
Total tackles /interceptions	60 vs 62	79 vs 147	108 vs 163	141 vs 130	109 vs 102	100 vs 95	71 vs 75
Total involvement	350 vs 334	360 vs 454	423 vs 424	351 vs 469	409 vs 343	312 vs 375	344 vs 322

Figure 1. Resting salivary measures (fasted state) across the investigation period (Mean \pm SD).

Figure 2. Tackles and interceptions over the 7 assessed matches (Mean \pm SD)





Figure 1. Resting salivary measures (fasted state) across the investigation period (Mean ± SD). 338x190mm (96 x 96 DPI)



†= significant difference to 1st and 7th matches (p<0.05); * significant difference to 4th and 5th matches (p<0.05); TTI = Total tackles/interceptions; STI = Successful tackles /interceptions