

LJMU Research Online

Eijsvogels, TM, George, KP and Thompson, PD

Cardiovascular benefits and risks across the physical activity continuum.

<http://researchonline.ljmu.ac.uk/id/eprint/3984/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Eijsvogels, TM, George, KP and Thompson, PD (2016) Cardiovascular benefits and risks across the physical activity continuum. Current Opinions in Cardiology, 31 (5). pp. 566-571. ISSN 0268-4705

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

Cardiovascular Benefits and Risks Across the Physical Activity Continuum

Thijs M.H. Eijsvogels, Ph.D^{1,2}

Keith P. George, Ph.D¹

Paul D. Thompson, M.D.³

Affiliations:

¹Research Institute for Sports and Exercise Sciences, Liverpool John Moores University, Liverpool, United Kingdom. ²Department of Physiology, Radboud University Medical Center, Nijmegen, The Netherlands. ³Division of Cardiology, Hartford Hospital, Hartford, Connecticut.

Manuscript word count: 2473

Funding: The work of T.M.H.E is supported by a European Commission Horizon 2020 grant (Marie Skłodowska-Curie Fellowship 655502).

Conflict of Interest: T.M.H.E and K.P.G. reports no conflicts. P.D.T. reports receiving research grants from the Aventis, Regeneron, Sanolfi, and Pfizer; serving as a consultant for Aventis, Regeneron, Merck, Genomas, Abbvie, Sanolfi, and Pfizer; receiving speaker honoraria from Regeneron, Sanolfi, Amgen, Aventis, and Merck; owning stock in General Electric, JA Wiley Publishing, J&J, and Abbvie, Abbott, Medtronic, and Cryolife.

Author for correspondence:

Dr. Thijs Eijsvogels, Research Institute for Sports and Exercise Sciences, Liverpool John Moores University, Tom Reilly Building, L3 3AF, Liverpool, United Kingdom
E-mail: T.M.Eijsvogels@lmu.ac.uk. Tel. +44 151 904 62 64

ABSTRACT

Purpose of review: Habitual physical activity can reduce the risk of future cardiovascular morbidity and mortality. This review evaluates recent publications that have assessed the impact of the dose of physical (in)activity on cardiovascular outcomes.

Recent findings: Sedentary behavior, characterized by prolonged sitting, is increasingly prevalent across the globe and increases the risk for cardiovascular events in a dose-dependent fashion. Similarly, the number of individuals performing endurance exercise events has tripled over the last 2 decades, and some studies suggest that the high volumes of exercise training and competition may attenuate the health benefits of a physically active lifestyle.

Summary: Breaking-up sitting time or replacing sitting by (light) physical activity are effective strategies to attenuate its detrimental health effects. Low doses of physical activity, preferably at a high-intensity, significantly reduce the risk for cardiovascular and all-cause mortality. Larger doses of exercise yield larger health benefits. Extreme doses of exercise neither increase nor decrease the risk for adverse outcomes. Athletes demonstrate a transient cardiac dysfunction and biomarker release directly post-exercise. Chronic exercise training may increase the risk for atrial fibrillation, but is also associated with a superior life expectancy compared to the general population.

Keywords: sedentary behavior, sitting, endurance exercise, athletes, lifestyle

INTRODUCTION

Physical activity and the ability to perform endurance exercise played an essential role in human evolution [1]. Our early ancestors combined long-distance running and walking to track and hunt animals on the African savannah. During so-called ‘persistence hunts’, distances >30 km were regularly covered [2]. In contrast to this intermittent but substantial exertion, it is believed that hunters were predominantly physically inactive during the remainder of the day [1]. This inactive behavior reduced their energy expenditure and was essential to maintain a proper balance between energy intake and expenditure.

During the past century, our lifestyle has changed dramatically and the role of physical exertion is minimized in our contemporary lives. Machines have taken over the majority of our physical efforts at work, home and during transportation. Consequently, the prevalence of sitting time has increased, whereas the time performing exercise has decreased. These changes in habitual physical activity patterns greatly impact the energy intake/expenditure balance, which have contributed to an alarming increase in the incidence of obesity and other chronic diseases [3, 4]. Hence, physical inactivity was recently recognized as a major threat to global health [5].

The World Health Organization recommends that adults engage in at least 150 min/week of moderate-intensity exercise or 75 min/week of vigorous-intensity exercise [6]. Currently, only 61% of the European population [7] and 44% of the North American population [8] perform sufficient physical activity to meet the WHO guidelines, percentages that have changed only slightly over the past 20 years [9]. Incongruously, an increasing number of amateur athletes are participating in endurance exercise events. In fact, the number of US running race participants has tripled over the past

two decades [10]. Although exercise training is believed to improve cardiovascular health [11], recent studies suggest that excessive volumes of physical activity may harm the heart [12].

The purpose of this review is to provide an overview of recent insights relating to the risks and benefits of physical activity. Given the increased prevalence of physical inactivity and the increasing popularity of endurance exercise activities, we will summarize the cardiovascular risks and benefits across the physical activity continuum: from sitting behavior to extreme volumes of exercise.

PHYSICAL INACTIVITY AND SITTING BEHAVIOR

Physical activity is defined as any bodily movement produced by skeletal muscles that results in energy expenditure beyond resting expenditure. Hence, physical inactivity represents sedentary behavior that does not involve muscle contraction, which is most prevalent during sitting and lying. Recent studies revealed that accelerometer-measured mean daily sedentary time was 8.2 hrs/day among New York City adults [13], whereas Australian desk workers reported an average of 9.0 hrs/day of sitting time [14]. Physical inactivity is not restricted to the general population; it can be observed in (half-)marathon runners as they have reported sitting 10.75 hrs/day on workdays and 8 hrs/day on non-workdays [15].

A recent meta-analysis combining the outcomes of 41 studies ($n=829,917$ participants) found that sedentary time was associated with cardiovascular mortality (HR: 1.18, 95%CI: 1.11-1.26), but also with cardiovascular disease incidence (HR: 1.14, 95%CI: 1.00-1.73), cancer mortality (HR: 1.17, 95%CI: 1.11-1.24), cancer incidence (HR: 1.13, 95%CI 1.05-1.21), and the incidence of type 2 diabetes (HR: 1.91, 95%CI: 1.64-2.22) [16]. The authors emphasized that the detrimental health effects of

sitting were independent of the physical activity patterns of study participants [16]. The population-attributable fraction for all-cause mortality associated with sitting time was explored in another paper and included data from 54 countries. Sitting time was responsible for 3.8% of all-cause mortality, but large differences were observed across countries (0.6-11.6%) [17]. The sitting-related mortality risk was the highest in Western Pacific countries (5.7%), followed by European (4.4%), Eastern Mediterranean (3.3%), American (3.2%), and Southeast Asian (2.0%) countries [17].

A potential strategy to reduce the harmful effects of prolonged sitting, is to limit the duration of sitting sessions [18]. Breaking-up prolonged sitting time with 2 minute bouts of walking reduced postprandial glucose and insulin levels [19] and lowered systolic and diastolic blood pressure [20]. A different strategy is to replace sitting time with exercise or non-exercise activities (i.e. household chores, lawn and garden work, and daily walking). In a cross-sectional analysis, less active individuals (<2 hrs/day, n=69,606) demonstrated a reduced risk for cardiovascular mortality when 1 hr/day of sitting was replaced by exercise (HR: 0.47, 95%CI: 0.40-0.56) or non-exercise activities (HR: 0.64, 95%CI: 0.57-0.71) [21]. Active individuals (≥ 2 hrs/day, n=85,008) also demonstrated a reduced risk for cardiovascular mortality when 1 hr/day of sitting was replaced by exercise (HR: 0.84, 95%CI 0.78-0.90) but no benefit was observed for non-exercise activities (HR: 1.00, 95%CI: 0.96-1.04) [21].

An observational study modeled the health benefits of replacing 2 hrs/day of sitting by standing or stepping. Sitting-to-standing reallocation was associated with lower levels of fasting glucose (~2%), total/HDL-cholesterol ratio (~6%), triglycerides (~11%), and a higher HDL-cholesterol (~0.06 mmol/L) [22]. Sitting-to-stepping reallocation was associated with a lower BMI (~11%) and waist circumference (~7.5 cm), and lower levels of post-load glucose (~12%), triglycerides (~14%) and a higher

HDL-cholesterol (~0.10 mmol/L) [22]. The benefits of low-intensity activities to reduce the detrimental effects of sitting were reinforced by a recent study in the UK Women's Cohort Study (n=12,778) [23]. Sitting ≥ 7 hrs/day significantly increased the risk for all-cause mortality compared to sitting <5 hrs/day. However, fidgeting behavior (small movements of hands and feet) appeared to modify the association between sitting time and all-cause mortality. The increased mortality risk associated with sitting was only observed in women reporting no fidgeting, whereas women reporting regular to frequent fidgeting demonstrated comparable mortality risks between high and low volumes of sitting [23].

Importantly, these observations demonstrate that small changes in sitting behavior can improve (cardiovascular) health. Consequently, policy documents from the United Kingdom and Australia already include statements about sitting behavior [24, 25]. Experts from the United Kingdom recommend including specific guidelines on sedentary behavior in future physical activity guidelines [24]. In Australia, minimizing time spent in prolonged sitting and breaking up long periods of sitting as often as possible are already included in the national physical activity guidelines [25]. These initiatives are likely to contribute to increased awareness of the detrimental health effects of sitting. Indeed, the time spent in sedentary behavior has not increased in European adults over the past decade [26]. More importantly, the prevalence of prolonged sitting (7.5 hrs/day) decreased from 23.1% in 2002 to 17.8 in 2013 [26]. Further reductions in sitting times may be achieved via workplace interventions such as sit-stand desks, but high quality intervention trials are needed to provide evidence for the cost-effectiveness and health benefits of such interventions [27].

THE OPTIMAL EXERCISE DOSE

Exercise is associated with risk reductions in at least 26 different diseases, including the metabolic syndrome, polycystic ovarian syndrome, type 1 and 2 diabetes, cancer, musculo-skeletal disorders and psychiatric, neurological, cardiovascular, pulmonary and metabolic diseases [28]. Furthermore, physically active individuals have a lower risk for all-cause and cardiovascular mortality and morbidity compared to sedentary peers [29, 30].

Several recent papers have explored the dose-response relationship between physical activity and adverse health outcomes [31-33]. A pooled analysis including 661,137 men and women from 6 large prospective American and European population studies found a 20% risk reduction for all-cause mortality in individuals performing moderate-intensity physical activity <100 min/week during 14 years of follow-up [34]. Increasing volumes of physical activity gradually decreased the mortality risk. The maximal benefit of an active lifestyle was found at an exercise dose representing 3 to 5 times the WHO physical activity recommendation (HR: 0.61, 95%CI: 0.59-0.62) [34]. Larger doses of exercise did not further decrease mortality risks, but did not increase it either. These findings align with a recent perspective document from the American College of Cardiology's (ACC's) Sports and Exercise Cardiology Leadership Council. They reported that the 'optimal' exercise dose to reduce the risk for cardiovascular events was established at 41 MET-hrs/week: i.e. 9.1 hrs/week of moderate-intensity exercise [35].

Interestingly, the dose-response relationship appears to be different for moderate *versus* vigorous intensity activities. Whereas a progressive decrease in the risk for cardiovascular mortality is observed for increasing volumes of moderate intensity physical activity, no further risk reduction is observed beyond a vigorous intensity exercise dose of 1.3 hrs/week (11 MET-hrs/week) [35]. These findings were

reinforced by an Australian prospective cohort study ($n=204,542$) that investigated the role of vigorous activities on all-cause mortality during 6.5 years of follow-up. Individuals performing <30% of their total exercise dose at a vigorous intensity had a significantly lower mortality risk (HR: 0.91, 95%CI 0.84-0.98) compared to the reference group that performed a similar exercise dose but refrained from vigorous activities [36]. Individuals reporting $\geq 30\%$ of their exercise dose to be vigorous demonstrated a comparable mortality risk reduction (HR: 0.87, 95%CI: 0.81-0.93) [36]. Thus, low doses of vigorous intensity physical activities seem to be extremely efficient at reducing the risk for adverse (cardiovascular) events.

Despite the undeniable health benefits of exercise, a substantial proportion of the population does not perform enough physical activity to derive a health benefit [37]. Therefore, novel strategies to improve active behavior are needed [38]. Activity trackers are available globally and these devices provide real-time quantification and insight into one's activity pattern. Hence, these trackers can assist an individual in reaching activity goals and adopting a physically active lifestyle. A randomized clinical trial found an increase of 970 steps/day in individuals receiving a wireless activity tracker, regardless of their initial activity level [39]. Personalized encouragement, social competition and effective feedback loops are other key factors needed to pursue a behavioral change towards an active lifestyle [40]. The 'setting' to stimulate individuals to become physically active is important. A randomized clinical trial compared 3 methods to frame financial incentives to increase physical activity among overweight and obese adults [41]. Participants were instructed to cover 7000 steps/day and were randomly allocated to a 1) control, 2) gain incentive (\$1.40/day if goal was achieved), 3) lottery incentive (daily eligibility for \$1.40 if goal was achieved), or 4) loss incentive (\$42 allocated monthly upfront and \$1.40/day was removed if goal was not achieved)

study groups. Only the loss incentive group demonstrated a larger proportion (0.45, 95%CI: 0.38-0.52) of participant-days achieving the 7000 steps/day goal compared to the control group (0.30, 95%CI: 0.22-0.37) [41]. These findings emphasize the difficulty in changing behavior, but also that a tailored intervention can increase activity patterns in a group at risk. Personalized exercise prescriptions are therefore warranted in the era of precision medicine.

TOO MUCH EXERCISE?

The volume of exercise performed during training and competition by amateur and professional athletes to improve cardiorespiratory fitness often exceeds the dose needed to optimize cardiovascular health. High volumes of exercise training improve cardiovascular risk factors [42], and cause an initial increase in left ventricular chamber size followed by an increase in wall thickness during chronic exercise training [43]. Remodeling also occurs in the right heart, allowing the right ventricle to tolerate the increased pulmonary artery pressures during exercise [44]. These adaptations are part of the ‘athlete’s heart’ and are believed to represent physiological remodeling.

Some studies suggest that performance of prolonged vigorous exercise (such as marathon running) may harm the heart acutely or chronically. For example, cardiac dysfunction of the left and right ventricles was observed immediately post-exercise in endurance athletes [45, 46]. Similarly, increased cardiac troponin levels have been reported following marathon running, with 69% of the population exceeding the upper limit of the normal threshold [47]. Both phenomena are transient, however, as cardiac function and biomarker levels typically recover within 24-48 hours post exercise [12]. The risk for acute adverse cardiac events during endurance exercise is therefore considered low, and data from a French registry demonstrated a prevalence of life threatening events of only 0.75 per 100 000 athletes [48].

Cardiac remodeling associated with chronic exercise exposure may also increase the risk for arrhythmias in athletes [49]. A previous athletic population study [50] and a systematic review [51] have demonstrated an increased risk for atrial fibrillation (AF) in endurance athletes. However, contrasting findings were reported in 2 recent studies. Data from the Henry Ford Exercise Testing (FIT) Project found a 7% risk reduction for AF with every increase of 1 MET in cardiorespiratory fitness [52]. Fit individuals had a substantially lower risk (HR: 0.44, 95%CI: 0.39-0.50) for incident AF compared to unfit individuals [52]. Similarly, data from the CARDIO-FIT Study found a 20% reduction in the risk of AF recurrence for each MET increase in cardiorespiratory fitness among overweight and obese individuals with symptomatic AF [53]. Differences in maximum exercise capacity and cardiorespiratory fitness between the FIT/CARDIO-FIT studies and previous observations in athletes may explain the conflicting outcomes [54]. It may well be that initial increases in cardiorespiratory fitness decrease the risk for AF, but that excessive exercise training and associated fitness increase the AF risk.

Finally, recent epidemiological studies have assessed the long-term outcomes of high volumes of exercise training. Data from the Million Women Study [55] and Copenhagen City Heart study [56] report a U-shaped curve for the relationship between exercise exposure and risk for morbidity and mortality. Thus, inactive individuals had the highest risks and physically active individuals demonstrated a reduced risk, but the benefits of exercise were attenuated in vigorous exercisers. These observations contradict pooled data from 6 European and American cohorts that noted that individuals performing exercise at a dose 5 – 10 times the international recommendations, had a 31% reduction in all-cause mortality risk (HR=0.69, 95%CI 0.59-0.78) compared to inactive peers [34]. Furthermore, a 50-year follow-up study among Finnish elite athletes demonstrated that endurance athletes (HR: 0.70, 95%CI:

0.61-0.79) and team sport athletes (HR: 0.80, 95%CI: 0.72-0.89) had lower mortality risks compared to controls [57]. Explanations for the different outcomes in the Million Women Study and Copenhagen City Heart study may relate to the characteristics of the most active individuals (high smoking prevalence and low socioeconomic status [31]) and definition of the control group (allowed to exercise <2h/week [58]). Therefore, we believe that there is currently no solid evidence for an increased risk for adverse outcomes in the most active individuals.

CONCLUSIONS

The priority for reducing cardiovascular burden should be mainly focused on the lower end of the physical activity continuum. Physical inactivity, characterized by too much sitting, is a serious health problem as it independently increases the risk for cardiovascular morbidity and mortality. Future physical activity guidelines should incorporate specific recommendations on strategies to reduce sedentary behavior. Habitual physical activity and exercise training are powerful strategies to reduce the risk for adverse cardiovascular outcomes and mortality in a dose dependent way. High intensity exercise produces larger health benefits compared to moderate intensity exercise. Personalized exercise programs and wireless devices with real-time feedback may help individuals meet the international guidelines for physical activity. Currently, there is no strong evidence that supports the ‘too much exercise hypothesis’. Individuals performing exercise at a multiple of the recommended dose live longer and have a comparable risk for cardiovascular morbidity and mortality as moderately active peers.

Key points

- Prolonged sitting is highly prevalent in the general population and increases the risk for cardiovascular mortality. Breaking-up of prolonged sitting time or replacement of sitting time by (light) physical activity can effectively reduce the detrimental effects of sitting.
- A curvilinear dose-response relationship between exercise and cardiovascular health is found. Low doses of exercise improve health, but higher doses give larger benefits. Also, high-intensity activities induce larger risk reductions compared to moderate-intensity activities of a similar volume.
- Exercise-induced cardiac remodeling of all cardiac chambers is present in athletes. Acute exercise can lead to transient cardiac dysfunction and cardiac biomarker release. Chronic exercise may increase the risk for atrial fibrillation. Nevertheless, there is strong evidence that athletes live longer compared to individuals from the general population.

REFERENCES

* of special interest

** of outstanding interest

* [1] Lieberman DE. Is Exercise Really Medicine? An Evolutionary Perspective. Current sports medicine reports 2015; 14:313-319.

An interesting review paper that summarizes the health benefits of exercise from an evolutionary angle.

[2] Liebenberg L. Persistence hunting by modern hunter-gatherers. Curr. Anthropol. 2006; 47:1017-1026.

[3] Piirtola M, Kaprio J, Waller K et al. Leisure-time physical inactivity and association with body mass index: a Finnish Twin Study with a 35-year follow-up. Int J Epidemiol 2016.

** [4] NCD Risk Factor Collaboration. Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 19·2 million participants. The Lancet 2016; 387 1377 - 1396.

Unique study that demonstrates global changes in BMI and prevalence of overweight and obesity across 40 years of follow-up.

[5] Lee IM, Shiroma EJ, Lobelo F et al. Effect of physical inactivity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. Lancet 2012; 380:219-229.

[6] World Health Organization W. Physical Activity and Adults. Global Strategy on Diet, Physical Activity and Health. In: 2010.

[7] Marques A, Sarmento H, Martins J, Saboga Nunes L. Prevalence of physical activity in European adults - Compliance with the World Health Organization's physical activity guidelines. Preventive medicine 2015; 81:333-338.

[8] Carlson SA, Fulton JE, Schoenborn CA, Loustalot F. Trend and prevalence estimates based on the 2008 Physical Activity Guidelines for Americans. Am J Prev Med 2010; 39:305-313.

[9] Moore LV, Harris CD, Carlson SA et al. Trends in no leisure-time physical activity--United States, 1988-2010. Res Q Exerc Sport 2012; 83:587-591.

[10] Lampka R, Yoder T. State of the Sport - Part III: U.S. Race Trends. In: Running USA; 2013.

* [11] Lavie CJ, Arena R, Swift DL et al. Exercise and the cardiovascular system: clinical science and cardiovascular outcomes. Circulation research 2015; 117:207-219.

Comprehensive overview about the impact of physical activity, exercise training, and cardiorespiratory fitness in the prevention and treatment of cardiovascular diseases

** [12] Eijsvogels TM, Fernandez AB, Thompson PD. Are There deleterious Cardiac Effects of Acute and Chronic Endurance Exercise? Physiological reviews 2016;

96:99-125.

Comprehensive review that summarizes the evidence that was published in the field of exercise cardiology over the past 30 years.

[13] Yi SS, Bartley KF, Firestone MJ et al. Self-reported sitting time in New York City adults, the Physical Activity and Transit Survey, 2010-2011. Prev Chronic Dis 2015; 12:E85.

[14] Bennie JA, Pedisic Z, Timperio A et al. Total and domain-specific sitting time among employees in desk-based work settings in Australia. Aust N Z J Public Health 2015; 39:237-242.

[15] Whitfield G, Pettee Gabriel KK, Kohl HW, 3rd. Sedentary and active: self-reported sitting time among marathon and half-marathon participants. J Phys Act Health 2014; 11:165-172.

** [16] Biswas A, Oh PI, Faulkner GE et al. Sedentary time and its association with risk for disease incidence, mortality, and hospitalization in adults: a systematic review and meta-analysis. Annals of internal medicine 2015; 162:123-132.

A timely paper summarizing the latest insight in the detrimental health effects of sitting.

* [17] Rezende LF, Sa TH, Mielke GI et al. All-Cause Mortality Attributable to Sitting Time: Analysis of 54 Countries Worldwide. Am J Prev Med 2016.

This paper provides insight in the consequences of sitting, and how reductions in sitting time may improve longevity.

[18] Benatti FB, Ried-Larsen M. The Effects of Breaking up Prolonged Sitting Time: A Review of Experimental Studies. Medicine and science in sports and exercise 2015; 47:2053-2061.

[19] Dunstan DW, Kingwell BA, Larsen R et al. Breaking up prolonged sitting reduces postprandial glucose and insulin responses. Diabetes care 2012; 35:976-983.

[20] Larsen RN, Kingwell BA, Sethi P et al. Breaking up prolonged sitting reduces resting blood pressure in overweight/obese adults. Nutr Metab Cardiovasc Dis 2014; 24:976-982.

[21] Matthews CE, Moore SC, Sampson J et al. Mortality Benefits for Replacing Sitting Time with Different Physical Activities. Medicine and science in sports and exercise 2015; 47:1833-1840.

** [22] Healy GN, Winkler EA, Owen N et al. Replacing sitting time with standing or stepping: associations with cardio-metabolic risk biomarkers. European heart journal 2015; 36:2643-2649.

This paper demonstrates that reductions in sitting time have beneficial effects on cardiovascular risk parameters.

* [23] Hagger-Johnson G, Gow AJ, Burley V et al. Sitting Time, Fidgeting, and All-Cause Mortality in the UK Women's Cohort Study. Am J Prev Med 2016; 50:154-160.

This paper suggests that frequent performance of low-intensity activities may attenuate the detrimental health effects of too much sitting.

[24] Bull FC, and the Expert Working Groups. Physical Activity Guidelines in the U.K.: Review and Recommendations. In: School of Sport, Exercise and Health Sciences, Loughborough University; May 2010.

[25] Brown WJ, Bauman AE, Bull FC, Burton NW. Development of Evidence-based Physical Activity Recommendations for Adults (18-64 years). In: Report prepared for the Australian Government Department of Health, August 2012.

* [26] Milton K, Gale J, Stamatakis E, Bauman A. Trends in prolonged sitting time among European adults: 27 country analysis. Preventive medicine 2015; 77:11-16.
This paper presents data on changes in sitting time across Europe during a decade of follow-up.

[27] Shrestha N, Ijaz S, Kukkonen-Harjula KT et al. Workplace interventions for reducing sitting at work. Cochrane Database Syst Rev 2015; 1:CD010912.

* [28] Pedersen BK, Saltin B. Exercise as medicine - evidence for prescribing exercise as therapy in 26 different chronic diseases. Scandinavian journal of medicine & science in sports 2015; 25 Suppl 3:1-72.

Comprehensive review paper summarizing the benefits of exercise in many chronic diseases

[29] Feldman DI, Al-Mallah MH, Keteyian SJ et al. No evidence of an upper threshold for mortality benefit at high levels of cardiorespiratory fitness. Journal of the American College of Cardiology 2015; 65:629-630.

* [30] Maessen MF, Verbeek AL, Bakker EA et al. Lifelong Exercise Patterns and Cardiovascular Health. Mayo Clinic proceedings 2016; 91:745-754.

This paper explores the dose-response relationship between lifelong exercise exposure and the prevalence of cardiovascular morbidity.

[31] Eijsvogels TM, Thompson PD. Exercise Is Medicine: At Any Dose? JAMA 2015; 314:1915-1916.

[32] Wen CP, Wai JP, Tsai MK et al. Minimum amount of physical activity for reduced mortality and extended life expectancy: a prospective cohort study. Lancet 2011; 378:1244-1253.

[33] Lee DC, Pate RR, Lavie CJ et al. Leisure-time running reduces all-cause and cardiovascular mortality risk. Journal of the American College of Cardiology 2014; 64:472-481.

** [34] Arem H, Moore SC, Patel A et al. Leisure time physical activity and mortality: a detailed pooled analysis of the dose-response relationship. JAMA Intern Med 2015; 175:959-967.

A large population study providing strong evidence for the benefit of low, moderate and high dosages of exercise on cardiovascular mortality risks.

** [35] Eijsvogels TM, Molossi S, Lee DC et al. Exercise at the Extremes: The Amount of Exercise to Reduce Cardiovascular Events. Journal of the American College of Cardiology 2016; 67:316-329.

An ACC council perspective paper summarizing the evidence for the optimal exercise dose and intensity for primary and secondary prevention purposes.

** [36] Gebel K, Ding D, Chey T et al. Effect of Moderate to Vigorous Physical Activity on All-Cause Mortality in Middle-aged and Older Australians. *JAMA Intern Med* 2015; 175:970-977.

A large population study providing strong evidence for the superior exercise-induced health benefits of vigorous *versus* moderate intensity exercise.

[37] Physical Activity Council. 2013 participation report. In: www.physicalactivitycouncil.com/PDFs/2013_PAC_Overview_Report_Final.pdf.: 2015.

[38] Kamerow D. Why don't people exercise, even a little? *BMJ (Clinical research ed* 2015; 350:h3024.

[39] Poirier J, Bennett WL, Jerome GJ et al. Effectiveness of an Activity Tracker- and Internet-Based Adaptive Walking Program for Adults: A Randomized Controlled Trial. *J Med Internet Res* 2016; 18:e34.

[40] Patel MS, Asch DA, Volpp KG. Wearable devices as facilitators, not drivers, of health behavior change. *JAMA* 2015; 313:459-460.

** [41] Patel MS, Asch DA, Rosin R et al. Framing Financial Incentives to Increase Physical Activity Among Overweight and Obese Adults: A Randomized, Controlled Trial. *Annals of internal medicine* 2016; 164:385-394.

An original study that compares different approaches to stimulate physically active behavior in a population at risk.

* [42] Zilinski JL, Contursi ME, Isaacs SK et al. Myocardial adaptations to recreational marathon training among middle-aged men. *Circulation. Cardiovascular imaging* 2015; 8:e002487.

This prospective study elegantly demonstrates the effects of high volumes of exercise training on cardiovascular parameters.

* [43] Weiner RB, DeLuca JR, Wang F et al. Exercise-Induced Left Ventricular Remodeling Among Competitive Athletes: A Phasic Phenomenon. *Circulation. Cardiovascular imaging* 2015; 8.

A prospective study demonstrating the acute and chronic effects of exercise training on cardiac remodeling.

[44] La Gerche A, Claessen G. Is exercise good for the right ventricle? Concepts for health and disease. *The Canadian journal of cardiology* 2015; 31:502-508.

* [45] Elliott AD, La Gerche A. The right ventricle following prolonged endurance exercise: are we overlooking the more important side of the heart? A meta-analysis. *British journal of sports medicine* 2015; 49:724-729.

A comprehensive overview about the impact of acute exercise on the structure and function of the right heart.

[46] Middleton N, Shave R, George K et al. Left ventricular function immediately following prolonged exercise: A meta-analysis. Medicine and science in sports and exercise 2006; 38:681-687.

[47] Eijsvogels TM, Hoogerwerf MD, Maessen MF et al. Predictors of cardiac troponin release after a marathon. Journal of science and medicine in sport / Sports Medicine Australia 2015; 18:88-92.

** [48] Gerardin B, Collet JP, Mustafic H et al. Registry on acute cardiovascular events during endurance running races: the prospective RACE Paris registry. European heart journal 2015.

This paper provides insight into the prevalence of adverse cardiac events in marathon athletes.

[49] Redpath CJ, Backx PH. Atrial fibrillation and the athletic heart. Current opinion in cardiology 2015; 30:17-23.

[50] Andersen K, Farahmand B, Ahlbom A et al. Risk of arrhythmias in 52 755 long-distance cross-country skiers: a cohort study. European heart journal 2013; 34:3624-3631.

[51] Sorokin AV, Araujo CG, Zweibel S, Thompson PD. Atrial fibrillation in endurance-trained athletes. British journal of sports medicine 2011; 45:185-188.

* [52] Qureshi WT, Alirhayim Z, Blaha MJ et al. Cardiorespiratory Fitness and Risk of Incident Atrial Fibrillation: Results From the Henry Ford Exercise Testing (FIT) Project. Circulation 2015; 131:1827-1834.

This paper demonstrates that higher cardiorespiratory fitness reduces the risk for atrial fibrillation.

[53] Pathak RK, Elliott A, Middeldorp ME et al. Impact of CARDIOrespiratory FITness on Arrhythmia Recurrence in Obese Individuals With Atrial Fibrillation: The CARDIO-FIT Study. Journal of the American College of Cardiology 2015; 66:985-996.

[54] Thompson PD. Physical Fitness, Physical Activity, Exercise Training, and Atrial Fibrillation: First the Good News, Then the Bad. Journal of the American College of Cardiology 2015; 66:997-999.

* [55] Armstrong ME, Green J, Reeves GK et al. Frequent physical activity may not reduce vascular disease risk as much as moderate activity: large prospective study of women in the United Kingdom. Circulation 2015; 131:721-729.

This paper suggests that daily sessions of physical activity may attenuate the health benefits of exercise.

* [56] Schnohr P, O'Keefe JH, Marott JL et al. Dose of jogging and long-term mortality: the Copenhagen City Heart Study. Journal of the American College of Cardiology 2015; 65:411-419.

This paper suggests that high volumes of exercise may reduce its health benefits.

* [57] Kettunen JA, Kujala UM, Kaprio J et al. All-cause and disease-specific mortality among male, former elite athletes: an average 50-year follow-up. British journal of

sports medicine 2015; 49:893-897.

This paper provides evidence for superior longevity in athletes *versus* controls.

[58] Maessen MF, Hopman MT, Verbeek AL, Eijsvogels TM. Dose of Jogging: Mortality Versus Longevity. Journal of the American College of Cardiology 2015; 65:2672-2673.