



LJMU Research Online

Mohammed, T, Murphy, MF, Lilley, F, Burton, DR and Bezombes, F

The effects of acoustic vibration on fibroblast cell migration

<http://researchonline.ljmu.ac.uk/4037/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Mohammed, T, Murphy, MF, Lilley, F, Burton, DR and Bezombes, F (2016)
The effects of acoustic vibration on fibroblast cell migration. Materials
Science and Engineering C, 69. pp. 1256-1262. ISSN 1873-0191**

LJMU has developed [LJMU Research Online](#) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

The effects of acoustic vibration on fibroblast cell migration

Taybia Mohammed^a, Mark F Murphy^a, Francis Lilley^a, David R Burton^a

^aGeneral Engineering Research Institute , Liverpool John Moores University, Liverpool, L3 3AF, UK.

Abstract

Cells are known to interact and respond to external mechanical cues and recent work has shown that application of mechanical stimulation, delivered via acoustic vibration, can be used to control complex cell behaviours. Fibroblast cells are known to respond to physical cues generated in the extracellular matrix and it is thought that such cues are important regulators of the wound healing process. Many conditions are associated with poor wound healing, so there is need for treatments/interventions, which can help accelerate the wound healing process. The primary aim of this research was to investigate the effects of mechanical stimulation upon the migratory and morphological properties of two different fibroblast cells namely; human lung fibroblast cells (LL24) and subcutaneous areolar/adipose mouse fibroblast cells (L929). Using a speaker-based system, the effects of mechanical stimulation (0-1600Hz for 5 minutes) on the mean cell migration distance (μm) and actin organisation was investigated. The results show that 100Hz acoustic vibration enhanced cell migration for both cell lines whereas acoustic vibration above 100Hz acoustic vibration was found to decrease cell migration in a frequency dependent manner. Mechanical stimulation was also found to promote changes to the morphology of both cell lines, particularly the formation of lamellipodia and filopodia. Overall lamellipodia was the most prominent actin structure displayed by the lung cell (LL24), whereas filopodia was the most prominent actin feature displayed by the fibroblast derived from subcutaneous areolar/adipose tissue. Mechanical stimulation at all the frequencies used here was found not to affect cell viability. These results suggest that low-frequency acoustic vibration may be used as a tool to manipulate the mechano-sensitivity of cells to manipulate cell migration and which may be used to aid wound repair.

Key words: Acoustic vibration; Mechanical stimulation; Fibroblast Cell Migration; Actin reorganisation; wound healing

34

35 **1. Introduction**

36 Cells, by their very nature, have evolved to respond to external physical cues and it is now
37 known that there is a complex interplay between the physical extracellular microenvironment
38 and cellular function [1],[2] and [3]. Cells sense their physical surroundings by converting
39 mechanical forces and distortions into biochemical signals, via the activation of diverse
40 intracellular signalling pathways, through a process known as mechanotransduction [4]. Little
41 is known about mechanotransduction, however work on eukaryotic cells is helping to unravel
42 the complexities of this process. For example, it is known that stretch-sensitive ion channels
43 [5] are key regulators of this process, along with an architectural control of
44 mechanotransduction, through a mechanochemical coupling between the cell surface and
45 nucleus [6]. We also know that changes in a cell's/tissue's ability to respond to forces are
46 associated with certain disease states, including; muscular dystrophies, cardiomyopathies,
47 cancer progression and metastasis [4] and [7].

48

49 Recent work has shown that mechanical stimulation at the whole organism level can have
50 therapeutic benefits, while at a cellular level it can be used to control stem cell differentiation.
51 For example, whole body mechanical stimulation (WBV), which is being increasingly used in
52 a clinical setting, has been shown to have therapeutic benefits. In a study carried out on
53 patients with Multiple Sclerosis, it was found that mechanical stimulation at frequencies
54 between 40-50Hz (delivered through a vibrating platform) in combination with exercise,
55 helped to improve patient muscle strength, functional ability and general wellbeing [8].
56 While, Weinheimer-Haus et al [9], found that low intensity mechanical stimulation (45Hz)
57 significantly increased the wound closure rate in diabetic mice, compared to those mice that
58 were not exposed to mechanical stimulation. Similarly, Wang et al [7] used a dynamic motion
59 platform to apply mechanical stimulations (32-37Hz) to mice and found that expression of
60 the mechanical stimulation-induced protein R-Spondin, which has the capacity to promote
61 bone formation, was enhanced. The authors suggest that some 'vibe proteins' may be
62 candidates for pre-clinical development as anabolic agents for treatments of osteoporosis [7].

63

64 At the cellular level Aryaei and Jayasuriya [10] found that mechanical stimulation can be
65 used to improve osteoblast attachment and proliferation [10]. Kulkarni et al [11], found that
66 mechanical stimulation can inhibit bone reabsorption. Similarly, Wu et al [12] found that

67 mechanical stimulation produced an anabolic effect on bone through the inhibition of
68 osteoclast differentiation and Kim et al [12], found that low-magnitude high-frequency
69 (LMHF) mechanical stimulation enables the osteogenic process of human mesenchymal
70 stromal cells (hMSCs). Such studies highlight the potential of using mechanical stimulation
71 as a novel therapeutic treatment for a range of conditions, particularly those associated with a
72 loss of bone mass. However, knowledge on the effects of mechanical stimulation on other
73 cell types is lacking.

74

75 We were interested to demonstrate whether mechanical stimulation delivered via acoustic
76 stimulation, could be used to enhance cell migration. For this work we used fibroblast cells,
77 as these cells play an important role in wound healing. We hypothesised that mechanical
78 stimulation would influence the migratory properties of fibroblast cells, as these cells have
79 been shown to be mechanosensitive [13] and respond to mechanical signals generated within
80 the extracellular matrix [14]. Results show that mechanical stimulation delivered to fibroblast
81 cells in culture, can enhance, or decrease, cell migration in a frequency-dependent manner.

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101 **2. Materials and Methods**

102

103 2.1. Mechanical stimulation via acoustic vibration

104

105 To mechanically stimulate cells we used a speaker-based system. The speaker-based system
106 was built using a 0.2W super-thin, waterproof Mylar speaker (45mm) and an Arduino
107 microcontroller board for control (Fig. 1). This system enables mechanical stimulation to be
108 applied (via a sinusoidal waveform) to the underside of the cell culture dish to which the cells
109 adhere, at frequencies ranging between 100-1600Hz.

110

111 2.2. Calibration of mechanical stimulation system

112

113 To calibrate the frequency and amplitude generated by the speaker a laser vibrometer
114 (Polytech Ltd.) was used. Briefly, a cell culture dish (35mm) containing cell culture growth
115 medium was rested upon the speaker (40mm diameter). The laser spot from the vibrometer
116 was focused, through a $\times 10$ microscope objective lens, onto the inner bottom surface of the
117 dish (the surface to which the cells would adhere). Next, the speaker was set to vibrate at
118 either 100, 200, 400, 800 or 1600Hz. Vibration frequency (Hz) and amplitude of
119 displacement (μm) were obtained through measurement of the displacement of the laser spot,
120 which is software driven in the Polytech system (see Figure 2 for results).

121

122 2.3. Cell culture

123

124 Human lung fibroblast cells (LL24) and mouse fibroblast cells (L929) were used in this study
125 as fibroblasts are involved in the wound healing process. The two different fibroblast cells
126 were used as they originate from different tissue types with mechanically different properties
127 and functions. The cells were obtained from the European Collection of Animal Cell Cultures
128 (ECACC). The cells were maintained in Dulbecco's Modified Eagles Medium (DMEM)
129 supplemented with 2mM L-glutamine (Invitrogen), 10% foetal bovine serum (FBS) (sigma,
130 UK) and 1% penicillin- streptomycin in T75 cell culture flasks, at 37°C in a humidified
131 atmosphere containing 95% air and 5% CO₂.

132

133

134

135 2.4 Cell Imaging

136

137 All images obtained during this study were from a Zeiss 510 Meta laser scanning microscope,
138 mounted on an Axiovert 200M BP computer-controlled inverted microscope. This
139 microscope is equipped with the following laser lines; blue diode 405nm, Argon ion 458,
140 477, 488 and 514nm, He-Ne 543nm. For cell migration studies, Differential Interference
141 Contrast (DIC) microscopy was carried out using the He-Ne 543nm laser. For imaging
142 filamentous actin the He-Ne 543nm laser was employed.

143

144 2.5. Mechanical stimulation of cells

145

146 Cells were seeded into 35mm cell culture dishes at a density of 2×10^4 cells/cm² and left to
147 attach to the dishes for 24 hours at 37°C in a humidified atmosphere, containing 95% air and
148 5% CO₂. After 24 hours the dish was removed from the incubator and mechanically
149 stimulated for 5 minutes at either 0 (control), 100, 200, 400, 800, or 1600Hz. Next, the dish
150 was immediately placed into the environmental chamber (37°C humidified atmosphere with
151 95% air and 5% CO₂) of the laser scanning microscope and imaged every 5 minutes for 4
152 hours using DIC time-lapse microscopy.

153

154 Each experiment was repeated three times and ImageJ software was used to manually track
155 the migration of single cells (n=30 cells from each frame in the time-lapse sequence) in each
156 population after 4 hours so as to determine mean migration distance (µm).

157

158 2.6. Mechanical stimulation and actin organisation

159

160 Given that cell morphology and the actin cytoskeleton undergo reorganisation prior to and
161 during cell migration, fluorescence microscopy was used to document the effects of
162 mechanical stimulation upon cell morphology and actin organisation. The LL24 and L929
163 cells were seeded into 35mm cell culture dishes at a density of 2×10^4 cells/cm² and left to
164 attach for 24 hours at 37°C in a humidified atmosphere containing 95% air and 5% CO₂.
165 Next, the cells were mechanically stimulated for 5 minutes at 0, 100, 200, 400, 800, or
166 1600Hz and then chemically fixed to preserve their morphology.

167

168 For fixation the cells were washed (×1) in phosphate buffered saline solution (PBS) for 5
169 minutes and then fixed at room temperature in 10% paraformaldehyde for 10 minutes.
170 Following fixation the cells were washed with PBS and permeabilised for 20 minutes at room
171 temperature using 0.5% Triton-X 100. The cells were then washed (x1) with PBS and the
172 actin filaments labelled by staining with Rhodamine-Phalloidin according to the
173 manufacturer's instructions (Cytoskeleton Inc.). After staining, the dishes were thoroughly
174 washed with PBS to eliminate any background staining and the cells were imaged using an
175 excitation wavelength of 543nm. Images were analysed to determine changes to the
176 morphology of cells. In total 200 cells from each sample (across ×10 images from each
177 sample), were counted and the percentage of cells displaying prominent rounding, filopodia
178 and lamellipodia were recorded and the data plotted in graphical form (Fig. 7 & 8).

179

180 2.7. MTT cell viability assay

181

182 To investigate the effects of mechanical stimulation on cell viability the MTT assay was
183 used. This is a colorimetric assay that relates absorbance to viable cell number. The LL24 and
184 L929 cells were seeded into 35mm cell culture dishes at 2×10^4 cells/cm² and left to attach for
185 24 hours at 37°C in a humidified atmosphere, containing 95% air and 5% CO₂. Next, the cells
186 were mechanically stimulated for 5 minutes at 0, 100, 200, 400, 800, or 1600Hz and the MTT
187 assay carried out. Briefly, MTT solution (5mg/ml) was added to each dish and the cells left to
188 incubate at 37°C, 5% CO₂ for 2 hours. Next, 10% dimethyl sulfoxide (DMSO) was added to
189 each dish in order to solubilize the resulting formazan crystals. The formazan solution was
190 then added to wells of a 96-well plate and the absorbance measured at 540nm using a 96-well
191 plate reader. Each experiment was carried out three times to ensure repeatability.

192

193 2.8 Statistical Analysis

194 Statistical analysis to test for significance between the mean of the control and individual
195 treatments was carried out using an unpaired, two-tailed Student's t-test at 95% confidence
196 limit.

197

198


199

200 **3. Results and discussion**

201 3.1. Speaker system calibration

202

203 To assess the effects of mechanical stimulation upon fibroblast cell migration, a system was
204 developed that can deliver low-frequency-low-amplitude acoustic mechanical stimulation, via
205 a sinusoidal waveform, to cells growing in a cell culture dishes (Fig. 1 left). This system
206 allows the frequency of mechanical stimulation to be controlled via an Arduino controller. In
207 order to determine that the vibration frequency was accurate and stable, a laser vibrometer
208 was used to calibrate the system. It was found that below 100Hz and above 1600Hz the
209 system was unstable (in terms of frequency), as laser vibrometry recorded multiple harmonics
210 outside of this frequency range (data not shown). Therefore, frequencies between 100 and
211 1600Hz were used, so as to accurately deliver stable, low-frequency-low-amplitude
212 mechanical stimulations to cells in a continuous manner. Laser vibrometry also recorded
213 amplitude of displacement (μm), which can be seen to decrease as frequency is increased
214 (Fig. 1 right).



215

216 **Fig. 1** Speaker-based system (left) and calibration of frequency versus amplitude of
217 displacement of the inside bottom surface of a 35mm cell culture dish to which the cells
218 adhere (right).

219


220 3.2. The effect of mechanical stimulation on cell migration.

221

222 In order to investigate if mechanical stimulation has any effect on the migration of individual
223 fibroblast cells, time-lapse microscopic imaging and subsequent cell tracking using ImageJ
224 software was carried out. Cells were mechanically stimulated using 100, 200, 400, 800, or
225 1600Hz for 5 minutes and migration was subsequently recorded over a 4-hour period. As can
226 be seen from Figure 2, mechanical stimulation for 5 minutes at 100Hz significantly enhanced

227 (approx. 10% increase) the mean migration distance (μm) of human lung fibroblast (LL24)
228 cells, when compared to the control population ($p < 0.05$). In contrast, mechanical stimulation
229 for 5 minutes for all frequencies above 100Hz resulted in a decrease in the mean cell
230 migration distance (μm), when compared to the control ($p < 0.05$). This decrease can be seen
231 to occur in a steady manner as mechanical stimulation frequency increased (and amplitude
232 decreased) resulting in a decrease in the mean migration distance (μm) of approximately 50%
233 (at 1600Hz), compared to the control population.

234



235


236 **Fig. 2** Total distance travelled (μm) over a 4-hour period for LL24 fibroblast cells ($n=30$)
237 versus frequency of mechanical stimulation (Hz). Error bars represent standard deviation. *
238 denotes Significance ($P < 0.05$).

239

240

241 The results show a similar trend for the LL29 cells, in that mechanical stimulation at 100Hz
242 resulted in a significant increase in the mean cell migration distance (μm) (approximately
243 6%) compared to the control population ($p < 0.05$) (Fig.3). Above 100Hz there was a steady
244 decrease in the mean cell migration distance (μm) up to 400Hz ($p > 0.05$). Stimulation at
245 800Hz resulted in an almost identical mean cell migration distance (μm) compared to the
246 control population, while stimulation at 1600Hz resulted in a lower mean migration distance
247 (μm) compared to the control, although this was not found to be statistically significant
248 ($p > 0.05$).

249



250

251 **Fig. 3** Total distance travelled (µm) over 4 hours for L929 fibroblast cells (n=30) versus
 252 frequency (Hz) following mechanical stimulation. Error bars represent standard deviation. *
 253 denotes Significance (P < 0.05).

254

255 These results show that fibroblast cell migration can be controlled using acoustic vibration
 256 and that migration distance is dependent upon vibration frequency. The trend observed was
 257 repeatable for both cell lines, particularly the increase in mean cell migration distance (µm)
 258 seen at 100Hz. On all occasions a steady decrease in mean cell migration was observed for
 259 the lung cells (LL24), with 1600Hz always resulting in lowest mean cell migration distance
 260 (µm). The results for the L929 cells somewhat mirrored those of the LL24 cells, however the
 261 trend did not always produce a steady decrease in mean cell migration. To our knowledge this
 262 is the first example of investigating the effects of mechanical stimulation delivered via
 263 acoustic vibration on cell migration. The results are somewhat consistent with similar work
 264 reported in the literature. For example, Aryaei and Jayasuriya [10] found that mechanical
 265 stimulation, in the form of shear stress, can be used to enhance osteoblast adhesion and
 266 proliferation. Similarly, Ito et al [15] found that mechanical stimulation (100-1000Hz) of
 267 L929, HeLa and human umbilical vein endothelial cells (HUVECs) had no effect on cell
 268 morphology, or adhesion. In contrast, mechanical stimulation of mouse embryonic fibroblasts
 269 at 1 kHz was found to increase cell adhesion and alter cell morphology. Such work supports
 270 the work presented here, in that cells respond to external physical cues and how they respond
 271 seems to depend on the cell type and nature and properties of the mechanical stimulation.

272 These results are interesting as they indicate that the use of mechanical/acoustic stimulation
273 may have potential therapeutic benefits for example, enhancing the wound healing process.
274 However, more work is required to investigate the underlying mechanisms as well as the
275 migratory response of other cell types, e.g. human dermal fibroblast and gingival fibroblast
276 cells to mechanical stimulation and the effects of mechanical stimulation on collective cell
277 migration. Our preliminary work in this area using wound healing assays indicates that the
278 enhanced migratory rate following 100Hz migration diminishes 2-3 hours post stimulation.
279 Therefore, we aim to investigate the effects of intermittent stimulation on cell migration e.g.
280 hourly exposure to the stimulus.


281

282 3.3 Effects of mechanical stimulation on cell viability

283


284 In order to determine if the vibration-induced decrease in cell migration was due to changes
285 in cell viability, an MTT assay was carried out. The MTT is a colorimetric assay that relates
286 absorbance to the number of viable cells. A reduction/increase in absorbance (relative to the
287 control) would indicate that the mechanical stimulation has had some effect on the viability
288 (or mitochondrial activity) of the cells. It was observed on all occasions that mechanical
289 stimulation had no effect on cell viability of LL24 or L929 cells at any of the frequencies
290 used here. Trypan blue cell viability assays were also carried out and the results support those
291 obtained from the MTT assays (data not shown).

292



293

294 **Fig. 4** MTT assay results showing absorbance versus frequency (Hz) following mechanical
295 stimulation of LL24 cells for 5 minutes. Error bars represent standard deviation.
296
297




298
299 **Fig. 5** MTT assay results showing absorbance versus frequency (Hz) following mechanical
300 stimulation of L929 cells for 5 minutes. Error bars represent standard deviation.
301

302 3.4. Effects of mechanical stimulation on actin organisation

303


304 Cell migration is associated with well characterised changes to the cell. Many of these
305 changes are macro-scale structural and morphological changes and include; the formation of
306 lamellipodia/membrane ruffling, membrane blebs and actin filopodia [16], the latter of which
307 has been shown to stimulate cell migration [17]. Given that fibroblast cells have been shown
308 to be mechanosensitive and that actin remodelling is associated with cell migration, the effect
309 of mechanical stimulation on actin organisation was investigated. Following mechanical
310 stimulation of both LL24 cells and L929 cells at 0-1600Hz, the cells were fixed and stained
311 using Phalloidin, which specifically labels actin filaments (F-actin). Confocal microscopy
312 revealed that mechanical stimulation encouraged actin remodelling in both cell types. In
313 particular, for LL24 cells it was observed that there was an increase in
314 lamellipodia/membrane ruffling (identified by the arrows in Fig. 6) and an increase in stress-

315 fiber formation/density, compared to the control. However, there were no notable differences
316 in the F-actin organisation as frequency increased (Fig. 6).



317
318 **Fig. 6** F-actin organisation in LL24 cells following mechanical stimulation (0Hz to 1600Hz)
319 for 5 minutes. Yellow arrows highlight lamellipodia/ruffling.


320
321 Similarly, mechanical stimulation of L929 cells also resulted in a reorganisation of F-actin,
322 however the response was markedly different when compared to that of the LL24 cells. The
323 control L929 cells generally appeared to be less spread compared to the LL24 cells. However,
324 when mechanically stimulated, at all frequencies, there were distinct morphological changes
325 to the L929 cells that were characterised by an increased level of cell spreading and actin
326 filopodia formation (Fig. 7).



327
 328 **Fig. 7** F-actin organisation in L929 cells following mechanical stimulation (0Hz to 1600Hz)
 329 for 5 minutes Yellow arrows highlight lamellipodia/ruffling.

330
 331 Following actin staining and confocal imaging a more detailed analysis was carried out to try
 332 and quantify changes to cell morphology. Morphological changes are highlighted in Fig.8 and
 333 are grouped into three categories; cells displaying a rounded morphology, cells displaying
 334 prominent filopodia and cells displaying prominent lamellipodia/ruffling.

335



336
 337
 338 **Fig. 8** Confocal microscopy showing (A) rounded shape morphology, (B) cells with filopodia
 339 and (C) cells displaying lamellipodia/membrane ruffling.


340

341 For the analysis a total of 200 cells were counted for each frequency (and control) and the
342 percentage of rounded cells, number of cells with prominent lamellipodia/ruffling and
343 number of cells with prominent filopodia was recorded. Fig. 9 displays the results for LL24
344 cells and Fig. 10 displays the results for L929 cells. As can be seen from Fig. 9, the control
345 population contained cells that displayed a predominantly rounded morphology with many
346 cells in this population also displaying some degree of lamellipodia and filopodia.
347 Mechanical stimulation at all frequencies resulted in changes to LL24 cell morphology. In
348 particular, mechanical stimulation increased the level of cell spreading, with none of the
349 mechanically stimulated cell populations observed to have cells with a rounded morphology.
350 Mechanical stimulation of LL24 cells also resulted in an increase in the percentage of cells
351 displaying prominent lamellipodia/ruffling and filopodia, with lamellipodia/ruffling being the
352 most prominent morphological feature for the LL24 cell. This increase in
353 lamellipodia/ruffling and filopodia formation can be seen to be somewhat dependent upon
354 mechanical stimulation frequency. For example, the percentage of cells displaying membrane
355 ruffling/lamellipodia increased steadily as mechanical stimulation frequency increased up to
356 400Hz and then was observed to decrease at 800 and 1600Hz, respectively. While the
357 percentage of cells displaying actin filopodia was found to increase markedly at 200, 400 and
358 800Hz.

359

360 Similarly, mechanical stimulation of L929 cells was found to enhance cell spreading and
361 increase the percentage of cells displaying lamellipodia/ruffling and filopodia. As can be seen
362 from Fig. 10, the control population of L929 cells contained a high percentage of cells
363 displaying a rounded morphology. However, mechanical stimulation was seen to reduce the
364 percentage of rounded cells in a frequency-dependent manner. Similarly, the percentage of
365 cells displaying lamellipodia/ruffling was observed to increase in a frequency-dependent
366 manner, while the percentage of cells displaying actin filopodia was found to increase
367 markedly above the control population.

368



369


370 **Fig. 9** Image analysis of LL24 cells stimulated for 5 mins showing the percentage of cells
 371 being round, membrane ruffling and filopodia.

372

373

374

375



376

377 **Fig. 10** Image analysis of L929 cells stimulated for 5 mins showing the percentage of cells
 378 being round, membrane ruffling and filopodia.

379 Two alternate forms of actin machinery coexist at the leading edge of most motile cells;
380 lamellipodia (persistent protrusion over a surface) and filopodia (sensory and exploratory
381 functions to steer cells depending on cues from the environment) [18]. Although most cells
382 in culture express both lamellipodia and filopodia, the levels of each structure is thought to be
383 cell-specific. The response of the LL24 and L929 cells to mechanical stimulation was found
384 to differ. Lamellipodia were the more prominent actin structure formed by the LL24 cells,
385 while filopodia were the most prominent actin structure formed by the L929 cells.

386

387 When adhesions to the substrate are weak, contraction of the actin filaments occurs, causing
388 the lamellipodium to bend upwards, resulting in ruffling and transient retraction [19]. Such
389 events are characterized by an extension of the cellular leading edge before retraction, or
390 ruffling, occurs. This type of pattern of migration has been noted in fibroblast cells [20] and
391 was observed here for both cell lines. Given that the formation of lamellipodia is associated
392 with cell migration and that mechanical stimulation of LL24 cells at 400Hz, resulted in the
393 highest percentage of cells displaying lamellipodia, one may expect the mean cell migration
394 distance (μm) to be greatest at 400Hz; this however was not the case as migration was
395 greatest at 100Hz. Similarly, for L929 cells both membrane ruffling and filopodia were found
396 to be greatest following mechanical stimulation at a frequency of 1600Hz, however the least
397 cell migration occurred at 1600Hz and the greatest at 100Hz (which consequently also
398 produced cells having the lowest percentage of lamellepodia). Thus it is clear from these
399 results that actin remodelling occurs due to mechanical stimulation and the extent to this
400 remodelling seems to be dependent, somewhat, on frequency of vibration. This observation
401 agrees with previous work, whereby it was shown that cortical actin remodelling is dependent
402 on the rate of applied stress [21].

403 It is unknown why such structural differences are observed between the different cells, but it
404 should be noted that although both are fibroblast cells, one cell is of human origin and one
405 sourced from a mouse. Also, they originate from distinctly different tissues (i.e. LL24 cells
406 are derived from the lung, while L929 cells are derived from subcutaneous areolar/adipose
407 tissue) and it is known that fibroblast cells, although structurally similar, can differ
408 genetically depending on where they originate within the body. However, although both cells
409 appeared to respond differently to mechanical stimulation, from a structural point of view,
410 their migratory response was similar, particularly at 100Hz.

411

412

413 This work set out to determine the effects of mechanical stimulation on fibroblast cell
414 migration and to our knowledge this is the first time this has been explored. The results show
415 that acoustic vibration can enhance (100Hz) and decrease (200-1600Hz) cell migration
416 without affecting cell viability. At present, the underlying mechanisms behind these results
417 are unknown and clearly more work needs to be done in this area so to decipher such
418 mechanisms and help us understand more about the mechanobiology of cells. However, the
419 work is promising and may open up new avenues of research that focus on exploiting
420 mechanical means for controlling cell behaviour. For example, the development of novel
421 wound care technologies that use mechanical means to accelerate the wound healing process
422 in patients with conditions associated with poor wound healing.

423

424 **4. Conclusion**

425

426 We have shown that mechanical stimulation (100-1600Hz) applied via acoustic vibration, can
427 affect cell fibroblast cell migration in a frequency-dependent manner. In particular,
428 mechanical stimulation at 100Hz significantly increased the mean cell migration distance for
429 both cell types studied here. For the LL24 lung fibroblast cells, mechanical stimulation at
430 frequencies above 100Hz resulted in a steady decrease in the mean cell migration distance
431 without affecting cell viability. A similar response was seen with the L929 cells, however the
432 decrease in migration was less uniform when compared to that of the LL24 lung cells.
433 Mechanical stimulation was also found to affect cell morphology and actin organisation, with
434 acoustic vibration increasing the formation of lamellipodia and filopodia in both cell lines.
435 Acoustic vibration promoted more lamellipodia in the LL24 lung fibroblast cells, while the
436 formation of filopodia was more prominent in the L929 cells.

437 The results obtained are interesting and may lead to developments in novel wound care
438 technologies that exploit mechanical means to stimulate wound repair/regeneration.

439

440 **Conflict of interest statement**

441 There are no conflicts of interest.

442

443

444

445

446

447

448

449 **References**

450 1. Sirio Dupont, L.M., Mariaceleste Aragona, Elena Enzo,Stefano Giulitti,Michelangelo
451 Cordenonsi, Francesca Zanconato, Jimmy Le Digabel, Mattia Forcato, Silvio Bicciato, Nicola
452 Elvassore & Stefano Piccolo., Role of YAP/TAZ in mechanotransduction. *Nature*, 2011. 474:
453 p. 179-183.

454 2. Janmey PA, M.R., Mechanisms of mechanical signaling in development and disease. *J Cell*
455 *Sci.*, 2011. 124(1): p. 9-18.

456 3. Callie Johnson Miller, L.D., The interplay between cell signaling and mechanics in
457 developmental processes. *Nat Rev Genet*, 2013. 14(10): p. 733-744.

458 4. Ingber, D.E., Cellular mechanotransduction: putting all the pieces together again. *FASEB*
459 *J*, 2006. 20(7): p. 811-27.

460 5. Martinac, B., Mechanosensitive ion channels: molecules of mechanotransduction. *J Cell*
461 *Sci.* , 2004. 117: p. 2449-60.

462 6. Wang N, T.J., Ingber DE., Mechanotransduction at a distance: mechanically coupling the
463 extracellular matrix with the nucleus. *Nat Rev Mol Cell Biol.*, 2009. 10(1): p. 75-82.

464 7. Wang H, B.T.A., Russell E, Kim J.H, Egan K.P, Chen Q, Israelite C, Schultz D.C, Johnson
465 F.B. and Pignolo R.J., R-Spondin 1 promotes vibration-induced bone formation in mouse
466 models of osteoporosis. *J Mol Med (Berl)*. 2013. 91(12): p. 1421-9.

467 8. Othmar Schuhfried, C.M., Tatjana Jovanovic, Karin Pieber, and Tatjana Paternostro-
468 Sluga., Effects of whole-body vibration in patients with multiple sclerosis: a pilot study. *Clin*
469 *Rehabil*, 2005. 19(8): p. 834-842.

470 9. Weinheimer-Haus E.M., J.S., Ennis WJ and Koh TJ., Low-Intensity Vibration Improves
471 Angiogenesis and Wound Healing in Diabetic Mice. *PLoS ONE.* , 2014. 9(3).

472 10. Ashkan Aryaeia, A.C.J., The effect of oscillatory mechanical stimulation on osteoblast
473 attachment and proliferation. *Materials Science and Engineering: C*, 2015. Volume 52: p.
474 129–134.

475 11. Kulkarni R.N., V.P.A.a.L.D., Mechanical vibration inhibits osteoclast formation by
476 reducing DC-STAMP receptor expression in osteoclast precursor cells. *Bone.*, 2013. 57(2): p.
477 493-498.

478 12. Wu S.H., Z.Z.M.a.C.J.T., Low-Magnitude High-Frequency Vibration Inhibits RANKL-
479 Induced Osteoclast Differentiation of RAW264.7 Cells. . *Int J Med Sci.* , 2012. 9(9): p. 801-
480 7.

- 481 13. Turner N.A and Potter, K.E., Function and fate of myofibroblasts after myocardial
482 infarction. *Fibrogenesis Tissue Repair* 2013. 6(5).
- 483 14. Matthias Chiquet, L.G., Roman Lutz, Silke Maier, From mechanotransduction to
484 extracellular matrix gene expression in fibroblasts. *Biochimica et Biophysica Acta (BBA) -*
485 *Molecular Cell Research*, 2009. 1793(5): p. 911–920.
- 486 15. Ito Y, K.T., Ago Y, Nam K, Hiraku K, Miyazaki K, Masuzawa T, Kishida A., Nano-
487 vibration effect on cell adhesion and its shape. *Biomed Mater Eng*, 2011. 21(3): p. 149-58.
- 488 16. David J. Barry, C.H.D., Jasmine V. Abella, and Michael Way, Open source software for
489 quantification of cell migration, protrusions, and fluorescence intensities. *JCB*, 2015. 209(1):
490 p. 163-180.
- 491 17. Antti Arjonen, R.K., and Johanna Ivaska, Filopodia and adhesion in cancer cell motility.
492 *Cell Adh Migr*, 2011. 5(5): p. 421-430.
- 493 18. Marisan R. Mejillano¹, et al., Lamellipodial Versus Filopodial Mode of the Actin
494 Nanomachinery: Pivotal Role of the Filament Barbed End. *Cell Mol Bioeng.*, 2004. 118(3):
495 p. 363–373.
- 496 19. Giannone G., D.-T.B., Rossier O., Cai Y., Chaga O., Jiang G., Beaver W., Döbereiner
497 HG., Freund Y., Borisy G., Sheetz MP. , Lamellipodial actin mechanically links myosin
498 activity with adhesion-site formation. *cell*, 2007. 128(3).
- 499 20. Bear JE., S.T., Krause M., Schafer DA., Loureiro JJ., Strasser GA., Maly IV., Chaga OY.,
500 Cooper JA., Borisy GG., Gertler FB., Antagonism between Ena/VASP proteins and actin
501 filament capping regulates fibroblast motility. *Cell*, 2002. 109(4): p. 509-21.
- 502 21. Priyanka Pravincumar, D.L.B., Martin M. Knight, Viscoelastic Cell Mechanics and Actin
503 Remodelling Are Dependent on the Rate of Applied Pressure. 10, 2012. 1371.

504

505