



LJMU Research Online

Holdsworth, DL, Rushton, MT, Bewsher, D, Walter, FM, Eyres, SPS, Hounsell, R and Darnley, MJ

STEREO/HI and optical observations of the classical nova V5583 Sagittarii

<http://researchonline.ljmu.ac.uk/id/eprint/4131/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Holdsworth, DL, Rushton, MT, Bewsher, D, Walter, FM, Eyres, SPS, Hounsell, R and Darnley, MJ (2014) STEREO/HI and optical observations of the classical nova V5583 Sagittarii. MONTHLY NOTICES OF THE ROYAL ASTRONOMICAL SOCIETY. 438 (4). pp. 3483-3489. ISSN 0035-8711

LJMU has developed [LJMU Research Online](http://researchonline.ljmu.ac.uk/) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

STEREO/HI and Optical Observations of the Classical Nova V5583 Sagittarii

Daniel. L. Holdsworth^{1,2}, M. T. Rushton¹, D. Bewsher¹, F. M. Walter³, S. P. S. Eyres¹, R. Hounsell⁴ and M. J. Darnley⁵

¹ *Jeremiah Horrocks Institute, University of Central Lancashire, Preston, PR1 2HE.*

² *Astrophysics Group, Keele University, Staffordshire, ST5 5BG.*

³ *Department of Physics & Astronomy, Stony Brook University, Stony Brook, NY 11794-3800, USA.*

⁴ *Space Science Institute, 3700 San Martin Dr, Baltimore, MD 21218, USA.*

⁵ *Astrophysics Research Institute, Liverpool John Moores University, Liverpool, L3 5RF.*

Version of September 2013

ABSTRACT

The classical nova V5583 Sgr (Nova Sagittarii 2009 N^o 3) has been observed during the rise phase and shortly after by NASA's *STEREO*/HI instruments, with later optical spectroscopy obtained with the R-C Spectrograph at CTIO, Chile. The time of peak in the *STEREO* passband has been constrained to within 4 hours, as a result of the high cadence data obtained by *STEREO*/HI. The optical spectra show the nova evolving from the permitted to the nebular phases. The neon abundance in the ejecta is $[\text{Ne}/\text{O}] \gtrsim +1.0$, which suggests that V5583 Sgr was most likely a neon nova.

Key words: stars: novae, cataclysmic variables

1 INTRODUCTION

Classical nova (CN) eruptions occur in close binaries containing a white dwarf (WD) primary and a low mass star (see Bode & Evans 2008, for recent reviews). In these systems, the secondary fills its Roche lobe and transfers hydrogen-rich material to the WD via an accretion disk. The outbursts occur when the mass accreted on the white dwarf reaches a critical value. Novae generate $\sim 10^{37} - 10^{39}$ J and eject $\sim 10^{-4} M_{\odot}$ into the Interstellar Medium (see Gehrz et al. 1998 for a review).

The Galactic nova rate has been estimated to be 35 ± 11 yr⁻¹ (Shafter 1997), and 34^{+15}_{-12} yr⁻¹ (Darnley et al. 2006). However, < 10 yr⁻¹ are well observed, because of our position within the galaxy and the interstellar extinction in the Galactic disc (Shafter 1997). It is thought that a further ~ 5 nova a year are missed because of the close proximity of the Sun on the sky (Hounsell et al. 2010).

Novae are usually discovered by amateurs and reported to the American Association of Variable Star Observers (AAVSO) or Variable Star NETwork (VSNET) databases. This means that novae tend to be discovered when they are brightest. The rise phase is rarely well observed, although it is known that at least some novae pause about 1-2 magnitudes below the peak, a phenomenon known as the pre-maximum halt (PMH; Warner 2008).

Hounsell et al. (2010) presented light curves of four nova

eruptions from an unlikely source: the Solar Mass Ejection Imager (SMEI) on board the *Coriolis* satellite (Eyles et al. 2003), an instrument designed to observe coronal mass ejections (CMEs). Because of the continuous coverage of the SMEI data, they detected the novae before the peak, and tracked them into the decline. Due to the high cadence of the observations (102 minutes), they were able to construct nova light curves at unprecedented temporal resolution. PMHs were detected in three of the novae.

The Heliospheric Imagers (HI; Eyles et al. 2009) on board the Solar TERrestrial RELations Observatory (*STEREO*; Kaiser et al. 2008) also have the potential to detect nova outbursts. They monitor the heliosphere for CMEs, observing the background stars. This data set, spanning nearly seven years, has recently been exploited for stellar work, by e.g. Wraight et al. (2011), Wraight et al. (2012).

V5583 Sagittarius (Nova Sagittarii 2009 N^o 3) was discovered in eruption by Nishiyama et al. (2009) on 2009 August 6.494 UT at a magnitude of $V = 7.7$. Optical spectroscopy showed broad Balmer and Fe II emission lines as expected for a classical nova (Maehara & Fujii 2009). This interpretation was supported by the observations of Hachisu et al. (2009), who found that the multi-band light curves were almost identical to those of the classical nova V382 Vel. In this paper, we present the light curve for V5583 Sgr, the first nova detected in the HI instruments.




Figure 1. HI magnitude light curves of V5583 Sgr over the first 5 years of HI-2A data. The outburst can clearly be seen in the 3rd epoch of data.

2 PHOTOMETRIC OBSERVATIONS

2.1 STEREO/HI light curve

The HI observations used in this paper were obtained by the Heliospheric Imagers (HI-2) on the STEREO-Ahead (STEREO-A) spacecraft.

The HI-2 images have a field of view (FOV) of $\sim 70^\circ \times 70^\circ$ with a pixel size of ~ 4 arcmin pixel $^{-1}$, centred 53° along the ecliptic plane from Sun centre. The HI-2 cameras have a broad spectral response ($\sim 4000 - 10000\text{\AA}$) with the aim of detecting as much white light from faint CMEs as possible (Eyles et al. 2009). Ninety-nine exposures of 50 seconds each, make up each image, with an image cadence of 120 minutes. The solar F-corona is the dominant large scale structure in the images, which is removed using a running daily background.

V5583 Sgr entered the FOV of the HI-2A camera on 2009 July 29 UT (8 days before its discovery), and left the FOV on 2009 October 5 UT. As this is the first stellar observation using data from HI-2A, we give a short overview of the data analysis.

The HI-2 instruments have baffles which prevent stray-light from entering the optical system, but result in a re-

duced portion of the FOV being used for science purposes. Therefore, in this work, we exclude data in the baffled regions.

At the time of writing, a post-launch calibrated largescale flatfield is unavailable for the HI-2A instrument. However, two stars located on the same part of the CCD as the nova have been used to estimate the large scale flatfield using the same methodology that Bewsher et al. (2010) used for the HI-1 instruments. This correction for the flatfield has been applied to the data.

In each HI-2A image, the position of the nova is determined using the pointing and optical calibration of Brown et al. (2009). An aperture with radius of 2.18 pixels is used to calculate the intensity of the nova. This is repeated on every HI-2A image which contains the nova location over a 5 year period, which corresponds to crossing the CCD five times. The nova outburst is observed during the third of the five crossings of the CCD. The aperture size was chosen so that the aperture was large enough to render any PSF variation negligible, yet avoid any contamination from nearby sources. Catalogues show a number of other sources within the aperture, but all are faint ($V > 10$) and below the sensitivity limit of the instrument. Because the number of counts




Figure 2. HI magnitude light curve of V5583 Sgr. A close up view of the rise to the time of maximum. The uncertainty is ~ 0.06 at magnitudes above 10th magnitude.

observed when the nova is not in outburst are very small, a background is not removed as part of the aperture photometry. Data from the first, second, fourth and fifth CCD crossings of data are used to estimate the background, which is then fitted with a polynomial, and subtracted from all five years of data. The large scale flatfield corrected and background subtracted light curves over the five year period can be seen in Figure 1.

The determination of the photometric response of the HI-2A instrument follows the method of Bewsher et al. (2010) for the HI-1 instruments. The HI magnitude, m_{HI} , scale is given by

$$m_{\text{HI}} = -2.5 \log \frac{I}{F_0},$$

where I is the intensity of the star

and F_0 is the intensity of Vega derived from spectral folding (see Bewsher et al. 2010 for a detailed discussion) and has a value of $71655.1 \text{ DN pixel}^{-1}$.

The optical maxima of novae are usually either missed or poorly constrained in time because of their unexpected nature. However, the 120 minute cadence of the HI-2A camera allows us to place a tight constraint on the time of maximum of V5583 Sgr. The time of maximum (t_0) determined from the HI data is $\text{JD } 2455050.54 \pm 0.17$, which translates to 00:09 UT on 2009 August 7th, with an error of ± 4 hours due to the flatness of the peak. Due to the limiting magnitude of the detector, the rise phase is not seen from the quiescent magnitude ($V > 13.3$, Nishiyama et al. 2009). However, *STEREO/HI* did detect the nova 1 – 2 days before the optical maximum. We note that the progenitor is not present in the UNSO-B1 catalogue, giving an outburst amplitude of $\Delta B \geq 13$, typical of a fast classical nova (Bode & Evans 2008).

Figure 2 shows in detail the rise phase of the outburst. It shows a relatively smooth rise, although several data points deviate from the rising curve. The PMHs identified by Hounsell et al. (2010) develop 0.49 – 2.19 days before maximum, have a duration of 0.14 – 0.42 days, and occur 0.63 – 1.74 m_{SMEI} below maximum, where m_{SMEI} is the unfiltered apparent magnitude derived from the SMEI images. The light curve in Figure 2 shows similar features at 1.2 days before peak, 1.2 m_{HI} below the maximum, 0.9 days before




Figure 3. Light curve generated from the AAVSO data (crosses) and VSNET data (open triangles). Additional data points from the IAU circulars are plotted in open diamonds. The vertical lines show the times of the spectroscopic observations discussed in §3.1.

peak, 0.9 m_{HI} below maximum, and 0.8 days before peak, 0.7 m_{HI} below. These are consistent with PMH seen in other novae. Furthermore, Hillman et al. (2013) have recently observed similar features in their models of nova light curves. They attribute the PMH to the recession of the convection zone and a decline in the opacity of the expanding envelope. Their models show considerable variation in the shape and amplitude of the PMH and some show more than one halt.

The eruption of V5583 Sgr was also detected by the SMEI instrument, but the light curve is noisy because the nova peaked at $m_{\text{SMEI}} = 6.49 \pm 0.05$, just one magnitude above limiting value; as a result only a small portion of the rise phase is seen. The time of maximum seen in the SMEI data is 2455050.6 (2009 Aug 7.08 UT).

2.2 AAVSO and VSNET Light Curve

The IAU 9061 observations of V5583 Sgr constrain the time of optical maximum to within approximately 7.6 days, between 29th July and 6th August (Nishiyama et al. 2009). *STEREO/HI* provides a much more precise time, to within just a few hours. The light curve produced from the AAVSO & VSNET data is shown in Figure 3. The first measurement was recorded at JD 2455051.17, which translates to 1603 UT on 7th August 2009, ~ 22 hours after the maximum detected in HI.

The initial decline of the nova, which can be seen in both light curves, is smooth and fast. The AAVSO & VSNET photometry shows the late decline phase, which is also smooth, implying that there was no dust formation in the ejecta. The behaviour of V5583 Sgr is consistent with a very fast nova (Payne-Gaposchkin 1957), with an A type light curve (Duerbeck 1981). We determine the times to decline by 2 (t_2) and 3 (t_3) magnitudes from maximum of $t_2 = 4.97 \pm 0.32$ days and $t_3 = 9.52 \pm 0.32$ days from the data in Figure 3; the errors reflect the scatter in the AAVSO & VSNET light curve (~ 0.2 mag) - this will dominate the uncertainty, being more significant than the fading in the visual that occurred in the ~ 22 hr following the t_0 determined from the *STEREO* data.

Using the MMRD relationship (Downes & Duerbeck 2000), we estimate a magnitude at maximum of -9.55 ± 0.50 from t_2 and -9.70 ± 1.14 from t_3 . The interstellar extinc-

tion has been estimated to be $A_V = 1.0$ ($E(B - V) = 0.33$; Hachisu et al. 2009), which is consistent with the dust maps of Schlegel et al. (1998) along the line of sight. These values give an estimate of the distance to the nova of 14.7 ± 3.4 kpc.

3 OPTICAL SPECTRA

3.1 Overview of the spectra

Optical spectra of V5583 Sgr were obtained with the RC Spectrograph, which is mounted on the 1.5m telescope at the Cerro Tololo Inter-American Observatory, Chile. Table 1 logs relevant information on each spectra, see Walter et al. (2012).

Figures 4 & 5 show the optical spectra of V5583 Sgr on given dates. The spectral lines were identified using an online Atomic Line List (<http://www.pa.uky.edu/~peter/atomic/>) and published spectra of fast novae (e.g. Williams et al. 1991, Augusto & Diaz 2003 and Iijima & Cassatella 2010). The times of the spectroscopic observations with respect to the visual light curve of V5583 Sgr are shown in Figure 3. Table 2 shows the line fluxes on selected dates of observation. The dereddened values have been derived using the extinction law of Seaton (1979) and a ratio of total-to-selective extinction of $R = 3.1$. The smooth light curves are indicative of a dust poor nova, with no significant circumstellar extinction.

The spectra are dominated by H I, He I, O I, [O III], and [Ne III] emission lines. The lack of Fe lines suggests that V5583 Sgr was most likely a ‘He/N’ type of classical nova. The later spectra are dominated by the [Ne III]₃₈₆₉ line, which is second in strength only to H α , consistent with a transition from the P o phase to the N $_{ne}$ phase in the Tololo classification scheme of nova spectra. The intermediate auroral phase is not observed in our data, although this phase is by-passed by novae with a ONeMg white dwarf (Williams et al. 1991). In these novae, the ejecta is enriched in neon, due to mixing of accreted mater with the white dwarf core material (Starrfield et al. 1985). The strong [Ne III] lines in the spectra of V5583 Sgr suggest that there may be an enhancement of this species.

The line profiles are flat or saddled-shaped and jagged, as is usually the case in ‘He/N’ novae (Williams 1992). The full width at zero intensity (FWZI) of the lines is 5000 km s^{-1} or greater, consistent with the speed classification derived in §2.2 (Williams 1992). The line profiles vary between species and ionisation states. The [Ne III], [N II] and [O III] lines show peaks at ~ -1150 , -300 , 450 , and 1220 km s^{-1} ; in most cases, the blue peak is stronger than the red peak. The He I, H I and O I also show these velocities, but the red peak is stronger than the blue peak. These line profiles are often seen in novae and are attributed to inhomogeneities and clumpiness in the ejecta (Gill & O’Brien 1999).

3.2 The neon abundance

To determine the abundance of neon in the ejecta of V5583 Sgr, we consider a line ratio from the species [Ne III] and [O III], which have similar ionisation potentials (41 eV

and 35 eV respectively). If the density is $n_e > 10^5 \text{ cm}^{-3}$, the line flux (F_{ij}) is given by the formula:

$$F_{ij} = N_i E_{ij} A_{ij} \frac{g_j}{g_i} \exp \frac{-E_{ij}}{kT_e},$$

where N_i is the number density of level i , E_{ij} is the energy of the transition, A_{ij} is the transition probability, g is the statistical weight, and T_e is the electron temperature. In principal, T_e can be determined from the flux ratios [O III] (4959 + 5007)/4363 and [N II] (6548 + 6584)/5755 (Osterbrock & Ferland 2006), but these are severely effected by blending with H I lines in the data. However, in Figure 6 we show the [Ne/O] abundance in V5583 Sgr for a range of electron temperatures from which we can make a meaningful constraint. We note that the H β /H γ line ratio is lower than expected, and that the [O III] 4363 line must contribute significantly to the H γ flux; this shows that collisional effects are affecting the population of the [O III] levels, which suggests $n_e > 10^5 \text{ cm}^{-3}$ (Osterbrock & Ferland 2006). Furthermore, we extract the [O III] 4363 Å line flux from the blend with H γ using the H β profile. This gives the flux ratio [O III] (4959 + 5007)/4363 = 2.92. The density is about $1.5 \times 10^7 \text{ cm}^{-3}$ at $T_e = 10,000 \text{ K}$, $2 \times 10^6 \text{ cm}^{-3}$ at $50,000 \text{ K}$ and $4 \times 10^6 \text{ cm}^{-3}$ at 10^6 K .

We consider the data on days 62, 71 and 90, in which the relevant lines are relatively free of contamination. Although [O III] 5007 forms a blend with [O III] 4959, we can extract its flux using the transition probabilities, since these lines share the same upper level; this gives a ratio of 5007/4959 = 2.9. In deriving the [Ne/O] abundance, we average the results from each dataset, which give similar values. We give the results with respect to solar, assuming the abundances in Asplund et al. (2009).

Figure 6 shows that we would find lower [Ne/O] abundances if we assume higher assumed electron temperatures, with the abundance asymptotically approaching a value of about [Ne/O] $\gtrsim +1.0$ for $T_e > 20 \times 10^4 \text{ K}$. We can therefore conclude that [Ne/O] is well in excess of the solar value in the ejecta of V5583 Sgr. Such a high abundance is usually attributed to a nova occurring on a ONeMg WD, in which some of the WD material has been dredged-up and ejected in the eruption, while novae occurring on CO WDs have [Ne/O] $< +0.11$ (Gehrz 2008). It is worth noting that [Ne/O] $> +1.0$ places V5583 Sgr among the extreme neon novae, along with V693 CrA and V1370 Aql (Andrea et al. 1994).

4 DISCUSSION

In this paper we have presented a light curve of the very fast classical nova V5583 Sgr, obtained by the HI-2A camera on board the *STEREO* spacecraft. The extracted data show a clear detection of the nova outburst, and include the rarely seen phase before maximum.

The cadence of the HI-2 instrument is 120 minutes, giving a tight constraint on the time of the maximum light. The derived t_0 is JD 2455050.54 \pm 0.17, two days earlier than that estimated by Hachisu et al. (2009), who assumed that V5583 Sgr was identical to another nova, V382 Vel. In the HI light curve of V5583 Sgr, a PMH may have occurred one day before maximum (as shown in Figure 2), similar to

Table 1. Log of spectra information. Classes shown in brackets are alternate classifications, see text for details.

| Date | JD (- 2455000) | Time since t_0 (Days) | Slit (arcsec) | Range (Å) | Resolution (Å) | Tololo Class |
|------------------|-------------------|----------------------------|------------------|--------------|-------------------|-----------------------|
| 07/09/2009T03:51 | 081 | 31 | 0.8 | 5630-6950 | 3.1 | P_{he} (P_o^o) |
| 16/09/2009T03:27 | 090 | 40 | 1.0 | 6000-9500 | 8.6 | P_o^o |
| 17/09/2009T23:40 | 091 | 41 | 1.0 | 3650-5450 | 4.1 | N_{ne} (P_o^o) |
| 08/10/2009T01:48 | 112 | 62 | 1.0 | 3650-5450 | 4.1 | N_{ne} |
| 17/10/2009T00:58 | 121 | 71 | 1.0 | 3650-5450 | 4.1 | N_{ne} |
| 19/10/2009T01:11 | 123 | 73 | 0.8 | 5630-6950 | 3.1 | P_{he} (N_{ne}) |
| 05/11/2009T00:23 | 140 | 90 | 1.0 | 3650-5450 | 4.1 | N_{ne} |




Figure 4. Red spectra of V5583 Sgr on 7th September 2009, $t_0 + 31$. The dotted line in the lower figure shows the profile of $H\alpha$ at a scale of one quarter.

those of RS Oph and KT Eri in the SMEI data (Hounsell et al. 2010).

The detection of PMHs is rare due to the unpredictable nature of novae. However, it is possible that other novae are present in the HI archive data. Furthermore, any nova detected with the HI-1 instruments, with their smaller pixel




Figure 5. Blue spectra of V5583 Sgr on 8th October 2009, $t_0 + 62$.

size and higher cadence, will be shown in unprecedented detail.

The optical spectra of V5583 Sgr show an emission line spectrum typical of a fast nova in the decline stage. Most of the spectra have been classified as N_{ne} in the Tololo system, with the remaining being P_o^o . None are consistent with the coronal or auroral phase, which suggests that V5583 Sgr is a neon nova (Williams 1992). The profiles of the spectral lines,

Table 2. Table of line fluxes on selected dates. Notes: 1) \diamond indicates a line with an absorption component.

| Line | Time since t_0 (Days) | JD (- 2455000) | Flux $\times 10^{-15}$ (W m $^{-2}$) | Extinction corrected (Flux) $\times 10^{-15}$ (W m $^{-2}$) | FWZI ± 500 (km s $^{-1}$) |
|--------------|----------------------------|-------------------|---|--|--------------------------------------|
| H δ | 62 | 112 | 2.43 \pm 0.13 | 8.89 \pm 2.05 | 7000 |
| H γ | 62 | 112 | 4.70 \pm 0.12 | 16.6 \pm 3.8 | 11000 |
| H β | 62 | 112 | 4.10 \pm 0.21 | 12.9 \pm 3.0 | 6000 |
| H α | 40 | 90 | 114 \pm 12 | 240 \pm 55 | 6000 |
| H 18865 | 40 | 90 | 1.40 \pm 0.12 | 2.18 \pm 0.50 | 5000 |
| H 19017 | 40 | 90 | 1.89 \pm 0.28 | 2.93 \pm 0.67 | 4700 |
| H 19231 | 40 | 90 | 5.20 \pm 0.39 | 8.06 \pm 1.85 | 8300 |
| He 14471 | 62 | 112 | 0.93 \pm 0.12 | 3.11 \pm 0.72 | 11200 |
| He 15876 | 73 \diamond | 123 | 3.35 \pm 0.27 | 8.04 \pm 1.85 | 5600 |
| He 16679 | 40 | 90 | 2.74 \pm 0.06 | 5.74 \pm 1.32 | 6600 |
| He 17065 | 40 | 90 | 6.67 \pm 0.19 | 13.2 \pm 3.0 | 6800 |
| C III197 | 40 | 90 | 0.71 \pm 0.02 | 1.24 \pm 0.29 | 4800 |
| N III4582 | 62 | 112 | 1.66 \pm 0.20 | 5.54 \pm 1.28 | 16500 |
| N II5681 | 73 | 123 | 0.51 \pm 0.05 | 1.32 \pm 0.30 | 5818 |
| [N II]5755 | 73 | 123 | 0.85 \pm 0.422 | 2.04 \pm 0.47 | 5400 |
| [O III]4959 | 62 | 112 | 1.22 \pm 0.09 | 3.61 \pm 0.83 | 5100 |
| [O III]5007 | 62 | 112 | 5.48 \pm 0.33 | 16.2 \pm 3.7 | 7300 |
| [O III]7325 | 40 | 90 | 6.27 \pm 1.00 | 12.4 \pm 2.9 | 7100 |
| O 17776 | 40 | 90 | 2.03 \pm 0.08 | 3.78 \pm 0.87 | 5100 |
| O 18221 | 40 | 90 | 19.5 \pm 0.7 | 34.2 \pm 7.9 | 4600 |
| O 18446 | 40 | 90 | 29.8 \pm 4.3 | 52.1 \pm 12.0 | 4000 |
| [Ne III]3869 | 62 | 112 | 8.22 \pm 0.34 | 32.5 \pm 7.5 | 8600 |
| [Ne III]3968 | 62 | 112 | 3.26 \pm 0.23 | 12.4 \pm 2.9 | 5200 |
| Ne II6168 | 40 | 90 | 0.73 \pm 0.02 | 1.65 \pm 0.38 | 10900 |
| Fe II5169 | 62 | 112 | 0.20 \pm 0.03 | 0.548 \pm 0.126 | 7000 |

and the FWZIs are also consistent with this interpretation, as is our conclusion that the object is a ‘He/N’ type nova (Williams 1992).

We have derived a neon abundance in the ejecta of [Ne/O] \gtrsim +1.0, which shows that V5583 Sgr was likely to be a neon nova, and that material has been dredged-up from the white dwarf and ejected in the explosion. We can therefore conclude that the primary is a massive ($M_{WD} \gtrsim 1.2M_{\odot}$)

ONeMg white dwarf (Gehrz 2008), which is also consistent with the speed classification.

V382 Vel, which was discovered in outburst in 1999, is, in many respects, similar to V5583 Sgr. It, too, was a very fast, neon-type nova, with a neon abundance of [Ne/O] = +0.66 and t_2 of 4.5 days, and showed little or no evidence of dust formation in the ejecta (Della Valle et al. 2002). Early spectroscopic observations (t_0+5) showed many Fe II lines in its spectra (Della Valle et al. 2002) which




Figure 6. The $[\text{Ne}/\text{O}]$ abundance in V5583 Sgr as a function of assumed electron temperature.

disappeared about one month later. The spectra following this epoch, when the nova was in the nebular phase, are almost identical to those presented here for V5583 Sgr over the same evolutionary period. It is possible that earlier spectra of V5583 Sgr would have also shown Fe II lines, similar to those seen in V382 Vel. The latest observations of V382 Vel, ~ 184 days after maximum, showed highly ionised Fe lines, which are not present in the V5583 Sgr data. From these spectra, Della Valle et al. (2002) classify the object as an ‘Fe IIb’ type nova. On the basis of the lines present in our spectra of V5583 Sgr, we classified the object as a ‘He/N’ type nova; however, given the similarity to V382 Vel, it is possible that V5583 Sgr is a ‘Fe IIb’ nova, and would have developed Fe lines at a later stage.

5 CONCLUSIONS

In this paper we have presented *STEREO/HI* observations of V5583 Sgr, which tightly constrains the time of maximum light at JD 2455050.54 \pm 0.17. The t_2 of the nova is 4.97 \pm 0.32 days, making V5583 Sgr a very fast nova.

Optical spectroscopy of V5583 Sgr show the nova evolving from the permitted phase to the nebula phase. The neon abundance in the ejecta is $[\text{Ne}/\text{O}] \gtrsim +1.0$, which makes V5583 Sgr one of the most neon enriched novae known.

ACKNOWLEDGEMENTS

We acknowledge with thanks the variable star observations from the AAVSO and VSNET international databases contributed by observers worldwide and used in this research. The Heliospheric Imager (HI) instrument was developed by a collaboration that included the Rutherford Appleton Laboratory and the University of Birmingham, both in the United Kingdom, and the Centre Spatial de Liège (CSL), Belgium, and the US Naval Research Laboratory (NRL), Washington DC, USA. The STEREO/SECCHI project is an international consortium of the Naval Research Laboratory (USA), Lockheed Martin Solar and Astrophysics Lab (USA), NASA Goddard Space Flight Center (USA), Rutherford Appleton Laboratory (UK), University of Birmingham (UK), Max-Planck-Institut für Sonnen- system-

forschung (Germany), Centre Spatial de Liège (Belgium), Institut d’Optique Théorique et Appliquée (France), and Institut d’Astrophysique Spatiale (France). This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France. The authors thank the anonymous reviewer for useful comments.

REFERENCES

- Andrea J., Drechsel H., Starrfield S., 1994, *A&A*, 291, 869
 Asplund M., Grevesse N., Sauval A. J., Scott P., 2009, *ARA&A*, 47, 481
 Augusto A., Diaz M. P., 2003, *AJ*, 125, 3349
 Bewsher D., Brown D. S., Eyles C. J., Kellett B. J., White G. J., Swinyard B., 2010, *Sol Phys*, 264, 433
 Bode M. F., Evans A., 2008, *Classical Novae*, 2 edn. Cambridge University Press
 Brown D. S., Bewsher D., Eyles C. J., 2009, *Solar Physics*, 254, 185
 Darnley M. J., Bode M. F., Kerins E., *et. al.* 2006, *MNRAS*, 369, 257
 Della Valle M., Pasquini L., Daou D., Williams R. E., 2002, *A&A*, 390, 155
 Downes R. A., Duerbeck H. W., 2000, *AJ*, 120, 2007
 Duerbeck H. W., 1981, *PASP*, 93, 165
 Eyles C. J., Harrison R. A., Davis C. J., Waltham N. R., Shaughnessy B. M., Mapson-Menard H. C. A., Bewsher D., Crothers S. R., Davies J. A., Simnett G. M., Howard R. A., Moses J. D., Newmark J. S., Socker D. G., Halain J., Defise J., Mazy E., Rochus P., 2009, *Sol Phys*, 254, 387
 Eyles C. J., Simnett G. M., Cooke M. P., Jackson B. V., Buffington A., Hick P. P., Waltham N. R., King J. M., Anderson P. A., Holladay P. E., 2003, *Sol Phys*, 217, 319
 Gehrz R. D., 2008, in Bode, M. F. and Evans, A. ed., *Classical Novae Infrared studies of classical novae*. Cambridge University Press, pp 167–193
 Gehrz R. D., Truran J. W., Williams R. E., Starrfield S., 1998, *PASP*, 110, 3
 Gill C. D., O’Brien T. J., 1999, *MNRAS*, 307, 677
 Hachisu I., Kato M., Kiyota S., Maehara H., 2009, *ArXiv e-prints*
 Hillman Y., Prialnik D., Kovetz A., Shara M. M., Neill J. D., 2013, *MNRAS*, Advance Access published November 19, 2013, doi: 10.1093/mnras/stt2027
 Hounsell R., Bode M. F., Hick P. P., Buffington A., Jackson B. V., Clover J. M., Shafter A. W., Darnley M. J., Mawson N. R., Steele I. A., Evans A., Eyres S. P. S., O’Brien T. J., 2010, *ApJ*, 724, 480
 Iijima T., Cassatella A., 2010, *A&A*, 516, A54
 Kaiser M. L., Kucera T. A., Davila J. M., St. Cyr O. C., Guhathakurta M., Christian E., 2008, *Space Science Reviews*, 136, 5
 Maehara H., Fujii M., 2009, *Central Bureau Electronic Telegrams*, 1900, 2
 Nishiyama K., Kabashima F., Pojmanski G., Szczygiel D., Pilecki B., Maehara H., Sakane Y., Yamaoka H., Elenin L., Kiyota S., Fujii M., Arai A., Komatsu T., Yamanaka M., Sasada M., Itoh R., 2009, *IAUCirc*, 9061, 1
 Osterbrock D. E., Ferland G. J., 2006, *Astrophysics of Gaseous Nebulae and Active Galactic Nuclei*, 4 edn. University Science Books

- Payne-Gaposchkin C., 1957, Book Reviews: The Galactic Novae. Amsterdam: North Holland Publishing Co.
- Schlegel D. J., Finkbeiner D. P., Davis M., 1998, ApJ, 500, 525
- Seaton M. J., 1979, MNRAS, 187, 73P
- Shafter A. W., 1997, ApJ, 487, 226
- Starrfield S., Sparks W. M., Truran J. W., 1985, ApJ, 291, 136
- Walter F. M., Battisti A., Towers S. E., Bond H. E., Stringfellow G. S., 2012, PASP, 124, 1057
- Warner B., 2008, in Bode, M. F. and Evans, A. ed., Classical Novae *Properties of novae: an overview*. Cambridge University Press, pp 16–33
- Williams R. E., 1992, AJ, 104, 725
- Williams R. E., Hamuy M., Phillips M. M., Heathcote S. R., Wells L., Navarrete M., 1991, ApJ, 376, 721
- Wraight K. T., Bewsher D., White G. J., Nowotny W., Norton A. J., Paladini C., 2012, MNRAS, 426, 816
- Wraight K. T., White G. J., Bewsher D., Norton A. J., 2011, MNRAS, 416, 2477