



LJMU Research Online

Broughton, LJ, Giuntini, F, Savoie, H, Bryden, F, Boyle, RW, Maraveyas, A and Madden, LA

Duramycin-porphyrin conjugates for targeting of tumour cells using photodynamic therapy

<http://researchonline.ljmu.ac.uk/id/eprint/5064/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Broughton, LJ, Giuntini, F, Savoie, H, Bryden, F, Boyle, RW, Maraveyas, A and Madden, LA (2016) Duramycin-porphyrin conjugates for targeting of tumour cells using photodynamic therapy. JOURNAL OF PHOTOCHEMISTRY AND PHOTOBIOLOGY B-BIOLOGY. 163. pp. 374-384.

LJMU has developed [LJMU Research Online](#) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

Duramycin-porphyrin conjugates for targeting of tumour cells using photodynamic therapy

Laura J. Broughton,^a Francesca Giuntini^b, Huguette Savoie^c, Francesca Bryden^c, Ross W. Boyle^c, Anthony Maraveyas^a, Leigh A. Madden^d

^a Hull York Medical School, University of Hull, Hull HU6 7RX, UK

^b School of Pharmacy and Biomolecular Sciences, Liverpool John Moores University, Liverpool L3 2AJ, UK

^c Department of Chemistry, University of Hull, Hull, HU6 7RX, UK

^d School of Life Sciences, University of Hull, Hull, HU6 7RX, UK

Duramycin, through binding with phosphatidylethanolamine (PE), has been shown to be a selective molecular probe for the targeting and imaging of cancer cells. Photodynamic therapy aims to bring about specific cytotoxic damage to tumours through delivery of a photosensitising agent and light irradiation. Conjugation to biological molecules that specifically target cancer has been shown to increase photosensitiser (PS) selectivity and decrease damage to surrounding normal tissue. The aim of this study was to target tumour cells with a PE-specific PS therefore duramycin was conjugated to a porphyrin based PS which was achieved via direct reaction with the ϵ -amino group on the lysine residue near duramycin's N-terminal. The compound was subsequently purified using RP-HPLC and confirmed using mass spectrometry. Binding of the conjugate to ovarian and pancreatic cancer cell lines was assessed by flow cytometry. Light irradiation with a light fluence of 7.5 J/cm² was delivered to conjugate treated cancer cells and cell proliferation analysed by MTT assay. The conjugate detected PE on all 4 cancer cell lines in a concentration dependent manner and conjugate plus irradiation effectively reduced cell proliferation at concentrations ≥ 0.5 μ M, dependent on cancer cell line. Reduction in cell proliferation by the irradiated conjugate was enhanced over unconjugated duramycin in A2780, AsPC-1 and SK-OV-3 ($p < 0.05$). In this study we have shown that a duramycin-porphyrin conjugate retained good binding affinity for its target and, following irradiation, reduced cell proliferation of pancreatic and ovarian cancer cell lines.

1. Introduction

Duramycin is a small tetracyclic peptide produced by the bacterium *Streptovorticillium cinnamoneus* [1]. Only 19 amino acids in length duramycin is one of very few identified, relatively, short peptides to have a defined 3-dimensional structure [2]. This structure is stable due to the presence of 3 thioether bridges formed via the unusual amino acids lanthionine and methyllanthionine. Further a covalent linkage between the cysteine at duramycin's N-terminal and an alanine residue produces lysinoalanine [3]. The strength of duramycin's structure is likely to confer its resistance to thermal and proteolytic degradation [2]. The presence of the covalent linkages also results in duramycin's stable binding site which selectively recognises the membrane phospholipid phosphatidylethanolamine (PE) through binding to its polar head group [4]. Duramycin binds to PE with high specificity with a binding molar ratio of 1:1 and a dissociation constant in the low nanomolar range [5]. The lantibiotic cinnamycin, which is structurally similar to duramycin, was shown to induce non-specific transbilayer membrane movement of phospholipids in model and cell membranes [6]. It is therefore possible that duramycin self-promotes its own binding to cells by inducing movement

of PE molecules from the inner to the outer membrane. PE is an abundant membrane phospholipid which is ubiquitous in eukaryotic cells. In quiescent cells PE is predominantly located on the inner cell membrane along with phosphatidylserine [7,8]. The asymmetrical distribution of PE and phosphatidylserine is enzymatically maintained by a number of ATP-dependent aminophospholipid translocases and floppases and ATP-independent scramblases [9,10]. On activation PE can become exposed on the outer cell membrane during a number of physiologically important processes such as apoptotic and necrotic cell death, reorganisation of the membrane during cytokinesis and initiation of the coagulation cascade [11–14]. Upregulation of cell surface PE expression has been observed in cancer cells, cancer de-rived microparticles and on the tumour endothelium of a variety of in vitro and in vivo tumours [1,15–17]. It is well documented that upregulation of cell surface exposure of phosphatidylserine occurs after chemotherapy treatment and that this can increase the procoagulant activity of the tumour [18–20]. On the premise that PE and phosphatidylserine are regulated by the same mechanisms and that both are exposed during apoptosis [21] and after irradiation [1] a number of studies investigated whether PE exposure was upregulated after chemotherapy. It was demonstrated that externalisation of PE, detected by duramycin, could potentially be a marker of early apoptosis and could be utilised as a detection method for the response to chemotherapy treatment [22–24].

Photodynamic therapy (PDT) is a minimally invasive treatment that can deliver selective cytotoxic activity towards cancer [25]. PDT requires three essential components; a photosensitising agent, light and oxygen. Accumulation of the photosensitiser (PS) at the site of cancer and activation with light at a specific wavelength causes the PS to transfer energy, via its excited triplet state, to molecular oxygen which is converted from its ground triplet state to an excited singlet state. This leads to the generation of cell damaging reactive oxygen species (ROS) which ultimately causes apoptotic or necrotic cell death [25,26]. As well as its direct cytotoxic effect on cancer, PDT can have anti-tumour effects via induction of the inflammatory response and damage to tumour vasculature [27,28]. PDT has been approved for treatment of a variety of cancer types including premalignant and malignant skin cancers [29,30], early carcinomas of the oral cavity, pharynx, and larynx [31], cancers of the GI tract [32], prostate cancer, bladder cancer, non-small cell lung cancer and brain tumours [25]. PDT has been used as a synergistic method combined with other major therapies e.g. chemotherapy, radiotherapy and surgery [33–35] as it does not compromise the effect of the therapies and therefore can be used in conjunction without effecting future treatment options for residual or recurrent malignancy [25]. A number of PS have been approved for use in clinical oncology including m-THCP (meta-tetrahydroxyphenylchlorin, Foscan®), ALA (5-aminolevulinic acid, Levulan®) and its methyl ester (Metvix®) and porfimer sodium (Photofrin®). There are a wide range of PS still in pre-clinical and clinical trials [36,37].

Porphyrins are highly conjugated heterocyclic tetrapyrroles that were first identified as having photodynamic properties in the 1970s [38,39]. A porphyrin PS (5-[Aminobutyl-N-oxycarbonyl]phenyl)phenyl]-10,15,20-tris(N-methyl-4-pyridinium) porphyrin trichloride) plus light irradiation was able to reduce cell survival and prevent colony forming in prostate epithelial cells at concentrations of 8.75 µM [40]. The porphyrins TTP (5,10,15,20-tetra-p-tolyl porphyrin) and THNP

(5,10,15,20-tetra-p-naphthyl-porphyrin) after a light irradiation dose of 4.5 J/cm² were able to cause cell death in a human melanoma cell line in a concentration dependent manner [37]. The main disadvantage of PDT is the low selectivity of PS to their target, which can lead to damage to surrounding normal tissue and sustained skin photosensitivity [25]. In an attempt to overcome this research in PDT in oncology has focused on developing PS's conjugated to tumour targeted biological molecules [28]. Porphyrins have been successfully conjugated to a variety of biomolecules including mAbs [26,41,42], lipoproteins [43,44] and nanoparticles [45] which allowed for more targeted delivery of PS, and therefore a more specific tissue response with limited damage to normal tissue. A number of porphyrins have been developed that conjugate to biomolecules under mild conditions through direct reaction with lysine residues [26,33,46].

The two primary amines, cysteine and lysine, at the N-terminal of duramycin offer potential sites for conjugation without interference with duramycin's binding site [3]. Using this theory a number of duramycin conjugates have been developed to take advantage of PE as a possible selective molecular probe. Fluorescent duramycin conjugates have been used to successfully target and image the lymphoma cancer cell line U937 [4], cancer derived microparticles [17], prostate tumour endothelium in rats [1] and the endothelium of aortic flow dividers [47]. Duramycin has been radiolabelled with gadolinium [3] and tech-netium-99 m where it was used to image cell death in a rat model of acute myocardial infarction [47-49]. The radiolabelled duramycin conjugate quickly accumulated in the PE expressing myocardium. Fast accumulation of a duramycin-PS conjugate could potentially provide an advantage over some of the current PS used in the clinical setting as a short drug-light interval could be achieved. Thus patients could receive PDT relatively quickly in the outpatient setting and would have a reduced treatment duration. In this proof of concept study we conjugated duramycin to a porphyrin based PS and assessed its ability to bind to cancer cells via cell surface PE. It has been previously shown that the level of cell surface PE expression, detected by duramycin, was increased on the pancreatic cancer cell lines AsPC-1 and CFPAC-1 when apoptotic (induced by camptothecin treatment) compared to untreated viable cells and was further increased when the cells were necrotic (induced by 30% H₂O₂ treatment) [15]. As duramycin can bind to quiescent, apoptotic and necrotic cancer cells it was theorised that a duramycin conjugate could potentially provide targeted treatment of tumours at all stages of progression. We also examined its cytotoxic activity on ovarian and pancreatic cancer cell lines following irradiation with visible light.

2. Materials and Methods

2.1. Cell Culture

The cancer cell lines AsPC-1 and MIA PaCa-2 were purchased from the American Type Culture Collection (UK). The cancer cell lines A2780 and SK-OV-3 were purchased from the European Collection of Cell Cultures (UK). All cells were grown using distributors' instructions. All cells were cultured in either IMDM, McCoy's 5a Medium Modified or RPMI (Lonza, UK) substituted with 10% FBS (15% for McCoy's 5a Medium Modified) (Bio-Sera, UK) and (v/v); 100 units/mL penicillin, 100 µg/mL streptomycin (P/S) (Lonza, UK). All serum was filtered using a 0.2 µm syringe filter prior to addition to media. When not in use all media was stored between 4 and 6 °C. All cells were incubated at 37 °C in a 5% CO₂ atmosphere. Cells were cultured in tissue culture flasks (Sarstedt, UK) and removed via scraping when cells were 70– 90% confluent.

2.2. Conjugation of Porphyrin to Duramycin

A solution of duramycin (Sigma-Aldrich, UK, 10 mg, 4.97×10^{-3} mmol) in DMSO (2 mL) plus trimethylamine was treated with 5-[4-(N-succinimidyl)oxycarbonyl]phenyl]-10,15,20-tris(4-methylpyridinium)porphyrin trichloride [50] (8 mg, 9.94×10^{-3} mmol) and the resulting solution was mixed on a rotating shaker at room temperature for 24 h. The crude material was recovered by addition of dichloromethane (5 mL) and filtration through paper. The solid was dissolved in 1 mL of 0.1% aqueous trifluoroacetic acid (TFA) and the 1:1 porphyrin/duramycin conjugate was isolated by semi-preparative HPLC and analysed by electron ionising mass spectrometry (ESI-MS) (Section 2.3). Recovery of 4.8 mg of conjugate was achieved equating to 34.4% yield. An unconjugated, “capped” porphyrin was used as a control in which the NHS ester group used for conjugation was reacted with butylamine to eliminate any non-specific conjugation [40]. All photosensitiser containing compounds were kept, protected from light, at -20 °C. Final concentrations of the re-suspended lyophilised powder of the conjugate and “capped” control were determined using UV-visible spectrophotometry.




Fig. 1. Structure of duramycin-porphyrin conjugate. A covalent amide bond was formed through the reaction with the NHS ester of the porphyrin 5-[4-(N-succinimidyl)oxycarbonyl]phenyl]-10,15,20-tris(4-methylpyridinium) porphyrin trichloride and the ϵ -amino group of the lysine residue at duramycin's N-terminal in the organic solvent DMSO

2.3. Analyses and Purification of Duramycin-Porphyrin Conjugate

RP-HPLC-PDA analyses were performed on an Agilent 1200 HPLC system (Agilent, Germany) equipped with a G1312B BinPump SL, G13798 degasser, G1367D HiP ALS SL plus autosampler, a G1316B column compartment (set at 35 °C), and a G1315C diode array detector (acquisition rate, 80 Hz; scan rate:

380–600 nm, step: 1 nm), using a Gemini C18 column, 5 μ , 150 \times 4.6 mm, 110 Å column (Phenomenex, UK), equipped with a SecurityGuard C18 (ODS) 4 \times 3.0 mm ID guard column (Phenomenex, UK). The injection volume was 20 μ L, and the flow rate was 1 mL/min. Mobile phase consisted of a mixture of 0.1% TFA in deionised water (18.2 M Ω , Elga Purelab Ultra ULXXXGEM2), solvent A, and 0.1% TFA in HPLC-grade acetonitrile, solvent B. The gradient for the elution was as follows: 0–2 min, 5% B; 2–7 min, B to 95%; 7–12 min, 95% B; 12–12.1 min, B to 5%. Duramycin R_t = 7.32, porphyrin R_t = 7.01, conjugate R_t = 6.20 min (See Supplementary Fig. S-1). RP-HPLC purifications were carried out on a Jasco HPLC system equipped with a PU-1580 LC pumps, HG-1580 degasser, AS-1555 autosampler and MD-1515 array UV–visible detector, using a Luna C18, 5 μ , 250 \times 10 mm, 110 Å (Phenomenex, UK), at a flow rate of 2.5 mL/min. Mobile phase consisted of a mixture of 0.1% TFA in deionised water (18.2 M Ω , Elga Purelab Ultra ULXXXGEM2), solvent A, and 0.1% TFA in HPLC-grade acetonitrile, solvent B. The gradient for the elution was as follows: 0–8.5 min, 5% to 51% B; 8.5–13 min, 51% B; 13–14 min, 51 to 95% B; 14–18 min, 95% B, 18–18.1 min, 95 to 5% B. Conjugate R_t = 6.20 min (See supplementary Fig. S-1). Q-TOF-MS data were acquired in positive mode scanning from 400 to 3000 m/z with and without auto MS/MS fragmentation. Ionisation was achieved with an Agilent JetStream electrospray source and infused internal reference masses. Agilent 6540 Q-TOF-MS parameters: gas temperature, 325 °C; drying gas, 10 L/min; and sheath gas temperature, 400 °C. ESI-MS (+): 675.66 [(M + 4H⁺)/4]⁴⁺ (exp: 675.67). Conjugate R_t = 7.34 (see Supplementary Fig. S-2).

2.4. Flow Cytometric Detection of Phosphatidylethanolamine

Cells (1×10^5) were re-suspended in 25 μ L PBS and added to a 5 mL polypropylene tube (Sarstedt, UK). Duramycin-porphyrin conjugate (5 μ L) at required concentration was added to the tubes and incubated at room temperature in the dark for 30 min. Cells were washed with PBS, centrifuged at 320g for 3 min and re-suspended in 300 μ L PBS and then analysed using a BD FACScalibur (BD Biosciences, UK).

2.5. Flow Cytometric Analysis of Forward Scatter and Side Scatter

Analysis of forward scatter (FSC) and side scatter (SSC) data was performed using BD CellQuest Pro software™ version 5.1 (BD Biosciences, UK) after cells were processed using the method described above. FSC and SSC data using the unconjugated duramycin was collected as part of a study to determine the effect of duramycin on cell viability over a 2 hour time course (unpublished data). These cells were processed using a flow cytometric cell viability analysis method de-scribed previously [16]. Briefly cells (2×10^5) were incubated with an appropriate concentration of unconjugated duramycin at room temperature in the dark for 30 min and analysed using Annexin V: FITC Apoptosis Detection Kit I (BD Biosciences, UK).

2.6. Competition Binding Assays

Cells (2×10^5) were re-suspended in 40 μ L PBS and added to a 5 mL polypropylene tube (Sarstedt, UK). Duramycin-porphyrin conjugate (10 μ L, final concentration 50 μ M) was added to the tubes and incubated at room temperature in the dark for 5 min. The cells were washed with PBS, centrifuged at 320g for 3 min and re-suspended in 40 μ L PBS except for the 0 minute tube which was re-suspended with 300 μ L PBS and analysed by flow cytometry. Unconjugated duramycin (10 μ L, final

concentration 50 μM) was added to the remaining tubes. After appropriate incubation time (2, 5, 15 or 30 min) 250 μL PBS was added to the tubes and the cells analysed using flow cytometry.

2.7. Photodynamic Light Treatment Cells

Cells (8×10^5) in serum free medium were added to 5 mL polypropylene tubes (Sarstedt, UK) containing duramycin-porphyrin conjugate at required concentration (0–10 μM) and incubated in the dark at 37 °C in a 5% CO_2 atmosphere for 5 min. Cells were then washed with 3 mL medium so as to remove any unbound conjugate and centrifuged at 300g for 10 min and re-suspended in 1 mL medium. Cells (8×10^4) were then plated into 96 well plates and incubated in the dark at 37 °C in a 5% CO_2 atmosphere for 5 min. The PDT light supply (9 \times 14 LED Array, wavelength 620–660 nm) was allowed to stabilise and irradiance reading taken using a R203 Macam radiometer (Irradian Ltd. UK). After incubation one plate was irradiated with a light fluence of 7.5 J/cm^2 red light whilst the other served as the dark control (i.e. not irradiated). After light treatment 5 μL FBS was added to each well and the plates incubated at 37 °C in a 5% CO_2 atmosphere for 24 h.

2.8. Cell Proliferation Assay

After 24 h incubation 10 μL MTT (3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide; thiazolyl blue) (Sigma-Aldrich, UK) (12 mM solution in PBS) was added to each well of the 96 well plates. The plates were incubated at 37 °C in a 5% CO_2 atmosphere for 4 h and then the crystals were dissolved by adding 150 μL of acid-alcohol mixture (0.04 mol/L HCl in absolute 2-propanol). The plates were then analysed at 570 nm using a Biotek ELX800 Universal Microplate Reader (Corgenix Ltd., UK). Percentage cell survival was expressed relative to control values.

2.9. Statistical Analysis

Statistical difference between the irradiated and non-irradiated conjugate conditions and between the duramycin-porphyrin conjugate and unconjugated duramycin conditions was performed using the t-test for unequal variances (heteroscedastic) and a p value of ≤ 0.05 was considered statistically significant.

3. Results

3.1. Phosphatidylethanolamine Detection

The duramycin-porphyrin conjugate ([Fig. 1](#)) was able to detect cell surface PE expression on the ovarian cancer cell lines, A2780 and SK-OV-3, and the pancreatic cancer cell lines, AsPC-1 and MIA PaCa-2 using flow cytometry. Log_{10} median PE fluorescence (635 nm laser, FL-4 flow cytometer channel) was used to express levels of detected PE. The duramycin-porphyrin conjugate detected levels of PE in a concentration dependent manner on all 4 cell lines ($n = 2$) ([Fig. 2](#)). Optimal PE detection was seen using the highest concentration (1 mM) of the conjugate. As an average of all 4 cell lines the log_{10} median PE fluorescence increased from 0.17 ± 0.007 ($n = 8$) when detected with 1 μM conjugate to 2.21 ± 0.192 ($n = 8$) when detected with 1 mM conjugate. The duramycin-porphyrin conjugate was detectable through the flow cytometer's FL-4 channel (flow cytometer's 635 nm laser) due to fluorescence of the porphyrin component of the conjugate at this wavelength.




Fig. 2. Detection of cell surface PE on ovarian and pancreatic cancer cell lines using duramycin-porphyrin conjugate. i) Median PE fluorescence (FL-4 flow cytometer channel, 635 nm laser) was used to express detected levels of PE (plotted on a logarithmic to base 10 scale). The conjugate was able to detect PE on the cell surface of all 4 cancer cell lines, A2780, AsPC-1, MIA PaCa-2 and SK-OV-3, in a concentration dependent manner. Error bars show standard deviation of 2 replicates. ii) Flow cytometry histograms showing detectable shift in fluorescence on a) A2780, b) AsPC-1, c) MIA PaCa-2 and d) SK-OV-3 cells (purple) using 0.1 µM (green) and 10 µM (pink) duramycin-porphyrin conjugate.

3.2. Competitive Binding for Phosphatidylethanolamine

To investigate whether the binding ability of the duramycin-porphyrin conjugate to PE was affected by the conjugation procedure a competition assay was performed using flow cytometry. The unconjugated duramycin was unable to compete for cell surface PE on the cancer cell lines which had been pre-incubated with duramycin-porphyrin conjugate (Fig. 3). The median PE fluorescence (FL-4 channel) of A2780 and MIA PaCa-2 was determined, using the pre-bound conjugate, as 10.9 ± 0.75 ($n = 2$) and 16.7 ± 5.9 ($n = 2$), respectively. Duramycin was incubated with the cancer cell lines for 0, 2, 5, 15 and 30 min to allow time to potentially displace the conjugate and competitively bind to PE. The median PE fluorescence of MIA PaCa-2 remained constant, within standard deviation, at each interval across the 30 minute time course. The median PE fluorescence of the A2780 however increased over time to the maximum of 12.8 ± 0.16 ($n = 2$) at 30 min. The unaffected binding affinity of the conjugate for PE after ligation to porphyrin provides evidence that the conjugation method used in this study does not disrupt PE binding. Alternatively, the assay was performed with the unconjugated duramycin pre-bound and the conjugate as the competing species, similarly over a 30 min time course. The median PE fluorescence (FL-1 channel) detected remained constant at 8.3 ± 1.8 ($n = 2$) for A2780 and 11.7 ± 1.4 ($n = 2$) for MIA PaCa-2 over the 30 min. However it cannot be certain that the FL-1 signal originated from the unconjugated duramycin (labelled with a tertiary FITC antibody detected at 488 nm) as the duramycin-porphyrin conjugate signal is also detectable on this channel.




Fig. 3. Competitive binding for PE between pre-bound duramycin-porphyrin conjugate and unconjugated duramycin. The unconjugated duramycin was unable to compete against the conjugate for cell surface PE on the cancer cell lines A2780 and MIA PaCa-2. The median PE fluorescence of MIA PaCa-2 remained constant, within standard deviation, at each interval across the 30 minute time course. However the median PE fluorescence of A2780 increased slightly over time. Error bars show standard deviation of 2 replicates.

3.3. Effect on Cell Size and Morphology

We have previously reported the effect that unconjugated duramycin has on the morphology of A2780, AsPC-1, MIA PaCa-2 and SK-OV-3 with relation to an increase in cell size (FSC) and internal cell complexity/granularity (SSC) [16]. FSC and SSC data collected via flow cytometry was analysed after the same 4 cancer cell lines were treated with the duramycin-porphyrin conjugate at concentrations between 1 μ M and 1 mM (Fig. 4). A2780 and SK-OV-3 increased in SSC when treated with 1 mM conjugate and AsPC-1 and MIA PaCa-2 increased in SSC when treated with N250 μ M (n = 2). A2780 and MIA PaCa-2 showed a slight increase in FSC when treated with 1 mM duramycin-porphyrin conjugate (n = 2).




Fig. 4. The effect of duramycin-porphyrin conjugate on the forward scatter and side scatter properties of ovarian and pancreatic cancer cell lines. The effect on FSC of a) unconjugated duramycin and b) duramycin-porphyrin conjugate and the effect on SSC of c) unconjugated duramycin and d) duramycin-porphyrin conjugate. Unconjugated duramycin has the ability to increase FSC (cell size) of AsPC-1 and SK-OV-3 cells at N50 μ M and increase SSC (cell granularity) of A2780, AsPC-1, MIA PaCa-2 and SK-OV-3 at N200 μ M. Duramycin-porphyrin conjugate slightly increased FSC of A2780 and MIA PaCa-2 at 1 mM. SSC was increased in A2780 and SK-OV-3 when treated with 1 mM conjugate and AsPC-1 and MIA PaCa-2 when treated with N250 μ M conjugate. Error bars show standard deviation of 2 replicates

3.4. Photodynamic Cytotoxicity Assays

The duramycin-porphyrin conjugate was used in the photodynamic treatment of the cancer cell lines A2780, AsPC-1, MIA PaCa-2 and SK-OV-3. Each concentration of either “capped” control porphyrin, unconjugated duramycin or duramycin-porphyrin conjugate was repeated 4 times as biological replicates and each assay performed 3 times as technical replicates. The irradiated duramycin-porphyrin conjugate was able to reduce the cell survival of the A2780 cancer cells at concentrations $\geq 5 \mu\text{M}$ (Fig. 5a). A significant difference ($p = 0.029$) in reduction in cell survival between the irradiated and non-irradiated conjugate treated cells occurred at $5 \mu\text{M}$ where cell survival was $17.7 \pm 26.9\%$ and $65.9 \pm 2.4\%$, respectively. The conjugate at the highest concentration of $10 \mu\text{M}$ was able to reduce cell survival to $3.2 \pm 3.7\%$. The unconjugated duramycin caused a slight reduction in cell survival at concentrations $\geq 5 \mu\text{M}$. The non-irradiated “capped” control porphyrin (porphyrin with no duramycin) showed no cytotoxicity. The irradiated “capped” control porphyrin however showed similar levels of cytotoxicity as the irradiated conjugate treated cells at concentrations $\geq 5 \mu\text{M}$. MIA PaCa-2 cells showed similar sensitivity to photodynamic light treatment as A2780. For conjugate treated MIA PaCa-2 cells the difference in reduction of cell survival between the irradiated and non-irradiated conditions also occurred at $5 \mu\text{M}$ (Fig. 5b) though this was not significantly different ($p = 0.32$). At $10 \mu\text{M}$ cell survival was reduced to $36.3 \pm 13.4\%$. At a concentration of $10 \mu\text{M}$ the cell survival for the irradiated and non-irradiated unconjugated duramycin was $40.3 \pm 16.3\%$ and $43.5 \pm 14\%$, respectively. The non-irradiated conjugate and “capped” control porphyrin showed no cytotoxicity. The irradiated “capped” control porphyrin reduced cell survival slightly at the highest concentration of $10 \mu\text{M}$. The cancer cell lines AsPC-1 (Fig. 5c) and SK-OV-3 (Fig. 5d) were more sensitive to photodynamic light treatment than A2780 and MIA PaCa-2. The significant reduction in cell survival between the irradiated and non-irradiated conjugate occurred at $2.5 \mu\text{M}$ ($p = 0.018$) for AsPC-1. The cell survival of non-irradiated AsPC-1 cells treated with $2.5 \mu\text{M}$ conjugate was $86.4 \pm 4.9\%$ compared to irradiated cells at $6.3 \pm 2.4\%$. For SK-OV-3 cells the significant reduction in cell survival occurred at $0.5 \mu\text{M}$ ($p = 0.039$) and was $95.5 \pm 3.9\%$ and $86.9 \pm 3.8\%$ for the non-irradiated and irradiated conditions, respectively. This reduction in cell survival became more pronounced at $2.5 \mu\text{M}$ where cell survival for non-irradiated conjugate treated cells was $91.2 \pm 7.7\%$ and only $11.7 \pm 4.4\%$ for irradiated cells. At the highest concentration of $10 \mu\text{M}$ duramycin-porphyrin conjugate reduced cell survival to $1.83 \pm 1\%$ in AsPC-1 and $2.76 \pm 2.2\%$ in SK-OV-3. The non-irradiated conjugate showed dark toxicity at concentrations $\geq 5 \mu\text{M}$ and the irradiated “capped” control porphyrin control slightly reduced cell survival at $10 \mu\text{M}$ for both cancer cell lines. Unconjugated duramycin showed a cytotoxic effect in a concentration dependent manner for both AsPC-1 and SK-OV-3. For example at $10 \mu\text{M}$ the irradiated unconjugated duramycin reduced cell survival to $48.7 \pm 20.2\%$ in AsPC-1 and $60.6 \pm 7.2\%$ in SK-OV-3. Indeed unconjugated duramycin reduced cell survival in a concentration dependent manner in all 4 cell lines. However this is to be expected due to the cytotoxic abilities of unconjugated duramycin at these concentrations as seen previously [16]. Yet, a significantly enhanced cytotoxic effect of the irradiated duramycin-porphyrin conjugate over unconjugated duramycin occurred at concentrations $\geq 1.25 \mu\text{M}$ for A2780 ($p = 0.045$) and AsPC-1 ($p = 0.03$) and at all concentrations ($0.5\text{--}10 \mu\text{M}$) for SK-OV-3 ($p = 0.046$). Reduction in cell survival of MIA PaCa-2 cells when treated with irradiation plus conjugate was not significantly different to the unconjugated duramycin treated cells ($p = 0.91$).




Fig. 5. PDT Treatment of ovarian and pancreatic cancer cell lines. The irradiated duramycin-porphyrin conjugate at concentrations $\geq 0.5 \mu\text{M}$ significantly reduced cell survival of AsPC-1, A2780 and SK-OV-3 cells when compared to the non-irradiated conjugate. a) The unconjugated duramycin caused a slight reduction in cell survival of A2780 cells at concentrations $\geq 5 \mu\text{M}$. The irradiated capped porphyrin treated cells showed similar levels of cytotoxicity as the irradiated conjugate treated cells at concentrations $\geq 5 \mu\text{M}$. b) The irradiated capped porphyrin reduced cell survival of MIA PaCa-2 cells slightly at the highest concentration of $10 \mu\text{M}$. Reduction in cell survival in conjugate treated cells was not significantly different to unconjugated duramycin treated cells and there was no significant difference between irradiated and non-irradiated conjugate conditions. c) and d) Both AsPC-1 and SK-OV-3 showed dark toxicity at concentrations with non-irradiated conjugate $\geq 5 \mu\text{M}$. The irradiated capped porphyrin control slightly reduced cell survival at $10 \mu\text{M}$. Unconjugated duramycin showed cytotoxic effect in a concentration dependent manner. Error bars show standard deviation of 3 independent experiments with 4 repetitions each.

4. Discussion

Conjugation of tetrapyrroles to peptides can be chemically challenging due to the requirement of bringing the two moieties together regioselectively whilst maintaining their separate functionalities [33].

In this study conjugation occurred by the formation of a covalent amide bond between an N-hydroxysuccinimide (NHS) ester of the porphyrin and the ϵ -amino group on the lysine residue at duramycin's N-terminal. The reaction conditions were chosen in order to target the desired lysine residue. NHS esters are highly amine-reactive and form stable amide bonds primarily with lysine's ϵ -amino group and the α -amino group of the N-terminal residue under physiological or slightly alkaline pH (pH 7.2–9) [51]. As the NHS ester was non-sulphonated, causing it to be water insoluble, the reaction was performed in the water miscible organic solvent DMSO. To produce a pH of pH 9, provide a base in the non-aqueous DMSO solution and catalyse the reaction between NHS and lysine trimethylamine (TEA) was added [50]. In addition to the reaction conditions used the structure of duramycin means it is reasonable to presume the desired location for conjugation was achieved. The only exposed reactive functionality on duramycin is the ϵ -amino group of the N-terminal lysine residue as the α -amino group of the lysine and the amino terminus itself is folded inside the peptide's structure. The conjugation of the porphyrin to duramycin at this chosen location should minimise interference with the PE binding site [3]. A number of biological molecules conjugated to porphyrin through lysine residues have been shown to retain their antimicrobial function [50,52, 53]. The scFv antibody fragment LAG-3 when conjugated through its lysine residues to a porphyrin PS was shown to bind to the colorectal cancer cell line Caco-2 and effectively induce cell death following 15 J/cm² light treatment [26].

In this study the duramycin-porphyrin conjugate was used to detect cell surface PE expression on the 4 cancer cell lines A2780, AsPC-1, MIA PaCa-2 and SK-OV-3 via flow cytometry. In this way the conjugate was tested to see whether the duramycin component's ability to bind the ethanolamine head group of PE had been altered during conjugation. It was also assessed herein whether the porphyrin had retained its fluorescent properties. The conjugate detected PE expression on all 4 cell

lines and was detected in a concentration dependent manner thus verifying the conjugates ability to bind its target and maintain detectable fluorescence. Though it was not shown directly in this study that there was an upregulation of PE on the cancer cell lines compared to their normal counterparts, we have previously shown that PE expression occurs on a broad range of cancer cell lines [16]. Also it has been recently shown that PE was present in much higher abundance in xenografts formed from a number of different tumour types (lung, colon, pancreatic and glioblastoma) than in their normal tissue counterparts [54]. It is well established that duramycin binds to PE when exposed on the surface of cells and on the luminal surface of the endothelium [1,15,47, 49,55]. Duramycin was used in conjunction with bavituximab to image the co-localisation of exposed PE and phosphatidylserine on irradiated endothelial cells [56]. The cells were stained 24 hour post irradiation and coincidental exposure of the two phospholipids was observed on the surface of the cells. Duramycin staining occurred only where PE was exposed on the surface of the cell, even when the exposure area moved due to cell movement across the dish, and this staining was maintained

on the surface over the 8 hour imaging period. Therefore we theorise that the binding of duramycin and thus the duramycin-porphyrin conjugate occurs only at the cell surface where PE is exposed and is likely not internalised. To further investigate whether the conjugate's PE binding ability was affected by the conjugation procedure a competition assay was performed via flow cytometry. Unconjugated duramycin did not have the ability to outcompete the conjugate for binding of cell surface PE. The median PE fluorescence detected remained constant for MIA PaCa-2 cells over the 30 minute time course and was seen to slightly increase for A2780.

We have previously reported that unconjugated duramycin has an effect on the FSC and SSC properties of cancer cell lines [16]. Duramycin has the ability to increase cell size (FSC) of AsPC-1 and SK-OV-3 cells at N50 μ M and increase cell granularity (SSC) of A2780, AsPC-1, MIA PaCa-2 and SK-OV-3 at N200 μ M. Necrotic cell death is associated with swelling of the cell followed by rapid release of intracellular content and an increase in granularity which have been linked to an increase in FSC and SSC, respectively [57]. An increase in SSC was seen to correlate with increased granularity in prostate cancer cells (LNCaP and PC3) [58]. Cells with an increased SSC in the renal proximal tubular epithelial cell line LLC-PK1 were identified as apoptotic by flow cytometry and those with an increased FSC as necrotic [59]. Heightened SSC was also seen in early apoptotic cells where condensation of the cell nucleus and cytoplasm during cell shrinkage lead to enhanced light refraction [60]. However it is important to state that morphological and cell size changes measured via light scatter should be considered as supportive rather than as definitive evidence of cell death. We have previously shown via confocal imaging that duramycin treated AsPC-1 cells lose cell membrane staining accompanied with an uptake of propidium iodide suggesting a loss in membrane integrity or cell death. An increase in cell size was also observed, which is indicative of cell swelling prior to cell death [16]. The duramycin-porphyrin conjugate also effected the FSC/SSC properties of the ovarian and pancreatic cancer cell lines. FSC was increased in A2780 and MIA PaCa-2 when treated with 1 mM conjugate and SSC increased in all 4 cell lines though AsPC-1 and MIA PaCa-2 at N250 μ M and A2780 and SK-OV-3 at 1 mM. The similar effects on FSC and SSC produced by duramycin and duramycin-porphyrin conjugate could provide additional evidence of duramycin's maintained functionality when conjugated to porphyrin. The higher concentration of conjugate required to exert this effect on the cell lines compared to duramycin alone is beneficial to PDT where minimal non-specific damage is a desired characteristic of any PS [36].

A number of studies in the literature have investigated the effect of PDT on ovarian and pancreatic cancer in a variety of models including cell lines [61], 3D cell culture [62–64], transplanted mice [65,66] and Phase I/II human trials [67,68]. A benzoporphyrin derivative (BPD) at 0.25 μ M was able to reduce cell viability and volume in the human ovarian carcinoma cell line OVCAR5 and penetrate well through OVCAR5 3D cell culture [62,63]. In another study in which OVCAR5 cells were grown in a culture film a fibre optic delivery system was used where the PS (pheophorbide) and oxygen gas was delivered separately [64]. Good diffusion of the PS to cells beyond the reach of the probe tip was observed. Balb/C mice transplanted with the ovarian cancer cell line OVCAR3 where treated with an immunoconjugated PS and irradiated with in vivo results showing targeted treatment and minimal damage to the surrounding vasculature [65]. A pancreatic cell line panel showed the cancer cell line AsPC-1

was particularly sensitive to PDT. Treatment of this cell line with verteporfin at 250 nM with a light dose of 10 J/cm² was effective enough to induce complete cell death [61]. Verteporfin has been used successfully in a Phase I/II trial against advanced pancreatic cancer where it accumulated quickly in the target tissue, induced necrotic cell death and had a fast clearance via excretion in bile [68].

In this study we used 2 ovarian and 2 pancreatic cancer cell lines to evaluate the cytotoxic extent of our newly developed PS, duramycin-porphyrin conjugate, taking advantage of duramycin's specificity for cell surface expressed PE. The duramycin-porphyrin conjugate was able to reduce cell survival of all 4 cancer cell lines after irradiation with a light dose of 7.5 J/cm² red light. The difference in cell survival between the non-irradiated and irradiated cells treated with the conjugate occurred at 0.05 µM for SK-OV-3, 2.5 µM for AsPC-1 and 5 µM for A2780 and MIA PaCa-2 (though not statistically significant for MIA PaCa-2). Treatment with 10 µM conjugate was effective enough to reduce cell survival to 6% in A2780, AsPC-1 and SK-OV-3. MIA PaCa-2 exhibited less sensitivity to the PDT light treatment as cell survival was only ever reduced to 36.3% at the highest concentration. Reduction in cell survival of MIA PaCa-2 in conjugate treated cells was not significantly different to unconjugated duramycin treated cells. The greater resistance to irradiation and the lack of increased cytotoxic effect of the conjugate over duramycin alone may suggest that MIA PaCa-2 cancer cells may not be a candidate for PDT using this PS. Out of the 2 pancreatic cancer cell lines used in this study MIA PaCa-2 is the less aggressive form. Samkoe et al. [66] discovered that aggressive, fast growing pancreatic cancer (AsPC-1) responded better to verteporfin PDT than the less aggressive cell line (PANC-1) in transplanted mice. In their vascular occlusion experiments 40 J/cm light dose was enough to induce complete necrosis in the tumours of AsPC-1 transplanted mice but not in PANC-1. They concluded that this difference in reaction to light treatment between cell lines could depend on tumour aggressiveness and other inherent biophysical properties. It has been observed in human trials that these differences in response to light treatment are of clinical importance where light dose can raise different reactions even in patients with the same type of malignancy [28,67].

The "capped" porphyrin control when irradiated reduced cell survival slightly in AsPC-1, SK-OV-3 and MIA PaCa-2 at 10 µM. In a recent study [40] this "capped" control porphyrin was used in PDT experiments against benign and malignant prostate cells to assess its cytotoxic potential. "Capped" porphyrin 8.75 µM was seen to induce necrosis and prevent colony forming in irradiated prostate cells. In the A2780 cancer cell line the capped porphyrin control produced similar levels of cytotoxicity as to that produced by the irradiated conjugate. An increased uptake and localisation of the "capped" porphyrin could potentially explain this increased sensitivity. However DNA damage in the prostate cells (suggesting localisation occurred in the nucleus) was induced after 1 hour incubation with the porphyrin [40]. This movement into the cell and localisation of the "capped" control porphyrin is unlikely to have occurred in the A2780 cells in the relatively short incubation of 5 min used in this study.

The differences in sensitivities to irradiation between cancer cell lines could potentially be explained by the cytoprotective mechanisms that cancer cells can employ to escape PDT induced cell death. Antioxidant molecules and detoxifying enzymes such as superoxide

dismutase (SOD) present in cancer cells have been implicated in the protection against PDT effects [25]. Other cytoprotective mechanisms have been demonstrated to be induced by PDT itself including the i) initiation of stress-related transcription factors e.g. hypoxia inducible factor (HIF) and nuclear factor erythroid 2-related factor 2 (NRF2), ii) increased heat shock protein (HSP) levels e.g. elevated HSP60 and HSP70 have been seen to be inversely correlated with PDT sensitivity and iii) inhibition of NF- κ B which can activate anti-apoptotic genes [25]. The literature reports at least some of these protective mechanisms are active in the ovarian and pancreatic cancer cell lines used in this study. An upregulation of HIF-1- α , an important factor in tumour metastasis and progression, was seen to accompany the hypoxic state of A2780 tumours and has also been observed in hypoxia induced MIA PaCa-2 cells and following oxidative stress-induced ROS production in SK-OV-3 cells [71]. SK-OV-3 cells also have an upregulation of the antioxidant gene SOD2 [72] which could protect against harmful PDT effects. This ovarian cancer cell line has also been shown to have enhanced NF- κ B activity, when stimulated by ROS, which lead to the activation of anti-apoptotic genes [73]. NRF2 is the principal transcription factor of many antioxidant and detoxification genes and is overexpressed in A2780 [74], SK-OV-3 [75] and AsPC-1 cells [76,77] and can be induced in the presence of ROS especially in pancreatic cancer [77]. HSPs can bind oxidatively damaged proteins and play a role in pro-survival path-ways [25]. HSP70 is overexpressed in MIA PaCa-2 cells [78] and is found in much higher quantities in tissue from patients with ovarian cancer than in normal ovarian tissue [79]. The effectiveness of PDT relies on an available and sustained oxygen supply to tissue, which may be reduced in hypoxic conditions [80]. A2780 xenografts in a murine model of ovarian cancer were found to be extremely hypoxic (2.0 ± 0.7 mmHg) when compared to RIF-1 xenografts (7.8 ± 1.4 mmHg); a typical hypoxic tumour model [69].

An important property of any good PS is a high intersystem crossing (ISC) yield i.e. a high possibility of conversion from the excited singlet state to the longer lived excited triplet state, from which energy transfer to ground state (triplet) oxygen occurs [28]. Tetrapyrroles, such as the porphyrin moiety used in this study, are very efficient at undergoing ISC [81]. The benefit of conjugating tetrapyrroles with peptides is that their small size allows for good tissue penetration and usually renders them non-immunogenic [33]. An advantage of the type of porphyrin used in this study is that the conjugation procedure does not produce reactive intermediates or by-products [26,46]. This porphyrin has previously been shown to exhibit antibacterial properties, be an efficient cytotoxic after 5 minute irradiation and be stable against photodegradation [50]. The duramycin-porphyrin conjugate reported here is an effective PS for the targeted PDT treatment of cancer cells. With a relatively low (in vitro) cytotoxic concentration of $\geq 0.5 \mu\text{M}$ (de-pendent on cell line) the conjugate offers benefits over unconjugated duramycin induced cell death. At these low concentrations duramycin does not cause necrosis but can reliably bind cell surface expressed PE

Unconjugated duramycin has been shown to bind PE expressed on the tumour vasculature [1]. Through the possibility of targeted irradiance the duramycin-porphyrin conjugate could allow for specific death of the tumour and therefore reduce the risk of collateral damage to surrounding normal tissue. Photodynamic effect usually occurs very close to the localisation of the PS [25] and when the PS's binding site is the plasma membrane necrotic cell death can very rapidly follow irradiation [82]. As the duramycin-porphyrin conjugate binds to cancer cell

membranes it is possible that cell death occurs via rapid necrosis which may reduce the risk of PDT-induced cytoprotective mechanisms.

PDT is evidently an effective anti-cancer therapy though only a limited number of PS have been approved by the FDA and EU for use in the clinical setting [28,37]. For PDT to be better utilised as a treatment strategy the development of PS that possess selectivity for binding sites overexpressed or enhanced on a wide range of cancer types is needed. In this proof of concept study we have shown that the PS duramycin-porphyrin conjugate retained good binding affinity for its target and, following irradiation, reduced cell survival of pancreatic and ovarian cancer cell lines in a concentration dependent manner. As we have previously shown, duramycin has the ability to bind to cell surface expressed PE on a range of cancer cell lines [16], and so we hypothesise that this PS could potentially be used against a number of different cancer types. Further work with duramycin-porphyrin conjugate in the in vivo environment could potentially confirm this conjugate as a promising new PS in the PDT treatment of cancer.

Conflict(s) of Interest

The authors declare that they have no conflicts of interest with the contents of this article

Supplementary data to this article can be found at

<http://dx.doi.org/10.1016/j.jphotobiol.2016.09.001>

References

- [1] J.H. Stafford, P.E. Thorpe, Increased exposure of phosphatidylethanolamine on the surface of tumor vascular endothelium, *Neoplasia* 13 (4) (2011) 299-IN2.
- [2] N. Zimmermann, S. Freund, A. Fredenhagen, G. Jung, Solution structures of the lantibiotics duramycin B and C, *Eur. J. Biochem.* 216 (2) (1993) 419–428.
- [3] M. Zhao, Lantibiotics as probes for phosphatidylethanolamine, *Amino Acids* 41 (5) (2011) 1071–1079.
- [4] Y. Aoki, T. Uenaka, J. Aoki, M. Umeda, K. Inoue, A novel peptide probe for studying the transbilayer movement of phosphatidylethanolamine, *J. Biochem.* 116 (2) (1994) 291–297.
- [5] F. Hayashi, K. Nagashima, Y. Terui, Y. Kawamura, K. Matsumoto, H. Itazaki, The structure of PA48009: the revised structure, *J. Antibiot.* 43 (11) (1990) 1421–1430.
- [6] A. Makino, T. Baba, K. Fujimoto, K. Iwamoto, Y. Yano, N. Terada, S. Ohno, S.B. Sato, A. Ohta, M. Umeda, K. Matsuzaki, T. Kobayashi, Cinnamycin (Ro 09-0198) promotes cell binding and toxicity by inducing transbilayer lipid movement, *J. Biol. Chem.* 278 (5) (2003) 3204–3209.
- [7] B. Fadeel, D. Xue, The ins and outs of phospholipid asymmetry in the plasma membrane: roles in health and disease, *Crit. Rev. Biochem. Mol. Biol.* 44 (5) (2009) 264–277.
- [8] G. van Meer, D.R. Voelker, G.W. Feigenson, Membrane lipids: where they are and how they behave, *Nat. Rev. Mol. Cell Biol.* 9 (2) (2008) 112–124.
- [9] D.L. Daleke, Phospholipid flippases, *J. Biol. Chem.* 282 (2) (2007) 821–825.
- [10] S.K. Sahu, S.N. Gummadi, N. Manoj, G.K. Aradhyam, Phospholipid scramblases: an overview, *Arch. Biochem. Biophys.* 462 (1) (2007) 103–114.
- [11] R.F.A. Zwaal, A.J. Schroit, Pathophysiologic implications of membrane phospholipid asymmetry in blood cells, *Blood* 89 (4) (1997) 1121–1132.
- [12] K. Emoto, N. Toyama-Sorimachi, H. Karasuyama, K. Inoue, M. Umeda, Exposure of phosphatidylethanolamine on the surface of apoptotic cells, *Exp. Cell Res.* 232 (2) (1997) 430–434.
- [13] R.F.A. Zwaal, P. Comfurius, E.M. Bevers, Lipid–protein interactions in blood coagulation, *Biochim. Biophys. Acta Rev. Biomembr.* 1376 (3) (1998) 433–453.

- [14] L.A. Falls, B. Furie, B.C. Furie, Role of phosphatidylethanolamine in assembly and function of the Factor IXa–Factor VIIIa complex on membrane surfaces†, *Biochemis- try* 39 (43) (2000) 13216–13222.
- [15] K.R. Yates, J. Welsh, N.O. Udegbunam, J. Greenman, A. Maraveyas, L.A. Madden, Duramycin exhibits antiproliferative properties and induces apoptosis in tumour cells, *Blood Coagul. Fibrinolysis* 23 (5) (2012) 396–401.
- [16] L.J. Broughton, C. Crow, A. Maraveyas, L.A. Madden, Duramycin-induced calcium re- lease in cancer cells, *Anti-Cancer Drugs* 27 (3) (2016) 173–182.
- [17] M.C. Larson, J.E. Woodliff, C.A. Hillery, T.J. Kearl, M. Zhao, Phosphatidylethanolamine is externalized at the surface of microparticles, *Biochim. Biophys. Acta – (BBA) Mol. Cell Biol. Lipids* 1821 (12) (2012) 1501–1507.
- [18] Y. Fu, J. Zhou, H. Li, F. Cao, Y. Su, S. Fan, Y. Li, S. Wang, L. Li, G.E. Gilbert, J. Shi, Dau- norubicin induces procoagulant activity of cultured endothelial cells through phosphatidylserine exposure and microparticles release, *Thromb. Haemost.* 104 (6) (2010) 1235–1241.
- [19] J.C. Boles, J.C. Williams, R.M. Hollingsworth, J.G. Wang, S.L. Glover, A.P. Owens 3rd, D.A. Barcel, R.S. Kasthuri, N.S. Key, N. Mackman, Anthracycline treatment of the human monocytic leukemia cell line THP-1 increases phosphatidylserine exposure and tissue factor activity, *Thromb. Res.* 129 (2) (2012) 197–203.
- [20] X. Dong, J. Shi, J. Zhou, X. Chen, Y. Jin, X. Zhang, X. Li, H. Dai, J. Wang, Chemotherapy induces enhanced procoagulant activity through phosphatidylserine exposure in acute lymphoblastic leukemia, *Thromb. Res.* 132 (5) (2013) 614–620.
- [21] N. Maulik, V.E. Kagan, V.A. Tyurin, D.K. Das, Redistribution of phosphatidylethanol- amine and phosphatidylserine precedes reperfusion-induced apoptosis, *Am. J. Phys.* 274 (1 Pt 2) (1998) H242–H248.
- [22] S. Yao, K. Hu, G. Tang, X. Liang, K. Du, D. Nie, S. Jiang, L. Zang, Positron emission to- mography imaging of cell death with [(18)F]FPDuramycin, *Apoptosis* 19 (5) (2014) 841–850.
- [23] F. Elvas, C. Vangestel, K. Pak, P. Vermeulen, B. Gray, S. Stroobants, S. Staelens, L. Wyffels, Early prediction of tumor response to treatment: preclinical validation of ^{99m}Tc-duramycin, *J. Nucl. Med.* 57 (5) (2016) 805–811.
- [24] R. Luo, L. Niu, F. Qiu, W. Fang, T. Fu, M. Zhao, Y.J. Zhang, Z.C. Hua, X.F. Li, F. Wang, Monitoring apoptosis of breast cancer xenograft after paclitaxel treatment with ^{99m}Tc-labeled duramycin SPECT/CT, *Mol. Imaging* 15 (2016).
- [25] P. Agostinis, K. Berg, K.A. Cengel, T.H. Foster, A.W. Girotti, S.O. Gollnick, S.M. Hahn, M.R. Hamblin, A. Juzeniene, D. Kessel, M. Korbelik, J. Moan, P. Mroz, D. Nowis, J. Piette, B.C. Wilson, J. Golab, Photodynamic therapy of cancer: an update, *CA Cancer J Clin.* 61 (4) (2011) 250–281.
- [26] C. Staneloudi, K.A. Smith, R. Hudson, N. Malatesti, H. Savoie, R.W. Boyle, J. Greenman, Development and characterization of novel photosensitizer: scFv conjugates for use in photodynamic therapy of cancer, *Immunology* 120 (4) (2007) 512–517.
- [27] M. Korbelik, PDT-associated host response and its role in the therapy outcome, *La- sers Surg. Med.* 38 (5) (2006) 500–508.
- [28] T.C. Zhu, J.C. Finlay, The role of photodynamic therapy (PDT) physics, *Med. Phys.* 35 (7) (2008) 3127.
- [29] M.S. Nestor, M.H. Gold, A.N. Kauvar, A.F. Taub, R.G. Geronemus, E.C. Ritvo, M.P. Goldman, D.J. Gilbert, D.F. Richey, T.S. Alster, R.R. Anderson, D.E. Bank, A. Carruthers, J. Carruthers, D.J. Goldberg, C.W. Hanke, N.J. Lowe, D.M. Pariser, D.S. Rigel, P. Robins, J.M. Spencer, B.D. Zelickson, The use of photodynamic therapy in dermatology: results of a consensus conference, *J. Drugs Dermatol.* 5 (2) (2006) 140–154.
- [30] L.R. Braathen, R.M. Szeimies, N. Basset-Seguín, R. Bissonnette, P. Foley, D. Pariser, R. Roelandts, A.M. Wennberg, C.A. Morton, D. International Society for Photodynamic Therapy, Guidelines on the use of photodynamic therapy for nonmelanoma skin cancer: an international consensus.

International Society for Photodynamic Therapy in Dermatology, 2005, *J. Am. Acad. Dermatol.* 56 (1) (2007) 125–143.

[31] W. Jerjes, T. Upile, S. Akram, C. Hopper, The surgical palliation of advanced head and neck cancer using photodynamic therapy, *Clin. Oncol. (R. Coll. Radiol.)* 22 (9) (2010) 785–791.

[32] H.C. Wolfsen, Carpe luz—seize the light: endoprevention of esophageal adenocarcinoma when using photodynamic therapy with porfimer sodium, *Gastrointest. Endosc.* 62 (4) (2005) 499–503.

[33] F. Giuntini, C.M. Alonso, R.W. Boyle, Synthetic approaches for the conjugation of porphyrins and related macrocycles to peptides and proteins, *Photochem. Photobiol. Sci.* 10 (5) (2011) 759–791.

[34] Q. Xie, Synergetic anticancer effect of combined gemcitabine and photodynamic therapy on pancreatic cancer in vivo, *World J. Gastroenterol.* 15 (6) (2009) 737.

[35] S.R. Lucena, N. Salazar, T. Gracia-Cazana, A. Zamarron, S. Gonzalez, A. Juarranz, Y. Gilaberte, Combined treatments with photodynamic therapy for non-melanoma skin cancer, *Int. J. Mol. Sci.* 16 (10) (2015) 25912–25933.

[36] D. Nowis, M. Makowski, T. Stoklosa, M. Legat, T. Issat, J. Golab, Direct tumor damage mechanisms of photodynamic therapy, *Acta Biochim. Pol.* 52 (2) (2005) 339–352.

[37] I. Baldea, R.M. Ion, D.E. Olteanu, I. Nenu, D. Tudor, A.G. Filip, Photodynamic therapy of melanoma using new, synthetic porphyrins and phthalocyanines as photosensitisers - a comparative study, *Clujul Med* 88 (2) (2015) 175–180.

[38] J.D. Spikes, Porphyrins and related compounds as photodynamic sensitizers, *Ann. N. Acad. Sci.* 244 (1975) 496–508.

[39] X. Yang, P. Palasuberniam, D. Kraus, B. Chen, Aminolevulinic acid-based tumor detection and therapy: molecular mechanisms and strategies for enhancement, *Int. J. Mol. Sci.* 16 (10) (2015) 25865–25880.

[40] F.M. Frame, H. Savoie, F. Bryden, F. Giuntini, V.M. Mann, M.S. Simms, R.W. Boyle, N.J. Maitland, Mechanisms of growth inhibition of primary prostate epithelial cells following gamma irradiation or photodynamic therapy include senescence, necrosis, and autophagy, but not apoptosis, *Can. Med.* 5 (1) (2016) 61–73.

[41] M.B. Vrouenraets, G.W. Visser, F.A. Stewart, M. Stigter, H. Oppelaar, P.E. Postmus, G.B. Snow, G.A. van Dongen, Development of meta-tetrahydroxyphenylchlorin-monoclonal antibody conjugates for photoimmunotherapy, *Cancer Res.* 59 (7) (1999) 1505–1513.

[42] M. Del Governatore, M.R. Hamblin, E.E. Piccinini, G. Ugolini, T. Hasan, Targeted photodestruction of human colon cancer cells using charged 17.1A chlorin e6 immunoconjugates, *Br. J. Cancer* 82 (1) (2000) 56–64.

[43] M.R. Hamblin, E.L. Newman, Photosensitizer targeting in photodynamic therapy. II. Conjugates of haematoporphyrin with serum lipoproteins, *J. Photochem. Photobiol. B.* 26 (2) (1994) 147–157.

[44] U. Schmidt-Erfurth, H. Diddens, R. Birngruber, T. Hasan, Photodynamic targeting of human retinoblastoma cells using covalent low-density lipoprotein conjugates, *Br. J. Cancer* 75 (1) (1997) 54–61.

[45] D.L. Vega, P. Lodge, J.L. Vivero-Escoto, Redox-responsive porphyrin-based polysilsesquioxane nanoparticles for photodynamic therapy of cancer cells, *Int. J. Mol. Sci.* 17 (1) (2016).

[46] J.M. Sutton, O.J. Clarke, N. Fernandez, R.W. Boyle, Porphyrin, chlorin, and bacteriochlorin isothiocyanates: useful reagents for the synthesis of photoactive bioconjugates, *Bioconjug. Chem.* 13 (2) (2002) 249–263.

[47] Z. Li, C.W. Wells, P.E. North, S. Kumar, C.B. Duris, J.A. McIntyre, Z. Ming, Phosphatidylethanolamine at the luminal endothelial surface—implications for hemostasis and thrombotic autoimmunity, *Clin. Appl. Thromb. Hemost.* 17 (2) (2011) 158–163.

[48] M. Zhao, Z. Li, S. Bugenhagen, ^{99m}Tc-labeled duramycin as a novel phosphatidylethanolamine-binding molecular probe, *J. Nucl. Med.* 49 (8) (2008) 1345–1352.

- [49] M. Zhao, Z. Li, A single-step kit formulation for the (99 m)Tc-labeling of HYNIC- Duramycin, *Nucl. Med. Biol.* 39 (7) (2012) 1006–1011.
- [50] J.P. Tome, M.G. Neves, A.C. Tome, J.A. Cavaleiro, M. Soncin, M. Magaraggia, S. Ferro, G. Jori, Synthesis and antibacterial activity of new poly-S-lysine-porphyrin conjugates, *J. Med. Chem.* 47 (26) (2004) 6649–6652.
- [51] G.T. Hermanson, Chapter 5 - homobifunctional crosslinkers, *Bioconjugate Tech- niques*, Third edition Academic Press, Boston 2013, pp. 275–298.
- [52] N.S. Soukos, L.A. Ximenez-Fyvie, M.R. Hamblin, S.S. Socransky, T. Hasan, Targeted an- timicrobial photochemotherapy, *Antimicrob. Agents Chemother.* 42 (10) (1998) 2595–2601.
- [53] L. Polo, A. Segalla, G. Bertoloni, G. Jori, K. Schaffner, E. Reddi, Polylysine-porphycene conjugates as efficient photosensitizers for the inactivation of microbial pathogens, *J. Photochem. Photobiol. B* 59 (1–3) (2000) 152–158.
- [54] R. Fernandez, J. Garate, S. Lage, S. Teres, M. Higuera, J. Bestard-Escalas, D.H. Lopez, F. Guardiola-Serrano, P.V. Escriba, G. Barcelo-Coblijn, J.A. Fernandez, Identification of biomarkers of necrosis in xenografts using imaging mass spectrometry, *J. Am. Soc. Mass Spectrom.* 27 (2) (2016) 244–254.
- [55] K. Iwamoto, T. Hayakawa, M. Murate, A. Makino, K. Ito, T. Fujisawa, T. Kobayashi, Curvature-dependent recognition of ethanolamine phospholipids by duramycin and cinnamycin, *Biophys. J.* 93 (5) (2007) 1608–1619.
- [56] A. Marconescu, P.E. Thorpe, Coincident exposure of phosphatidylethanolamine and anionic phospholipids on the surface of irradiated cells, *Biochim. Biophys. Acta* 1778 (2008) 2217–2224.
- [57] C. Dive, C.D. Gregory, D.J. Phipps, D.L. Evans, A.E. Milner, A.H. Wyllie, Analysis and discrimination of necrosis and apoptosis (programmed cell death) by multiparam- eter flow cytometry, *Biochim. Biophys. Acta, Mol. Cell Res.* 1133 (3) (1992) 275–285.
- [58] M.K. Haynes, J.J. Strouse, A. Waller, A. Leitao, R.F. Curpan, C. Bologa, T.I. Oprea, E.R. Prossnitz, B.S. Edwards, L.A. Sklar, T.A. Thompson, Detection of intracellular granu- larity induction in prostate cancer cell lines by small molecules using the HyperCyt high-throughput flow cytometry system, *J. Biomol. Screen.* 14 (6) (2009) 596–609.
- [59] E. Healy, M. Dempsey, C. Lally, M.P. Ryan, Apoptosis and necrosis: mechanisms of cell death induced by cyclosporine A in a renal proximal tubular cell line, *Kidney Int.* 54 (6) (1998) 1955–1966.
- [60] D. Wlodkowic, W. Telford, J. Skommer, Z. Darzynkiewicz, Apoptosis and beyond: cy- tometry in studies of programmed cell death, *Methods Cell Biol.* 103 (2011) 55–98.
- [61] J.P. Celli, N. Solban, A. Liang, S.P. Pereira, T. Hasan, Verteporfin-based photodynamic therapy overcomes gemcitabine insensitivity in a panel of pancreatic cancer cell lines, *Lasers Surg. Med.* 43 (7) (2011) 565–574.
- [62] I. Rizvi, S. Anbil, N. Alagic, J. Celli, L.Z. Zheng, A. Palanisami, M.D. Glidden, B.W. Pogue, Hasan, PDT dose parameters impact tumoricidal durability and cell death pathways in a 3D ovarian cancer model, *Photochem. Photobiol.* 89 (4) (2013) 942–952.
- [63] S. Anbil, I. Rizvi, J.P. Celli, N. Alagic, B.W. Pogue, T. Hasan, Impact of treatment re- sponse metrics on photodynamic therapy planning and outcomes in a three-dimen- sional model of ovarian cancer, *J. Biomed. Opt.* 18 (9) (2013) 098004.
- [64] D. Bartusik, D. Aebisher, A. Ghogare, G. Ghosh, I. Abramova, T. Hasan, A. Greer, A fiberoptic (photodynamic therapy type) device with a photosensitizer and singlet oxygen delivery probe tip for ovarian cancer cell killing, *Photochem. Photobiol.* 89 (2013) 936–941.
- [65] B.A. Goff, J. Blake, M.P. Bamberg, T. Hasan, Treatment of ovarian cancer with photo- dynamic therapy and immunoconjugates in a murine ovarian cancer model, *Br. J. Cancer* 74 (8) (1996) 1194–1198.

- [66] K.S. Samkoe, A. Chen, I. Rizvi, J.A. O'Hara, P.J. Hoopes, S.P. Pereira, T. Hasan, B.W. Pogue, Imaging tumor variation in response to photodynamic therapy in pancreatic cancer xenograft models, *Int. J. Radiat. Oncol. Biol. Phys.* 76 (1) (2010) 251–259.
- [67] S.G. Bown, A.Z. Rogowska, D.E. Whitelaw, W.R. Lees, L.B. Lovat, P. Ripley, L. Jones, P. Wyld, A. Gillams, A.W.R. Hatfield, Photodynamic therapy for cancer of the pancreas, *Gut* 50 (4) (2002) 549–557.
- [68] M.T. Huggett, M. Jermyn, A. Gillams, R. Illing, S. Mosse, M. Novelli, E. Kent, S.G. Bown, T. Hasan, B.W. Pogue, S.P. Pereira, Phase I/II study of verteporfin photo-dynamic therapy in locally advanced pancreatic cancer, *Br. J. Cancer* 110 (7) (2014) 1698–1704.
- [69] K. Selvendiran, A. Bratasz, M.L. Kuppusamy, M.F. Tazi, B.K. Rivera, P. Kuppusamy, Hypoxia induces chemoresistance in ovarian cancer cells by activation of signal transducer and activator of transcription 3, *Int. J. Cancer* 125 (9) (2009) 2198–2204.
- [70] Z.A. Sibenaller, J.L. Welsh, C. Du, J.R. Witmer, H.E. Schrock, J. Du, G.R. Buettner, P.C. Goswami, J.A. Cieslak 3rd, J.J. Cullen, Extracellular superoxide dismutase suppresses hypoxia-inducible factor-1alpha in pancreatic cancer, *Free Radic. Biol. Med.* 69 (2014) 357–366.
- [71] Y. Wang, J. Ma, H. Shen, C. Wang, Y. Sun, S.B. Howell, X. Lin, Reactive oxygen species promote ovarian cancer progression via the HIF-1alpha/LOX/E-cadherin pathway, *Oncol. Rep.* 32 (5) (2014) 2150–2158.
- [72] Y. Wang, L. Dong, H. Cui, D.H. Shen, Y. Wang, X.H. Chang, T.Y. Fu, X. Ye, Y.Y. Yao, Up-regulation of mitochondrial antioxidation signals in ovarian cancer cells with aggressive biologic behavior, *J. Zhejiang Univ. Sci. B* 12 (5) (2011) 346–356.
- [73] J.A. Saunders, L.C. Rogers, C. Klomsiri, L.B. Poole, L.W. Daniel, Reactive oxygen species mediate lysophosphatidic acid induced signaling in ovarian cancer cells, *Free Radic. Biol. Med.* 49 (12) (2010) 2058–2067.
- [74] L.J. Bao, M.C. Jaramillo, Z.B. Zhang, Y.X. Zheng, M. Yao, D.D. Zhang, X.F. Yi, Nrf2 induces cisplatin resistance through activation of autophagy in ovarian carcinoma, *Int. J. Clin. Exp. Pathol.* 7 (4) (2014) 1502–1513.
- [75] J.M. Cho, S. Manandhar, H.R. Lee, H.M. Park, M.K. Kwak, Role of the Nrf2-antioxidant system in cytotoxicity mediated by anticancer cisplatin: implication to cancer cell resistance, *Cancer Lett.* 260 (1–2) (2008) 96–108.
- [76] Y.B. Hong, H.J. Kang, S.Y. Kwon, H.J. Kim, K.Y. Kwon, C.H. Cho, J.M. Lee, B.V. Kallakury, I. Bae, Nuclear factor (erythroid-derived 2)-like 2 regulates drug resistance in pancreatic cancer cells, *Pancreas* 39 (4) (2010) 463–472.
- [77]
- [78] Y.W. Yi, S. Oh, Comparative analysis of NRF2-responsive gene expression in AcPC-1 pancreatic cancer cell line, *Genes Genomics* 37 (2015) 97–109.
- [79] N. Walsh, A. Larkin, N. Swan, K. Conlon, P. Dowling, R. McDermott, M. Clynes, RNAi knockdown of Hop (Hsp70/Hsp90 organising protein) decreases invasion via MMP-2 down regulation, *Cancer Lett.* 306 (2) (2011) 180–189.
- [80] H.J. Kang, H.S. Moon, H.W. Chung, The expression of FAS-associated factor 1 and heat shock protein 70 in ovarian cancer, *Obstet. Gynecol. Sci.* 57 (4) (2014) 281–290.
- [81] J. Golab, D. Olszewska, P. Mroz, K. Kozar, R. Kaminski, A. Jalili, M. Jakobisiak, Erythropoietin restores the antitumor effectiveness of photodynamic therapy in mice with chemotherapy-induced anemia, *Clin. Cancer Res.* 8 (5) (2002) 1265–1270.
- [82] L.B. Josefsen, R.W. Boyle, Photodynamic therapy and the development of metal-based photosensitisers, *Metal-Based Drugs* 2008 (2008) 276109.
- [83] D. Kessel, R.D. Poretz, Sites of photodamage induced by photodynamic therapy with a chlorin e6 triacetoxymethyl ester (CAME), *Photochem. Photobiol.* 71 (1) (2000) 94–96.

