



LJMU Research Online

Lardo, C, Pancino, E, Mucciarelli, A and Milone, AP

The discovery of an anomalous RGB in M 2

<http://researchonline.ljmu.ac.uk/id/eprint/5320/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Lardo, C, Pancino, E, Mucciarelli, A and Milone, AP (2013) The discovery of an anomalous RGB in M 2. Memorie Della Società Astronomica Italiana, 84 (1). ISSN 0037-8720

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>



The discovery of an anomalous RGB in M 2

C. Lardo¹, E. Pancino^{2,3}, A. Mucciarelli¹, and A. P. Milone^{4,5}

¹ Department of Physics and Astronomy – University of Bologna, Via Ranzani 1, I-40127 Bologna, Italy e-mail: carmela.lardo2@unibo.it

² Istituto Nazionale di Astrofisica – Osservatorio Astronomico di Bologna, Via Ranzani 1, I-40127 Bologna, Italy

³ ASI Science Data Center, I-00044 Frascati, Italy

⁴ Instituto de Astrofísica de Canarias, E-38200 La Laguna, Tenerife, Canary Islands, Spain

⁵ Department of Astrophysics – University of La Laguna, E-38200 La Laguna, Tenerife, Canary Islands, Spain

Abstract. Using *UV* images taken with the Telescopio Nazionale Galileo, we discovered an anomalous sequence in the color-magnitude diagram of M 2. This feature appears as a narrow poor-populated red giant branch, which extends down to the sub giant branch region. We speculate that this new feature could be the extension of the faint component of the split sub giant branch recently discovered by Piotto et al. We identified in our *U, V* images two CH stars detected in previous studies. These stars, which are both cluster members, fall on this redder sequence, suggesting indeed that the anomalous RGB should have a peculiar chemical pattern. Unfortunately, no additional spectra were obtained for stars in this previously unknown substructure.

Key words. stars: abundances – stars: red giant branch – GCs: individual (M 2) – C-M diagrams

1. Introduction

In the last decade, a large collection of new spectroscopic and photometric data had conclusively determined that Globular Clusters (GCs) are neither coeval nor monometallic, reopening the issue of the formation of such systems. The formation of GCs is now schematized as a two-step process, during which the enriched matter from the more massive stars – both intermediate mass asymptotic giant branch stars (Ventura & D’Antona 2008) or fast rotating massive stars (Decressin et al. 2007) – from a first generation gives birth, in

the cluster innermost regions, to a subset of stars with the characteristic signature of fully CNO-processed material (see Gratton et al. 2012, and references therein). Variations in the light element abundances have been associated with the multiple sequences observed in the color-magnitude diagram (CMD), demonstrating that these two phenomena are intimately linked (e.g., Marino et al. 2008; Yong et al. 2008). It is not surprising that the abundance variations should have large effects on photometry, particularly in the *UV*-blue bands where CN, NH, and CH molecular bands can be dominant. Here, we exploit the property of *U*-band observation of tracing light-element spreads

Send offprint requests to: C. Lardo

and thus revealing multiple populations in the case of the poorly studied cluster M 2.

2. Observations and data reduction

We obtained images of the cluster in the standard Johnson U and V filters for a total of 540 s shifted in 3 single exposures in each filter with the DOLORES camera¹. The choice of passbands is due to the ability of separating photometric sequences at different evolutionary stages along the CMD (see Sects. 1 and 3). The raw frames were processed using the standard tasks in IRAF. Point spread function (PSF) fitting photometry was hence carried out with the DAOPHOT II and ALLSTAR packages. The photometric calibration was done using stars in common with Stetson Photometric standard field². Stars within 1' and outside of 4' from the cluster center are excluded from the CMD to reduce blending effects and the field star contamination, respectively. To select a sample of well-measured stars we have followed the procedure given in Lardo et al. (2012a), Sect. 5.1³.

3. The anomalous RGB in M 2

Besides the remarkable exception of ω Centauri, (see Johnson & Pilachowski 2010, and references therein), variations in the heavy element content have been detected only for few clusters. NGC 1851 and M 22 are among the best studied of all these. For these clusters, a bimodal distribution of s -process elements abundance have been identified (Yong & Grundahl 2008; Marino et al. 2012). The chemical inhomogeneity reflects itself in a complex CMD: multiple stellar groups in M 22 and NGC 1851 are also clearly manifest by a split in the sub giant branch (SGB) region (Piotto 2009; Milone et al. 2008) which appears to be related to chemical variations observed among red giant branch (RGB) stars

¹ DOLORES is a low resolution spectrograph and camera installed at Telescopio Nazionale Galileo (TNG) located in La Palma, Canary Islands.

² <http://www3.cadc-ccda.hia-ihp.nrc-cnrc.gc.ca/community/STETSON/standards/>

³ We imposed also the selection limits of $\text{CHI} < 2.0$ and $-1 < \text{SHARP} < 1$ photometric parameters.




Fig. 1. (a) U, V CMD from DOLORES images is shown in gray. Selected RGB stars are plotted as black dots, while the red continuous line is the fiducial obtained in the way described in the text. Panels (b) and (c) show the rectified RGB in function of the color difference and the histogram color distribution respectively (see the text for details).

(Marino et al. 2012; Lardo et al. 2012a). Indeed, carefully constructed CMDs —based on colors which include a blue filter (Han et al. 2009; Marino et al. 2012)— clearly reveal that the bright SGB is connected to the blue RGB, while red RGB stars are linked to the faint SGB. The split of the RGB discovered in the $U-I$ and $U-V$ colors for NGC 1851 and M 22 respectively, would not be detected in usual optical colors.

M 2 DOLORES photometry (see Fig. 1) displays an *anomalous* branch beyond the red edge of the main body of the RGB. The difference in color between stars belonging to this structure and *normal* RGB stars is quite large (of the order of 0.2-0.3 mags, well above the typical measurement errors) and extends down to the SGB region. There may be a second group of stars that are 0.3 mags redder with respect to this sequence and can possibly be more, anomalous RGB stars. We eliminated unphysical effects (i.e., differential reddening and field contamination) as the cause of this observed additional RGB (for details refer to




Fig. 2. (a) $u, (u-g)$ CMD from An et al. (2008) corrected photometry zoomed in around the RGB. Stars selected as red in Fig. 1 are plotted as red circles, while the red continuous line is the fiducial obtained in the way described in the text. Panels (b) show the color distribution in the $(u-g)$ color, while panel (c) show spatial distribution of the selected red stars.

Lardo et al. 2012b). To take photometric errors into due account, we follow the method described in Anderson et al. (2009). We considered the two independent CMDs obtained from DOLORES and An et al. (2008) photometry. In Fig. 1 we selected the portion of the RGB sequence with magnitudes between $14.5 \leq V \leq 18.5$ mag. In addition, we defined *bona fide* RGB members as the stars closer to the main RGB locus in the corrected DOLORES CMD (showed as black dots in panel (a) of Fig 1). The red continuous line in the same panel is the RGB fiducial obtained following as described in Milone et al. (2008). In brief, we drew a ridge line by putting a best-fit spline through the average color computed in successive short (0.2 mag) magnitude intervals. In panel (b) we subtracted from the color of each star the color of the fiducial at the same magnitude and plotted the V magnitude in function of this color difference; $\Delta(U-V)$. The histogram color distribution in logarithmic scale in panel (c) presents a clear substructure at the red end of the RGB, and we arbitrarily isolated RGB




Fig. 3. CMD for M2. The location of the carbon stars in the CMD is indicated by the open stars.

stars with $\Delta(U-V) > 0.15$ (red shaded region). These stars are plotted as red crosses in panel (b).

If the red branch we see was due to photometric errors, then a star redder than the RGB ridge line in the $V, (U-V)$ diagram would have the same probability of being bluer or redder in a different CMD obtained with different data. To this purpose, we identified the selected stars in u, g photometry (An et al. 2008) in Fig. 2 (red dots). The (a) panel shows a zoom around the RGB, again the red line is the fiducial defined as discussed above. In the following analysis, we considered only stars in common with the DOLORES photometry and, for the sake of homogeneity, we kept only stars between $1' < R < 4'$ from the cluster center. The fact that the histogram distributions of the selected RGB stars systematically have red colors demonstrates that we are seeing a *real* feature: no random or systematic errors can explain why the two distribution remain confined in the CMDs obtained from independent data-sets. Similar spatial distributions of stars on the bluer and redder RGBs also indicate that the differential reddening, if any, is not likely the cause of the double RGBs (see panel (c) in Fig. 2).

We found that the stars located on the anomalous red substructure account for only $\approx 4\%$ of the whole RGB population in this range of magnitude ($14.5 \leq V \leq 18.5$ mag). For comparison, $\approx 5\%$ of stars turn out to belong to the faint SGB if the two subpopulations have the same age and twice C+N+O and $\approx 3\%$ if they have the same C+N+O but differ of $\approx 1-2$ Gyr in age (Piotto et al. 2012). Moreover Piotto et al. (2012) claimed for this cluster the presence of a split SGB, with a fainter component less populous than the brighter one. We tentatively speculate that, also for M 2, this newly discovered double RGB might be photometrically connected to the split SGB, in close analogy to the case of NGC 1851 and M 22.

3.1. CH stars along the anomalous RGB

M 2 contains two CH stars, as discovered by Zinn (1981) and Smith & Mateo (1990). These stars show abnormally high CH absorption, together with deep CN bands, compared to other cluster giants. They are seen in dSph galaxies, and in the Galactic halo, but they are relatively rare within globular clusters. In Fig. 3 we identified the two CH stars in our $V, U - V$ photometry. Interestingly enough, both stars belong to the additional RGB, pointing out the anomalous chemical nature of this redder branch. Regardless of the exact classification of I-240 and I-451, it is apparent that the anomalous RGB contains a populations of giants which exhibit both a strong CN and strong G band. These stars may be the analogous to other CN and CH-strong RGB stars found in ω Cen, M 22 and NGC 1851 (Hesser et al. 1982). Given the peculiarity of other clusters that contain CH stars, it is of extreme interest to investigate the chemical pattern of stars in this red substructure.

4. Conclusions

Among the GCs with photometric evidence of multiple populations only NGC 1851 and M 22 display a bimodal SGB which is photometrically connected to the split RGB (Lardo et al. 2012a; Marino et al. 2012). The apparent similarity of M 2 to NGC 1851 and M 22 calls for a

deeper and complete spectroscopic characterization of stars in this anomalous RGB: only accurate measurements of metal abundances for a representative sample of stars will shed light on the origin of this poorly studied cluster.

Acknowledgements. Support for this work has been provided by the IAC (grant 310394), and the Education and Science Ministry of Spain (grants AYA2007-3E3506, and AYA2010-16717).

References

- An, D., Johnson, J. A., Clem, J. L., et al. 2008, *ApJS*, 179, 326
- Anderson, J., Piotto, G., King, I. R., Bedin, L. R., & Guhathakurta, P. 2009, *ApJ*, 697, L58
- Decressin, T., Charbonnel, C., & Meynet, G. 2007, *A&A*, 475, 859
- Gratton, R. G., Carretta, E., & Bragaglia, A. 2012, *A&A Rev.*, 20, 50
- Han, S.-I., Lee, Y.-W., Joo, S.-J., et al. 2009, *ApJ*, 707, L190
- Hesser, J. E., Bell, R. A., Harris, G. L. H., & Cannon, R. D. 1982, *AJ*, 87, 1470
- Johnson, C. I. & Pilachowski, C. A. 2010, *ApJ*, 722, 1373
- Lardo, C., Milone, A. P., Marino, A. F., et al. 2012a, *A&A*, 541, A141
- Lardo, C., Pancino, E., Mucciarelli, A., & Milone, A. P. 2012b, *A&A*, 548, A107
- Marino, A. F., Milone, A. P., Sneden, C., et al. 2012, *A&A*, 541, A15
- Marino, A. F., Villanova, S., Piotto, G., et al. 2008, *A&A*, 490, 625
- Milone, A. P., Bedin, L. R., Piotto, G., et al. 2008, *ApJ*, 673, 241
- Piotto, G. 2009, in *IAU Symposium*, Vol. 258, *IAU Symposium*, ed. E. E. Mamajek, D. R. Soderblom, & R. F. G. Wyse, 233–244
- Piotto, G., Milone, A. P., Anderson, J., et al. 2012, *ApJ*, 760, 39
- Smith, G. H. & Mateo, M. 1990, *ApJ*, 353, 533
- Ventura, P. & D’Antona, F. 2008, *MNRAS*, 385, 2034
- Yong, D. & Grundahl, F. 2008, *ApJ*, 672, L29
- Yong, D., Grundahl, F., Johnson, J. A., & Asplund, M. 2008, *apj*, 684, 1159
- Zinn, R. 1981, *ApJ*, 251, 52