



## LJMU Research Online

**Banat, IM, Diaz De Rienzo, MA and Quinn, GA**

**Microbial biofilms: biosurfactants as antibiofilm agents.**

<http://researchonline.ljmu.ac.uk/id/eprint/7241/>

### Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Banat, IM, Diaz De Rienzo, MA and Quinn, GA (2014) Microbial biofilms: biosurfactants as antibiofilm agents. Applied Microbiology and Biotechnology, 98 (24). pp. 9915-9929. ISSN 0175-7598**

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)

<http://researchonline.ljmu.ac.uk/>

# Microbial biofilms: biosurfactants as antibiofilm agents

Ibrahim M Banat<sup>1\*</sup>, Mayri A. Díaz De Rienzo and Gerry A. Quinn<sup>3</sup>

<sup>1</sup> School of Biomedical Sciences, University of Ulster, Coleraine, BT52 1SA, Northern Ireland, UK.

<sup>2</sup> School of Chemical Engineering and Analytical Science, University of Manchester, Manchester, UK.

<sup>3</sup> College of Medicine, Swansea University, Swansea, SA2 8PP, Wales, UK.

**Key words:** Biosurfactants, biofilm, disruption, antimicrobial, antibiofilm, dispersal agents, adjuvants.

\* Corresponding author email: [im.banat@ulster.ac.uk](mailto:im.banat@ulster.ac.uk)

## Abstract

1  
2 Current microbial inhibition strategies based on planktonic bacterial physiology have been  
3  
4 known to have limited efficacy on the growth of biofilms communities. This problem can be  
5  
6 exacerbated by the emergence of increasingly resistant clinical strains. All aspects of biofilm  
7  
8 measurement, monitoring, dispersal, control and inhibition are becoming issues of increasing  
9  
10 importance. Biosurfactants have merited renewed interest in both clinical and hygienic  
11  
12 sectors due to their potential to disperse microbial biofilms in addition to many other  
13  
14 advantages. The dispersal properties of biosurfactants have been shown to rival that of  
15  
16 conventional inhibitory agents against bacterial and yeasts biofilms. This makes them  
17  
18 suitable candidates for use in new generations of microbial dispersal agents and for use as  
19  
20 adjuvants for existing microbial suppression or eradication strategies. In this review we  
21  
22 explore aspects of biofilms characteristics and examine the contribution of biologically  
23  
24 derived surface-active agents (biosurfactants) to the disruption or inhibition of microbial  
25  
26  
27  
28  
29 biofilms.  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Introduction

1  
2 Microorganisms in general gravitate towards solid surfaces forming biofilms as a strategy to  
3 protect themselves from environmental challenges. Such deposition and subsequent biofilm  
4 formation is a phenomenon that happens naturally and is usually part of the microorganisms'  
5 strategy to protect themselves from external toxic factors (Pereira et al. 2007). They have  
6 the ability to sense their own cell density, communicate and behave as a population through  
7 cell to cell signalling, a phenomenon known as quorum-sensing (Liu et al. 2012). This  
8 behaviour has been documented for some time in microbial biofilm formations (Davies 2003)  
9 and is dependent on the nutritional/environment and the maturation stage of development of  
10 the microorganisms. Microbial biofilms represent a distinct bacterial physiology characterised  
11 by a multicellular phenotype that is fundamentally different from planktonic bacteria. They  
12 have been implicated in chronic and recalcitrant healthcare associated infections (Dowd et  
13 al. 2008), the dissemination of community acquired diseases (Stewart et al. 2012), effective  
14 hygienic processing, increased failure rate of anti-infective therapy (Bueno, 2014) and  
15 marine water and electronics environments (Lourenco et al. 2011). Biofilms that are  
16 composed of one species are relatively rare in the majority of the natural environment; rather  
17 microorganisms tend to be found in complex multispecies communities associated with  
18 surfaces (Stoodley et al. 2002).

19  
20 Until recently the differences between planktonic and biofilm physiologies seemed  
21 inconsequential. Standard bacterial inhibition tests were almost exclusively based on  
22 planktonic bacterial physiology and not the biofilm physiology even though these conditions  
23 were not readily observed in the natural environment. The standard planktonic bacterial  
24 physiology is typically exemplified by free-living single bacteria with optimal nutrition, gas  
25 exchange and agitation (typically 250rpm) (Bueno 2014; Kotulova and Slobodnikova, 2010).

26  
27 In contrast, the biofilm physiology has multicellular differentiation, multicellular  
28 communication, internal architecture and rudimentary fluid transport systems (Girard et al.  
29 2010; Leis et al. 2005). More importantly for in-vitro testing procedures, biofilms have  
30 variable levels of nutrients, gas exchange, little or no agitation and therefore slower growth.

1 This difference in bacterial physiology can be critical especially in clinical situations where  
2 there is a higher production of virulence factors in pathogens such as *Pseudomonas*  
3 *aeruginosa* (Croda-García et al. 2011). In the biofilm physiology these pathogens can be one  
4 to three orders of magnitude more resistant to dispersal/inhibition by conventional  
5 chemotherapy than their planktonic counterparts of the same species (Girard et al. 2010;  
6 Olson et al. 2002; Sepandj et al. 2004). This has been demonstrated in recent experiments  
7 on biofilm formation during peritoneal dialysis, where all the antibiotics tested were effective  
8 in laboratory MIC tests but (with the exception of gentamicin) lost their efficacy against  
9 *Staphylococcus aureus* biofilms (Girard et al. 2010). Globally, methicillin resistant  
10 *Staphylococcus aureus* (MRSA) is a serious problem due to limited efficacy of antibiotic  
11 options, hospital hygiene and the resistance of biofilm associated clinical strains (Samadi et  
12 al. 2012). Some biofilms also undergo phenotypic change as a result of chemotherapy  
13 resulting in increased resistance.

14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

New insights into biofilm physiology have now enabled researchers to design more effective bacterial inhibition/dispersal strategies. There are two main inhibitory strategies, based on the formulation of new antibiofilm compounds and the construction of biofilm resistant surfaces (Villa and Cappitelli, 2013).

Some of the most promising candidates for the inhibition of bacterial biofilms have come from biological surface-active agents (biosurfactants) (Kiran et al. 2010; Pradhan et al. 2013). Many of these have been reported to have anti-adhesive, antimicrobial and biofilm disruption properties (Rodrigues et al., 2006a,b,c; Rodrigues et al., 2007). Enzymatically synthesized surfactants such as lauryl glucose have also been reported to be effective against fungal and bacterial biofilms (Dusane et al. 2010).

Biosurfactants are a heterogeneous group of amphiphilic compounds produced mainly by microorganisms that accumulate at the interface between liquid phases and therefore reduce surface and interfacial tension. They have been recognised for some time in a diverse array of potential applications in a wide range of industries including agriculture, food, cosmetic, pharmaceutical and petroleum industries (Banat et al. 2010). The surface

1 and interfacial tension reducing properties of surfactants provide excellent detergency,  
2 emulsification, foaming and dispersing traits, making them some of the most versatile  
3 products in chemical processes (Desai and Banat 1997). They are highly sought after  
4 molecules due to their specificity, low toxicity, high biodegradability, widespread applicability  
5 and effectiveness at extremes of pH and temperature (Muthusamy et al. 2008).  
6  
7  
8  
9

10  
11 Several strands of research have demonstrated that under certain testing conditions  
12 biosurfactants can be more effective than many traditional biofilm inhibition and or disruption  
13 strategies (Epstein et al. 2011). There have been many reviews of biosurfactants and their  
14 potential applications in environmental and biomedical related areas (Neu 1996; Banat et al.  
15 2010; Banat et al. 2000). There has been however renewed interest in biosurfactants in  
16 relation to healthcare associated infections (Krasowska 2010). In addition, the rapid pace of  
17 advances in biofilm inhibition, control/disruption and the emergence of biofilms as potential  
18 reservoirs for the dissemination of disease have necessitated a review of the current state of  
19 the art on biofilms measurements and potentially effective biosurfactants against microbial  
20 biofilms.  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31

32  
33 To our knowledge the area of biofilms and role of biosurfactants within is becoming  
34 an increasingly important topic of research yet has not been the subject of a review article. In  
35 this review therefore we examine biofilms characteristics, monitoring and quantification and  
36 the main classes of current biosurfactants in use, their contribution to the dispersal or  
37 inhibition of biofilms, their scope and efficiency, quantification of this dispersal/inhibition and  
38 the sources and limitations of their uses.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48

### 49 **The nature and functions of biosurfactants**

50  
51 Biosurfactants are amphiphilic compounds of biological origin containing a hydrophilic region  
52 (polar or non-polar) and a hydrophobic region (lipid or fatty acid). The hydrophilic group is  
53 the base of the International Union of Pure & Applied Chemistry nomenclature i.e. those  
54 biosurfactants containing rhamnose are described as rhamnolipids; while those containing  
55 sophorose are sophorolipids and those generally containing a carbohydrate moiety including  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 the previously mentioned two types are described as glycolipids. Other lipopeptide  
2 biosurfactants contain a lipophilic hydrocarbon chain described as hydrophobic and a polar  
3 or hydrophilic part which is usually composed of a string of amino acids.  
4  
5  
6  
7

### 8 *Function* 9

10 Biosurfactants have been identified in many biological processes as the components of  
11 cellular metabolism, motion and defence. They are found in great abundance in bacteria, in  
12 biofilms, as quorum sensing molecules, lubricants, promoting the uptake of poorly soluble  
13 substrates, as immune modulators, virulence factors, secondary metabolites and  
14 antimicrobial compounds (Fracchia et al. 2012). In a review by Neu (1996) it has also been  
15 proposed that biosurfactants act as important molecules for interfacial processes,  
16 conditioning the microbial cell surface, interfaces and surfaces with which the  
17 microorganisms interact. These biosurfactants can be found in greater concentrations in the  
18 layers of cells associated with movement and hydration although they can have an  
19 intracellular location.  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31

32  
33 Biosurfactants' also have important roles in the dissolution and accessibility of oil  
34 molecules especially for oil degrading microorganisms; adhesion to hydrocarbons as a result  
35 of the emulsification of water-insoluble substrate compounds; the de-adhesion from  
36 interfaces; facilitating the in gliding of bacteria through wetting interfaces. Such surface  
37 active molecules can also have a role in enhancing the interaction between microorganisms  
38 and all the natural organic hydrophobic compounds interfaces including plant and animal  
39 derived polymeric compounds and microbial exopolysaccharides (Neu 1996). The role of  
40 bacterial biosurfactants has been extensively studied in *Pseudomonas* where they are  
41 known to promote colonisation and migration-dependent structural development (Pamp and  
42 Tolker-Nielsen 2007).  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53

54  
55 Other roles for biosurfactants including biocidal activity have been reported. This is  
56 mainly related to the effects of the lipidic moiety against eucaryotic cells. This has also been  
57 reported to lead to toxicity, lysis, pyrogenicity, mitogenicity and immunogenicity among other  
58  
59  
60  
61  
62  
63  
64  
65

1 effects (Wicken and Knox 1980). Lysis of red blood cells has been used as a selection  
2 criterion for microorganisms producing biosurfactants (Satpute et al. 2009). Finally human  
3 derived biosurfactants have recently received increased attention because of their role in  
4 immunity and defence (Gakhar et al. 2010).  
5  
6  
7  
8  
9

### 10 *Measurements of biosurfactants physical properties*

11 There are many methods employed to test physico-chemical properties of biosurfactants.  
12

13 These are very important for base line comparisons. The standard tests are based on the  
14 physical properties of biosurfactants such as measurement of reduced surface tension.  
15

16 Other tests measure the critical micelle concentration (CMC) which is the concentration of  
17 surfactants above which micelle formation occurs. The CMC for example of sodium dodecyl  
18 sulphate in water (with no other additives or salts) at 25°C and, atmospheric pressure, is  
19 8x10<sup>-3</sup>mol/l. The emulsification index (E24 or EI24) is another method used to characterize  
20 a biosurfactants' ability to form a stable emulsion with a hydrophobic phase. The hydrophilic  
21 phase in this instance is usually water which, can be mixed with Kerosene and the  
22 biosurfactant, shaken vigorously and allowed to stand for 24 hours. The percentage  
23 emulsion of the water solution in kerosene is reported as the E24 or EI24 (Desai and Banat  
24 1997). Other characterisation methods in use are the oscillating jets and the maximum  
25 bubble pressures measured in the presence of the surface active compounds.  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43

### 44 *Conditions for monitoring biofilm formation*

45 There is no standard laboratory method for quantifying biofilms though there are preferred  
46 methods.. In the past, planktonic bacterial inhibition assays have had to have strictly defined  
47 experimental criteria in order to reduce variation in results and increase confidence in  
48 antibiotic comparisons. However, these tests do not adequately represent different bacterial  
49 growth physiologies such as that in biofilms. The first biofilm tests were very similar to these  
50 planktonic experiments and created the impression that biosurfactants were weak  
51 counterparts of conventional inhibitory agents. Later, research into biofilm inhibition showed  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1 that these tests did not give an accurate reflection of the efficacy of biosurfactants. Today's  
2 biosurfactant tests are more accurate and try to represent the in-situ environment as much  
3 as possible. Many of these tests are based on pre-coating a surface with a known amount of  
4 biosurfactant overlaid with microbial biofilm (O'Toole 2011). This can be alternated with  
5 overlaying pre-existing biofilms with the test substance.  
6  
7  
8  
9

10  
11 Since the biofilm physiology is distinct from the planktonic physiology, biofilm  
12 experimental conditions have had to be adjusted accordingly. In terms of temperature the  
13 biofilm cultivation is carried out at the optimal temperature for biofilm growth of a particular  
14 species which may not be the same as the optimal temperature for planktonic growth, this  
15 could mean that biofilm cultivation may be at 20°C (even for clinical specimens) whilst others  
16 may be at 10°C in the cases of some environmental biofilms (Quinn et al. 2012).  
17  
18  
19  
20  
21  
22  
23

24 In terms of nutrition, it is common practice for biofilms to be cultivated in a dilution of the  
25 media that is used for planktonic cultivation, this is usually ½ to 1/5th of standard  
26 concentrations reflecting the sub-optimal conditions of biofilm growth, however this practice  
27 is not universally applicable (Stepanovic et al. 2004).  
28  
29  
30  
31  
32

33 Since biofilms also grow slower than optimised planktonic conditions, the typical cultivation  
34 period for biofilms can vary from 4 hours to 3-4 days or even 7-10 days in the case of slower  
35 growing environmental biofilms (Quinn et al. 2012; Stepanovic et al. 2007). Agitation  
36 considerations are equally important. In the earliest biofilm growth assays it was thought that  
37 environments of high shear stress were necessary. However more recent research has  
38 shown that environments of high agitation are not necessary for all biofilm growth and these  
39 growth conditions can be considered strain specific. Rather biofilm tests are typically  
40 conducted in almost static environments or environments of minimum perturbation (O'Toole  
41 2011; Stepanovic et al. 2007).  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

53 The standard inoculation density of microorganisms also differs greatly from standard  
54 planktonic tests. For planktonic MIC tests organisms are seeded at a density of  $1 \times 10^6$  /ml of  
55 fresh cells taken from the logarithmic stage of growth. In biofilm cultivation seeding densities  
56 are typically a 1/100 dilution of a stationary phase culture (McLaughlin and Hoogewerf 2006;  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Quinn et al. 2012). Some researchers use biofilm induction agents such as high glucose or alcohol to aid biofilm formation but these may add unknown variables to the assay making the final biofilm data difficult to interpret.

Recently, Lourenco and co-workers (2014) published the results of an initiative to establish “minimum information about a biofilm experiments” (MIABiE) which is a project partly funded by EU grants to find a scientifically adequate procedures to document biofilm-related data. They asserted that this could be achieved through ensuring a set of minimum information that should be reported to guarantee the independent verification and interpretation of experimental results in a way that would allow their integration with biofilm related information generated by other fields.

### **Surfaces for the quantification of biofilm growth**

The physiochemical properties of substrates used for biosurfactant evaluations can affect the nature of biofilm adhesion, the subsequent biofilm architecture in the case of monocultures or the selection of the microbial species which colonise in the case of mixed and environmental biofilms. Biofilms also express different repertoires of proteins or adhesion characteristics depending on the surface characteristics of the substrate they are attached to (Stoodley et al. 2002). Hence the choice of surface for biofilm cultivation is very important and must be taken into account even when comparing the results of inhibitory tests. .

The different surfaces used in biofilm tests range from animate/inanimate, rough/smooth, hydrophobic/ hydrophilic and liquid/air/liquid etc. Laboratory cultivation of biofilms can be conducted on many surfaces including glass, plastic, metal, silicone and tissue models (O'Toole 2011). In more comprehensive assessments of the inhibition potential biosurfactants can be applied to a broad range of surfaces especially in clinical environments. Research into the efficacy of Pseudofactin II (a newly characterised biosurfactant) used many different surfaces such as glass, polystyrene and silicone to cultivate biofilms in combination with different bacterial strains in order to demonstrate its

1 wide efficacy (Janek et al. 2010 & 2012). In other research on biofilms of *Salmonella*,  
2 investigators used PVC and silicone (urethral catheters) as biofilm substrates to demonstrate  
3 the applicability of biosurfactants in the reduction of biofilm formation/attachment (Mireles et  
4 al. 2001).  
5  
6  
7  
8  
9

## 10 **Quantification of biofilm inhibition/dispersal**

### 11 *The Calgary Biofilm Device*

12  
13 One of the first devices employed to measure biofilm inhibition/dispersal was the Calgary  
14 biofilm device (CBD) (Olson et al. 2002). This technique is widely used in flow tests for  
15 microbial biofilms (Rivardo et al. 2009; Girard et al. 2010; Rivardo et al. 2011). The  
16 cultivation chamber consists of a 96-well plate together with a lid that contains 96 peg  
17 projections (Figure 1). These pegs provide a maximum surface area for the growth of  
18 biofilms. The CBD has a typical seeding density of  $1 \times 10^4$  to  $1 \times 10^6$  bacteria per well or  
19 McFarland standard 1, a cultivation speed of  $\geq 10$  rpm and an incubation period of 4-24 h  
20 depending on species and conditions (Girard et al. 2010).  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

33 Microbial biofilms are cultivated on test pegs projecting into a growth media, removed  
34 after a given time, washed and then inserted into wells containing an inhibitory/test  
35 substance. Mature biofilms can be subsequently detached from the pegs by ultrasonic  
36 treatment. The detached microbes can be enumerated by standard cultivation techniques or  
37 quantified by measuring their optical density at 650nm. The amount of bacterial inhibition of  
38 the biofilms is referred to as the minimal biofilm eradication concentration (MBEC). The  
39 MBEC represents the lowest dilution of inhibitory substance. If cultivation conditions require  
40 a greater circulation of media the lid of this plate can be modified to accommodate 12 media  
41 channels into which the 96 pins are extended.. In this manner 96 pins can be simultaneously  
42 exposed to a given culture (Ceri et al 1999)  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54

55 Although the CBD was a welcome departure from planktonic based testing regimes  
56 and a step towards a more accurate portrayal of biofilm physiology; the method still relied on  
57 the final detection of viable planktonic microorganisms rather than directly measuring the  
58  
59  
60  
61  
62  
63  
64  
65

1 whole biofilm biomass. It also assumes that bacteria from viable biofilms can immediately  
2 rejuvenate on agar or directly into culture broth. This is an important point in biofilm  
3 physiology since studies on the resuscitation of bacterial cells have shown that some  
4 microorganisms may still be viable in the biofilm but not immediately cultivatable especially  
5 after prolonged chemotherapy (Rollet et al. 2009). This is also important when considering  
6 the negative impacts of the selective pressure of chemotherapy on biofilm forming  
7 pathogens. In some cases it has been shown that severe chemotherapy can induce a viable  
8 but dormant pathogen that can resuscitate in more favourable conditions to contribute to the  
9 chronic character of a biofilm infection (Zhang 2014).

10  
11 Finally, the CBD measures the amount of cells in a biofilm and not the biofilm  
12 biomass i.e. the biofilm + extra polymeric substances (EPS). However biofilm substances  
13 that are not cells can constitute a significant proportion of biofilms (Decho 2013).  
14  
15  
16  
17  
18  
19

#### 20 *Biofilm growth within flow-through devices*

21  
22 Biofilms can be analysed under flow conditions by a variety of methods including the CBD.  
23  
24 However another flow system currently used to test biofilms is the BioFlux 200 system  
25 (Fluxion Biosciences Inc., South San Francisco, CA) (Benoit et al. 2010; Ding et al 2014;  
26 Chabane et al. 2014). One of the benefits of such a system is that it is amenable to real-time  
27 analysis of the biofilm through automated image acquisition within specialized multi-well  
28 plates. In order to cultivate biofilms, microfluidic channels are primed with the culture  
29 medium at a specific rate. Each channel is seeded with an overnight culture with a cell  
30 density of  $10^7$  CFU. The biofilms are subsequently incubated at specific time and  
31 temperature levels in order for the bacterial cells to adhere. Once the biofilms have formed,  
32 planktonic cells are removed, and washed. The biofilm growth can then be recorded using a  
33 phase contrast or fluorescence microscope (Ding et al. 2014).  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57

#### 58 *In-vitro Biofilm formation in a 8 well chamber*

59  
60  
61  
62  
63  
64  
65

1  
2 Another variation of biofilm chamber growth is the use of an 8-well chamber slide. This  
3 method uses 200µL aliquots of mid-logarithmic cells diluted in fresh medium (1:2500 (v/v)).  
4 The medium can be replaced every 12h if the biofilm takes longer than 24h to grow or as  
5 needed to maintain bacterial viability (O'Toole 2011).  
6  
7

8  
9 The resulting biofilms can be visualized by aspirating the medium and washing with  
10 saline. The viability of the biofilm cells is typically assessed by the addition of BacLight  
11 Live/Dead stain (O'Toole 2011). Additionally, EPS or pili in the biofilms can be visualized  
12 under SEM by dehydrating the sample in a graded series of alcohols and adding  
13 hexamethyldisilazane (HMDS) (Araujo et al 2003).  
14  
15  
16  
17  
18  
19  
20  
21

## 22 **Crystal violet quantification of biofilm growth**

23  
24 One of the most commonly used methods to assess the effectiveness of biosurfactants and  
25 biofilm inhibitory agents is the crystal violet quantification of biofilm growth (O'Toole 2011).  
26 The technique involves the cultivation of a microbial biofilm in a 96-well (high bind PVC)  
27 plate, a rinsing step and final staining with 1% crystal violet. Biofilms are quantified by  
28 assessing the proportion of crystal violet bound to the biofilm biomass in control and test  
29 cultivations. The surfaces of high-bind 96 well plates were originally designed for ELISA  
30 tests and hence contain organically compatible high protein binding plastic (other types of  
31 PVC have different binding properties). This type of surface allows the binding of large  
32 molecules with ionic groups or large hydrophobic regions and permits a wide diversity of  
33 bacteria to form biofilms.  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45

46 The advantages of this method of biofilm quantification is that dispersal/inhibition can  
47 be measured directly in-situ rather than extrapolated from viable planktonic microorganisms.  
48 The crystal violet stains the total biofilm biomass which includes EPS and extracellular  
49 proteins rather than just its component cells. There may be some variability in the results  
50 obtained from this test but this can be rectified by a higher number of replicates which is  
51 afforded by the 96-well plate.  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Quantification of biofilm inhibition by direct analysis

One of the simplest methods of biofilm quantification is by direct measurements of bacterial viability as directly proportional to biofilm dispersal (Rodrigues et al. 2004). This technique does not measure total biofilm biomass or biofilm adhesion, however it is a useful validation step for other methods. This quantification becomes more problematic for mixed bacterial populations and viable but non-cultivable microorganisms.

### *Bacterial Viability Quantification*

There are several viability dyes that are used to quantify biofilm. Most of these are based on DNA binding. These include two of the most widespread fluorescent dyes, propidium iodide which binds to DNA when the cell nuclear membrane is damaged fluorescing red and syto 9 green which binds to DNA when the nuclear membrane is intact (Lehtinen et al. 2004). In the case of biofilms this quantification can be complicated by extracellular DNA but this might only apply in very dense biofilms.

### *Digital quantification*

Fluorescent stains are easily quantified by digital technologies. This makes it easier to assess biofilm growth/dispersal. As mentioned above although this technique may be directly quantitative for bacterial monolayers or biofilms of several layers thick, there are still technical issues however with proportional measurements of complex multi-layered biofilms with all the associated dead spaces and channels.

### *Other microscopic quantification*

Scanning electron microscopy (SEM) has proved to be a useful technique for pictorial representations of biofilms, however the preparation methods involved including successive dehydrations in alcohol and gold sputtering can fundamentally alter the composition and biofilm architecture of biofilms. More promising results have recently been obtained by the use of cryo-SEM (Alhede et al. 2012). As previously stated, the biofilm substrates used in

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

microscopic techniques have to be quite robust such as glass; however this may also have a role in determining the formation of the biofilm and cannot always be used in direct comparisons to the same biofilms growth on plastic or silicone.

### **Biosurfactants as antibiofilm molecules**

One of the most common questions posed on the effects of biosurfactants on biofilms is why are there still biofilms when biosurfactants are powerful molecules mostly leading to biofilm inhibition? The current hypothesis is that surface active molecules play a major role in the development and maintenance of biofilms partly through the maintenance of water channels through the biofilm which enhances nutrients movements and gaseous exchange, and which ultimately leads to the dissociation of parts of the biofilm into planktonic mobile forms (Marchant and Banat 2012). However the current focus of research is the ability of biosurfactants to disrupt established biofilms and prevention of the development of new ones. Although there are diverse arrays of biosurfactants, this review focuses on those in current use or known for the ability to disrupt biofilms in-vitro (Table 1).

#### *Lipopeptides biosurfactant as disruptor molecules*

Lipopeptides are one of the largest groups of biosurfactants that can effectively disperse microbial biofilms. These generally referred to by their group name although they can be composed of three or more varieties of homologous or congeners molecules. This group includes surfactins, polymixins, fengycins and fusaricidins (Krupovic et al. 2007; Pecci et al. 2010; Raza et al. 2009; Rivardo et al. 2009). Structurally lipopeptides are composed of a hydrophilic peptide attached to hydrophobic lipid or fatty acid. The peptides can either be aliphatic, branched or cyclic. Similarly, the lipids chains can vary in length and conformations ensuring a wide diversity of structures. Many of the current lipopeptides reported to inhibit/disperse biofilms originate from *Bacillus* or *Paenibacillus* (Kim et al. 2009; Price et al. 2007; Quinn et al. 2012).

## Polymyxins

1  
2 Polymyxins, are a class of non-ribosomally synthesized cyclic lipopeptides. They are  
3  
4 generally produced as secondary metabolites of *Bacillus* or similar species (Price et al.  
5  
6 2007). Their typical structure is that of a cyclic polypeptide attached to a fatty acid tail. They  
7  
8 can also contain exotic bacterial amino acids such as 2, 4, diaminobutyric acid (DAB) (Figure  
9  
10 2A). Polymyxins are known have a limited clinical spectrum of inhibition in the treatment of  
11  
12 Gram negative infections. There are several commercially available formulations of  
13  
14 polymyxins including Colistin (polymyxin E) (Falagas and Kasiakou 2005), Neosporin and  
15  
16 Polymyxin B which can be supplied as polymyxin B sulphate (a mixture of polymixins) (He et  
17  
18 al. 2010). Polymyxin can also be combined with trimethoprim for eye conditions (Polytrim)  
19  
20 and with neomycin and bacitracin to make triple antibiotic ointment Neosporin.  
21  
22  
23

24  
25 Polymyxins are the last drug of choice in some infections and are often prescribed  
26  
27 with caution due to fears of their toxicity however this estimation has been reappraised in the  
28  
29 light of more rigorous testing (Arnold et al. 2007). Polymyxins are prescribed intestinally or  
30  
31 topically as cream or powders in most cases of multi drug resistant *Acinetobacter baumannii*,  
32  
33 *Klebsiella pneumoniae*, and *Pseudomonas aeruginosa* (Falagas and Kasiakou 2006; Milletti  
34  
35 Sezgin et al. 2012). Polymyxin has been reported to reduce biofilms of *Pseudomonas*  
36  
37 *aeruginosa* at concentrations of 20µg/ml by 99% in a 12 hour time period and almost  
38  
39 completely over 24 hours (Jass and Lappin-Scott 1996). However these results are based  
40  
41 on the viability of bacteria and not their dispersal; although it was noted that bacterial cells  
42  
43 displayed an altered morphology. Polymyxin E (colistin) is recommended as an early  
44  
45 aggressive therapy to delay the onset of chronic *P. aeruginosa* infection (which frequently  
46  
47 forms biofilms) or intermittent colonization in cystic fibrosis patients, a combination of oral  
48  
49 ciprofloxacin with colistin inhalation (Doring et al. 2000).  
50  
51  
52

53  
54 Polymyxin D1 has been shown to be effective against mixed bacterial biofilms,  
55  
56 however our earlier work has shown this compound was found in combination with  
57  
58 fusaricidin and surfactin in undefined ratio's (Quinn et al. 2012). This complex of  
59  
60 biosurfactants was also reported to inhibit the formation of biofilms of both Gram positive  
61  
62  
63  
64  
65



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

bacteria such as *Staphylococcus aureus*, *Streptococcus bovis*, *Bacillus subtilis* and *Micrococcus luteus* and Gram negative bacteria such as *Pseudomonas aeruginosa*. Most interestingly the biosurfactants were able to inhibit the formation of mixed species biofilms such as self-assembling marine biofilm (SAMB) in co-incubation assays by 99.3% and disrupt previously established mixed SAMB by 72.4% (Quinn et al. 2012).

The mechanism of action of polymyxins on bacterial biofilms remains largely undefined. However the mechanism of action on planktonic bacteria is proposed to be related to their high affinity for lipopolysaccharide (LPS) (Domingues et al. 2012). This induces LPS aggregation increasing the surface charge of LPS leading to internalization and binding to the bacterial phosphatidylglycerol-rich membrane leaflets which in turn induces leakage of cellular contents (Domingues et al. 2012).

#### *Fengycin-like lipopeptides*

Fengycin-like lipopeptides are derived from *Bacillus subtilis* and *Bacillus licheniformis*. These are cyclic lipopeptides containing 8-10 amino acids linked to a beta hydroxy fatty acid (Figure 2B). Fengycin-like peptides have also been reported to be involved in the inhibition of biofilms (Xu et al. 2013) causing up to 90% dispersion of Gram positive *S. aureus* biofilms and up to 97% dispersion of Gram negative *E. coli* biofilm (Rivardo et al. 2009) .

#### *Putisolvin*

Putisolvin is a cyclic lipodepsipeptide isolated from *Pseudomonas putida*. This has been characterised in two forms, putisolvin I and putisolvin II. This biosurfactant has a four member cyclic peptide; the valine residue in putisolvin I being substituted by a leucine or isoleucine in putisolvin II. (Figure 2C ) (Dubern et al. 2006). Although Putisolvin is involved in biofilm formation by *Pseudomonas putida* these surfactants have also been shown to be effective dispersal agents in pre- and post-addition to biofilms of other *Pseudomonas* sp. strains (Kuiper et al. 2004).

### *Pseudofactin*

1  
2 Pseudofactin is a cyclic lipodepsipeptide derived from *P. fluorescens*. The structure of  
3  
4 Pseudofactin is based on that of a palmitic acid attached to the terminal amino group of an  
5  
6 eight amino acid peptide chain. The C-terminal carboxylic group of the last amino acid forms  
7  
8 a lactone with the hydroxyl of third amino acid which is a threonine (Figure 2D). Pseudofactin  
9  
10  
11 It has been reported to be 36-90% effective against the adhesion of five species of bacterial  
12  
13 biofilms on glass, polystyrene and silicone substrates. These strains include *Enterococcus*  
14  
15 *faecalis*, *Escherichia coli*, *Staphylococcus epidermidis*, *Enterococcus hirae* and *Proteus*  
16  
17 *mirabilis*. Similar inhibition of adhesion (92-99%) was reported on yeast biofilms of *Candida*  
18  
19 *albicans* at concentrations of 0.5 mg/ml (Janek et al. 2012).  
20  
21

22 Pseudofactin has been documented to produce an effective dispersal of 26-70% on  
23  
24 pre-existing biofilms grown on untreated surfaces and has been shown to cause a marked  
25  
26 inhibition of the initial adhesion of *E. hirae*, *E. coli*, *E. faecalis* and *C. albicans* to silicone  
27  
28 urethral catheters. Total growth inhibition of *S. epidermidis* has been observed at the  
29  
30 highest concentration tested (0.5 mg/ml), which causes a partial (18-37%) inhibition of other  
31  
32 bacteria, a 8-9% inhibition of *C. albicans* yeast growth and a 99% prevention of adhesion  
33  
34 (Janek et al. 2012).  
35  
36  
37  
38  
39

### *Surfactin*

40  
41  
42 Surfactins are one of the most powerful biosurfactants originally isolated from *Bacillus*  
43  
44 *subtilis* and consist of a cyclic peptide heptamer connected to a 13-15 carbon, beta-hydroxy  
45  
46 fatty acid chain (Figure 3A). Unfortunately surfactins can also be indiscriminately cytotoxic  
47  
48 with haemolytic activities due to its interactions with cellular membranes (D'Auria et al.  
49  
50 2013). They have been reported to inhibit the growth of biofilms of *Salmonella* sp. cultivated  
51  
52 on PVC microtitre wells and urethral catheters (Mireles et al. 2001). They have been  
53  
54 observed to cause a rippling effect in lipid bilayers perhaps indicating a clue to the  
55  
56 mechanism of biosurfactant action or biofilm permeability or integrity (Brasseur et al. 2007)  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 most likely through the formation of some kind of channels within the biofilm increasing  
2 penetrability.  
3  
4  
5

### 6 *Complexes of lipopeptides*

7  
8 Although many lipopeptides have been characterised for experimental purposes as pure  
9 compounds they are in fact usually associated with groups of similar compounds. This is  
10 reflected in their availability as minimally purified preparations. Siram and co-workers (2011)  
11 reported on one such complex of lipopeptide biosurfactants produced by a heavy metal  
12 tolerant strain of *Bacillus cereus*. This surfactant effectively dispersed biofilms at an active  
13 dose of 0.150µg and was noted to be very tolerant of fluxes in pH, temperature and NaCl, in  
14 addition to being resistant to high levels of iron, lead and zinc whilst maintaining  
15 antimicrobial and biofilm dispersal activity. Another complex of surfactants isolated from  
16 *Paenibacillus polymyxa*. PPE was found to consist of polymyxin D1, fusaricidin B and traces  
17 of surfactin (Deng et al. 2011; Quinn et al. 2012) (Figure 3 B & C).  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

31 A preparation containing 2mg/ml of such lipopeptides tested in one of our  
32 laboratories inhibited (87-98%) the formation of many Gram positive bacterial biofilms such  
33 as *Staphylococcus aureus*, *Streptococcus bovis*, *Micrococcus luteus*, *Bacillus subtilis* and  
34 also some Gram negative bacteria such as *Pseudomonas aeruginosa* (Quinn et al. 2012).  
35 More uniquely in terms of biofilm experiments, this combination of lipopeptides was effective  
36 against mixed environmental strains' biofilms formation (99% inhibition) and up to 74% in  
37 pre-established biofilms.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

### 48 *Synergy of lipopeptides with other inhibitors*

49 Lipopeptide biosurfactants have been combined with conventional antibiotics in an effort to  
50 produce synergistic inhibition effects. Lipopeptides isolated from *Bacillus licheniformis*  
51 (strain V9T14) were reported by Rivardo and co-workers (2011) to have a synergistic effect  
52 against a mature 24-h uropathogenic *E. coli* (CFT073) biofilms when combined with  
53 ciprofloxacin, cefazolin, piperacillin, ceftriaxone, ampicillin, tobramycin and  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 trimethoprim/sulfamethoxazole. They concluded that some combinations led to total  
2 eradication of biofilm; however the antibiotics on their own had poor inhibitory activity  
3  
4 (Rivardo et al 2011).  
5  
6  
7

### 8 *Glycolipid biosurfactants as antibiofilm molecules*

9

10 Glycolipids consist of a carbohydrate attached to aliphatic or hydroxy-aliphatic acid. These  
11 are one of the most studied groups of biosurfactants in other fields although they are  
12 underrepresented as agents of biofilm dispersal.  
13  
14  
15  
16  
17

### 18 *Rhamnolipids*

19

20 Rhamnolipids consist of di- or mono-rhamnose sugars attached to a fatty acid chain (Figure  
21 3 D & E). Originally isolated from *Pseudomonas aeruginosa*, analogues are also produced  
22 by isolates of *Burkholderia* (Costa et al. 2011), *Renibacterium salmoninarum*, *Cellulomonas*  
23 *cellulans*, *Nocardioides* and *Tetragenococcus koreensis* (Abdel-Mawgoud et al. 2010).  
24 Rhamnolipids have been reported as a potential replacement to chemical surfactants for  
25 many uses in the oil and petroleum industries and in use for the bioremediation of oil  
26 contaminated environments (Marchant and Banat 2012a & b). They are frequently cited as  
27 inhibitors of bacterial growth although their capacity to inhibit biofilms however has not been  
28 as extensively documented.  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41

42 Rhamnolipids are involved in biofilm formation in *Pseudomonads* sp. through the  
43 promotion of motility, the inhibition of attachment and degradation of the matrix maintaining  
44 channels throughout the biofilm for movement of water and oxygen (Boles et al. 2005; Davey  
45 et al. 2003). These biosurfactants were previously reported as antibacterial against  
46 *Staphylococcus aureus*, *Mycobacterium* sp, *Bacillus* sp, *Serratia marsecens*, *Enterobacter*  
47 *aerogenes*, *Klebsiella pneumonia* and against fungi such as *Chaetomium globosum*,  
48 *Aureobacidium pullulans*, *Gliocladium virens*, *Botryhs cinerea* and *Rhizoclonia solanii*  
49 (Benincasa et al. 2004; Haba et al. 2003).  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2 Rhamnolipids have also been shown to be effective against biofilms of *Bordetella*  
3 *bronchiseptica* (Irie et al. 2005). The mechanism of biofilm inhibition is thought to be by the  
4 detachment of cells; however some unattached cells may still be viable. They have been  
5 reported to disrupt pre-formed biofilms such as *Bacillus pumilus* from the marine  
6 environment (on polystyrene microplates) resulting in a dispersal at sub-MIC concentrations  
7 and confirming ability to remove pre-formed biofilms (Dusane et al. 2010). This was  
8 corroborated by scanning electron microscopy which showed that rhamnolipids removed  
9 biofilm-matrix components (Dusane et al. 2010). The effects of rhamnolipids on pre-formed  
10 biofilms of *P. aeruginosa* PAO1 generated in our laboratory can be seen here in Figure 4.  
11  
12  
13  
14  
15  
16  
17  
18  
19

20 Rhamnolipids were also been tested on devices such as voice prostheses and have  
21 been noted to reduce the initial deposition rates of biofilm after 4h (Rodrigues et al. 2006a).  
22 A maximum reduction of adhesion ( $\approx 66\%$ ) was observed when the surfaces such as silicone  
23 rubber had been preconditioned with rhamnolipids using biofilms of *Streptococcus salivarius*  
24 and *Candida tropicalis*. The number of cells adhering after 4h was reduced to  $\approx 48\%$  for  
25 *Staphylococcus epidermidis*, *Streptococcus salivarius*, *Staphylococcus aureus* and *C.*  
26 *tropicalis* in comparison to controls. This group managed to optimise the actions of this  
27 biosurfactant on the detachment of microorganisms adhering to silicone rubber by perfusing  
28 the flow chamber with a biosurfactant containing solution followed by passage at the liquid-  
29 air interface. By this method they were able to achieve a high detachment (96%) for most of  
30 the microbial cells.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43

44 Rhamnolipids have also been shown to be effective dispersal agents for fungi  
45 disrupting pre-formed biofilms of *Yarrowia lipolytica* on glass surfaces by  $\approx 67\%$  which was  
46 more effective in comparison to the surfactants cetyl-trimethyl ammonium bromide (CTAB)  
47 and sodium dodecyl sulphate (SDS) (Dusane et al. 2012).  
48  
49  
50  
51  
52

53 It is important to note that although rhamnolipids can effectively disrupt biofilm  
54 formation and integrity which we observed through phase contrast microscopy where thick  
55 dense cellular biofilm (Figure 4A) of microcolonies structures on glass coverslips stained  
56 with crystal violet was much reduced in after treatment with rhamnolipids biosurfactants  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 (Figure 4B). These molecules are also known to be extracellular virulence factors and  
2 related to the pathogenesis (infection procedure) in *Pseudomonas aeruginosa*. It has been  
3  
4 noted that rhamnolipids are also linked to increased lung epithelial permeability, rapid  
5  
6 necrotic killing of polymorphonuclear leukocytes and the malfunction of normal tracheal  
7  
8 ciliary motion in the respiratory system of infected patients (Read et al. 1992).  
9

### 10 11 12 *Sophorolipids*

13  
14 Sophorose lipids are typical glycolipids biosurfactants consisting of a dimer of sophorose  
15  
16 sugar and a long-chain fatty acid that are produced by yeasts belonging to the genus  
17  
18 *Candida*.  
19

20  
21  
22 The synergy between sophorolipids and antibiotics has been studied as potential  
23  
24 strategy to disrupt biofilms using The LIVE/DEAD BacLight Bacterial Viability Kits as a  
25  
26 method for detection . This method employs two nucleic acid stains — the green-fluorescent  
27  
28 SYTO 9® stain and the red-fluorescent propidium iodide stain. These stains differ in their  
29  
30 ability to penetrate healthy bacterial cells. When used alone, SYTO 9 stain labels both live  
31  
32 and dead bacteria. In contrast, propidium iodide penetrates only bacteria with damaged  
33  
34 membranes, reducing SYTO 9 fluorescence when both dyes are present. Thus, live bacteria  
35  
36 with intact membranes fluoresce green, while dead bacteria with damaged membranes  
37  
38 fluoresce red. Joshi-Navare and Prabhune (2013) reported the the effect of sophorolipids in  
39  
40 the disruption of biofilms from *Escherichia coli*. Figure 5 illustrated the examination of cells  
41  
42 of *Bacillus subtilis* attached to coverslips after 48h and stained with LIVE/DEAD BacLight  
43  
44 showing the presence of individual bacteria, small clusters of cells (microcolonies), and  
45  
46 extended areas of the glass surface covered with large numbers of microcolonies of active  
47  
48 cells (Figure 5A), as well as, those which their membrane was damage due to the effect of  
49  
50 sophorolipids 5% (v/v) concentration after 30min of treatment (Figure 5B).  
51  
52  
53  
54  
55  
56  
57

### 58 *Other Glycolipids as antibiofilm molecules*

59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Dusane et al. (2012) reported a glycolipid based on glucose and palmitic acid produced by a tropical marine *Serratia marcescens* was effective in inhibiting biofilms of the marine biofouling bacterium *Bacillus pumilus* and the adhesion of *C. albicans* and *P. aeruginosa* PAO1. This effect was also observed with preformed biofilms of these cultures on microtitre plate tests. Other complexes of glycolipids from *Brevibacterium casei* MSA19 have been reported to disrupt and significantly inhibit individual and mixed culture biofilms of human and fish at concentrations of 30 mg/ml (Kiran et al. 2010).

Antibiofilm glycolipids have also been isolated from *Lactobacillus* (Tahmourespour et al. 2011; Zakaria Gomaa 2013). In this case *L. paracasei* A20 produced biosurfactants that inhibit Gram-positive and Gram-negative bacteria, yeasts and filamentous fungi (Gudina et al. 2010). The biosurfactant also showed anti-adhesive activity against pathogenic *Candida albicans*, *Escherichia coli*, *Staphylococcus aureus*, *Staphylococcus epidermidis* and *Streptococcus agalactiae*. Glycolipids derived from plants have also been reported to inhibit biofilms. These include a novel hydroxyproline rich glycopeptide from the pericarp of *Datura stramonium* known as datucin which is also reported to eradicate biofilms of antifungal resistant *Candida albicans* (Mandal 2012).

### *Complex surfactant mixtures*

Biosurfactants are seldom found in pure form or isolation and are often associated together with isomers or congeners that share similar physiochemical characteristics which makes the process of purification either exhaustive or uneconomical. However these complexes of biosurfactants may have the advantage of a broader applicability than pure compounds. The same is true of complexes of compounds in other environments; this can be illustrated by the large diversity of antimicrobial peptides and surfactants found on the skin of amphibians (Bevins and Zasloff 1990). Similarly in innate human defence, antimicrobial peptides such as human beta defensins 1 , 2 and 3 and related human neutrophil peptides (Ganz et al. 1985) are found in homogenous groups. .

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Combinations of biosurfactants have also been extracted from *Robinia pseudoacacia* and *Nerium oleander*. These secretions inhibited attachment of biofilms of *Candida albicans* on silicon and denture prosthesis at concentrations of 78µg/ml and 156µg/ml (Cochis et al. 2012). Other biosurfactants obtained from probiotic bacteria *Lactococcus lactis* 53 and *Streptococcus thermophilus* greatly reduced microbial numbers on preconditioned voice prostheses in an artificial throat model and induced a decrease in the airflow resistance that occurs on voice prostheses after biofilm formation (Gakhar et al. 2010).

### *Biosurfactants from fungi*

Biosurfactants that inhibit biofilms have been found in fungi such as *Candida bombicola*. This produces sophorolipids that inhibit biofilms of *V. cholerae* (Mukherji and Prabhune 2014). Other strains of yeast such as *Candida sphaerica* have also been reported to produce biosurfactants such as lunasan (Luna et al. 2011). This inhibits the adhesion of *Pseudomonas aeruginosa*, *Streptococcus agalactiae* and *Streptococcus sanguis* to levels between 80-92%. Similarly rufisan from *Candida lipolytica* inhibits biofilm formation at concentrations greater or equal to 0.75 µg/ml against *S. aureus*, *Streptococcus agalactae*, *S. mutans* NS (Rufino et al. 2011).

### *Mammalian surface active secretions*

From a chemotherapeutical perspective, the most interesting groups of biosurfactants are those produced by humans. Not much is known about these molecules however it has recently been reported that PLUNC ("Palate, lung, nasal epithelium clone") protein has anti-biofilm activity (Gakhar et al. 2010). These molecules are mainly produced as a secretory product of epithelia lining the airways tubes within mammals including humans. They are evolutionarily related to the lipid transfer/lipopolysaccharide binding protein (LT/LBP) family. PLUNC are believed to have novel biologically relevant surface active properties as they significantly reduce surface tension at the air-liquid interface within aqueous solutions they



1  
2 also inhibited biofilm formation in the airways colonising potential pathogen *Pseudomonas*  
3 *aeruginosa in vitro* at physiologically relevant concentrations (Gakhar et al. 2010).  
4  
5

## 6 **Conclusions**

7  
8  
9

10 It has been acknowledged that microbial biofilms lie at the heart of many recalcitrant patient  
11 infections in the clinical environment, the dissemination of airborne pathogens and the  
12 fouling of industrial surfaces. These problems are increasingly exacerbated by the rise of  
13 resistant biofilm populations and the paucity of alternative eradication solutions.  
14  
15 Biosurfactants represent an emerging therapy which has inherent anti-bacterial, fungal and  
16 viral properties with an ability to effectively disperse or disrupt such biofilms. Their use  
17 therefore either on their own or as adjuvants to other antimicrobial chemotherapies may  
18 represent a potential way forward in tackling biofilms in the future.  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## References

- 1  
2  
3 Abdel-Mawgoud AM, Lepine F, Deziel E (2010) Rhamnolipids: diversity of structures,  
4 microbial origins and roles. *Appl Microbiol Biotechnol* 86:1323-1336  
5  
6  
7 Alhede M, Qvortrup K, Liebrechts R, Hoiby N, Givskov M, Bjarnsholt T (2012) Combination  
8 of microscopic techniques reveals a comprehensive visual impression of biofilm  
9 structure and composition. *FEMS Immunol Med Microbiol* 65: 335-342  
10  
11  
12 Araujo JC, Téran FC, Oliveira RA, Nour EA, Montenegro MA, Campos JR, Vazoller RF  
13 (2003) Comparison of hexamethyldisilazane and critical point drying treatments for SEM  
14 analysis of anaerobic biofilms and granular sludge. *J Electron Microsc (Tokyo)* 52:429-  
15 33  
16  
17  
18  
19  
20 Arnold TM, Forrest GN, Messmer KJ (2007) Polymyxin antibiotics for gram-negative  
21 infections. *American J Health-system Pharmacy* 64: 819-826  
22  
23  
24 Banat IM, Franzetti A, Gandolfi I, Bestetti G, Martinotti MG, Fracchia L, Smyth TJ, Marchant  
25 R, (2010) Microbial biosurfactants production, applications and future potential. *Appl*  
26 *Microbiol Biotechnol* 87:427-444  
27  
28  
29  
30 Banat IM, Makkar RS, Cameotra SS (2000) Potential commercial applications of microbial  
31 surfactants. *Appl Microbiol Biotechnol* 53:495-508  
32  
33  
34 Benoit MR, Conant CG, Lonescu-Zanetti C, Schwartz M, Martin A. (2010) New Device for  
35 High-Throughput Viability Screening of Flow Biofilms. *App Environ Microbiol* 76: 4136-  
36 42  
37  
38  
39  
40 Benincasa M, Abalos A, Oliveira I, Manresa A (2004) Chemical structure, surface properties  
41 and biological activities of the biosurfactant produced by *Pseudomonas aeruginosa* LBI  
42 from soapstock. *Antonie Van Leeuwenhoek* 85:1-8  
43  
44  
45  
46 Bevins CL, Zasloff M (1990) Peptides from frog skin. *Annu Rev Biochem* 59:395-414  
47  
48  
49 Boles BR, Thoendel M, Singh PK (2005) Rhamnolipids mediate detachment of  
50 *Pseudomonas aeruginosa* from biofilms. *Mol Microbiol* 57:1210-1223  
51  
52  
53 Brasseur R, Braun N, El Kirat K, Deleu M, Mingeot-Leclercq MP, Dufrene YF (2007) The  
54 biologically important surfactin lipopeptide induces nanoripples in supported lipid  
55 bilayers. *Langmuir* 23:9769-9772  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- Bueno J (2014) Anti-Biofilm Drug Susceptibility Testing Methods: Looking for New Strategies against Resistance Mechanism. *J Microbial Biochem Technol* S3: 004. doi: 10.4172/1948-5948.S3-004
- Ceri H, Olson ME, Stremick C, Read RR, Morck D, Buret A (1999) The Calgary Biofilm Device: new technology for rapid determination of antibiotic susceptibilities of bacterial biofilms. *J Clin Microbiol.* 37:1771-1776
- Chabane NY, Mlouka MB, Alexandre S, Nicol M, Marti S, Pestel-Caron M, Vila J, Jouenne T, Dé (2014) Virstatin inhibits biofilm formation and motility of *Acinetobacter baumannii*. *BMC Microbiol* 14:62
- Cochis A, Fracchia L, Martinotti MG, Rimondini L (2012) Biosurfactants prevent in vitro *Candida albicans* biofilm formation on resins and silicon materials for prosthetic devices. *Oral Surg Oral Med Oral Pathol Oral Radiol* 113:755-761
- Croda-García G, Grosso-Becerra V, Gonzalez-Valdez A, Servín-González L, Soberón-Chávez G (2011) Transcriptional regulation of *Pseudomonas aeruginosa* rhlR: role of the CRP orthologue Vfr (virulence factor regulator) and quorum-sensing regulators LasR and RhIR. *Microbiol.* 157:2545-55
- Costa SG, Deziel E, Lepine F (2011) Characterization of rhamnolipid production by *Burkholderia glumae*. *Lett Appl Microbiol* 53:620-627
- D'Auria L, Deleu M, Dufour S, Mingeot-Leclercq MP, Tyteca D (2013) Surfactins modulate the lateral organization of fluorescent membrane polar lipids: a new tool to study drug : membrane interaction and assessment of the role of cholesterol and drug acyl chain length. *Biochim Biophys Acta* 1828:2064-2073
- Davey ME, Caiazza NC, O'Toole GA (2003) Rhamnolipid surfactant production affects biofilm architecture in *Pseudomonas aeruginosa* PAO1. *J Bacteriol* 185:1027-1036
- Davies D (2003) Understanding biofilm resistance to antibacterial agents. *Nat Rev Drug Discov* 2:114-122
- Decho AW (2013) The EPS matrix as an adaptive bastion for biofilms: introduction to special issue. *Int J Mol Sci* 14:23297-23300
- Deng Y, Lu Z, Bi H, Lu F, Zhang C, Bie X (2011) Isolation and characterization of peptide antibiotics LI-F04 and polymyxin B6 produced by *Paenibacillus polymyxa* strain JSa-9. *Peptides* 32:1917-1923

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- Desai JD, Banat IM (1997) Microbial production of surfactants and their commercial potential. *Microbiol Mol Biol Rev* 61:47-64
- Ding Y, Wang W, Fan M, Tong Z, Kuang R, Jiang W, Ni L (2014) Antimicrobial and anti-biofilm effect of Bac8c on major bacteria associated with dental caries and *Streptococcus mutans* biofilms. *Peptides* 52:61-7
- Domingues MM, Inacio RG, Raimundo JM, Martins M, Castanho MA, Santos NC (2012) Biophysical characterization of polymyxin B interaction with LPS aggregates and membrane model systems. *Biopolymers* 98:338-344
- Doring G, Conway SP, Heijerman HGM, Hodson ME, Hoiby N, Smyth A, Touw DJ (2000) Antibiotic therapy against *Pseudomonas aeruginosa* in cystic fibrosis: a European consensus, *Eur Respiratory J.* 16: 749-767
- Dowd SE, Wolcott RD, Sun Y, McKeehan T, Smith E, Rhoads D (2008) Polymicrobial Nature of Chronic Diabetic Foot Ulcer Biofilm Infections Determined Using Bacterial Tag Encoded FLX Amplicon Pyrosequencing (bTEFAP). *PLoS ONE* 3:e3326
- Dubern JF, Lugtenberg BJ, Bloemberg GV (2006) The ppul-rsaL-ppuR quorum-sensing system regulates biofilm formation of *Pseudomonas putida* PCL1445 by controlling biosynthesis of the cyclic lipopeptides putisolvins I and II. *J Bacteriol* 188:2898-2906
- Dusane DH, Dam S, Nancharaiah YV, Kumar AR, Venugopalan VP, Zinjarde SS (2012) Disruption of *Yarrowia lipolytica* biofilms by rhamnolipid biosurfactant. *Aquat Biosyst* 8:17
- Dusane DH, Nancharaiah YV, Zinjarde SS, Venugopalan VP (2010) Rhamnolipid mediated disruption of marine *Bacillus pumilus* biofilms. *Colloids Surf B Biointerfaces* 81:242-248
- Dusane DH, Pawar VS, Nancharaiah YV, Venugopalan VP, Kumar AR, Zinjarde SS (2011) Anti-biofilm potential of a glycolipid surfactant produced by a tropical marine strain of *Serratia marcescens*. *Biofouling* 27:645-654
- Epstein AK, Pokroy B, Seminara A, Aizenberg J (2011) Bacterial biofilm shows persistent resistance to liquid wetting and gas penetration. *Proc Natl Acad Sci U S A* 108:995-1000
- Falagas ME, Kasiakou SK (2005) Colistin: the revival of polymyxins for the management of multidrug-resistant gram-negative bacterial infections. *Clin Infect Dis* 40:1333-1341

- 1  
2  
3  
4 Falagas ME, Kasiakou SK (2006) Toxicity of polymyxins: a systematic review of the  
5 evidence from old and recent studies. Crit Care 10:R27  
6  
7  
8  
9  
10 Fracchia L, Cavallo M, Martinotti MG, Banat IM (2012) Biosurfactants and Bioemulsifiers  
11 Biomedical and Related Applications – Present Status and Future Potentials. In:  
12 Biomedical Science, Engineering and Technology. pp 325-370  
13  
14  
15  
16 Gakhar L, Bartlett JA, Penterman J, Mizrachi D, Singh PK, Mallampalli RK, Ramaswamy S,  
17 McCray PB (2010) PLUNC is a novel airway surfactant protein with anti-biofilm activity.  
18 PLoS ONE 5:e9098  
19  
20  
21 Ganz T, Selsted ME, Szklarek D, Harwig SS, Daher K, Bainton DF, Lehrer RI (1985)  
22 Defensins. Natural peptide antibiotics of human neutrophils. J Clin Invest 76:1427-1435  
23  
24  
25  
26 Girard LP, Ceri H, Gibb AP, Olson M, Sepandj F (2010) MIC versus MBEC to determine the  
27 antibiotic sensitivity of *Staphylococcus aureus* in peritoneal dialysis peritonitis. Perit Dial  
28 Int 30:652-656  
29  
30  
31  
32 Gudina EJ, Rocha V, Teixeira JA, Rodrigues LR (2010) Antimicrobial and antiadhesive  
33 properties of a biosurfactant isolated from *Lactobacillus paracasei* ssp. paracasei A20.  
34 Lett Appl Microbiol 50:419-424  
35  
36  
37  
38 Haba E, Pinazo A, Jauregui O, Espuny MJ, Infante MR, Manresa A (2003) Physicochemical  
39 characterization and antimicrobial properties of rhamnolipids produced by  
40 *Pseudomonas aeruginosa* 47T2 NCBIM 40044. Biotechnol Bioeng 81:316-322  
41  
42  
43  
44 He J, Ledesma KR, Lam WY, Figueroa DA, Lim TP, Chow DS, Tam VH (2010) Variability of  
45 polymyxin B major components in commercial formulations. Int J Antimicrob Agents  
46 35:308-310  
47  
48  
49 Irie Y, O'Toole G A, Yuk MH (2005) *Pseudomonas aeruginosa* rhamnolipids disperse  
50 *Bordetella bronchiseptica* biofilms. FEMS Microbiol Lett 250:237-243  
51  
52  
53  
54 Janek T, Lukaszewicz M, Krasowska A (2012) Antiadhesive activity of the biosurfactant  
55 pseudofactin II secreted by the Arctic bacterium *Pseudomonas fluorescens* BD5. BMC  
56 Microbiol 12:24  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- Jass J, Lappin-Scott HM (1996) The efficacy of antibiotics enhanced by electrical currents against *Pseudomonas aeruginosa* biofilms. J Antimicrob Chemother 38:987-1000
- Joshi-Navare K, Prabhune A (2013) A biosurfactant sophorolipid acts in synergy with antibiotics to enhance their efficiency BioMed Res Int; 1-8
- Kim KM, Lee JY, Kim CK, Kang JS (2009) Isolation and characterization of surfactin produced by *Bacillus polyfermenticus* KJS-2 . Arch Pharm Res 32:711-715
- Kiran GS, Sabarathnam B, Selvin J (2010) Biofilm disruption potential of a glycolipid biosurfactant from marine *Brevibacterium casei*. FEMS Immunol Med Microbiol 59:432-438
- Kotulova D, Slobodnikova L (2010) Susceptibility of *Staphylococcus aureus* Biofilms to Vancomycin, Gentamicin and Rifampin. Epidemiologie Mikrobiologie Imunologie 59: 80-87
- Krasowska A (2010) Biomedical activity of biosurfactants. Postepy Hig Med Dosw) 64:310-313
- Krupovic M, Daugelavicius R, Bamford DH (2007) Polymyxin B induces lysis of marine *Pseudoalteromonads* Antimicrob Agents. Chemother 51:3908-3914
- Kuiper I, Lagendijk, EL, Pickford, R, Derrick, JP, Lamers, GEM, Thomas-Oates, JE, Lugtenberg, BJJ, Bloemberg, GV (2004) Characterization of two *Pseudomonas putida* lipopeptide biosurfactants, putisolvin I and II, which inhibit biofilm formation and break down existing biofilms. Mol Microbiol 51:97-113
- Lehtinen J, Nuutila J, Lilius, EM (2004) Green fluorescent protein-propidium iodide (GFP-PI) based assay for flow cytometric measurement of bacterial viability. Cytometry 60A: 165-172
- Leis AP, Schlicher S, Franke H, Strathmann M (2005) Optically transparent porous medium for nondestructive studies of microbial biofilm architecture and transport dynamics. Appl Environ Microbiol 71:4801-4808
- Liu L, Tan X, Jia A (2012) Relationship between bacterial quorum sensing and biofilm formation--a review. Acta microbiologica Sinica 52: 271-278.
- Lourenco A, Machado H, Brito L (2011) Biofilms of *Listeria monocytogenes* produced at 12 degrees C either in pure culture or in co-culture with *Pseudomonas aeruginosa* showed reduced susceptibility to sanitizers. J Food Sci 76: M143-148

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- Lourenco A, Coenye T, Goeres DM, Donelli G, Azevedo AS, Ceri H, Coelho FL, Flemming HC, Juhna T, Lopes SP, Oliveira R, Oliver A, Shirliff ME, Sousa AM, Stoodley P, Pereira MO, Azevedo NF (2014) Minimum information about a biofilm experiment (MIABIE): standards for reporting experiments and data on sessile microbial communities living at interfaces. *Pathogens and Disease* 70: 250-256
- Luna JM, Rufino RD, Sarubbo LA, Rodrigues LR, Teixeira JA, de Campos-Takaki GM (2011) Evaluation antimicrobial and antiadhesive properties of the biosurfactant Lunasan produced by *Candida sphaerica* UCP 0995. *Curr Microbiol* 62:1527-1534
- Mandal SM (2012) A novel hydroxyproline rich glycopeptide from pericarp of *Datura stramonium*: proficiently eradicate the biofilm of antifungals resistant *Candida albicans*. *Biopolymers* 98:332-337
- Marchant R, Banat IM (2012a) Biosurfactants: a sustainable replacement for chemical surfactants? *Biotechnol Lett* 34:1597-1605
- Marchant R, Banat IM (2012b) Microbial biosurfactants: challenges and opportunities for future exploitation. *Trends Biotechnol* 30:558-565
- McLaughlin RA, Hoogewerf AJ (2006) Interleukin-1beta-induced growth enhancement of *Staphylococcus aureus* occurs in biofilm but not planktonic cultures. *Microb Pathog* 41:67-79
- Milletli Sezgin F, Coban AY, Gunaydin M (2012) Investigation of biofilm formation in *Acinetobacter baumannii* isolates and their colistin susceptibilities in biofilm. *Int J Antimicrob Agents* 41:199
- Mireles JR, 2nd, Toguchi A, Harshey RM (2001) *Salmonella enterica* serovar *typhimurium* swarming mutants with altered biofilm-forming abilities: surfactin inhibits biofilm formation. *J Bacteriol* 183:5848-5854
- Mukherji R, Prabhune A (2014) Novel glycolipids synthesized using plant essential oils and their application in quorum sensing inhibition and as antibiofilm agents. *Scientific World Journal* 2014: 890709
- Muthusamy K, Gopalakrishnan S, Ravi TK, Sivachidambaram P (2008) Biosurfactants: Properties, commercial production and application. *Current Sci.* 94: 736-747
- Neu TR (1996) Significance of bacterial surface-active compounds in interaction of bacteria with interfaces. *Microbiological Reviews* 60: 151-166

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- O'Toole GA (2011) Microtiter dish biofilm formation assay. *J Vis Exp* Jan 30;(47). pii: 2437. doi: 10.3791/2437.
- Olson ME, Ceri H, Morck DW, Buret AG, Read RR (2002) Biofilm bacteria: formation and comparative susceptibility to antibiotics. *Can J Vet Res* 66:86-92
- Pamp SJ, Tolker-Nielsen T (2007) Multiple roles of biosurfactants in structural biofilm development by *Pseudomonas aeruginosa*. *J Bacteriol* 189:2531-2539
- Pecci Y, Rivardo F, Martinotti MG, Allegrone G (2010) LC/ESI-MS/MS characterisation of lipopeptide biosurfactants produced by the *Bacillus licheniformis* V9T14 strain. *J Mass Spectrom* 45:772-778
- Pereira MO, Machado I, Simões M, Vieira MJ (2007) Preventing biofilm formation using surfactants. *BiofilmClub*©. 167-74. <http://hdl.handle.net/1822/7534> accessed May 2014.
- Pradhan AK, Pradhan N, Mall G, Panda HT, Sukla LB, Panda PK, Mishra BK (2013) Application of lipopeptide biosurfactant isolated from a halophile: *Bacillus tequilensis* CH for inhibition of biofilm. *Appl Biochem Biotechnol* 171:1362-1375
- Price NP, Rooney AP, Swezey JL, Perry E, Cohan FM (2007) Mass spectrometric analysis of lipopeptides from *Bacillus* strains isolated from diverse geographical locations. *FEMS Microbiol Lett* 271:83-89
- Quinn GA, Maloy AP, McClean S, Carney B, Slater JW (2012) Lipopeptide biosurfactants from *Paenibacillus polymyxa* inhibit single and mixed species biofilms. *Biofouling* 28:1151-1166
- Raza W, Yang X, Wu H, Wang Y, Xu Y, Shen Q (2009) Isolation and characterisation of fusaricidin-type compound-producing strain of *Paenibacillus polymyxa* SQR-21 active against *Fusarium oxysporum* f.sp.nevium. *Eur J Plant Pathol* 125:471-483
- Read RC, Roberts P, Munro N, Rutman A, Hastie A, Shryock T, Hall R, McDonald-Gibson W, Lund V, Taylor G (1992) Effect of *Pseudomonas aeruginosa* rhamnolipids on mucociliary transport and ciliary beating. *J Appl Physiol* 72: 2271–2277
- Rivardo F, Martinotti MG, Turner RJ, Ceri H (2011) Synergistic effect of lipopeptide biosurfactant with antibiotics against *Escherichia coli* CFT073 biofilm. *Int J Antimicrob Agents* 37:324-331



- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- Rivardo F, Turner RJ, Allegrone G, Ceri H, Martinotti MG (2009) Anti-adhesion activity of two biosurfactants produced by *Bacillus* spp. prevents biofilm formation of human bacterial pathogens. *Appl Microbiol Biotechnol* 83:541-553
- Rodrigues L, Banat IM, Teixeira J, Oliveira R (2006a) Biosurfactants: potential applications in medicine. *J Antimicrob Chemother* 57:609-618
- Rodrigues L, Banat IM, van der Mei HC, Teixeira JA, Oliveira R (2006b) Interference in adhesion of bacteria and yeasts isolated from explanted voice prostheses to silicone rubber by rhamnolipid biosurfactants. *J Appl Microbiol* 100:470-480
- Rodrigues L, van der Mei HC, Banat IM, Teixeira J, Oliveira R (2006c) Inhibition of microbial adhesion to silicone rubber treated with biosurfactant from *Streptococcus thermophilus* A. *FEMS Immunology and Medical Microbiology* 46, 107-112
- Rodrigues L, van der Mei HC, Teixeira J, Oliveira R (2004) Influence of biosurfactants from probiotic bacteria on formation of biofilms on voice prostheses. *Appl Environ Microbiol* 70:4408-4410
- Rodrigues L, Banat IM, Teixeira J, Oliveira R (2007) Strategies for the prevention of microbial biofilm formation on silicone rubber voice prostheses. *J Biomed Materials Res Part B- Applied Biomaterials* 81B: 358-370
- Rollet C, Gal L, Guzzo J (2009) Biofilm-detached cells, a transition from a sessile to a planktonic phenotype: a comparative study of adhesion and physiological characteristics in *Pseudomonas aeruginosa*. *FEMS Microbiol Lett* 290:135-142
- Rufino RD, Luna JM, Sarubbo LA, Rodrigues LR, Teixeira JA, Campos-Takaki GM (2011) Antimicrobial and anti-adhesive potential of a biosurfactant Rufisan produced by *Candida lipolytica* UCP 0988. *Colloids Surf B Biointerfaces* 84:1-5
- Satpute SK, Banpurkar AG, Dhakephalkar PK, Banat IM, Chopade BA (2009) Methods for investigating biosurfactants and bioemulsifiers: A review. *Critical Reviews in Biotechnol* 30: 127-144
- Samadi N, Abadian N, Ahmadkhaniha R, Amini F, Dalili D, Rastkari N, Safaripour E, Mohseni FA (2012) Structural characterization and surface activities of biogenic rhamnolipid surfactants from *Pseudomonas aeruginosa* isolate MN1 and synergistic effects against methicillin-resistant *Staphylococcus aureus*. *Folia Microbiologica* 57: 501-508

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- Sepandj F, Ceri H, Gibb A, Read R, Olson M (2004) Minimum inhibitory concentration (MIC) versus minimum biofilm eliminating concentration (MBEC) in evaluation of antibiotic sensitivity of gram-negative bacilli causing peritonitis. *Perit Dial Int* 24:65-67
- Sriram MI, Kalishwaralal K, Deepak V, Gracerosept R, Srisakthi K, Gurunathan S (2011) Biofilm inhibition and antimicrobial action of lipopeptide biosurfactant produced by heavy metal tolerant strain *Bacillus cereus* NK1. *Colloids Surf B Biointerfaces* 85:174-181
- Stepanovic S, Cirkovic I, Ranin L, Svabic-Vlahovic M (2004) Biofilm formation by *Salmonella* spp. and *Listeria monocytogenes* on plastic surface. *Lett Appl Microbiol* 38:428-432
- Stepanovic S, Vukovic D, Hola V, Di Bonaventura G, Djukic S, Cirkovic I, Ruzicka F (2007) Quantification of biofilm in microtiter plates: overview of testing conditions and practical recommendations for assessment of biofilm production by *staphylococci* . *APMIS* 115:891-899
- Stewart CR, Muthye V, Cianciotto NP (2012) *Legionella pneumophila* Persists within Biofilms Formed by *Klebsiella pneumoniae*, *Flavobacterium* sp., and *Pseudomonas fluorescens* under Dynamic Flow Conditions *PLoS ONE* 7:e50560
- Stoodley P, Sauer K, Davies DG, Costerton JW (2002) Biofilms as complex differentiated communities. *Annu Rev Microbiol* 56:187-209
- Tahmourespour A, Salehi R, Kasra Kermanshahi R (2011) *Lactobacillus acidophilus*-Derived Biosurfactant Effect on GTFB and GTFC Expression Level in *Streptococcus mutans* Biofilm Cells. *Braz J Microbiol* 42:330-339
- Velraeds MMC, van de Belt-Gritter B, Busscher HJ, Reid G, van der Mei HC (2000) Inhibition of uropathogenic biofilm growth on silicone rubber in human urine by *lactobacilli* - a teleologic approach. *World J Urology* 18: 422-426
- Villa, F, Cappitelli, F (2013) Plant-derived bioactive compounds at sub-lethal concentrations: towards smart biocide-free antibiofilm strategies. *Phytochem Rev* 12: 245-254
- Wicken AJ, Knox KW (1980) Bacterial cell surface amphiphiles. *Biochim. Biophys. Acta* 604: 1–26. Wu ZY, Ye CS, ; Guo F, Zhang SH, Yu X (2013) Evidence for Broad-Spectrum Biofilm Inhibition by the Bacterium *Bacillus* sp Strain SW9. *Appl Environ Microbiol* 79: 1735-1738

1 Xu Z, Shao J, Li B, Yan X, Shen Q, Zhang R (2013) Contribution of Bacillomycin D in  
2 *Bacillus amyloliquefaciens* SQR9 to Antifungal Activity and Biofilm Formation. Appl  
3 Environ Microbiol 79: 808-815  
4

5 Zakaria Gomaa E (2013) Antimicrobial and anti-adhesive properties of biosurfactant  
6 produced by *Lactobacilli* isolates, biofilm formation and aggregation ability. J Gen Appl  
7 Microbiol. 2013;59 :425-36  
8  
9

10 Zhang Y (2014) Persisters, persistent infections and the Yin-Yang model. Emerg Microbes  
11 Infect 3: e3. Published online: doi:10.1038/emi.2014.3  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

**Table 1** Selected biosurfactants reported in literature with antibiofilm/microbial activities.

---

Biosurfactant class	Name	Source	Reference	Effectiveness
Lipopeptide	Putisolvin I & II	<i>Pseudomonas putida</i>	Kuiper et al. 2004	Biofilm inhibition of <i>Pseudomonas</i> spp.
Lipopeptide	Pseudofactin II		Janek et al. 2010	Effective against <i>E. coli</i> , <i>Enterococcus faecalis</i> , <i>Proteus mirabilis</i> and <i>Candida</i> sp.
Lipopeptide	NS	<i>Bacillus subtilis</i>	Mireles et al. 2001	Biofilm inhibition of <i>S. entrica</i> on urethral catheter
Lipopeptide	Fengycin	<i>B. subtilis</i> & <i>B. licheniformis</i>	Rivardo et al. 2009	Inhibition of pathogenic <i>E. coli</i> & <i>S. entrica</i>
Lipopeptide	NS	Heavy metal tolerant strain of <i>Bacillus</i>	Sriram et al. 2011	Inhibits Gram positive and negative bacteria and fungi
Lipopeptide	NS	<i>Bacillus</i> sp. strain SW9	Wu et al. 2013	Inhibits biofilm formation in a wide range of bacteria
Lipopeptide	NS	<i>Bacillus tequilensis</i>	Pradhan et al. 2013	Biofilm inhibition of <i>E. coli</i> & <i>Streptococcus mutans</i>
Lipopeptide	L. fermentum B54	<i>Lactobacillus</i>	Velraeds et al. 2000	Inhibits uropathogens
Glycolipids	NS	<i>Brevibacterium casei</i> .	Kiran et al. 2010	Inhibits mixed pathogenic biofilm bacteria
Mixture of biosurfactants	Lunasan	<i>Candida sphaerica</i>	Luna et al. 2011	Inhibition of <i>P. aeruginosa</i> and <i>S. agalactae</i>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

NS	NS	<i>Lactobacillus paracasei</i> A20	Gudina et al. 2010	Biofilm inhibition for a range of bacteria, yeasts & filamentous fungi.
Glycolipid	Rhamnolipid	<i>P. aeruginosa</i>	Rodrigues et al. 2006b	Inhibits biofilms in <i>S. aureus</i> <i>Candida tropicalis</i>
Glycolipid	Rhamnolipid	<i>P. aeruginosa</i>	Dusane et al. 2010	Inhibits <i>B. pumilus</i>
Mixed biosurfactants		<i>Lactococcus lactis</i> / <i>Strep thermophilus</i>	Rodrigues et al. 2004	Effective against <i>Staphylococcus</i> , <i>Streptococcus</i> , <i>Rothia</i> and <i>Candida sp.</i>
NS	NS	<i>Robinia pseudocacia</i> / <i>Nerium oleander</i>	Cochis et al. 2012	Effective against <i>C. albicans</i>
Glycolipids	Rhamnolipid	<i>P. aeruginosa</i>	Dusane et al. 2012	Effective against <i>Yarrowia sp.</i>
NS	Rufisan	<i>Candida lypolytica</i>	Rufino et al. 2011	Effective against <i>Streptococcus sp</i>
Glycolipid	Glucose + palmitic acid	<i>Serratia Marsecens</i>	Dusane et al. 2011	Effective against <i>C. albicans</i> , <i>P. aeruginosa</i> and <i>B. pumilus</i>

---

NS= Not specified

## Figures legends

**Figure 1.** Calgary Biofilm Device (CBD) measures the minimum biofilm eradication concentration (MBEC). (1) Biofilms cultivated on pegs in 1/10th Muller Hinton broth (2) Pegs rinsed with PBS (3) Pegs exposed to test substances in new wells (4) Pegs rinsed in PBS (5) Biofilm removed by sonicating pegs into sterile media (6) Remaining viable bacteria in wells is proportional to the biofilm biomass.

**Figure 2.** Biosurfactants: (A) Polymyxin B2, (B) Fengycin-like peptide, (C) Putisolvin II and (D) Pseudofactin II:

**Figure 3.** Biosurfactants : (A) Surfactin, (B) - Polymyxin D1, (C) Fusaricidin B1, (D) Rhamnolipids: mono rhamnolipid, (l-rhamnosyl- $\beta$ -hydroxydecanoyl- $\beta$ -hydroxydecanoate)(RL-1) and (E) di-rhamnolipid, (l-rhamnosyl l-rhamnosyl- $\beta$ -hydroxydecanoyl- $\beta$ -hydroxydecanoate (RL-2), DAB = diaminobutyric acid.

**Figure 4.** Representative images depicting the effect of rhamnolipids on pre-formed biofilms of *P. aeruginosa* PAO1 on cover slips. Cells were stained with crystal violet 1%, and observed using a phase contrast microscope at 40x. (A) *P. aeruginosa* PAO1 Biofilms after 48h. (B) After 30min treatment with rhamnolipids (5%) v/v on 48h biofilms.

**Figure 5.** Biofilm formation by *Bacillus subtilis* BBK006 on coverslips. Cells were stained with Syto9®, and observed using a fluorescence microscope at 40x. The bar represent 100 $\mu$ m . (A) *Bacillus subtilis* BBK006 biofilms after 48h as a control. (B) After 30min treatment in the presence of Sophorolipids 5% v/v on 48h preformed biofilms.

**Figure 1**



Biofilms cultivated in wells with pegs



Biofilms mature



Non-adherent bacteria rinsed off



Biofilms incubated with test substance



Rinse off inhibitor



Sonicate pegs into media growth = biofilm survival

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Figure 2








Figure 3


A




B




C



D



E



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Figure 4**





Figure 5

