



LJMU Research Online

Murray, AT, Dowley, MJH, Pradaux-Caggiano, F, Baldansuren, A, Fielding, AJ, Tuna, F, Hendon, CH, Walsh, A, Lloyd-Jones, GC, John, MP and Carbery, DR

Catalytic Amine Oxidation under Ambient Aerobic Conditions: Mimicry of Monoamine Oxidase B

<http://researchonline.ljmu.ac.uk/8146/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Murray, AT, Dowley, MJH, Pradaux-Caggiano, F, Baldansuren, A, Fielding, AJ, Tuna, F, Hendon, CH, Walsh, A, Lloyd-Jones, GC, John, MP and Carbery, DR (2015) Catalytic Amine Oxidation under Ambient Aerobic Conditions: Mimicry of Monoamine Oxidase B. *Angewandte Chemie-*

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

Catalytic Amine Oxidation under Ambient Aerobic Conditions: Mimicry of Monoamine Oxidase B**

Alexander T. Murray, Myles J. H. Dowley, Fabienne Pradaux-Caggiano, Amgalanbaatar Baldansuren, Alistair J. Fielding, Floriana Tuna, Christopher H. Hendon, Aron Walsh, Guy C. Lloyd-Jones, Matthew P. John, and David R. Carbery*

Abstract: The flavoenzyme monoamine oxidase (MAO) regulates mammalian behavioral patterns by modulating neurotransmitters such as adrenaline and serotonin. The mechanistic basis which underpins this enzyme is far from agreed upon. Reported herein is that the combination of a synthetic flavin and alloxan generates a catalyst system which facilitates biomimetic amine oxidation. Mechanistic and electron paramagnetic (EPR) spectroscopic data supports the conclusion that the reaction proceeds through a radical manifold. This data provides the first example of a biorelevant synthetic model for monoamine oxidase B activity.

Monoamine oxidase (MAO) is a mitochondrial flavin-dependent oxidoreductase enzyme which oxidizes a range of important amines to imines, for example, the neurotransmitters serotonin, histamine, and noradrenaline.^[1] With such an integral role in the neurochemical network, MAO function has been pinpointed as an underlying rationale for a range of behavioral, evolutionary, and physiological observations. For example, variations in the MAO A gene can lead to increased

aggression, known as the “warrior gene”, ultimately impacting human evolution.^[2] Inhibition of MAO has been an important area for medicinal chemistry with MAO inhibitors (MAOIs) acting as potent antidepressants and having potential applications as neuroprotective agents.^[3] Mechanistic studies have also helped in understanding the role of lysine-specific demethylase 1 (LSD1), a key epigenetic modulator, with MAOIs impacting a number of key biological processes.^[4]

It is remarkable that no consensus has been reached with respect to a mechanism of action, despite over 45 years of investigation.^[5] There are two isozymes of MAO: MAO-A and MAO-B. While the flavin active sites are identical, each form displays a different substrate and inhibitor profile, and the mechanistic basis of this selectivity unknown.^[6]

²H primary kinetic isotope (KIE) effects have been observed for the C–H bond cleavage step(s) with both MAO A and B. In principle, rate-contributing cleavage may be envisaged as proceeding by either H⁺-, H⁻-, or H[•]-transfer mechanisms (Scheme 1). These options have been widely discussed,^[5] with rate-contributing C–H cleavage by H⁺ transfer being the most prevalent mechanistic description. Two mechanistic postulates have been developed to account for the requisite increase in acidity of the relevant α -amino C–H bond: the formation of a covalent flavin–amine conjugate,^[7] and the formation of an aminium radical cation^[8] after single-electron transfer from amine to flavin. As both mechanisms require discrete steps prior to the rate-contributing C–H cleavage, it is notable that no intermediates accumulate to observable populations. C–H cleavage in the context of a direct hydride transfer has also been suggested.^[9] However, such a synchronous event would not be consistent with the ¹⁵N KIE measured for amine oxidation by MAO B, thus pointing to an absence of synchronicity between C–H cleavage and sp²→sp³ nitrogen atom re-hybridization.^[10] Finally, H[•] transfer from the substrate to the flavin has been

[*] A. T. Murray, M. J. H. Dowley, Dr. F. Pradaux-Caggiano, C. H. Hendon, Prof. A. Walsh, Dr. D. R. Carbery
 Department of Chemistry, University of Bath
 Claverton Down, Bath (UK)
 E-mail: d.carbery@bath.ac.uk

Dr. A. Baldansuren, Dr. A. J. Fielding, Dr. F. Tuna
 EPSRC National EPR Facility, Photon Science Institute
 School of Chemistry
 University of Manchester, Oxford Road, Manchester (UK)


Prof. G. C. Lloyd-Jones
 School of Chemistry, Joseph Black Building
 West Mains Road, Edinburgh EH9 3 JJ (UK)

Dr. M. P. John
 GlaxoSmithKline Research and Development
 Gunnels Wood Road, Stevenage (UK)

[**] We acknowledge funding of a studentship (A.T.M.) from GSK, EPSRC, and the University of Bath. The Bath, Bristol, and Cardiff Catalysis alliance is acknowledged for support (F.P.-C.). C.H.H. and A.W. are supported by the ERC (Grant 277757) and the computing was enabled by access to ARCHER through membership of the UK's HPC Materials Chemistry Consortium, which is funded by EPSRC (Grant No. EP/L00202).

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201503654>.

© 2015 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.



Scheme 1. MAO-catalyzed oxidation of amines and qualitative overview of possible modes of C–H bond cleavage.

suggested.^[11] This possibility was discarded on the grounds that no hydrogen-atom abstracting moiety, which was reactive enough to overcome relevant α -amino C–H bond dissociation energies, could be identified in the enzyme active site.^[12]

Studies using synthetic flavins have played a crucial role in elucidating flavoenzyme mechanisms.^[13] Accordingly, insight gained from studying model cofactors is a valid strategy to unlocking mechanistic problems in flavoenzymology. Pioneering work on primary amines by various groups supported the polar, proton-transfer mechanism, but the low turnover, tendency of catalysts to decompose, and requirement of heating in an enriched O₂ atmosphere for several days meant that they are perhaps of limited relevance to biological processes.^[14] We,^[15] and others,^[16] have previously applied cationic flavin catalysts in biomimetic monooxygenase contexts, as well as donor–acceptor chemistry,^[17] and now report the oxidation of biologically pertinent amines as a vehicle to understanding MAO mechanism.

Initial exploratory studies demonstrated catalytic aerobic oxidation of benzylamine, with formation of the imine **4a** being consistent with oxidase rather than monooxygenase-like reactivity (Table 1).^[18] Excellent yields of **4a** were obtained if a thioether additive (Me₂S) and a cocatalyst, alloxan (**3a**), were used (Table 1). Initially **3a** was present as an undetected by-product from the synthesis of **2a**, however, was found to be crucial for this transformation. *N,N*-dimethylalloxan (**3b**) was found to be inactive (entry 3) despite possessing structural similarity to **3a**. Additionally, cobalamin synthase, BluB, has been implicated in the cannibalization of flavin mononucleotide to form alloxan, which acts as a crucial multifunctional redox catalyst in the biosynthesis of vitamin B12.^[19] A series of substituted benzylamines, typical substrates for MAO-B, have been examined.

Table 1: Flavin-organocatalyzed amine oxidation.^[a]

Entry	Substrate	R ¹	Product	Yield [%]
1	1a	Ph	4a	94
2 ^[b]	1a	Ph	4a	49
3 ^[c]	1a	Ph	–	0
4	1b	4-MeC ₆ H ₄	4b	99
5	1c	4-MeOC ₆ H ₄	4c	95
6	1d	4- <i>t</i> BuC ₆ H ₄	4d	92
7 ^[d]	1e	4-FC ₆ H ₄	4e	98
8 ^[d]	1f	4-ClC ₆ H ₄	4f	77
9 ^[d,e]	1g	4-CF ₃ C ₆ H ₄	4g	37
10	1h	3-MeC ₆ H ₄	4h	91
11	1i	3-OMeC ₆ H ₄	4i	72
12	1j	2-MeC ₆ H ₄	4j	96
13	1k	2-MeOC ₆ H ₄	4k	68
14 ^[d]	1l	2-ClC ₆ H ₄	4l	66
15 ^[d]	1m	2-furyl	4m	70
16	1n	2-thiophenyl	4n	87
17	1o	1-naphthyl	4o	72

[a] Reaction conditions: **2a** (2 mol %), **3a** (2 mol %), Me₂S (10 equiv), 5 h. [b] **2b** used. [c] **3b** used. [d] 18 h. [e] **2a** (4 mol %), **3a** (4 mol %) used.

Generally, high yields of imine products are attainable, although substrates with a strongly electron-withdrawing *para*-substituent group (entry 9) are less reactive, thus mirroring MAO B reactivity trends.

Upon attempted in situ ¹H NMR analysis, the inability to locate the lock signal suggested paramagnetic behavior. Accordingly, EPR studies at the X-band were initiated. Mixing **2a** and Me₂S generated the flavin radical cation **2a'** (Figure 1). The structure was further confirmed by pulsed




Figure 1. EPR spectra of **2a'** and **4a'**, and DFT-calculated spin densities measured from solutions of **2a** + Me₂S (top) and **2a** + Me₂S + **3a** + **1a** (bottom).

EPR studies. In particular, the protonation state of **2a'** was assessed by electron spin echo envelope modulation (ESEEM), and is a rare example of an aerobically generated flavin semiquinone, having demonstrable relevance to catalysis, observed by EPR spectroscopy.^[20] The use of a strong hydrogen-bonding solvent, trifluoroethanol, may aid the stabilization of the semiquinone formation, as discussed by Massey and co-workers, for flavins with amino acids.^[21] Upon sequential addition of alloxan and amine, a new EPR spectrum was observed and characterized as the radical **4a'**, and is consistent with charge-transfer-initiated hydrogen-atom abstraction from **4a**. Hybrid-DFT and post-Hartree Fock calculations were performed on **2a'** and **4a'** and the spin density isosurfaces are shown in Figure 1.^[22] Importantly, the theoretical calculations quantify the local spin density distribution, thus further corroborating the simulations of the continuous-wave EPR spectra.^[23]

Kinetic studies provided additional important mechanistic information with the transformation being first order in benzylamine^[24] and showing a KIE of $k_H/k_D = 1.9$ when using PhCD₂NH₂ (**7**), thus supporting rate-contributing C–H bond cleavage (Figure 2). A range of studied *para*-substituted benzylamines provided a negative Hammett correlation ($\rho = -2$).

The observed rates of reaction were found to be independent of the Me₂S concentration. Kinetic analysis for




Figure 2. Hammett and kinetic isotope effect study carried out by HPLC analysis of imine formation from amines against an internal standard of naphthalene.


3a did not demonstrate a simple reaction order, with saturation behavior observed over the concentrations examined (see the Supporting Information). The kinetic order in **2a** was probed by means of a $\ln(k_{\text{obs}})$ versus $\ln([\text{flavin}])$ plot, which was linear with a slope of 0.25 and consistent with de-aggregation of a higher order resting state, but with a monomeric semiquinone being catalytically active. Significantly, the less oxidizing flavin **2b** also mediates this reaction ($k_{\text{obs}2a}/k_{\text{obs}2b} = 2.96$) with an electrochemical reduction potential of +66 mV vs. SHE, which parallels MAO-B at +40 mV (Table 1, entry 2)^[25] Therefore, the flavin catalysts **2a,b** offer themselves as realistic mimics of MAO through the neutral N(5)-H semiquinone.^[26]

A mechanism that accounts for EPR and kinetic data is underpinned by the realization that rate-determining C–H cleavage is mediated by **2a'** (Scheme 2). The radical cation **2a'** is formed by a proton-coupled electron transfer from Me_2S , as observed by EPR. BnNH_2 promotes the formation of **2a'** by mediating the de-aggregation and deprotonation of **2a'**, thus generating the neutral semiquinone **2a''**, with subsequent $\text{H}\cdot$ transfer, initiated by a charge-transfer event, from **1a** to **2a''**. An α -amino radical is formed (**1a'**), and it acts as a potent reductant,^[27] thus reducing alloxan and forming **1a''**. Electron transfer from α -amino radicals to vicinal dicarbonyl compounds is regarded as one of the fastest reactions between a radical and a neutral closed-shell organic molecule.^[28] Alloxan (**3a**) reacts as an amide tautomer, thus allowing stabilization of a developing oxyanion character, a feature which is impossible for the inactive **3b** (Table 1, entry 3). This captodative-stabilized radical^[29] subsequently reacts with O_2 , thus generating **5**. The peroxy radical **5** oxidizes **2a'''** to **2a'**, thus forming the hydroperoxide **5'** and completing the catalytic cycle. Formation of stoichiometric DMSO is observed. Therefore Me_2S mediates the reduction of **5'** to

alloxan. Additionally, a purple by-product, consistent with the dye murexide (**6**; UV/vis $\lambda_{\text{max}} = 521 \text{ nm}$; lit = 520 nm),^[30] is observed to accumulate from **3a''**, **3a**, and ammonia. This observation is consistent with a two-electron over-reduction of **3a**, thus leading to catalyst deactivation and suggesting that **3a''** is not a catalytically active species.

This model study supports a homolytic C–H bond cleavage mediated by a flavin semiquinone, and with a substrate preference for benzylamines, it has prompted us to ask whether any reasonable insight into the enzymatic mechanism of MAO B can be achieved through consideration of this currently presented model system. A linear correlation exists between the substrate $\text{p}K_{\text{a}}$ value and steady-state k_{cat} for MAO B (Figure 3),^[31] and is consistent with a neutral amine substrate. It is significant that Hammett electronic correlations for MAO B are only apparent at pH 9.0. The similarity of the model's KIE and Hammett profiles to the equivalent B isozyme data, when the enzyme kinetics are measured at pH 9.0, which is similar to this unbuffered system, is notable, (MAO B: $k_{\text{H}}/k_{\text{D}} = 2.25$, $\rho = -0.9$ at pH 9.0).^[32]

Our proposal for the MAO B mechanism is informed by the presented data, the substrate reactivity trends, and the pH sensitivity of MAO B.^[34] This mechanistic suggestion centers upon a charge-transfer event promoted by the free-base substrate interacting with an electron-rich phenol of Y398 near the flavin acceptor, as demonstrated by Scrutton and co-workers.^[35] This acceptor is itself activated by the H_2O -K296 hydrogen-bonding motif. The neutral semiquinone thus formed can mediate hydrogen-atom transfer from the substrate, with the tyrosinyl radical cation now able to accept the second substrate electron, in direct analogy to the role played



Scheme 2. Proposed reaction mechanism showing amine oxidation mediated by the key flavin semiquinone **2a''**.




Figure 3. a) MAO B steady-state k_{cat} relationship to substrate pK_{a} ; relevant k_{cat} values from Ref. [31]. b) Crystal structure of MAO B with 4-nitrobenzylamine inhibitor.^[31]

by alloxan in the currently discussed model. Indeed, both components can be viewed as redox-active hydroxylated units.

In summary, an aerobic, catalytic oxidation of benzylamines which mimics MAO B activity proceeding through charge-transfer-initiated substrate H[•] abstraction has been developed. EPR spectroscopy has revealed the operation of an aerobically generated flavin semiquinone. KIE and Hammett studies have demonstrated a pH-dependent kinetic parallel to MAO B activity. This model system has opened up an additional mechanistic model of MAO B activity, that is, a charge-transfer event is harnessed to access a reactive neutral flavin semiquinone as the C–H abstracting species in MAO B.

Keywords: amines · enzymes · EPR spectroscopy · oxidation · reaction mechanisms

How to cite: *Angew. Chem. Int. Ed.* **2015**, *54*, 8997–9000
Angew. Chem. **2015**, *127*, 9125–9128

- [1] C. M. McEwen, *J. Biol. Chem.* **1965**, *240*, 2003–2010.
[2] R. McDermott, D. Tingley, J. Cowden, G. Frazzetto, D. D. P. Johnson, *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 2118–2123.
[3] M. B. H. Youdim, D. Edmondson, K. F. Tipton, *Nat. Rev. Neurosci.* **2006**, *7*, 295–309.
[4] K. Helin, D. Dhanak, *Nature* **2013**, *502*, 480–488.
[5] D. E. Edmondson, C. Binda, J. Wang, A. K. Upadhyay, A. Mattevi, *Biochemistry* **2009**, *48*, 4220–4230.
[6] L. Pisani, M. Barletta, R. Soto-Otero, O. Nicolotti, E. Mendez-Alvarez, M. Catto, A. Introcaso, A. Stefanachi, S. Cellamare, C. Altomare, A. Carotti, *J. Med. Chem.* **2013**, *56*, 2651–2664.
[7] D. E. Edmondson, C. Binda, A. Mattevi, *Arch. Biochem. Biophys.* **2007**, *464*, 269–276.
[8] R. B. Silverman, *Acc. Chem. Res.* **1995**, *28*, 335–342.
[9] a) P. F. Fitzpatrick, *Arch. Biochem. Biophys.* **2010**, *493*, 13–25; b) R. Vianello, M. Repič, J. Mavri, *Eur. J. Org. Chem.* **2012**, 7057–7065.
[10] S. MacMillar, D. E. Edmondson, O. Matsson, *J. Am. Chem. Soc.* **2011**, *133*, 12319–12321.
[11] M. C. Walker, D. E. Edmondson, *Biochemistry* **1994**, *33*, 7088–7098.
[12] T. L. Cottrell, *The Strength of Chemical Bonds*, 2nd ed., Butterworth and co., London, **1958**.
[13] a) C. Kemal, T. C. Bruice, *Proc. Natl. Acad. Sci. USA* **1976**, *73*, 995–999; b) C. Kemal, T. C. Bruice, *J. Am. Chem. Soc.* **1977**, *99*, 7064–7067; c) T. C. Bruice, J. B. Noar, S. S. Ball, U. V. Venkataram, *J. Am. Chem. Soc.* **1983**, *105*, 2452–2463.
[14] a) G. A. Hamilton, L. E. Brown, *J. Am. Chem. Soc.* **1970**, *92*, 7225–7227; b) J. M. Kim, I. S. Cho, P. S. Mariano, *J. Org. Chem.* **1991**, *56*, 4943–4955; c) J. M. Kim, M. A. Bogdan, P. S. Mariano, *J. Am. Chem. Soc.* **1993**, *115*, 10591–10595; d) W.-S. Li, L. M. Sayre, *Tetrahedron* **2001**, *57*, 4523–4536.
[15] a) B. J. Marsh, D. R. Carbery, *Tetrahedron Lett.* **2010**, *51*, 2362–2365; b) A. T. Murray, P. Matton, N. W. G. Fairhurst, M. P. John, D. R. Carbery, *Org. Lett.* **2012**, *14*, 3656–3659.
[16] a) Y. Imada, H. Iida, S. Ono, S.-I. Murahashi, *J. Am. Chem. Soc.* **2003**, *125*, 2868–2869; b) Y. Imada, H. Iida, S.-I. Murahashi, T. Naota, *Angew. Chem. Int. Ed.* **2005**, *44*, 1704–1706; *Angew. Chem.* **2005**, *117*, 1732–1734; c) S. Chen, M. S. Hossain, F. W. Foss, *Org. Lett.* **2012**, *14*, 2806–2809; d) S. Chen, F. W. Foss, *Org. Lett.* **2012**, *14*, 5150–5153; e) J. Žurek, R. Cibulka, H. Dvořáková, J. Svoboda, *Tetrahedron Lett.* **2010**, *51*, 1083–1086; f) H. Kotoučová, I. Strnadová, M. Kovandová, J. Chudoba, H. Dvořáková, R. Cibulka, *Org. Biomol. Chem.* **2014**, *12*, 2137–2142.
[17] a) A. Niemz, J. Imbriglio, V. M. Rotello, *J. Am. Chem. Soc.* **1997**, *119*, 887–892; b) J. B. Carroll, B. J. Jordan, H. Xu, B. Erdogan, L. Lee, L. Cheng, C. Tiernan, G. Cooke, V. M. Rotello, *Org. Lett.* **2005**, *7*, 2551–2554; c) S. T. Caldwell, L. J. Farrugia, S. G. Hewage, N. Kryvokhyzha, V. M. Rotello, G. Cooke, *Chem. Commun.* **2009**, 1350–1352.
[18] S. Oae, K. O. Asada, T. Yoshimura, K. Fujimori, *Heterocycles* **1992**, *33*, 189–194.
[19] X.-L. Wang, J.-M. Quan, *J. Am. Chem. Soc.* **2011**, *133*, 4079–4091.
[20] T. Akiyama, F. Simeno, M. Murakami, F. Yoneda, *J. Am. Chem. Soc.* **1992**, *114*, 6613–6620.
[21] D. Xu, R. M. Kohli, V. Massey, *Proc. Natl. Acad. Sci. USA* **1999**, *96*, 3556–3561.
[22] C. H. Hendon, D. R. Carbery, A. Walsh, *Chem. Sci.* **2014**, *5*, 1390–1395.
[23] For full discussion of EPR spectroscopy and calculations, see the Supporting Information.
[24] The reaction surprisingly becomes apparently zero order in amine after 1 h of reaction time, perhaps because oxygen becomes rate contributing.
[25] a) W.-S. Li, N. Zhang, L. M. Sayre, *Tetrahedron* **2001**, *57*, 4507–4522; b) D. P. H. M. Heuts, N. S. Scrutton, W. S. McIntire, M. W. Fraaije, *FEBS J.* **2009**, *276*, 3405–3427.
[26] ES-MS monitoring did not reveal any catalyst–amine adducts that would be expected from polar mechanisms.
[27] I. R. Gould, J. R. Lenhard, A. A. Muentner, S. A. Godleski, S. Farid, *J. Am. Chem. Soc.* **2000**, *122*, 11934–11943.
[28] J. C. Scaiano, *J. Phys. Chem.* **1981**, *85*, 2851–2855.
[29] H. G. Viehe, Z. Janousek, R. Merenyi, L. Stella, *Acc. Chem. Res.* **1985**, *18*, 148–154.
[30] J. Torrelles, M.-C. Guerin, M.-L. Carrie, *Biochimie* **1989**, *71*, 1231–1234.
[31] A. K. Tan, R. R. Ramsay, *Biochemistry* **1993**, *32*, 2137–2143.
[32] a) T. Jonsson, D. E. Edmondson, J. P. Klinman, *Biochemistry* **1994**, *33*, 14871–14878; b) R. Orru, M. Aldeco, D. E. Edmondson, *J. Neural Transm.* **2013**, *120*, 847–851.
[33] M. Li, C. Binda, A. Mattevi, D. E. Edmondson, *Biochemistry* **2006**, *45*, 4775–4784.
[34] a) L. De Colibus, M. Li, C. Binda, A. Lustig, D. E. Edmondson, A. Mattevi, *Proc. Natl. Acad. Sci. USA* **2005**, *102*, 12684–12689; b) C. Binda, P. Newton-Vinson, F. Hubalek, D. E. Edmondson, A. Mattevi, *Nat. Struct. Biol.* **2002**, *9*, 22–26.
[35] R. V. Dunn, A. W. Munro, N. J. Turner, S. E. J. Rigby, N. S. Scrutton, *ChemBioChem* **2010**, *11*, 1228–1231.

Received: April 21, 2015

Published online: June 18, 2015