

LJMU Research Online

Shaw, TA, Plater, AJ, Kirby, JR, Roy, K, Holgate, S, Tutman, P, Cahill, N and Horton, BP

Tectonic influences on late Holocene relative sea levels from the centraleastern Adriatic coast of Croatia

http://researchonline.ljmu.ac.uk/id/eprint/9541/

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Shaw, TA, Plater, AJ, Kirby, JR, Roy, K, Holgate, S, Tutman, P, Cahill, N and Horton, BP (2018) Tectonic influences on late Holocene relative sea levels from the central-eastern Adriatic coast of Croatia. Quaternary Science Reviews. 200. pp. 262-275. ISSN 0277-3791

LJMU has developed LJMU Research Online for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

http://researchonline.ljmu.ac.uk/

1	Tectonic influences on late Holocene relative sea levels from the central-
2	eastern Adriatic coast of Croatia.
3	
4	Timothy A. Shaw ^{a, *} , Andrew J. Plater ^b , Jason R. Kirby ^c , Keven Roy ^a , Simon Holgate ^d , Pero
5	Tutman ^e , Niamh Cahill ^f , Benjamin P. Horton ^{a, g,}
6	^a Asian School of the Environment, Nanyang Technological University, Singapore 639798,
7	Singapore.
8	^b Department of Geography and Planning, School of Environmental Sciences, University of

- 9 Liverpool, Roxby Building, Liverpool, Merseyside L69 7ZT, U.K.
- 10 ^c School of Natural Sciences and Psychology, Liverpool John Moores University, Liverpool,
- 11 Merseyside, L3 3AF, U.K.
- ^d Sea Level Research Ltd, Studio D, Baltic Creative Campus, Liverpool, Merseyside, L1 0AH,
 U.K.
- ^e Institute of Oceanography and Fisheries, Šetalište Ivana Meštrovića 63, 21000, Split, Croatia.
- ¹⁵ ^f School of Mathematics and Statistics, University College Dublin, Dublin, Ireland.
- 16 g Earth Observatory of Singapore, Nanyang Technological University, Singapore 639798,
- 17 Singapore.
- 18 * Corresponding author email: tshaw@ntu.edu.sg
- 19

20 Abstract

Differential tectonic activity is a key factor responsible for variable relative sea-level (RSL)
changes during the late Holocene in the Adriatic. Here, we compare reconstructions of RSL from

23 the central-eastern Adriatic coast of Croatia with ICE-7G NA (VM7) glacial-isostatic model RSL 24 predictions to assess underlying driving mechanisms of RSL change during the past ~ 2700 years. Local standardized published sea-level index points (n=23) were combined with a new salt-marsh 25 26 RSL reconstruction and tide-gauge measurements. We enumerated fossil foraminifera from a short 27 salt-marsh sediment core constrained vertically by modern foraminiferal distributions, and 28 temporally by radiometric analyses providing sub-century resolution within a Bayesian age-depth 29 framework. We modelled changes in RSL using an Errors-In-Variables Integrated Gaussian 30 Process (EIV-IGP) model with full consideration of the available uncertainty. Previously 31 established index points show RSL rising from -1.48 m at 715 BCE to -1.05 m by 100 CE at 0.52 32 mm/yr (-0.82-1.87 mm/yr). Between 500 and 1000 CE RSL was -0.7 m below present rising to -33 0.25 m at 1700 CE. RSL rise decreased to a minimum rate of 0.13 mm/yr (-0.37-0.64 mm/yr) at \sim 1450 CE. The salt-marsh reconstruction shows RSL rose ~ 0.28 m since the early 18^{th} century at 34 35 an average rate of 0.95 mm/yr. Magnitudes and rates of RSL change during the twentieth century 36 are concurrent with long-term tide-gauge measurements, with a rise of ~ 1.1 mm/yr. Predictions 37 of RSL from the ICE-7G NA (VM7) glacial-isostatic model (-0.25 m at 715 BCE) are consistently 38 higher than the reconstruction (-1.48 m at 715 BCE) during the Late Holocene suggesting a 39 subsidence rate of 0.45 ± 0.6 mm/yr. The new salt-marsh reconstruction and regional index points 40 coupled with glacial-isostatic and statistical models estimate the magnitude and rate of RSL change 41 and subsidence caused by the Adriatic tectonic framework.

42

43 Keywords: Sea Level Changes; Adriatic; Croatia; late Holocene; Glacial isostatic adjustment;
44 Tectonic subsidence; Salt marsh; Foraminifera.

46 **1. Introduction**

47 Significant efforts have been made towards understanding Holocene relative sea-level (RSL) 48 changes in the Mediterranean (e.g., Flemming, 1969; Pirazzoli, 1976, 1991, 1996; Flemming and 49 Webb, 1986; Zerbini et al., 1996, 2017; Woodworth, 2003; Lambeck et al., 2004a; Marcos and 50 Tsimplis, 2008; Vacchi et al., 2016). During the Holocene, geological records illustrate eustatic 51 and glacio-hydro-isostatic changes (e.g., Lambeck and Purcell, 2005; Stocchi and Spada, 2007, 52 2009; Roy and Peltier, 2018) superposed by tectonic and local processes (e.g., Pirazzoli, 2005; 53 Antonioli et al., 2009, 2011; Vacchi et al., 2016). Indeed, tectonic effects on late Holocene RSL 54 histories in the northern Adriatic are particularly important, attesting to variable subsidence and 55 uplift rates (e.g., Benac et al., 2004, 2008; Furlani et al., 2011; Surić et al., 2014; Fontana et al., 56 2017). The effect of tectonics on RSL histories in the central-eastern Adriatic is, however, less well constrained (e.g. Faivre et al., 2013). Anthropogenic forcings since the mid to late 19th century 57 58 have contributed towards sea-level changes (e.g. Jevrejeva et al., 2009; Dangendorf et al., 2015; 59 Kopp et al., 2016). In the Adriatic and wider Mediterranean region, tide-gauge stations document 60 coherent RSL trends, simultaneously recording large inter-annual and inter-decadal variability (Orlić and Pasarić, 2000; Tsimplis and Baker, 2000; Tsimplis and Josey, 2001; Tsimplis et al., 61 62 2012). Comparing independent RSL datasets with differing resolution and time periods is, 63 therefore, problematic and restricts our understanding of RSL changes in the Adriatic.

64

Here, we reconstruct late Holocene RSL using geological and tide-gauge data coupled with a new salt-marsh based reconstruction from the central-eastern coast of Croatia that bridges the gap between late Holocene and modern sea-level data. Salt-marsh environments afford a unique ability providing near continuous, decimeter vertical (Scott and Medioli, 1978, 1980; Horton and

69 Edwards, 2006) and sub-century temporal resolution (Törnqvist et al., 2015; Corbett and Walsh, 70 2015; Marshall, 2015). Their use in reconstructing RSL is well established across regions in 71 Northern (Gehrels et al., 2005; Kemp et al., 2013; Barlow et al., 2014; Saher et al., 2015) and 72 Southern Hemispheres (Gehrels et al., 2008; 2012; Strachan et al., 2014). Salt-marsh based 73 reconstructions have aided our understanding of climate-sea-level connections (Kemp et al., 2011; Kopp et al., 2016); the onset of increases in the rate of RSL rise in the mid to late 19th century 74 75 (Kopp et al., 2016); and tectonic (van de Plassche et al., 2014), compaction (Brain et al., 2017) 76 and tidal range (Horton et al., 2013) influences on local RSL change. The Adriatic and wider 77 Mediterranean region, however, have evaded similar high-resolution RSL studies.

78

To better understand driving mechanisms of RSL change in the central-eastern Adriatic, we compare the composite RSL record with ICE-7G_NA (VM7) glacio-isostatic model predictions (Roy and Peltier, 2017) for the last ~ 2700 years. We show the magnitude of RSL change during this period is offset to model predictions by more than 1 m, implying an overarching influence of tectonic subsidence on RSL changes. We demonstrate the utility of the salt-marsh reconstruction in deriving similar magnitudes and rates of RSL change to long-term tide-gauges.

85

86 **2.** Study area

87 2.1. Tectonic setting

88 Tectonism in the western Mediterranean region is the consequence of the collision boundary 89 between the major tectonic plates of Africa and Eurasia (Fig. 1). This convergence zone results in 90 a number of microplates, including the Adriatic (McKenzie, 1972; Anderson and Jackson, 1987; 91 D'Agostino et al., 2008). The Adriatic microplate, which shows movements independent to Africa

92 and Eurasia (Grenerczy et al., 2005; Altiner et al., 2006; Serpelloni et al., 2013), is subdivided into 93 northern and southern sectors with the southern sector moving counterclockwise in a N-NW 94 direction at 5-10 mm/yr (Oldow et al., 2002; Herak et al., 2005; Marjanović et al., 2012). Tectonic 95 activity predominately occurs along the coasts and a through a number of fault lines that pass 96 through the region (Herak et al., 1996; 2017; Korbar, 2009). The distribution of earthquake 97 epicenters in the Adriatic between the Ancona-Zadar and Gargano-Dubrovnik lines (Fig. 2) 98 suggests this region is seismically more intense compared to the north with four $M_L = \ge 5.5$ events 99 recorded since the twentieth century (Herak et al., 2005). Most recently, a sequence of earthquakes peaking at M_L = 5.5 occurred in 2003 at Jabuka, some ~ 90 km west of Vis (Fig. 2) in the central 100 101 Adriatic Sea (Herak et al., 2005).

102

103 Modern measurements from Global Positioning System (GPS) stations reveal both lateral and 104 vertical land movements in the Adriatic region (Buble et al., 2010; Weber et al., 2010; Serpelloni et al., 2013; Devoti et al., 2017). Vertical velocities from GPS stations in the north-western Adriatic 105 106 show significant subsidence rates up to $\sim 8 \text{ mm/yr}$ near the Po River Delta, reflecting crustal 107 movements and also compaction of sediments (Carminati et al., 2003; Antonioli et al., 2009). 108 While the density of observations along the eastern Adriatic are limited, vertical motions in 109 northern and central Croatia are close to 0 mm/yr with minor subsidence up to 1 mm/yr recorded 110 in the south near to Dubrovnik (Fig. 2).



Fig. 1. Western Mediterranean showing location of major tectonic boundaries modified after
Faccenna et al. (2014) with ICE-7G_NA (VM7) (Roy and Peltier, 2017) model predictions of
present-day RSL change rate (mm/yr). Square outline depicts Adriatic study region presented in
Fig. 2.

116

117 **2.2. Oceanographic setting**

The Adriatic Sea is a relatively shallow elongated basin communicating with the Mediterranean Sea through the Strait of Otranto. The bathymetry is subdivided with a shallow (average ~ 35 m water depth) northern section near the Gulf of Trieste, progressively deepening to ~ 1200 m towards the south near Dubrovnik (Ciabatti et al., 1987; Orlić et al., 1992). Tidal ranges in the region are microtidal, increasing as water depth decreases to the north (Cushman-Roisin and Naimie, 2002). The influence of strong north-easterly Bora and south-easterly Sirocco winds can significantly alter the tidal regime (Orlić et al., 1994; Vilibić, 2006; Ferla et al., 2007) and
meteorological tsunamis associated with prolonged low atmospheric pressure systems are a
relatively common occurrence (Vilibić and Šepić, 2009; Vilibić et al., 2017).

127

Instrumental observations of RSL change from long-term (>50 years) tide-gauge stations in the Adriatic are restricted to the northern and eastern coastline (Fig. 2). The tidal station at Trieste provides an inference of RSL change since the late 19th century while Bakar extends (discontinuously) to 1930 CE. The tidal stations at Split and Dubrovnik extend to the mid-1950s. By comparison to the rest of the Adriatic, high rates of RSL change are observed in the north in Venice; however, this is attributable to anthropogenic influences exacerbating subsidence in the region (Woodworth, 2003) as illustrated by the GPS network.

135

136 **2.3. Study site**

137 We investigated the salt-marsh environments located near Jadrtovac, along the central-eastern 138 Adriatic coastline of Croatia (Fig. 2). Our focus on this region was motivated by the availability 139 of pristine salt marshes (Pandža et al., 2007) and nearby long-term tide-gauge stations. In this 140 context, tide gauges can provide a means of self-evaluation for proxy-based reconstructions, 141 permitting independent comparison of RSL changes (e.g. Donnelly et al., 2004; Gehrels et al., 142 2005; Kemp et al., 2009). Shaw et al. (2016) previously documented the vertical zonation of 143 contemporary foraminiferal assemblages at Jadrtovac, underpinning their potential to reconstruct 144 RSL change. The microtidal regime, with a mean tidal range of 0.23 m (Hydrographic Institute, 145 1955; Vilibić et al., 2005), also helps limit vertical uncertainties (Barlow et al., 2013). The salt-146 marsh environment is located at the head of a ~ 2.5 km channel in the Morinje Bay, northwest of Split and is a typical karstic environment with limited vegetation and poor soils on the surrounding slopes. The bay was infilled during the Holocene marine transgression, resulting in ~ 4.5 m of sediment (Bačani et al., 2004; Šparica et al., 2005). The main salt-marsh surface is ~ 130 m wide on the eastern side and gradually thins moving north around the bay.



Fig. 2. (A) Adriatic study area showing the location of long-term (>50 years) tide-gauge stations (stars), the vertical land movements recorded by GPS stations (dots) modified after Serpelloni et al. (2013) and the simplified tectonic setting of the Croatian coastline modified after Korbar (2009), together with the location of Island of Vis and the Ancona-Zadar and Gargano-Dubrovnik lines (Herak et al., 2005) referred to in text. (B) Sample site at Jadrtovac within the Morinje Bay showing stratigraphic transects and sample core location. (C) Tide-gauge measurements of relative sea-level (RSL) change from stations highlighted in panel A.

160 **3. Methodology**

161 We investigated the depositional history of the salt-marsh environment, describing the underlying 162 lithostratigraphy according to the Troëls-Smith (1955) classification of coastal sediments. Core 163 transects were established capturing the full range of sub-environments from the landward high 164 salt-marsh (hereafter termed 'high-marsh') edge to open water boundary. Following this, a short 165 42 cm core (43.6803 N, 15.9570 E) was selected from the high-marsh and extracted using a 1 m-166 long Eijkelkamp hand gouge corer with a diameter of 50 mm. The relative thinness of organic salt-167 marsh deposits in the Morinje Bay most likely reflects low biological productivity and suspended 168 sediment concentrations in the tidal waters related to the impoverished soils in the limestone 169 catchment area. Nonetheless, the shallow core depth helps minimize the effects of post-170 depositional lowering through sediment compaction (e.g. Brain et al., 2011) because of the limited depth of overburden (e.g. Törnqvist et al., 2008; Horton and Shennan, 2009) and was drilled onto 171 172 the limestone bedrock. The outer surface of the core was carefully cleaned to prevent 173 contamination prior to sub-sampling of the undisturbed internal section and samples were kept 174 refrigerated until ready for analysis. We surveyed core sample altitudes using Real Time Kinetic (RTK) satellite navigation and Leica Na820 optical leveling equipment relative to Croatian 175 176 national geodetic datum (m HVRS71).

177

178 Core samples were prepared at 1 cm intervals for all subsequent analyses. Samples for 179 foraminiferal analysis followed procedures outlined in Horton and Edwards (2006), enumerating 180 foraminiferal tests from sediments between sieve fractions 500 µm and 63 µm transferred to a wet 181 splitter (Scott and Hermelin, 1993) and analyzed wet under a binocular microscope. Our taxonomic 182 identification follows Shaw et al. (2016) where fossil foraminiferal assemblages mirrored those observed in the contemporary environments and are typical of intertidal environments (Edwards
and Wright, 2015). Calcareous taxa *Ammonia*, *Elphidium* and *Quinqueloculina* were recorded as
generic groups (Horton and Edwards, 2006) and followed contemporary studies (Shaw et al.,
2016).

187

We determined the organic matter content of core sediments through Loss-On-Ignition (LOI)
(Ball, 1964), combusting sediment samples at 450°C for four hours to provide supplementary
evidence for intertidal environmental change (Plater et al., 2015).

191

192 **3.1. Chronology**

193 We established sedimentation rates for the salt-marsh core using a composite chronology 194 combining Accelerator Mass Spectrometry (AMS) ¹⁴C dating coupled with short-lived radionuclides (²¹⁰Pb, ²²⁶Ra, ¹³⁷Cs and ²⁴¹Am) for sediments deposited in the past 100 years or so 195 196 (Corbett and Walsh, 2015). Radionuclide activities were analyzed by direct gamma assay at the 197 University of Liverpool Environmental Radioactivity Laboratory. Prior to gamma assay, we 198 determined the dry bulk density of the sediment samples by freeze-drying and weighing. Samples 199 were then lightly disaggregated before being stored for three weeks to allow radioactive 200 equilibration. Samples were analyzed using Ortec HPGe GWL series well-type coaxial low 201 background intrinsic germanium detectors (Appleby et al., 1986). We corrected for the effect of self-absorption of low energy γ -rays within the sample (Appleby and Oldfield, 1992) and ²¹⁰Pb 202 203 ages calculated using the Constant Rate of Supply (CRS) model (Appleby and Oldfield, 1978; Appleby et al., 1979). To further constrain ages obtained via ²¹⁰Pb dating, ¹³⁷Cs activities 204

205 referenced to nuclear weapons testing and the Chernobyl disaster were used as a chronological
206 marker (Appleby, 2001).

207

We selected plant macrofossils for AMS ¹⁴C dating as opposed to bulk sediment dating for 208 209 improved accuracy and reduced uncertainties (Törnquist et al., 2015). Following preparation 210 methods outlined in Kemp et al. (2013b), we identified a varying abundance of Scirpus 211 holoschoenus seeds, a common high-marsh plant of the eastern Adriatic seaboard (Pandža et al., 2007). Three, closely spaced intervals were selected for analysis by AMS ¹⁴C at the NERC 212 213 Radiocarbon Facility, U.K (Table 2). Using a priori knowledge of their stratigraphic position (i.e. 214 the assumption that the lowest most sample was deposited before those above), conventional 215 radiocarbon ages were calibrated using the INTCAL13 calibration curve (Reimer et al., 2013) 216 within a Bayesian age-depth framework using Bchron (Haslett and Parnell, 2008; Parnell et al., 217 2008) to provide 2σ age distributions.

218

219 **3.2. Reconstructing relative sea level**

Our assessment of late Holocene RSL changes in the central-eastern Adriatic are derived from
salt-marsh, tide-gauge and sea-level index points extracted from the quality controlled
Mediterranean Holocene RSL database of Vacchi et al. (2016). Indicative meanings of the proxy
RSL data are detailed in Table 1.

224

Our reconstruction of RSL changes from the salt-marsh environment uses indicative ranges from contemporary foraminiferal distributions (Shaw et al., 2016) to provide estimates of the paleomarsh elevation (PME) from fossil counterparts. We used stratigraphic markers of

228 environmental change (e.g., sediments and organic matter content) as supporting evidence. In 229 microtidal environments such as the Adriatic, using indicative ranges can derive an estimate of 230 PME with equivalent or improved precision over statistically more vigorous techniques (e.g. 231 transfer functions) (Kemp et al., 2017). We identified clusters in fossil foraminiferal assemblages 232 using Partitioning Around Medoids (PAM) cluster analysis (Kaufman and Rousseeuw, 1990) and 233 used contemporary distributions to provide an indicative range (i.e. vertical uncertainty) over 234 which the sample formed relative to mean tide level (MTL). To determine the most statistically 235 representative number of clusters, PAM produces silhouette widths providing a measure of the 236 samples classification. Our RSL reconstruction is restricted to the agglutinated assemblages only 237 within which chronologies and contemporary distributions are constrained. To attain RSL we 238 subtracted PME from surveyed sample elevations related to MTL (Shennan and Horton, 2002), 239 coupled with age estimations provided by the Bchron age-depth model.

240

We analyzed annual measurements from the Split Gradska tide-gauge spanning the period 1955 to 2009 CE. Tide-gauge measurements were analyzed relative to 2009 CE to directly compare RSL changes with the core extraction date of the salt-mash reconstruction. Vertical uncertainties of the tide gauge data were calculated from the standard deviation of annual measurements (\pm 0.03 m) and a temporal uncertainty of \pm 0.5 years follows that of Kemp et al. (2015).

246

We extracted Holocene sea-level index points (n = 23) from Vacchi et al. (2016) for the nearby Island of Vis, central-eastern Adriatic (Fig. 2). These RSL data are based on fossil rims of *Lithophyllum byssoides* (a precise fixed biological indicator of past RSL) and archaeological evidence recorded by Faivre et al. (2013). Temporal uncertainties of the RSL data ranged from \pm

- 251 50 to 244 years with vertical uncertainties of \pm 0.3 m. No reinterpretation of the RSL data was
- applied after Vacchi et al. (2016).
- 253
- **Table 1.** Indicative meanings of proxy RSL data used in RSL reconstruction.

Sea level	Description	Indicative		
indicator		meaning		
Salt-marsh	Organic sediment dominated by salt-marsh plant	MTL-HAT		
	macrofossils and agglutinated foraminifera (e.g. <i>Entzia macrescens</i>).			
*Lithophyllum	Fixed biological fossil rims of Lithophyllum byssoides	MTL-HAT		
byssoides	recorded by Faivre et al. (2013).			
*Archaeological	Functional interpretation of harbour structure (pier and	$MTL\pm0.25$		
	dolia) recorded by Faivre et al. (2103).			
3 (1977)				

255 MTL = mean tide level; HAT = highest astronomical tide. *Lithophyllum byssoides and archaeological evidence

- extracted from the Vacchi et al. (2016) Mediterranean RSL database.
- 257

258 We quantified RSL changes from the salt-marsh, instrumental and a composite RSL record using 259 an Error-In-Variables Integrated Gaussian Process (EIV-IGP) model (Cahill et al., 2015). The 260 EIV-IGP model takes an unevenly distributed RSL time series, prone to vertical and temporal 261 uncertainties, as input and produces estimates of RSL and rates of RSL with 95% credible 262 intervals. The EIV-IGP models rates of RSL change using a Gaussian process (GP) (Williams and Rasmussen, 1996) and models RSL as the integral of the GP (IGP) plus (measured and estimated) 263 264 vertical uncertainty. Temporal uncertainties are accounted for by setting the IGP model in an 265 errors-in-variables (EIV) framework (Dey et al., 2000).

267 **3.3. Glacial-isostatic model predictions**

268 The Adriatic and wider Mediterranean has been a region of great interest in the study of the glacial-269 isostatic adjustment (GIA) process. The large array of biological, archaeological and geological 270 indicators of past sea level have provided an opportunity to tune, test and/or validate GIA models 271 (e.g. Lambeck et al. 2004b; Lambeck & Purcell 2005; Stocchi & Spada 2009; Spada et al. 2009; 272 Lambeck et al. 2011; Vacchi et al. 2016). The recent availability of a standardized Holocene RSL 273 database covering the western Mediterranean basin (Vacchi et al., 2016) has enabled the 274 community to further test global models of the GIA process against RSL data, such as the ICE-275 7G NA (VM7) model (Roy & Peltier 2017; Roy & Peltier 2018).

276

277 Here, we compared the composite RSL record with glacial-isostatic model predictions in the 278 central-eastern Adriatic using the ICE-7G NA (VM7) model (Roy and Peltier, 2017). Our model 279 choice is motivated by the ability of the ICE-7G NA (VM7) model to explain a wide range of 280 geophysical observables related to the GIA process coming from geographically disparate regions 281 (covering formerly glaciated areas, forebulge regions and far-field sites) using a single, simple 282 rheological structure. This independence is important to understand patterns of sea-level evolution in the context of complex local effects, such as tectonic activity (Antonioli et al., 2011). Indeed, 283 284 the ICE-7G NA (VM7) model has been shown to fit a large proportion of the geographically and 285 temporally extensive RSL data set from Vacchi et al. (2016).

286

The ICE-7G_NA (VM7) model is an update to the precursor ICE-6G_C (VM5a) model of Peltier et al. (2015). It includes a modified spherically-symmetric viscosity structure and an updated North American ice-loading history, described in detail in Roy and Peltier (2017). 290

291 4. Results

292 4.1. Salt-marsh stratigraphy

293 Boreholes drilled across the salt-marsh showed an overall increase in sediment depth with distance 294 towards open water (Fig. 3). The lithostratigraphy revealed five main stratigraphic units where 295 sediment accumulation appeared relatively uniform across the site. An unrecoverable (i.e. overly 296 saturated) unit was found between 40 m and 110 m along the transect and overlain by varying silt 297 and clay units (often containing shells fragments) which become progressively more organic 298 towards the surface. An organic salt-marsh peat was restricted to the landward 20 m of the transect 299 where the sample core was extracted. The sediments in the sampled core were comprised of a silty 300 clay bottom unit with low organic content (LOI $\sim 8\%$) between 42 cm and 20 cm. This was overlain 301 by an increasingly organic clay (LOI 10-40%) up to 11 cm and an organic humified peat deposit 302 towards the surface (LOI > 50%) (Fig. 4).



Fig. 3. Simplified cross-sectional profile of the salt-marsh stratigraphy at Jadrtovac showing
location of sample core (see Fig. 2 for transect location).

306

4.2. Indicative meaning based on foraminiferal assemblages

308 The biostratigraphy shows fossil foraminifera preserved throughout the entire sediment core 309 sequence (Fig. 4). An up-core transition from a calcareous-dominated assemblage to agglutinated 310 types is broadly coincident with a shift in sedimentation regime with progressively increasing 311 organic matter content at ~ 27 cm depth. Indeed, for a bundance was significantly higher 312 in line with this sedimentation change, with a mean abundance of 1310 per 5 cm³. Between 42 cm 313 and 32 cm, Ammonia spp., Elphidium spp. and Haynesina germanica dominate before agglutinated 314 types Entzia macrescens, Miliamminia fusca and Trochammina inflata increase in relative 315 abundance. A decrease in the relative abundance of *M. fusca* from 19 cm (71%), corresponds with 316 an increase in E. macrescens and T. inflata towards the surface within the organic peat deposits 317 (LOI >40%).

318

319 Cluster analysis identified two foraminiferal assemblage groups in the fossil environment, 320 essentially discriminating between agglutinated and calcareous dominated assemblages, reflecting 321 the transition from intertidal muds and clays to organic salt-marsh sediments. Two broad indicative 322 meanings are appropriate given the current understanding of contemporary foraminiferal 323 distributions from the central-eastern Adriatic coast (e.g., Shaw et al., 2016). The fossil 324 foraminiferal assemblages mirrored those dominating the contemporary environment. The 325 contemporary distribution of agglutinated types (dominated by E. macrescens and T. inflata) 326 across the salt-marsh platform extends from $0.17 \text{ m} \pm 0.12 \text{ m}$ MTL. Current vertical uncertainties

327 of \pm 0.12 m using salt-marsh sediments are comparable to other RSL studies adopting different 328 sea-level indicators from the central-eastern Adriatic (Vacchi et al., 2016).



Fig. 4. Sample core sediment profile from Jadrtovac salt-marsh core showing lithology (following
that displayed in Fig. 3), relative abundance (%) of the most abundant agglutinated (shaded green)
and calcareous (shaded blue) foraminiferal taxa, organic matter content (LOI %) and results from
cluster analysis. Foraminiferal taxa from left to right; *Entzia macrescens*; *Miliammina fusca*; *Trochammina inflata*; (*Am*) *Ammonia* spp., (*Ep*) *Elphidium* spp., *Haynesina germanica*.

335

336 4.3. Chronology

We established age-depth relationships in the core through short-lived radionuclide analyses and AMS ¹⁴C dating of three intervals between depths 25-30 cm where *Scirpus holoschoenus* seeds were observed within the calcareous-agglutinated foraminiferal assemblage transition (Table 2). The upper ~ 20 cm were constrained from downcore profiles of ²¹⁰Pb and ¹³⁷Cs, respectively (Fig.

341	5). Total ²¹⁰ Pb activity reaches equilibrium with the supporting ²²⁶ Ra at \sim 20 cm depth.
342	Unsupported ²¹⁰ Pb concentrations record a minor discontinuity between 10-13 cm, below which
343	they decline exponentially with depth. Analysis of ¹³⁷ Cs activity shows a relatively well-defined
344	maximum at 9-12 cm (69.9 Bq kg ⁻¹). Its double peak reflects the same event that affected 210 Pb
345	concentrations at this depth. As a result, the 137 Cs/ 210 Pb activity ratio can be a more accurate marker
346	(Plater and Appleby, 2004) to show a well-defined peak between 10-12 cm that reflects peak
347	fallout from the atmospheric testing of nuclear weapons (1963 CE). A second, more recent peak
348	at 5-6 cm (57.1 Bq kg ⁻¹), is interpreted as fallout from the Chernobyl reactor accident (1986 CE).
349	

Table 2. Results from AMS ¹⁴C analyses.

Depth	Laboratory	¹⁴ C Year	$\delta^{13}C$	Modelled $(2\sigma)^{-14}C$	Material dated
(cm)	code	BP (±)	(‰)	ages (CE)*	
25-26	SUERC45020	256 (37)	-25.3	1764-1805	Scirpus
					holoschoenus seeds
26-27	SUERC45021	213 (37)	-26.2	1742-1800	Scirpus
					holoschoenus seeds
28-30	SUERC45022	112 (37)	-26.8	1692-1784	Scirpus
					holoschoenus seeds

351 *¹⁴C ages calibrated within Bchron age-depth modelling software (Haslett and Parnell, 2008; Parnell et al., 2008).
352

Peaks in ¹³⁷Cs broadly correspond to those found from previous research in the Morinje Bay environment where maximum ¹³⁷Cs activity occurs within the upper 20 cm (Mihelčič et al., 2006). The CRS dating model place 1963 at 11.5 cm and 1986 at 5.5 cm, in good agreement with the depths suggested by the ¹³⁷Cs record. The results are relatively unambiguous down to ~ 16 cm, dated to 1920 CE beyond which the uncertainty of age estimates increases. The stratigraphic position of ¹⁴C ages was used to constrain calibrated age distributions within Bchron. The composite chronologies were modelled to provide age estimates with 95% credible intervals for sediments in the upper 30 cm, with an average temporal uncertainty of \pm 19 years (Fig. 5).



Fig. 5. Down-core profiles from short-lived radionuclide analyses (A-D) described in text. Error bars from analyses are smaller than data point symbols used. (E) Bchron age-depth model with 95% credible interval incorporating short-lived radionuclide and AMS ¹⁴C dating.

365 4.4. Late Holocene relative sea-level trends

- 366 Application of the EIV-IGP model to the salt-marsh RSL reconstruction showed a magnitude of
- 367 RSL change of ~ 0.28 m since 1733 CE (Fig. 6) with an average rate of RSL change of 0.95 mm/yr
- 368 over the whole record. Rates of RSL increase from 0.71 mm/yr (-0.67-2.09 mm/yr) to 0.93 mm/yr
- 369 (0.39-1.47 mm/yr) at 1850 CE when RSL was at -0.19 m below present level. Since 1900 CE, RSL
- 370 rose ~ 0.14 m at an average rate of 1.09 mm/yr, increasing to 1.16 mm/yr (-0.08-2.42 mm/yr) at
- 371 2009 CE.



372

Fig. 6. (A) Reconstruction of relative sea-level (RSL) from salt-marsh core at Jadrtovac. (B) ErrorIn-Variables Integrated Gaussian Process (EIV-IGP) model showing mid-points from the RSL
reconstruction with mean, 68% and 95% credible intervals. (C) Rates of RSL (mm/yr) with mean,
68% and 95% credible intervals.

378 Annual measurements from the Split tide-gauge since 1955 CE show a magnitude RSL change of 379 ~ 0.09 m (Fig. 7), concurrent with that recorded by salt-marsh sediments for the same period (~

380 0.08 m; Fig. 6). The average rate of RSL change was 0.60 mm/yr increasing from 0.52 mm/yr (-



381 0.15-1.2 mm/yr) at 1955 CE to 0.67 mm/yr (0.01-1.33 mm/yr) at 2009 CE.

Fig. 7. (A) Annual mean relative sea-level (RSL) trends from the Split Gradska tide-gauge (see Fig. 2 for location). (B) EIV-IGP model showing mid-points from tide-gauge measurements with mean, 68% and 95% credible intervals. (C) Rates of RSL (mm/yr) with mean, 68% and 95%

386 credible intervals. Annual RSL data accessed from the Permanent Service for Mean Sea Level on
 387 13/11/2017 (http://www.psmsl.org/products/trends/trends.txt).

388

389 Late Holocene RSL data from the Island of Vis show a magnitude of RSL change of -1.48 m since 390 715 BCE increasing to -1.05 m by 100 CE at 0.52 mm/yr (-0.82-1.87 mm/yr) (Fig. 8). Between 391 500 and 1000 CE, RSL was at around -0.7 m below present increasing to -0.25 m at 1700 CE, 392 similar to that recorded by the salt-marsh reconstruction for the same time period (-0.28 m). 393 Acknowledging temporal paucity of earlier data, rates of RSL are relatively stable up to 800 CE 394 increasing to 0.77 mm/yr (-0.02-1.57 mm/yr) and then decreasing to 0.13 mm/yr (-0.37-0.64 395 mm/yr) at 1450 CE. The inclusion of the salt-marsh reconstruction and tide-gauge measurements 396 shows the gradual increase in RSL change towards the present.

397

Comparison of the late Holocene composite RSL record with ICE-7G_NA (VM7) model predictions for the central-eastern Adriatic coast reveals a significant offset between the RSL data and predicted results (Fig. 8d). At ~ 700 BCE, the ICE-7G_NA (VM7) model predicts RSL at -0.25 m below present, compared to -1.48 ± 0.3 m suggested by the RSL data. Indeed, this offset is manifest throughout the late Holocene towards the present, with the GIA model predicting magnitudes of RSL changes lower than RSL reconstructions in the central-eastern Adriatic.



405 Fig. 8. (A) Late Holocene relative sea-level (RSL) change from fixed biological (*Lithophyllum*)
406 littoral rims and archaeological evidence from the Island of Vis recorded by Faivre et al. (2013),
407 the salt-marsh RSL reconstruction from Jadrtovac and Split Gradska tide-gauge measurements.
408 (B) Application of EIV-IGP model to the composite RSL showing mid-points with mean, 68%
409 and 95% credible intervals. (C) Rate of RSL (mm/yr) with mean, 68% and 95% credible intervals.
410 (D) Comparison of the composite RSL reconstruction against glacial-isostatic adjustment model
411 predictions of RSL from ICE-7G_NA (VM7) for the study region.

412

413 **5. Discussion**

414 Eustatic and glacio-hydro-isostatic processes have been important driving mechanisms of RSL 415 change in the Mediterranean (e.g., Lambeck and Purcell, 2005; Stocchi and Spada, 2007, 2009; 416 Roy and Peltier, 2018). At more local scales, particularly in the northern Adriatic, geological 417 evidence from geomorphological, sedimentological and archeological sea-level indicators have 418 been utilized to illustrate the importance of differential tectonic movements and local processes 419 (e.g. sediment compaction) affecting late Holocene RSL histories (e.g., Pirazzoli, 2005; Antonioli 420 et al., 2009, 2011; Marriner et al., 2014; Surić et al., 2014; Benjamin et al., 2017; Fontana et al., 421 2017). Furthermore, understanding RSL changes from the late Holocene to the modern period have 422 been restricted by the temporal offset between geological and tide-gauge RSL records (Vacchi et 423 al., 2016), which record large inter-annual and inter-decadal variability (Tsimplis et al., 2012). Our 424 salt-marsh RSL reconstruction overcomes this limitation.

426 **5.1. Late Holocene relative sea levels in the Adriatic**

427 In the northwestern Adriatic, geomorphological and geoarchaeological evidence shows RSL was 428 at -2.0 ± 0.6 m between 1250-1110 BCE, increasing to -1.1 ± 0.3 m at ~ 50 BCE (Fontana et al., 429 2017). In the northeastern Adriatic, archeological evidence shows RSL was between -1.75 and -430 1.4 m ~ 0 CE (Vacchi et al., 2016). More recent RSL data from the Venice and Fruili lagoons 431 shows RSL was at -0.4 ± 0.6 m at ~ 1350 CE and below -0.3 m at 1650 CE (Vacchi et al., 2016). 432 These results compare well with our RSL data from the central-eastern Adriatic (Faivre et al., 433 2013), which show a magnitude of RSL change of -1.48 ± 0.3 m since 715 BCE increasing to -434 1.05 ± 0.3 m by ~ 100 CE and at -0.3 ± 0.3 m between 1350 and 1750 CE. 435 436 We compared the composite RSL record for the central-eastern Adriatic with glacio-isostatic 437 model predictions from ICE-7G NA (VM7) (Roy and Peltier, 2017) (Fig. 8). The magnitude of 438 RSL change from the RSL data (1.48 ± 0.3 m), however, is significantly greater than ICE-7G NA 439 (VM7) model predictions (0.25 m). If we assume the viscosity profile in the GIA model to be 440 accurate, the disparity between the RSL reconstructions and model predictions of RSL could be 441 due to eustatic input and/or tectonics in the absence of local processes (e.g., sediment compaction). 442 For example, the eustatic contribution to sea-level change during the late Holocene from the 443 Antarctic ice sheet may be underestimated in the ICE-7G NA (VM7) model. Reconstructions of 444 RSL from the Mediterranean basin can play a crucial role in understanding the response of the 445 cryosphere to deglacial warming, in particular with respect to the late melting history of the 446 Antarctic ice sheet (Stocchi et al., 2009; Roy and Peltier, 2018). Indeed, one of the key distinctions

448 component of sea-level change largely driven by the Antarctic and Greenlandic ice sheets.

between various reconstructions of ice sheet deglaciation history lies in the late Holocene eustatic

449 Whereas the ICE-6G C and ICE-7G NA models of ice sheet loading history show around 2 m of 450 global mean sea level (GMSL) rise since 7ka, with no substantial increase after 4ka, other models 451 have inferred up to 6 m of GMSL rise since 7ka, with more than 80 cm of this change having 452 occurred after 4ka (Lambeck et al., 2014). It is important to place this observation in the broader 453 context of the quality of the fit provided by the ICE-7G NA (VM7) to RSL data in the rest of the 454 western Mediterranean basin (Fig. 9). Roy and Peltier (2018) found their model to perform very 455 well in France, around the Ligurian Sea, in Corsica and in Sardinia. However, the authors identified 456 regions of sustained misfits between the model predictions and the Vacchi et al. (2016) database, 457 notably in central Spain, southwest Italy and in the Adriatic (Fig. 9). The GIA model is able to fit 458 the majority of RSL reconstructions between 3ka and 1ka in the western Mediterranean basin to 459 within two standard deviations. However, the RSL data that misfit with model predictions by 460 greater than two standard deviations are concentrated in the Adriatic.

461

462 The misfit between RSL data and model predictions provide support for the existence of tectonic 463 effects in the Adriatic, rather than an issue in the rate of GMSL rise included in the ICE-7G NA 464 model. The difference between the reconstruction and the model suggest a tectonic subsidence of 465 0.45 ± 0.6 mm/yr. Although a full assessment of the influence of tectonics operating in the region 466 is challenging, the large variations in GPS vertical velocities (Fig. 2) observed around the Adriatic 467 Sea (Serpelloni et al., 2013) supports the idea of substantial tectonic motion. It should also be noted 468 that the presence of a complex tectonic setting should also be studied in the context of potential 469 local lateral heterogeneity in the viscosity of the mantle. Due to the small scale of the Adriatic Sea, 470 any sensitivity to such variations would be expected to be limited to the uppermost layers of the

- 471 mantle. Nonetheless, any rigorous determination of the geological evolution throughout the region
- 472 will need to consider these effects.



473

Fig. 9. Quality of the fit to the late Holocene RSL data from the western Mediterranean (Vacchi et al., 2016) provided by the model predictions of the ICE-7G_NA (VM7) model, centered on 0 CE (\pm 1000 years). Green dots represent an agreement between the model prediction of RSL and the local RSL reconstruction within 2 standard deviations (SD), while red dots indicate locations where the difference between the model predictions and the RSL reconstruction is greater than 2 SD. Contours represent ICE-7G_NA (VM7) model predictions of RSL (m) at 0 CE.

The influence of local processes including paleotidal-range change and sediment compaction to Holocene RSL histories (e.g. Horton et al., 2013) may also contribute to differences between glacial-isostatic model predictions and RSL data. A lack of Mediterranean based studies currently restricts the assessment of paleotidal-range changes (e.g. Hill et al., 2011; Griffiths and Hill, 2015)

to Mediterranean Holocene RSL data (e.g. Vacchi et al., 2016). However, given the local micro tidal range and time period involved, we can consider the influence of paleotidal-range changes to be negligible. Furthermore, the influence of sediment compaction, is also negligible due to the limited depth of overburden (e.g. Törnqvist et al., 2008; Horton and Shennan, 2009) of the relatively thin organic salt-marsh peat deposits at Jadrtovac. Nonetheless, future RSL studies in the Adriatic and Mediterranean region accounting for these processes, would inherently provide more accurate predictions of RSL change.

492

493 **5.2.** Centennial scale relative sea level variability

494 Climate-driven centennial sea-level variability superposed on late Holocene RSL are expressed at 495 the global scale (Kopp et al., 2016) with the transition from the Medieval Climate Anomaly (MCA) 496 to the Little Ice Age (LIA) coinciding with a reduction in air and ocean temperatures (Mann et al., 497 2009; Marcott et al., 2013; Rosenthal et al., 2017). Application of the EIV-IGP model to the 498 geological data from Vis shows a (subtle) increase and decrease in RSL rate, occurring at 800 CE 499 and 1450 CE, respectively. This broadly coincides with the MCA to LIA transition which Faivre 500 et al. (2013) suggested a response of central-eastern Adriatic sea levels similar to the North Atlantic 501 based on comparisons of RSL trends with salt-marsh based RSL reconstructions from North 502 America (Kemp et al., 2011). While the temporal coverage and vertical resolution of late Holocene 503 RSL data from the western Mediterranean currently restricts more local interpretations (e.g. 504 Vacchi et al., 2016), variability of late Holocene RSL in the eastern Mediterranean has been 505 reported (Sivan et al., 2004). Archaeological and biological proxy data from Israel support 506 inferences for sea-level variability between 900 and 1300 CE (Toker et al., 2012). Indeed, the

507 climatic deterioration during the LIA has also been associated with a period of increased 508 storminess throughout the Mediterranean region (Marriner et al., 2017).

509

510 **5.3. Modern sea-level rise**

511 Empirical modelling of proxy and instrumental RSL records has enabled inferences regarding 512 timing the onset of modern sea-level rise (Kopp et al., 2016). At the global scale, sea levels began 513 rising around 1860 CE (Kemp et al., 2011; Kopp et al., 2016), synchronous with sustained 514 industrial-era warming of the tropical oceans and Northern Hemisphere continents (Abram et al., 515 2016). Here, we applied the EIV-IGP model to our salt-marsh reconstruction, which captures the 516 dynamic evolution of sea-level change with robust consideration of sources of uncertainty (Cahill 517 et al., 2015). Importantly, the EIV-IGP model shows a subtle but constant increase in the mean 518 rate of RSL rise from 0.71 mm/yr (-0.67-2.09 mm/yr) to 1.16 mm/yr (-0.08-2.42 mm/yr) between 519 1733 CE and 2009 CE (Fig. 6c). Indeed, this subtle increase in mean RSL rate stems from the 520 deviation of sea-level trends recorded up to ~ 1450 CE (Fig. 8c).

521

522 Uncertainties in constraining age-depth relationships in the salt-marsh reconstruction during the 523 nineteenth century may preclude important inferences regarding timing of the onset of modern 524 RSL rise in the Mediterranean. Records of RSL change from twentieth century tide-gauge stations 525 in the Mediterranean show RSL rising at a rate of 1.1 to 1.3 mm/yr (Tsimplis and Baker, 2000; 526 Orlić and Pasarić, 2000; Marcos and Tsimplis, 2008). Zerbini et al. (2017) also report a rising RSL 527 trend of 1.2 to 1.3 mm/yr (\pm 0.2 to 0.5 mm/yr) from their analyses. The salt-marsh RSL 528 reconstruction supports these findings with an average RSL rate of ~ 1.1 mm/yr between 1900 and 529 2009. While a lower average rate of RSL change was recorded by the Split tide-gauge (0.60

530 mm/yr), this reflects a deviation of sea levels recorded by Adriatic and Mediterranean tide-gauge 531 stations during the latter half of the twentieth century. A decrease in RSL rate between 1960-1993 532 (Tsimplis and Baker, 2000; Marcos and Tsimplis, 2008) coincided with a period of higher 533 atmospheric pressure and evaporation over the basin driven by the high state of the North Atlantic 534 Oscillation (Tsimplis and Josey, 2001).

535

536 6. Conclusions

537 Reconstructions of RSL change along the central-eastern coast of Croatia offer new insight to the 538 late Holocene sea-level history of the central-eastern Adriatic region. We reconstructed RSL using 539 salt-marsh sediments and foraminifera that underpins their under-utilized potential to derive RSL 540 changes in the Mediterranean. Fossil foraminifera enumerated from a short sediment core were 541 constrained vertically by contemporary foraminiferal distributions, and temporally, by radiometric 542 dating techniques within a Bayesian age-depth framework. The reconstruction shows RSL rose \sim 543 0.28 m since \sim 1733 CE, with a magnitude RSL change of \sim 0.14 m comparable to tide-gauge 544 records during the twentieth century. We modelled RSL changes using the EIV-IGP model (Cahill 545 et al., 2015) showing rates of RSL change increasing from 0.71 mm/yr (-0.67-2.09 mm/yr) at \sim 546 1733 CE to 0.93 mm/yr (0.39-1.47 mm/yr) at 1850 CE. Average rates of RSL during the twentieth 547 century, rising at ~ 1.1 mm/yr, are analogous with the instrumental measurements.

548

We compared a composite RSL record combining the tide-gauge and salt-marsh reconstruction with local published sea-level index points (n=23) (Faivre et al., 2013) against ICE-7G_NA (VM7) glacio-isostatic model predictions (Roy and Peltier, 2017) for the last ~ 2700 years. The magnitude of RSL change from the RSL reconstruction (1.48 m) differs from the glacio-isostatic model prediction by more than 1 m, supporting subsidence rates driven by the Adriatic tectonic framework of 0.45 ± 0.6 mm/yr. Application of the EIV-IGP model supports evidence for late Holocene sea-level variability with rates of RSL rise decreasing from 0.77 mm/yr (-0.02-1.57 mm/yr) at 800 CE to 0.13 mm/yr (-0.37-0.64 mm/yr) at ~ 1450 CE. The temporal coverage of the salt-marsh reconstruction bridging RSL changes from the late Holocene to the modern instrumental period shows the gradual increase in mean RSL rate towards the present.

559

560 Acknowledgements

561 This research was supported by the National Environment Research Council (NERC) U.K. PhD 562 studentship award SR1686482, Singapore Ministry of Education Academic Research Fund Tier 1 563 RG119/17, National Research Foundation Singapore and the Singapore Ministry of Education 564 under the Research Centres of Excellence initiative. We gratefully acknowledge the NERC 565 radiocarbon steering committee for granting radiocarbon dating support (Allocation Number 566 1678.1012) and thank Peter Appleby, University of Liverpool, for the radionuclide analyses. This 567 article is a contribution to PALSEA2 (Palaeo-Constraints on Sea-Level Rise), International 568 Geoscience Program (IGCP) Project 639, "Sea Level Change from Minutes to Millennia," and 569 INQUA Project 1601P "Geographic variability of HOLocene relative SEA level (HOLSEA)." 570 This work is Earth Observatory of Singapore contribution no. 204. We dedicate this paper to Peter 571 John Shaw.

573 **References**

- 574 Abram, N.J., McGregor, H.V., Tierney, J.E., Evans, M.N., McKay, N.P., Kaufman, D.S.,
- 575 Consortium, the P. 2k, Thirumalai, K., Martrat, B., Goosse, H., Phipps, S.J., Steig, E.J., Kilbourne,
- 576 K.H., Saenger, C.P., Zinke, J., Leduc, G., Addison, J.A., Mortyn, P.G., Seidenkrantz, M.-S., Sicre,
- 577 M.-A., Selvaraj, K., Filipsson, H.L., Neukom, R., Gergis, J., Curran, M.A.J., Gunten, L. von, 2016.
- 578 Early onset of industrial-era warming across the oceans and continents. Nature 536, 411.
 579 https://doi.org/10.1038/nature19082
- Altiner, Y., 2006. Present-day tectonics in and around the Adria plate inferred from GPS measurements. Special Paper of the Geological Society of America 409. https://doi.org/10.1130/2006.2409(03)
- Anderson, H., Jackson, J., 1987. Active tectonics of the Adriatic Region. Geophysical Journal of
 the Royal Astronomical Society 91, 937–98;3. https://doi.org/10.1111/j.1365246X.1987.tb01675.x
- 586 Antonioli, F., Faivre, S., Ferranti, L., Monaco, C., 2011. Tectonic contribution to relative sea level
- 587 change. Quaternary International 232, 1–4. https://doi.org/10.1016/j.quaint.2010.10.003
- 588 Antonioli, F., Ferranti, L., Fontana, A., Amorosi, A., Bondesan, A., Braitenberg, C., Dutton, A.,
- 589 Fontolan, G., Furlani, S., Lambeck, K., Mastronuzzi, G., Monaco, C., Spada, G., Stocchi, P., 2009.
- 590 Holocene relative sea-level changes and vertical movements along the Italian and Istrian
- 591 coastlines. Quaternary International 206, 102–133. https://doi.org/10.1016/j.quaint.2008.11.008
- 592 Appleby, P.G., 2001. Chronostratigraphic Techniques in Recent Sediments, in: Last, W., Smol, J.
- 593 (Eds.), Tracking Environmental Change Using Lake Sediments, Developments in
- 594 Paleoenvironmental Research. KluwerAcademic Publishers, Dordrecht, pp. 171–203.

- 595 Appleby, P.G., Nolan, P.J., Gifford, D.W., Godfrey, M.J., Oldfield, F., Anderson, N.J., Battarbee,
- 596 R.W., 1986. 210Pb dating by low background gamma counting. Hydrobiologia 143, 21-27.
- 597 https://doi.org/10.1007/bf00026640
- 598 Appleby, P.G., Oldfield, F., 1978. The calculation of lead-210 dates assuming a constant rate of
- supply of unsupported 210Pb to the sediment. CATENA 5, 1–8. https://doi.org/10.1016/s03418162(78)80002-2
- Appleby, P.G., Oldfield, F., Thompson, R., Huttunens, P., Tolone, K., 1979. 210Pb dating of
 annually laminated lake sediments from Finland. Nature 280, 53–55.
- Bačani, A., Koch, G., Bergant, S., Šparica, M., Viličić, D., Dolenec, T., Vreča, P., Ibrahimpašić,
- 604 H., 2004. Origin of Recent Organic-Rich Sediments from Morinje Bay (Northern Dalmatia,
- 605 Croatia): Aspects of Hydrological and Hydrogeological Impact. Abstracts, Scientific Sessions,
 606 Part 2, 32nd International Geological Congress, Florence.
- Ball, D.F., 1964. Loss-on-ignition as an estimate of organic matter and organic carbon in noncalcareous soils. Journal of Soil Science 15, 84–92. https://doi.org/10.1111/j.13652389.1964.tb00247.x
- 610 Barlow, N.L.M., Long, A.J., Saher, M.H., Gehrels, W.R., Garnett, M.H., Scaife, R.G., 2014. Salt-
- 611 marsh reconstructions of relative sea-level change in the North Atlantic during the last 2000 years.
- 612 Quaternary Science Reviews 99, 1–16. https://doi.org/10.1016/j.quascirev.2014.06.008
- Barlow, N.L.M., Shennan, I., Long, A.J., Gehrels, W.R., Saher, M.H., Woodroffe, S.A., Hillier,
- 614 C., 2013. Salt marshes as late Holocene tide gauges. Global and Planetary Change 106, 90–110.
- 615 https://doi.org/10.1016/j.gloplacha.2013.03.003

- 616 Benac, Č., Juračić, M., Bakran-Petricioli, T., 2004. Submerged tidal notches in the Rijeka Bay NE
- 617 Adriatic Sea: indicators of relative sea-level change and of recent tectonic movements. Marine
- 618 Geology 212, 21–33. https://doi.org/10.1016/j.margeo.2004.09.002
- 619 Benac, C., Juracic, M., Blaskovic, I., 2008. Tidal notches in Vinodol Channel and Bakar Bay, NE
- 620 Adriatic Sea: Indicators of recent tectonics. Marine Geology 248, 151–160.
 621 https://doi.org/10.1016/j.margeo.2007.10.010
- 622 Benjamin, J., Rovere, A., Fontana, A., Furlani, S., Vacchi, M., Inglis, R.H., Galili, E., Antonioli,
- 623 F., Sivan, D., Miko, S., Mourtzas, N., Felja, I., Meredith-Williams, M., Goodman-Tchernov, B.,
- 624 Kolaiti, E., Anzidei, M., Gehrels, R., 2017. Late Quaternary sea-level changes and early human
- 625 societies in the central and eastern Mediterranean Basin: An interdisciplinary review. Quaternary
- 626 International 449, 29–57. https://doi.org/10.1016/j.quaint.2017.06.025
- 627 Brain, M.J., Kemp, A.C., Hawkes, A.D., Engelhart, S.E., Vane, C.H., Cahill, N., Hill, T.D.,
- 628 Donnelly, J.P., Horton, B.P., 2017. Exploring mechanisms of compaction in salt-marsh sediments
- 629 using Common Era relative sea-level reconstructions. Quaternary Science Reviews 167, 96–111.
- 630 https://doi.org/10.1016/j.quascirev.2017.04.027
- Brain, M.J., Long, A.J., Petley, D.N., Horton, B.P., Allison, R.J., 2011. Compression behaviour of
- 632 minerogenic low energy intertidal sediments. Sedimentary Geology 233, 28-41.
- 633 https://doi.org/10.1016/j.sedgeo.2010.10.005
- Buble, G., Bennett, R.A., Hreinsdóttir, S., 2010. Tide gauge and GPS measurements of crustal
- motion and sea level rise along the eastern margin of Adria. Journal of Geophysical Research 115,
- 636 B02404. https://doi.org/10.1029/2008jb006155

- 637 Cahill, N., Kemp, A.C., Horton, B.P., Parnell, A.C., 2015. Modeling sea-level change using errors-
- 638 in-variables integrated Gaussian processes. Ann. Appl. Stat. 9, 547-571.
- 639 https://doi.org/10.1214/15-AOAS824
- 640 Carminati, E., Martinelli, G., Severi, P., 2003. Influence of glacial cycles and tectonics on natural
- 641 subsidence in the Po Plain (Northern Italy): Insights from 14C ages. Geochemistry, Geophysics,
- 642 Geosystems 4. https://doi.org/10.1029/2002GC000481
- 643 Ciabatti, M., Curzi, P., Ricci Lucchi, F. 1987. Quaternary sedimentation in the Central Adriatic
 644 Sea. Giornale di Geologia 49, 113–125.
- 645 Corbett, D.R., Walsh, J. p., 2015. 210Lead and 137Cesium, in: Shennan, I., Long, A.J., Horton,
- 646 B.P. (Eds.), Handbook of Sea-Level Research. John Wiley & Sons, Ltd, pp. 361-372.
- 647 https://doi.org/10.1002/9781118452547.ch24
- 648 Cushman-Roisin, B., Naimie, C.E., 2002. A 3D finite-element model of the Adriatic tides. Journal
- of Marine Systems 37, 279–297. https://doi.org/10.1016/S0924-7963(02)00204-X
- 650 D'Agostino, N., Avallone, A., Cheloni, D., D'Anastasio, E., Mantenuto, S., Selvaggi, G., 2008.
- 651 Active tectonics of the Adriatic region from GPS and earthquake slip vectors. J. Geophys. Res.
- 652 113, B12413. https://doi.org/10.1029/2008JB005860
- Dangendorf, S., Marcos, M., Müller, A., Zorita, E., Riva, R., Berk, K., Jensen, J., 2015. Detecting
- 654 anthropogenic footprints in sea level rise. Nature Communications 6, 7849.
- 655 https://doi.org/10.1038/ncomms8849
- 656 Devoti, R., D'Agostino, N., Serpelloni, E., Pietrantonio, G., Riguzzi, F., Avallone, A., Cavaliere,
- A., Cheloni, D., Cecere, G., D'Ambrosio, C., Franco, L., Selvaggi, G., Metois, M., Esposito, A.,
- 658 Sepe, V., Galvani, A., Anzidei, M., 2017. A Combined Velocity Field of the Mediterranean
- 659 Region. Ann. Geophys. 60, S0215. https://doi.org/10.4401/ag-7059

- 660 Devy, D.K., Ghosh, S.K., Mallick, B.K., 2000. Generalized Linear Models: A Bayesian
 661 Perspective. Marcel Dekker, New York.
- Donnelly, J.P., Cleary, P., Newby, P., Ettinger, R., 2004. Coupling instrumental and geological
- 663 records of sea-level change: Evidence from southern New England of an increase in the rate of
- 664 sea-level rise in the late 19th century. Geophysical Research Letters 31, L05203.
- 665 https://doi.org/10.1029/2003gl018933
- Edwards, R., Wright, A., 2015. Foraminifera, in: Shennan, I., Long, A.J., Horton, B.P. (Eds.),
 Handbook of Sea-Level Research. John Wiley & Sons, Ltd, pp. 191–217.
- 668 Engelhart, S.E., Horton, B.P., 2012. Holocene sea level database for the Atlantic coast of the
- 669 United States. Quaternary Science Reviews, Coastal Change during the Late Quaternary 54, 12–
- 670 25. https://doi.org/10.1016/j.quascirev.2011.09.013
- 671 Faccenna, C., Becker, T.W., Auer, L., Billi, A., Boschi, L., Brun, J.P., Capitanio, F.A., Funiciello,
- 672 F., Horvàth, F., Jolivet, L., Piromallo, C., Royden, L., Rossetti, F., Serpelloni, E., 2014. Mantle
- dynamics in the Mediterranean. Rev. Geophys. 52, 2013RG000444.
 https://doi.org/10.1002/2013RG000444
- 675 Faivre, S., Bakran-Petricioli, T., Horvatinčić, N., Sironić, A., 2013. Distinct phases of relative sea
- 676 level changes in the central Adriatic during the last 1500 years influence of climatic variations?
- 677 Palaeogeography, Palaeoclimatology, Palaeoecology 369, 163–174.
 678 https://doi.org/10.1016/j.palaeo.2012.10.016
- 679 Ferla, M., Cordella, M., Michielli, L., Rusconi, A., 2007. Long-term variations on sea level and
- tidal regime in the lagoon of Venice. Estuarine, Coastal and Shelf Science 75, 214-222.
- 681 https://doi.org/10.1016/j.ecss.2007.03.037

- Flemming, N.C., 1969. Archaeological Evidence for Eustatic Change of Sea Level and Earth
 Movements in the Western Mediterranean During the Last 2,000 Years. Geological Society of
- America Special Papers 109, 1–98. https://doi.org/10.1130/SPE109-p1
- Flemming, N.C., Webb, C.O., 1986. Tectonic and eustatic coastal changes during the last 10,000
- 986 years derived from archaeological data. Zeitschrift für Geomorphologie, Supplementband 62, 1–
 987 29.
- 688 Fontana, A., Vinci, G., Tasca, G., Mozzi, P., Vacchi, M., Bivi, G., Salvador, S., Rossato, S.,
- Antonioli, F., Asioli, A., Bresolin, M., Di Mario, F., Hajdas, I., 2017. Lagoonal settlements and
- 690 relative sea level during Bronze Age in Northern Adriatic: Geoarchaeological evidence and
- 691 paleogeographic constraints. Quaternary International, Quaternary coastal and marine studies in
- 692 Central Mediterranean 439, 17–36. https://doi.org/10.1016/j.quaint.2016.12.038
- Furlani, S., Biolchi, S., Cucchi, F., Antonioli, F., Busetti, M., Melis, R., 2011. Tectonic effects on
 Late Holocene sea level changes in the Gulf of Trieste (NE Adriatic Sea, Italy). Quaternary
 International 232, 144–157. https://doi.org/10.1016/j.quaint.2010.06.012
- 696 Gehrels, W.R., Callard, S.L., Moss, P.T., Marshall, W.A., Blaauw, M., Hunter, J., Milton, J.A.,
- 697 Garnett, M.H., 2012. Nineteenth and twentieth century sea-level changes in Tasmania and New
- 698 Zealand. Earth and Planetary Science Letters 315–316, 94–102.
- 699 https://doi.org/10.1016/j.epsl.2011.08.046
- 700 Gehrels, W.R., Hayward, B.W., Newnham, R.M., Southall, K.E., 2008. A 20th century
- 701 acceleration of sea-level rise in New Zealand. Geophysical Research Letters 35, L02717.
- 702 https://doi.org/10.1029/2007gl032632

- 703 Gehrels, W.R., Kirby, J.R., Prokoph, A., Newnham, R.M., Achterberg, E.P., Evans, H., Black, S.,
- 704 Scott, D.B., 2005. Onset of recent rapid sea-level rise in the western Atlantic Ocean. Quaternary
- 705 Science Reviews 24, 2083–2100. https://doi.org/10.1016/j.quascirev.2004.11.016
- 706 Grenerczy, G., Sella, G., Stein, S., Kenyeres, A., 2005. Tectonic implications of the GPS velocity
- 707 field in the northern Adriatic region. Geophys. Res. Lett. 32, L16311.
 708 https://doi.org/10.1029/2005GL022947
- 709 Griffiths, S.D., Hill, D.F., 2015. Tidal modeling, in: Shennan, I., Long, A.J., Horton, B.P. (Eds.),
- 710 Handbook of Sea-Level Research. John Wiley & Sons, Ltd, pp. 438–451.
- 711 Grinsted, A., Moore, J.C., Jevrejeva, S., 2009. Reconstructing sea level from paleo and projected
- 712 temperatures 200 to 2100. Clim Dyn 34, 461–472. https://doi.org/10.1007/s00382-008-0507-2
- 713 Haslett, J., Parnell, A., 2008. A simple monotone process with application to radiocarbon-dated
- depth chronologies. Journal of the Royal Statistical Society: Series C (Applied Statistics) 57, 399–
- 715 418. https://doi.org/10.1111/j.1467-9876.2008.00623.x
- Herak, D., Herak, M., Prelogović, E., Markušić, S., Markulin, Ž., 2005. Jabuka island (Central
 Adriatic Sea) earthquakes of 2003. Tectonophysics 398, 167–180.
 https://doi.org/10.1016/j.tecto.2005.01.007
- 719 Herak, D., Sović, I., Cecić, I., Živčić, M., Dasović, I., Herak, M., 2017. Historical Seismicity of
- the Rijeka Region (Northwest External Dinarides, Croatia)—Part I: Earthquakes of 1750, 1838,
- and 1904 in the Bakar Epicentral Area. Seismological Research Letters 88, 904–915.
- 722 https://doi.org/10.1785/0220170014
- Herak, M., Herak, D., Markušić, S., 1996. Revision of the earthquake catalogue and seismicity of
- 724 Croatia, 1908–1992. Terra Nova 8, 86–94. https://doi.org/10.1111/j.1365-3121.1996.tb00728.x

- Hill, D.F., Griffiths, S.D., Peltier, W.R., Horton, B.P., Törnqvist, T.E., 2011. High-resolution
 numerical modeling of tides in the western Atlantic, Gulf of Mexico, and Caribbean Sea during
- 727 the Holocene. J. Geophys. Res. 116, C10014. https://doi.org/10.1029/2010JC006896
- 728 Horton, B.P., Edwards, R.J., 2006. Quantifying Holocene Sea Level Change Using Intertidal
- Foraminifera: Lessons from the British Isles. Cushman Foundation for Foraminiferal Research,
 Special Publication 40, 97.
- 731 Horton, B.P., Engelhart, S.E., Hill, D.F., Kemp, A.C., Nikitina, D., Miller, K.G., Peltier, W.R.,
- 732 2013. Influence of tidal-range change and sediment compaction on Holocene relative sea-level
- change in New Jersey, USA. J. Quaternary Sci. 28, 403–411. https://doi.org/10.1002/jqs.2634
- 734 Horton, B.P., Shennan, I., 2009. Compaction of Holocene strata and the implications for relative 735 sealevel change on the east of England. Geology 37, 1083–1086. coast 736 https://doi.org/10.1130/G30042A.1
- Hydrographic Institute, 1955. Report on sea level measurement along the eastern Adriatic coast.Split.
- Jevrejeva, S., Grinsted, A., Moore, J.C., 2009. Anthropogenic forcing dominates sea level rise
 since 1850. Geophysical Research Letters 36. https://doi.org/10.1029/2009GL040216
- 741 Kaufman, L., Rousseeuw, P.J., 1990. Partitioning Around Medoids (Program PAM), in: Finding
- 742 Groups in Data. John Wiley & Sons, Inc., pp. 68–125. https://doi.org/10.1002/9780470316801.ch2
- 743 Kemp, A.C., Hawkes, A.D., Donnelly, J.P., Vane, C.H., Horton, B.P., Hill, T.D., Anisfeld, S.C.,
- Parnell, A.C., Cahill, N., 2015. Relative sea-level change in Connecticut (USA) during the last
- 745 2200 yrs. Earth and Planetary Science Letters 428, 217–229.
 746 https://doi.org/10.1016/j.epsl.2015.07.034

- 747 Kemp, A.C., Horton, B.P., Culver, S.J., Corbett, D.R., van de Plassche, O., Gehrels, W.R.,
- 748 Douglas, B.C., Parnell, A.C., 2009. Timing and magnitude of recent accelerated sea-level rise
- 749 (North Carolina, United States). Geology 37, 1035–1038. https://doi.org/10.1130/g30352a.1
- 750 Kemp, A.C., Horton, B.P., Donnelly, J.P., Mann, M.E., Vermeer, M., Rahmstorf, S., 2011. Climate
- related sea-level variations over the past two millennia. Proceedings of the National Academy of
- 752 Sciences. https://doi.org/10.1073/pnas.1015619108
- 753 Kemp, A.C., Horton, B.P., Vane, C.H., Bernhardt, C.E., Corbett, D.R., Engelhart, S.E., Anisfeld,
- S.C., Parnell, A.C., Cahill, N., 2013. Sea-level change during the last 2500 years in New Jersey,
- USA. Quaternary Science Reviews 81, 90–104. https://doi.org/10.1016/j.quascirev.2013.09.024
- 756 Kemp, A.C., Kegel, J.J., Culver, S.J., Barber, D.C., Mallinson, D.J., Leorri, E., Bernhardt, C.E.,
- 757 Cahill, N., Riggs, S.R., Woodson, A.L., Mulligan, R.P., Horton, B.P., 2017. Extended late
- Holocene relative sea-level histories for North Carolina, USA. Quaternary Science Reviews 160,
- 759 13–30. https://doi.org/10.1016/j.quascirev.2017.01.012
- 760 Kemp, A.C., Nelson, A.R., Horton, B.P., 2013. 14.31 Radiocarbon Dating of Plant Macrofossils
- 761 from Tidal-Marsh Sediment, in: Shroder, J.F. (Ed.), Treatise on Geomorphology. Academic Press,
- 762 San Diego, pp. 370–388.
- 763 Kopp, R.E., Kemp, A.C., Bittermann, K., Horton, B.P., Donnelly, J.P., Gehrels, W.R., Hay, C.C.,
- Mitrovica, J.X., Morrow, E.D., Rahmstorf, S., 2016. Temperature-driven global sea-level
 variability in the Common Era. PNAS 113, E1434–E1441.
 https://doi.org/10.1073/pnas.1517056113
- 767 Korbar, T., 2009. Orogenic evolution of the External Dinarides in the NE Adriatic region: a model
- constrained by tectonostratigraphy of Upper Cretaceous to Paleogene carbonates. Earth-Science
- 769 Reviews 96, 296–312. https://doi.org/10.1016/j.earscirev.2009.07.004

- T70 Lambeck, K., Anzidei, M., Antonioli, F., Benini, A., Esposito, A., 2004a. Sea level in Roman time
- in the Central Mediterranean and implications for recent change. Earth and Planetary Science
- 772 Letters 224, 563–575. https://doi.org/10.1016/j.epsl.2004.05.031
- T73 Lambeck, K., Antonioli, F., Purcell, A., Silenzi, S., 2004b. Sea-level change along the Italian coast
- 774 for the past 10,000 yr. Quaternary Science Reviews 23, 1567–1598.
 775 https://doi.org/10.1016/j.quascirev.2004.02.009
- Lambeck, K., Purcell, A., 2005. Sea-level change in the Mediterranean Sea since the LGM: model
- predictions for tectonically stable areas. Quaternary Science Reviews 24, 1969–1988.
 https://doi.org/10.1016/j.quascirev.2004.06.025
- 179 Lambeck, K., Antonioli, F., Anzidei, M., Ferranti, L., Leoni, G., Scicchitano, G., Silenzi, S., 2011.
- 780 Sea level change along the Italian coast during the Holocene and projections for the future.
- 781 Quaternary International 232, 250–257. https://doi.org/10.1016/j.quaint.2010.04.026
- Lambeck, K., Rouby, H., Purcell, A., Sun, Y., Sambridge, M., 2014. Sea level and global ice
 volumes from the Last Glacial Maximum to the Holocene. PNAS 111, 15296–15303.
 https://doi.org/10.1073/pnas.1411762111
- 785 Mann, M.E., Zhang, Z., Rutherford, S., Bradley, R.S., Hughes, M.K., Shindell, D., Ammann, C.,
- 786 Faluvegi, G., Ni, F., 2009. Global Signatures and Dynamical Origins of the Little Ice Age and
- 787 Medieval Climate Anomaly. Science 326, 1256–1260. https://doi.org/10.1126/science.1177303
- Marcos, M., Tsimplis, M.N., 2008. Coastal sea level trends in Southern Europe. Geophysical
 Journal International 175, 70–82.
- 790 Marcott, S.A., Shakun, J.D., Clark, P.U., Mix, A.C., 2013. A Reconstruction of Regional and
- 791 Global Temperature for the Past 11,300 Years. Science 339, 1198–1201.
- 792 https://doi.org/10.1126/science.1228026

- 793 Marriner, N., Morhange, C., Faivre, S., Flaux, C., Vacchi, M., Miko, S., Dumas, V., Boetto, G.,
- 794 Radic Rossi, I., 2014. Post-Roman sea-level changes on Pag Island (Adriatic Sea): Dating
- 795 Croatia's "enigmatic" coastal notch? Geomorphology 221, 83–94.
 796 https://doi.org/10.1016/j.geomorph.2014.06.002
- 797 Marriner, N., Kaniewski, D., Morhange, C., Flaux, C., Giaime, M., Vacchi, M., Goff, J., 2017.
- 798 Tsunamis in the geological record: Making waves with a cautionary tale from the Mediterranean.
- 799 Science Advances 3, e1700485. https://doi.org/10.1126/sciadv.1700485
- 800 Marjanović, M., Bačić, Ž., Bašić, T., 2012. Determination of Horizontal and Vertical Movements
- 801 of the Adriatic Microplate on the Basis of GPS Measurements, in: Kenyon, S., Pacino, M.C., Marti,
- 802 U. (Eds.), Geodesy for Planet Earth, International Association of Geodesy Symposia. Springer
- 803 Berlin Heidelberg, pp. 683–688.
- 804 Marshall, W., 2015. Chronohorizons, in: Shennan, I., Long, A.J., Horton, B.P. (Eds.), Handbook
- 805 of Sea-Level Research. John Wiley & Sons, Ltd, pp. 373–385.
 806 https://doi.org/10.1002/9781118452547.ch25
- McKenzie, D., 1972. Active Tectonics of the Mediterranean Region. Geophysical Journal
 International 30, 109–185. https://doi.org/10.1111/j.1365-246X.1972.tb02351.x
- 809 Oldow, J.S., Ferranti, L., Lewis, D.S., Campbell, J.K., D'Argenio, B., Catalano, R., Pappone, G.,
- 810 Carmignani, L., Conti, P., Aiken, C.L.V., 2002. Active fragmentation of Adria, the north African
- 811 promontory, central Mediterranean orogen. Geology 30, 779–782. https://doi.org/10.1130/0091-
- 812 7613(2002)030<0779:AFOATN>2.0.CO;2
- 813 Orlić, M., Gačić, M., La Violette, P.E., 1992. The currents and circulation of the Adriatic Sea.
- 814 Oceanologica Acta 15, 109–124.

- 815 Orlić, M., Kuzmić, M., Pasarić, Z., 1994. Response of the Adriatic Sea to the bora and sirocco
- 816 forcing. Continental Shelf Research 14, 91–116. https://doi.org/10.1016/0278-4343(94)90007-8
- 817 Orlić, M., Pašarić, M., 2000. Sea-level changes and crustal movements recorded along the east
- 818 Adriatic coast. Nuovo cimento della Società italiana di fisica. C 23, 351–364.
- 819 Pandža, M., Franjić, J., Škvorc, Ž., 2007. The salt marsh vegetation on the East Adriatic coast.
- 820 Biologia 62, 24–31. https://doi.org/10.2478/s11756-007-0003-x
- 821 Parnell, A.C., Haslett, J., Allen, J.R.M., Buck, C.E., Huntley, B., 2008. A flexible approach to
- 822 assessing synchroneity of past events using Bayesian reconstructions of sedimentation history.
- 823 Quaternary Science Reviews 27, 1872–1885. https://doi.org/10.1016/j.quascirev.2008.07.009
- 824 Peltier, W.R., Argus, D.F., Drummond, R., 2015. Space geodesy constrains ice age terminal
- deglaciation: The global ICE-6G_C (VM5a) model. J. Geophys. Res. Solid Earth 120,
 2014JB011176. https://doi.org/10.1002/2014JB011176
- Pirazzoli, P., 1996. Sea Level Changes. The Last 20 000 Years. John Wiley & Sons Ltd,
 Chichester, New York.
- 829 Pirazzoli, P.A., 2005. A review of possible eustatic, isostatic and tectonic contributions in eight
- 830 Late-Holocene relative sea-level histories from the Mediterranean area. Quaternary Science
- 831 Reviews, Quaternary coastal morphology and sea-level changes 24, 1989–2001.
- 832 https://doi.org/10.1016/j.quascirev.2004.06.026
- 833 Pirazzoli, P.A., 1991. World Atlas of Holocene Sea-Level Changes, 1st ed. Elsevier.
- 834 Pirazzoli, P.A., 1976. Sea Level Variations in the Northwest Mediterranean During Roman Times.
- 835 Science 194, 519–521. https://doi.org/10.1126/science.194.4264.519

- 836 Plater, A.J., Kirby, J.R., Boyle, J.F., Shaw, T., Mills, H., 2015. Loss on ignition and organic
- 837 content, in: Shennan, I., Long, A.J., Horton, B.P. (Eds.), Handbook of Sea-Level Research. John
- 838 Wiley & Sons, Ltd, pp. 312–330.
- 839 Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Bronk Ramsey, C., Buck, C.E.,
- 840 Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Haflidason, H., Hajdas,
- 841 I., Hatté, C., Heaton, T.J., Hoffmann, D.L., Hogg, A.G., Hughen, K.A., Kaiser, K.F., Kromer, B.,
- 842 Manning, S.W., Niu, M., Reimer, R.W., Richards, D.A., Scott, E.M., Southon, J.R., Staff, R.A.,
- 843 Turney, C.S.M., van der Plicht, J., 2013. IntCal13 and Marine13 Radiocarbon Age Calibration
- 844 Curves 0–50,000 Years cal BP. Radiocarbon; Vol 55, No 4 (2013).
- 845 Rosenthal, Y., Kalansky, J., Morley, A., Linsley, B., 2017. A paleo-perspective on ocean heat
- 846 content: Lessons from the Holocene and Common Era. Quaternary Science Reviews 155, 1–12.
- 847 https://doi.org/10.1016/j.quascirev.2016.10.017
- 848 Roy, K., Peltier, W.R., 2017. Space-geodetic and water level gauge constraints on continental
- 849 uplift and tilting over North America: regional convergence of the ICE-6G C (VM5a/VM6)
- 850 models. Geophys J Int 210, 1115–1142. https://doi.org/10.1093/gji/ggx156
- 851 Roy, K., Peltier, W.R., 2018. Relative sea level in the Western Mediterranean basin: A regional
- test of the ICE-7G NA (VM7) model and a constraint on late Holocene Antarctic deglaciation.
- 853 Quaternary Science Reviews 183, 76–87. https://doi.org/10.1016/j.quascirev.2017.12.021
- 854 Saher, M.H., Gehrels, W.R., Barlow, N.L.M., Long, A.J., Haigh, I.D., Blaauw, M., 2015. Sea-
- 855 level changes in Iceland and the influence of the North Atlantic Oscillation during the last half
- 856 millennium. Quaternary Science Reviews 108, 23–36.
- 857 https://doi.org/10.1016/j.quascirev.2014.11.005

- 858 Scott, D.B., Hermelin, J.O.R., 1993. A Device for Precision Splitting of Micropaleontological
- 859 Samples in Liquid Suspension. Journal of Paleontology 67, 151–154.
 860 https://doi.org/10.2307/1305976
- 861 Scott, D.B., Medioli, F.S., 1980. Quantitative studies of marsh foraminiferal distributions in nova-
- 862 scotia canada implications for sea level studies. Cushman Foundation for Foraminiferal Research
- 863 Special Publication 1–58.
- Scott, D.B., Medioli, F.S., 1978. Vertical zonations of marsh foraminifera as accurate indicators
 of former sea-levels. Nature 272, 528–531. https://doi.org/10.1038/272528a0
- Serpelloni, E., Faccenna, C., Spada, G., Dong, D., Williams, S.D.P., 2013. Vertical GPS ground
 motion rates in the Euro-Mediterranean region: New evidence of velocity gradients at different
 spatial scales along the Nubia-Eurasia plate boundary. J. Geophys. Res. Solid Earth 118,
- 869 2013JB010102. https://doi.org/10.1002/2013JB010102
- 870 Shaw, T.A., Kirby, J.R., Holgate, S., Tutman, P., Plater, A.J., 2016. Contemporary salt-marsh
- 871 foraminiferal distribution from the Adriatic coast of Croatia and its potential for sea-level studies.
- Journal of Foraminiferal Research 46, 314–332. https://doi.org/10.2113/gsjfr.46.3.314
- 873 Sivan, D., Lambeck, K., Toueg, R., Raban, A., Porath, Y., Shirman, B., 2004. Ancient coastal
- 874 wells of Caesarea Maritima, Israel, an indicator for relative sea level changes during the last 2000
- 875 years. Earth and Planetary Science Letters 222, 315–330.
 876 https://doi.org/10.1016/j.epsl.2004.02.007
- 877 Spada, G., Stocchi, P., Colleoni, F., 2009. Glacio–isostatic Adjustment in the Po Plain and in the
- 878 Northern Adriatic Region. Pure and Applied Geophysics 166, 1303–1318.
- 879 https://doi.org/10.1007/s00024-004-0498-9

- 880 Šparica, M., Bačani, A., Koch, G., Anda, A., Miko, S., Viličić, D., Galovć, I., Šparica, M., M.,
- Ibrahimpšarić, H., Bargant, S., Dolenec, T., 2005. Ecosystem of Morinje Bay (Adriatic Sea,
 Croatia): Aspects of the sediment/water interface. RMZ Materials and Geoenvironment 52, 115–
 118.
- 884 Stocchi, P., Spada, G., 2007. Glacio and hydro-isostasy in the Mediterranean Sea: Clark's zones
- and role of remote ice sheets. Annals of Geophysics 50, 741–761.
- Stocchi, P., Spada, G., 2009. Influence of glacial isostatic adjustment upon current sea level
 variations in the Mediterranean. Tectonophysics 474, 56–68.
 https://doi.org/10.1016/j.tecto.2009.01.003
- 889 Strachan, K.L., Finch, J.M., Hill, T.R., Barnett, R.L., 2014. A late Holocene sea-level curve for
- the east coast of South Africa. South African Journal of Science 110, 9 Pages.
 https://doi.org/10.1590/ sajs.2014/20130198
- Surić, M., Korbar, T., Juračić, M., 2014. Tectonic constraints on the late Pleistocene-Holocene
 relative sea-level change along the north-eastern Adriatic coast (Croatia). Geomorphology 220,
- 894 93–103. https://doi.org/10.1016/j.geomorph.2014.06.001
- 895 Toker, E., Sivan, D., Stern, E., Shirman, B., Tsimplis, M., Spada, G., 2012. Evidence for centennial
- scale sea level variability during the Medieval Climate Optimum (Crusader Period) in Israel,
- 897 eastern Mediterranean. Earth and Planetary Science Letters, Sea Level and Ice Sheet Evolution: A
- 898 PALSEA Special Edition 315–316, 51–61. https://doi.org/10.1016/j.epsl.2011.07.019
- 899 Törnqvist, T.E., Rosenheim, B.E., Hu, P., Fernandez, A.B., 2015. Radiocarbon dating and
- 900 calibration, in: Shennan, I., Long, A.J., Horton, B.P. (Eds.), Handbook of Sea-Level Research.
- 901 John Wiley & Sons, Ltd, pp. 347–360.

- 902 Törnqvist, T.E., Wallace, D.J., Storms, J.E.A., Wallinga, J., van Dam, R.L., Blaauw, M., Derksen,
- 903 M.S., Klerks, C.J.W., Meijneken, C., Snijders, E.M.A., 2008. Mississippi Delta subsidence
- 904 primarily caused by compaction of Holocene strata. Nature Geosci 1, 173–176.
- 905 https://doi.org/10.1038/ngeo129
- 906 Troëls-Smith, J., 1955. Characterization of unconsolidated sediments. Danmarks Geologiske
 907 Undersøgelse Series 4, 1–73.
- Tsimplis, M.N., Baker, T.F., 2000. Sea level drop in the Mediterranean Sea: An indicator of deep
 water salinity and temperature changes? Geophysical Research Letters 27, 1731–1734.
- 910 https://doi.org/10.1029/1999gl007004
- 911 Tsimplis, M.N., Josey, S.A., 2001. Forcing of the Mediterranean Sea by atmospheric oscillations
 912 over the North Atlantic. Geophysical Research Letters 28, 803–806.
- 913 https://doi.org/10.1029/2000gl012098
- 914 Tsimplis, M.N., Raicich, F., Fenoglio-Marc, L., Shaw, A.G.P., Marcos, M., Somot, S.,
- 915 Bergamasco, A., 2012. Recent developments in understanding sea level rise at the Adriatic coasts.
- 916 Physics and Chemistry of the Earth, Parts A/B/C 40-41, 59-71.
 917 https://doi.org/10.1016/j.pce.2009.11.007
- Vacchi, M., Marriner, N., Morhange, C., Spada, G., Fontana, A., Rovere, A., 2016. Multiproxy
 assessment of Holocene relative sea-level changes in the western Mediterranean: Sea-level
 variability and improvements in the definition of the isostatic signal. Earth-Science Reviews 155,
- 921 172–197. https://doi.org/10.1016/j.earscirev.2016.02.002
- 922 Van De Plassche, O., Wright, A.J., Horton, B.P., Engelhart, S.E., Kemp, A.C., Mallinson, D.,
- 923 Kopp, R.E., 2014. Estimating tectonic uplift of the Cape Fear Arch (south-eastern United States)

- 924 using reconstructions of Holocene relative sea level. J. Quaternary Sci. 29, 749–759.
 925 https://doi.org/10.1002/jqs.2746
- 926 Vilibić, I., 2006. The role of the fundamental seiche in the Adriatic coastal floods. Continental
- 927 Shelf Research 26, 206–216. https://doi.org/10.1016/j.csr.2005.11.001
- 928 Vilibić, I., Orlić, M., Čupić, S., Domijan, N., Leder, N., Mihanović, H., Pasarić, M., Pasarić, Z.,
- 929 Srdelić, M., Strinić, G., 2005. A new approach to sea level observations in Croatia. Geofizika 22,
 930 21–57.
- 931 Vilibić, I., Šepić, J., 2009. Destructive meteotsunamis along the eastern Adriatic coast: Overview.
- 932 Physics and Chemistry of the Earth, Parts A/B/C 34, 904–917.
 933 https://doi.org/10.1016/j.pce.2009.08.004
- 934 Vilibić, I., Šepić, J., Pasarić, M., Orlić, M., 2017. The Adriatic Sea: A Long-Standing Laboratory
- 935 for Sea Level Studies. Pure Appl. Geophys. 174, 3765–3811. https://doi.org/10.1007/s00024-017936 1625-8
- 937 Weber, J., Vrabec, M., Pavlovcic-Preseren, P., Dixon, T., Jiang, Y., Stopar, B., 2010. GPS-derived
- 938 motion of the Adriatic microplate from Istria Peninsula and Po Plain sites, and geodynamic
- 939 implications. Tectonophysics 483, 214–222. https://doi.org/10.1016/j.tecto.2009.09.001
- Williams, C.K.I., Rasmussen, C.E., 1996. Gaussian Processes for Regression. MIT Press,Cambridge.
- Woodworth, P.L., 2003. Some Comments on the Long Sea Level Records from the Northern
 Mediterranean. Journal of Coastal Research 19, 212–217.
- 244 Zerbini, S., Plag, H.-P., Baker, T., Becker, M., Billiris, H., Bürki, B., Kahle, H.-G., Marson, I.,
- 945 Pezzoli, L., Richter, B., Romagnoli, C., Sztobryn, M., Tomasi, P., Tsimplis, M., Veis, G., Verrone,
- 946 G., 1996. Sea level in the Mediterranean: a first step towards separating crustal movements and

- 947 absolute sea-level variations. Global and Planetary Change 14, 1–48.
 948 https://doi.org/10.1016/0921-8181(96)00003-3
- 949 Zerbini, S., Raicich, F., Prati, C.M., Bruni, S., Del Conte, S., Errico, M., Santi, E., 2017. Sea-level
- 950 change in the Northern Mediterranean Sea from long-period tide gauge time series. Earth-Science
- 951 Reviews 167, 72–87. https://doi.org/10.1016/j.earscirev.2017.02.009