



LJMU Research Online

Silva, AC, Silva, A, Edwards, B, Tod, D, de Souza Amarala, A, de Alcântara Borba, D, Grade, I and de Mello, MT

Sleep extension in athletes: What we know so far – a systematic review

<http://researchonline.ljmu.ac.uk/id/eprint/14111/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Silva, AC, Silva, A, Edwards, B, Tod, D, de Souza Amarala, A, de Alcântara Borba, D, Grade, I and de Mello, MT (2020) Sleep extension in athletes: What we know so far – a systematic review. *Sleep Medicine*, 77. pp. 128-135. ISSN 1389-9457

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

1 **Sleep extension in athletes: What we know so far – a systematic review**

2
3
4 Aldo Coelho Silva^a, Andressa Silva^a; Ben J Edwards^b, David Tod^b, Adriana de Souza Amaral^a,
5 Diego de Alcântara Borba^a, Isadora Grade^a, Marco Túlio de Mello^a

6
7 ^aUniversidade Federal de Minas Gerais, Escola de Educação Física, Fisioterapia e Terapia
8 Ocupacional, Departamento de Esportes, Belo Horizonte/Brazil

9 ^bResearch Institute for Sport and Exercise Sciences, Liverpool John Moores University,
10 Liverpool, UK

11
12 ***Corresponding author:** Marco Túlio de Mello, PhD, Escola de Educação Física,
13 Fisioterapia e Terapia Ocupacional, Universidade Federal de Minas Gerais, Av. Antônio Carlos, 6627,
14 Pampulha, CEP: 31270-901, Belo Horizonte, MG, Brasil. Telephone: +55 (31) 3409-2347, E-mail:
15 tmello@demello.net.br

16
17
18
19 **Declarations of interest:** none

20
21 Word count:

22 Main file:3997

23 Abstract: 248

24 Number of tables: 2

25 Number of supplementary material: 1

29 **Abstract**

30 Objective: This study reviewed systematically the effects of sleep extension on sports
31 performance.

32 Design: systematic review

33 Method: The systematic review was conducted in November 2020. Articles published in
34 English were searched in PubMed, Virtual Health Library, SPORTDiscus, and Web of Science and
35 Scopus databases. The search terms used were “sleep extension” AND athlete. The measures of
36 interest were sports performance. Studies were included if they were a) original articles, b) published
37 in English and peer-reviewed article, c) had only athletes as participants, d) experimental protocol
38 whose objective was to investigate the effects of sleep extension on sports performance, including
39 randomized (RCT) and non-randomized controlled trial (nRCT), and e) at least a sports performance
40 measure as a dependent variable.

41 Results: The primary search revealed that a total of 5 out of 74 articles were considered
42 eligible and 2 studies were subsequently included. The studies used different strategies to extend time
43 in bed or total sleep time (extending 26 to 106 min). From fifteen sports measures, six presented a
44 large effect size, and the others ranged from trivial to medium. Overall, the risk of bias was high to
45 RCT and low to nRCT and the quality of evidence ranged from very low quality to moderate quality in
46 ten outcomes.

47 Conclusions: The limited evidence suggests that sleep extension interventions may be
48 beneficial to improve sports performance in athletes where the magnitude is dependent on the variable
49 assessed, although such conclusions are tentative because of the quality of the evidence and risk of
50 bias.

51 Keywords: athlete, extra sleep, performance, and sport.

52

53 1. Introduction

54 Sleep is particularly important to athletes as sleep helps in the body restoration imposed by the
55 fatigue of the waking period, with a restorative and repairing process in the energy of different
56 physiological systems. Consequently, the athlete's body becomes prepared for both training and
57 competition. In addition, sleep provides cognitive recovery and optimal decision-making capacity,
58 contributing to an optimal mental state.¹ For an athlete to achieve the expected training results, recover
59 for the next training session, and increase performance, sleep must be restorative, accomplished by
60 getting adequate sleep per night.² The National Sleep Foundation (NSF) recommends different sleep
61 durations according to the age group. For most healthy adults (18 – 64 years old), the NSF
62 recommends between 7 and 9 h of sleep per night. Although some individuals might sleep longer or
63 shorter than the recommended times without deleterious effects, getting sleep duration continually
64 outside the normal range may harm his or her health and well-being. It is important to note that this
65 recommendation referring to sleep duration instead of time in bed. Typically, actual sleep duration is
66 lower than time in bed, which makes it important to know which term is related to in the studies with
67 athletes.³ Although there is no consensus on the ideal amount of sleep per night for athletes, especially
68 during a competitive period,⁴ higher amount of sleep per night than that recommended for non-athletes
69 has been suggested.^{5,6}

70 Unlike sleep loss, sleep extension has ambiguous effects on sports performance. In one of the
71 first studies to observe the effect of sleep extension on sports performance, Mah, Mah ⁷ extended the
72 sleeping time of swimmers (from 6–8 h per night to 10 h per night) for 6–7 weeks. The authors
73 observed an improvement in sprint time, reaction speed, and mood. The same group of researchers
74 conducted a sleep extension program for college basketball players. They found that the players could
75 achieve a minimum of 10 h of sleep for 5–7 weeks. In addition to improved sprinting, the authors
76 observed an improvement in accuracy in free-throw or 3-point throws.⁸ Likewise, Schwartz and Simon
77 Jr ⁹ observed an improvement in the accuracy of service among college tennis players following sleep
78 extension for 2 weeks. On the contrary, Fullagar, Skorski ¹⁰ did not observe the effect of sleep
79 extension on physical performance (countermovement jump and intermittent yoyo recovery test),
80 blood component measurements (creatinine kinase, urea, and C-reactive protein), stress markers, or

81 perceived recovery. These contradictory results can be attributed, at least in part, to the different
82 research methods and designs used.

83 Thus, interestingly, there are methodological issues that limit the application of the previous
84 findings, and consequently, development of a standard for recommending sleep extension in athletes.
85 For example, were the athletes with accumulated sleep debt or chronic sleep restriction? Would
86 extending sleep only benefit individuals with sleep debt or sleep restriction? The length of sleep may
87 have been an opportunity to achieve enough sleep, with no extension in the ideal amount of sleep. The
88 literature reveals that individuals with sleep debt or reduced amount of sleep, benefit when provided
89 with the opportunity to achieve adequate sleep. Another factor to be considered is the minimum
90 intervention period necessary to experience the benefits of sleep extension. Moreover, there is a need
91 to verify the effect of sleep extension programs on athletes and on what measures these interventions
92 can be beneficial. This could contribute to the standardization of the sleep extension intervention
93 recommendation for athletes.

94 To date, there are no systematic reviews on the use of sleep extension in athletes. Recently,
95 sleep extension intervention has been recommended as beneficial for subsequent performance
96 measures.¹¹ This article provides information from two articles that used intervention to extend sleep
97 and its results in sports performance. However, it does not provide critical information about the
98 quality of the studies that were used, the effect size of intervention, neither their risks of bias. In a
99 recent narrative review on athlete's sleep,¹² the authors described sleep extension as a strategy that
100 should be looked at carefully, with a brief introduction to the analyzes that should be done in sleep
101 extension studies. Therefore, this study aimed to conduct a systematic review of the literature, perform
102 a critical analysis of the sleep extension intervention designs used in athletes, and present the effects of
103 sleep extension on sports performance of athletes when compared to habitual sleep patterns.

104

105 **2. Methods**

106 Two researchers screened the relevant published articles from PubMed, Web of Science,
107 Scopus, SPORTDiscus, and Virtual Health Library. Moreover, we have researched on grey literature
108 (OpenGrey, New York Academy of Medicine Grey Literature Report, ClinicalTrials, EThOS: UK E-

109 Theses Online Service) in February 2020 were searched and updated in November 2020. The search
110 terms used were: “sleep extension” AND athlete and their Medical Subject Heading terms. Our review
111 was performed in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-
112 Analyses (PRISMA).¹³ The search strategy and PRISMA checklist are available as supplementary
113 material. Studies were eligible if they investigated the effect of sleep extension on athletes’ sports
114 performance. Sports performance was considered any activity whose outcome of interest required
115 physical performance (e.g., strength, endurance, and speed), or specific sports skills (e.g., tennis serve
116 and 3-point throw in basketball) in athletes. Following were the inclusion criteria: a) original article; b)
117 published in English and peer-reviewed article; c) only athletes as participants, regardless of the
118 modality practiced or age; d) experimental protocol (randomized and non-randomized controlled
119 trials) whose objective was to investigate the effects of sleep extension on sports performance, without
120 limit for the intervention time; and e) existence of a measure of sports performance as a dependent
121 variable. For this review, sleep extension was defined as an increase in habitual total sleep time, either
122 in nighttime sleep or addition of naps during the day. Studies were excluded if they did not meet at
123 least one inclusion criterion. Reviews, systematic or critical, short-communication, and editorial
124 articles were excluded. The screening and data extraction were performed by two researchers.
125 Disagreements in any article or result were discussed among the authors and resolved by additional
126 authors when necessary. We used the *k* statistic to describe the level of agreement between the
127 reviewers in this phase.¹⁴ After removing duplicate articles, the titles and abstracts were read by an
128 author. The two authors then examined the articles in full to confirm the inclusion and eligibility
129 criteria.

130 A total of 74 articles were found to be eligible. Of these articles, 16 were extracted from
131 PubMed, 17 from Web of Science, 15 from Scopus, 13 from SPORTDiscus, 11 from Virtual Health
132 Library and 2 from grey literature. After excluding the duplicate articles, 32 articles remained. After
133 reading the titles and abstracts, 17 articles were excluded for not meeting the eligibility criteria (e.g.,
134 reviews and editorials), leaving 15 articles to be examined. A second reviewer also read the titles,
135 abstracts and the full text. In case of discrepancy in any study, a third reviewer was invited to arrive at
136 a final decision. After full reading of the selected articles, 10 were excluded such that articles that did

137 not include athletes and articles whose outcome measure was not sports performance. Two articles
138 were added from the reference lists of included studies. Therefore, a total of seven articles were
139 included for the final review (Figure 1). Unpublished articles were not included for the analysis. There
140 were good agreements between the reviewers during the screening after excluding duplicates ($k= 0.86$,
141 $95\% \text{ CI}[0.66,1.05]$; $p= 0.00$; agreement percentage= 97.2%) and during the screening of included
142 articles in this review ($k= 1.0$, $95\% \text{ CI}[1.0,1.0]$; $p= 0.00$; agreement percentage= 100%).

143 Information regarding the population, intervention, comparisons, outcomes, study design
144 (PICOS), was extracted from each study. This Data concerned characteristics of the participants (age,
145 sex, level of athlete, and habitual sleep pattern), type of intervention (number of hours of sleep
146 prescribed, duration of intervention), and outcome measures (sports performance). Sports performance
147 measures were recorded using numerical information from the results of each study. The Cochrane
148 Collaboration tools assessed the risk of bias in randomized trials (RoB 2.0) and non-randomized trials
149 (ROBINS-I). The RoB 2.0¹⁵ assesses the risk of bias in five distinct domains and the judgments
150 within each domain lead to overall risk-of-bias. The ROBINS-I¹⁶ assesses the risk of bias in seven
151 domains and shows an overall risk-of-bias. The items of this evaluation were classified as low,
152 moderate, or high risk of bias. The plots obtained from these analyses¹⁷ are available as
153 supplementary material. All studies with matching eligibility were included in the review regardless of
154 their risk of bias or quality. We used the k statistic to describe the level of agreement between the
155 reviewers. The Grades of Recommendations, Assessment, Development, and Evaluation (GRADE)
156 rated the overall quality of evidence for each outcome. This tool evaluates the level of confidence for
157 each outcome and provides an overall summary of quality with 1 of 4 classifications: high, moderate,
158 low, and very low.

159 Due to the large variation in intervention programs and the different measurement tools, it was
160 not possible to perform a meta-analysis of the data. Hence, synthesis without meta-analysis guidelines
161 was used to report the results.¹⁸ The synthesis without meta-analysis guidelines (SWiM) checklist is
162 available as supplementary material. The mean and standard deviation of the outcomes were collected
163 from the three studies and were used to calculate the effect size and confidence interval of the

164 intervention. The effect size was determined by Cohen's d ($d = \frac{M2 - M1}{\sqrt{(SD2^2 - SD1^2)/2}}$), with $d > 0.2$ and $d <$
165 0.5 considered as small effect; $d > 0.5$ and $d < 0.8$ as moderate effect; and $d > 0.8$ as large effect¹⁹.
166 The following formula was used to calculate the percentage difference: $(X2 - X1)/X1 \times 100$, where $X1$
167 is pre-intervention, and $X2$ is post-intervention.

168 **3. Results**

169 The characteristics of the included studies are presented in Table 1. All the included articles
170 were interventional studies published between 2011 and 2020 (randomized and non-randomized
171 controlled trial). The study sample comprised 9–24 athletes aged 14–30 years playing eight different
172 sports (tennis, basketball, cycling, handball, rugby, shooting, soccer and triathlon). Of the seven
173 studies analyzed, five included only male athletes^{8, 10, 20} and two included both male and female
174 athletes.^{9, 21} Two studies included university athletes,^{8, 9} one included high-level student-athletes,²¹ one
175 included trained athletes,²⁰ one included highly trained athletes,²² one included amateurs athletes,²³ and
176 one included highly trained amateurs.¹⁰

177 To describe the sleep-wake cycle, five studies used actigraphy either in the pre-intervention
178 period or during the intervention^{8, 10, 20, 21, 23} and one used the polysomnography.²² Two studies used
179 actigraphy after the end of the intervention period, describing the sleep-wake cycle up to two days. Of
180 the studies that used actigraphy to describe the sleep-wake cycle, two studies described the activity
181 thresholds, considering the value of 40 and 60 for movement detection.^{20, 23} This pre-interventional
182 evaluation for sample characterization lasted from 1 day to 4 weeks. Roberts, Teo²⁰ observed the
183 amount of sleep for 4 days before the intervention period, and there were other periods like 1 day,^{10, 22,}
184 ²³ 2 days,²¹ 1 week,⁹ and 2-4 weeks.⁸ The total pre-interventional sleep time ranged from 06:54 to
185 08:45 h:min per night.

186 The duration of the sleep extension intervention in the studies ranged from 1 day to 7 weeks.
187 The strategies for sleep extension intervention included a) extension in the habitual time in bed, b)
188 provision of behavioral advice with sleep hygiene tips, and (c) a combination of both. There different
189 recommendations like getting 9 to 10 h of sleep per night, an increase 30 % in habitual time in bed,
190 and take 20, 40, or 90-min of nap. All studies reported an increase in the sleep parameter assessed with

191 the intervention. The smallest difference observed between the control condition and the experimental
192 condition was 00:26 h:min and the biggest difference observed was 01:48 h:min.

193 Performance measures varied between studies, from specific skills of the sports modality to
194 measures of physical performance, totaling fifteen performance measures analyzed. The percentage
195 difference in the measures evaluated in sleep extension intervention studies is provided in Table 2.
196 Compared to baseline, Mah, Mah⁸ reported improvements in free throw (11.4 %, $d = 0.918$), 3-point
197 field goals (13.7 %, $d = 0.757$), and 282 feet sprint (-4.3 %, $d = 1.215$). Similarly, Schwartz and
198 Simon Jr⁹ reported improvements in tennis serving accuracy (17.7 %, $d = 0.418$) after sleep extension
199 intervention compared to baseline. Roberts, Teo²⁰ reported significant decreases in time-trial
200 performance after the sleep extension intervention on day 4 compared to normal sleep condition (-3.2
201 %, $d = 0.583$). Boukhris, Trabelsi²³ reported improvements in maximal voluntary isometric
202 contraction and shuttle run test ranging from +5.7 to 12.6 % ($d = 0.44$ to 1.70). Two studies
203 demonstrated no improvement in performance measures^{10, 21, 22} despite the extension in total sleep
204 time, and one study described an association between hours of sleep and sports performance.²¹
205 Fullagar, Skorski¹⁰ reported percentage changes from -4.6 to +2.8 % and Cohen'd from 0.10 to 0.25,
206 and Petit, Mougin²² reported percentage changes of +0.6 and 0.9 % ($d = 0.050$ and 0.049,
207 respectively). Lastly, Suppiah, Low²¹ did not report the values of central tendency and dispersion of
208 sports performance variables in their results.

209 The three randomized controlled trial showed a high risk of bias. Of the four non-randomized
210 studies, two were classified as being low risk of bias, one as being the moderate risk of bias, and one
211 as a serious risk of bias. All non-randomized studies showed bias due to the selection of the
212 participants. Overall RoB for non-randomized controlled trials was graded as 50 % showing a low risk
213 of bias, 25 % showing a moderate risk of bias, and 25 % showing a serious risk of bias. The results of
214 GRADE rating showed that one outcome provided evidence with very low quality (1/15, 6 %), five
215 outcomes provided evidence with low quality (5/15, 33 %), four outcomes provided evidence with
216 moderate quality (4/15, 27 %), and five outcomes provided evidence with high quality (5/15, 33 %).
217 The risk of bias was found in all included articles. The RoB assessments and the quality of evidence
218 are available as supplementary material.

219

220 **4. Discussion**

221 This study aimed to conduct a systematic review to present the effects of sleep extension on
222 sports performance of athletes (9-10 h over 1-49 days who habitually sleep 6-9 h a day). To the best of
223 our knowledge, this is the first study to review the effect of sleep extension in athletes. We present the
224 initial results of a new strategy to improve sports performance and indicate that extending sleep in
225 athletes without sleep disorders or sleep debt may be beneficial, although the quality of the research
226 prevents firm conclusions. To apply a sleep extension intervention, it is important to consider two
227 factors: sleep characteristics of the athletes and the sleep extension intervention.

228 The most important factor is the individual amount of sleep habitually taken by the athletes
229 participating in the studies. Some factors must be considered while identifying the athlete's normal
230 sleep pattern, such as the analysis period (number of observation days) and the amount of pre-
231 intervention sleep. Concerning the analysis period, the wide range of the evaluation period (from 1 day
232 to 4 weeks) observed in this review can cause diversity in the athlete's sleep pattern submitted to
233 extension intervention. The Society of Behavioral Sleep Medicine recommends the use of actigraphy
234 for a period of 7 to 14 days, including at least one weekend to estimate sleep-wake pattern.²⁴ In
235 addition, the description of the activity count used to define the sleep-wake cycle should be
236 considered, as there is a recommendation to use a higher threshold in studies involving athletes.²⁵

237 The amount of sleep can vary between nights. A previous study reported that nightly
238 variability was greater than yearly variability. Thus, the analysis of only 1 or 2 days of sleep may not
239 represent the individual's sleep characteristics.²⁶ For a reliable measure of total sleep time, the
240 minimum recommended period is 7 days.²⁷ Interestingly, only 2 studies observed an amount of sleep
241 for least 7 days. Thus, other studies may fail to estimate the volunteers' habitual sleep time. Another
242 important point is the amount of sleep in the pre-intervention moment. To show that athletes did not
243 return from a period of sleep loss, three studies presented the amount of sleep referring to 2-3 days
244 before the beginning of the study.^{10, 20, 21} Although two days are sufficient to restore the daytime
245 sleepiness, fatigue, cortisol level, and IL-6 level, after a period of sleep loss, evidence shows that
246 cognitive performance may need a longer period to restore pre-debt sleep values.^{28, 29} Consequently,

247 athletes may have started the study period with a cognitive pattern different from a normal period of
248 sleep, compromising the results of the research.

249 Thus, standardization in the observation of the total sleep time before the intervention can help
250 describe whether the athlete has sleep debt or restriction or a period with an ideal amount of sleep.
251 Long-term sleep restriction can have deleterious cognitive and performance effects.³⁰ After a period of
252 chronic restraint or sleep deprivation, as long as there is a free amount of sleep, there is a rebound
253 effect in the period of sleep recovery (period after the end of the intervention to reduce the amount of
254 sleep) with an increase in the total sleep time. Thus, if the athlete has sleep debt, the sleep extension
255 intervention will only offer the ideal amount of sleep for recovery.

256 Knowledge of the athlete's sleep pattern in terms of quantity helps estimate whether or not the
257 athlete is a short, intermediate, or long sleeper. Although most of the population get a satisfactory
258 amount of 7.5 h of sleep a night, there are individuals with a lower ideal amount of sleep.³¹ Short
259 sleepers can achieve satisfactory sleep with only 3 h of sleep per night. Long sleepers may require >10
260 h of sleep per night.³² Thus, sleepers are categorized into three classes: short sleepers, with <6 h of
261 sleep per night; intermediate sleepers, with 7–8 h of sleep per night, and long sleepers, with >9 h of
262 sleep per night.^{31, 33}

263 An important aspect is that the ability to extend the amount of sleep beyond normal appears to
264 be a psychophysiological feature.³⁴ Normal sleepers sleep for 7–9 h per night and can extend the
265 amount of sleep to 10–11 h even without a previous period of sleep deprivation or restriction.
266 Individuals with this ability are called sleep extenders.^{35, 36} On the other hand, 5 consecutive nights
267 with an approximately 25 % reduction in total sleep time affects short and long sleepers than
268 intermediate sleepers.³⁷ In general, sleep extension intervention studies in athletes use an index on
269 PSQI less than 5 (good sleeper), and a sleep amount greater than 6 and less than 9 hours as the
270 inclusion criteria for participants. Thus, there is a margin of approximately 2 h in the usual sleep of
271 these athletes, which can affect the results of the studies. There are reports that individuals were able
272 to achieve 10 h of sleep per night after an acute recommendation for the length of sleep, although this
273 ability may be characteristic of some individuals.³⁶ On comparing the ability of single twins to extend

274 the sleep, Gagnon, De Koninck ³⁸ reported that 8 out of 10 normal sleepers were able to extend sleep
275 between 12 and 15 h a night.

276 A critical question to describe whether the subject is short or long sleeper is to define the sleep
277 pattern and this question has been addressed for several decades.³² As noted earlier, there is an intra-
278 individual variation in sleep nights. besides the variation that occurs between the weekdays, there is a
279 difference of approximately 30 minutes in sleep time on weekdays compared to weekends.³² Several
280 kinds of researches have been used the subjective description of the sleep time duration and then, the
281 volunteer is observed for a certain time to describe the sleep pattern. This period for describing the
282 sleep pattern can vary from two weeks to one year of follow-up.^{31, 39} Thus, research shows that
283 volunteers have a sleep pattern and classifies them according to the typology.

284 Only Mah, Mah ⁸ and Schwartz and Simon Jr ⁹ monitored the athletes' sleep at least 7 days
285 before the intervention period. The other studies did not monitor sleep for long enough to describe the
286 sleep patterns. Thus, it is not certain whether these articles used athletes coming from a chronic period
287 of sleep restriction or whether the athletes were short or long sleepers, or that they would impact the
288 sleep extension intervention; another important fact is that most studies did not describe the usual
289 schedule for training sessions for athletes. Training sessions started before 07:00 h are known to
290 negatively affect athletes' sleep. Thus, the authors may have monitored and described the opportunity
291 that athletes have to sleep more than the amount of sleep that athletes usually need.

292 We emphasize that few studies considered the chronotype of the athletes and the time of day
293 to start the activities. Chronotype is an individual characteristic that determines the propensity to be
294 more alert and more active at a certain time of the day, establishing a preference to be more active or
295 to sleep at a certain time. ⁴⁰ In the studies used in this review, only 3 articles analyzed the chronotype.
296 ^{10, 20, 22} Of these 3 studies, only Roberts, Teo ²⁰ considered the chronotype for the prescription of the
297 sleep extension program and the usual training schedule for performing the performance analysis.
298 Individuals who practice sports for their chronotype (example: evening type training at night), train
299 and compete better than at times opposite to their chronotype ⁴¹. Regarding the time of day, five
300 studies applied the tests between 12:00-18:00 h, one study applied the tests between 06:00-09:00 h,
301 and one study applied tests at 10:00 and 16:00 h. In a recent systematic review, Vitale and Weydahl ⁴²

302 observed that the chronotype influences the perceived exertion and fatigue scores, with morning types
303 feeling less fatigued in the early hours of the day than intermediate types and night types. Thus, it is
304 essential that future studies regarding sleep extension describe the athlete's chronotype, establishing
305 relationships between the chronotype and the time when the performance test is performed.

306 The magnitude of the sleep extension effects varies between the studies. The current review
307 shows that using sleep extension programs leads to trivial to larger effects on sports performance.
308 Regarding the percentage difference, Mah, Mah⁸ observed an 11 % increase in pitch, and Schwartz
309 and Simon Jr⁹ observed an increase of approximately 20 % in the accuracy of service, with moderate
310 and small effects, respectively. Roberts, Teo²⁰ have observed a better aerobic performance (~3 %) on
311 day 3 in the sleep extension group, compared to the normal group. It is important to note that even
312 small effects can be decisive in sports performance. Boukhris, Trabelsi²³ described improvement
313 ranging from +5.6 to +9.6 % in maximal voluntary isometric contraction and improvements ranging
314 +7.8 to 12.6 % in shuttle run test with large effect size. On the contrary, Petit, Mougin²² reported no
315 improvement in Wingate test, Suppiah, Low²¹ did not observe an increase in shooting performance,
316 after a period of unrestricted sleep, with at least 9 h of sleep, and Fullagar, Skorski¹⁰ describe that
317 there was no change in performance in the countermovement jump or in the yoyo test after an acute
318 sleep hygiene protocol that increased the total sleep time.

319 However, the results favorable to the extension of sleep in these studies should be viewed with
320 caution. As mentioned earlier, Boukhris, Trabelsi²³ assessed the total sleep time for only one night
321 and Roberts, Teo²⁰ assessed the total sleep time just 4 days before starting the intervention. This
322 period is not enough to describe the habitual sleep and the 2 days of free sleep before starting the study
323 may be insufficient to restore cognitive performance. Mah, Mah⁸ and Schwartz and Simon Jr⁹ studies
324 are non-randomized quasi-experimental trials. In both studies, the subjects served as their control, a
325 pre-post intervention design. Thus, the positive results of the intervention period may be due to the
326 previous trial, also known as serial order carryover effects.⁴³ That is, the improvement in performance
327 may be due to a dependence on previous testing.⁴⁴ This dependence may be because of learning from
328 the test performed or training adjustments. One way to try to mitigate the effect of dependence on the

329 previous testing would be to try to counterbalance the testing order of different groups. Another fact to
330 highlight is the lack of information regarding sports training developed during the study. Thus, the
331 increase in performance may have been due to a peak period of performance previously established by
332 the athletes' staff, more than an increase due to the period of sleep extension. It is important to note
333 that almost 50 % of the studies used in this review had a moderate or serious risk of bias and almost 70
334 % of the studies were between very low and moderate quality. All six outcomes that showed a large
335 effect size showed methodological bias that may have affected the results. Therefore, it is understood
336 that the positive results of sleep extension on athletes' performance are overestimated, considering the
337 methodological flaws presented in the articles cited.

338 The effect of sleep restriction on physical performance is not the scope of this review,
339 however, we would like to emphasize that research related to banking sleep is scarce in sports
340 sciences. In one of the first studies on banking sleep, Rupp, Wesensten ⁴⁵ found that the effect of sleep
341 restriction on cognitive performance is dependent on the amount of sleep prior to the restriction
342 period. In a randomized crossover study, Arnal, Sauvet ⁴⁶ confirmed this hypothesis and observed that
343 the sleep extension for one week attenuated the effects of sleep deprivation on psychomotor
344 performance. It is important to highlight that studies on this theme did not use athletes and
345 performance measures as dependent variables. Considering that athletes may have their sleep impacted
346 by competition, using banking sleep prior to competition may be an area of interest in future studies.
347 Moreover, all outcomes were evaluated in-laboratory settings, with control exerted on several
348 parameters. Considering that sports performance can be affected by the environment that it is
349 performed,^{47, 48} it would be prudent that future research evaluate the effects of sleep extension on the
350 real-life condition, especially, during a real sports competition.

351 The results observed in this review cannot be extrapolated to all athletes. In this review, most
352 studies included only men and few sports, introducing a risk of bias in generalizing all information.
353 Moreover, the included studies differed widely with respect to the instrument used to describe sleep,
354 sample size and outcome measures. It is likely that indirect measures may have values that are not in
355 accordance with direct measures. Finally, there is a risk of bias in the interpretation of the results when
356 considering whether the athletes submitted to the intervention were short or long sleepers or were

357 sleep restricted. Moreover, there is a risk of language bias because we only included studies written in
358 English. Research, however, reveals that language bias does not necessarily influence the results of a
359 systematic review.⁴⁹

360

361 **5. Conclusion**

362 Before starting a sleep extension program, athletic trainers and medical staff should analyze
363 the chronotype and sleep pattern, and if the athlete is getting adequate sleep for their needs. The
364 current review highlights that coaches and staff should be careful to use sleep extension as a measure
365 to improve sports performance. Rather, establishing good sleeping habits and meeting the required
366 sleep length and duration should be a priority. Evidence from this systematic review indicates that
367 sleep extension interventions may be beneficial for athletes in different sports but should be viewed
368 with caution due to the risks of bias and the quality of the studies. Future research should consider
369 training status of athletes, order of testing and methodological flaws to validate the current results
370 observed.

371

372 **Acknowledgment**

373 This study is supported by Fundação de Amparo à Pesquisa do Estado de Minas Gerais
374 (FAPEMIG), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq),
375 Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Centro de Estudos
376 em Psicobiologia e Exercício (CEPE), Centro Multidisciplinar em Sonolência e Acidentes
377 (CEMSA), FEPE/UFMG, EEFFTO/UFMG and Ministério da Cidadania do Governo Federal
378 (Brasília, Brazil).

379 **REFERENCES**

- 380 1. Halson SL, Juliff LE. Sleep, sport, and the brain. *Progress in Brain Research*. 234:
381 Elsevier; 2017. p. 13-31.
- 382 2. Fullagar HHK, Duffield R, Skorski S, et al. Sleep and recovery in team sport: current
383 sleep-related issues facing professional team-sport athletes. *Int J Sport Physiol*
384 2015;10(8):950-7.doi:10.1123/ijsp.2014-0565.
- 385 3. Hirshkowitz M, Whiton K, Albert SM, et al. National Sleep Foundation’s sleep time
386 duration recommendations: methodology and results summary. *Sleep Health* 2015;1(1):40-
387 3.doi:10.1016/j.sleh.2014.12.010.
- 388 4. Kirschen GW, Jones JJ, Hale L. The impact of sleep duration on performance among
389 competitive athletes: a systematic literature review. *Clin J Sport Med*
390 2020.doi:10.1097/JSM.0000000000000622.
- 391 5. Bird SP. Sleep, recovery, and athletic performance: a brief review and
392 recommendations. *Strength Cond J* 2013;35(5):43-7.doi:10.1519/SSC.0b013e3182a62e2f.
- 393 6. Simpson NS, Gibbs E, Matheson GO. Optimizing sleep to maximize performance:
394 implications and recommendations for elite athletes. *Scand J Med Sci Spor* 2017;27(3):266-
395 74.doi:10.1111/sms.12703.
- 396 7. Mah CD, Mah KE, Dement WC. Extended sleep and the effects on mood and athletic
397 performance in collegiate swimmers. *Sleep* 2008;31.
- 398 8. Mah CD, Mah KE, Kezirian EJ, et al. The effects of sleep extension on the athletic
399 performance of collegiate basketball players. *Sleep* 2011;34(7):943-
400 50.doi:10.5665/SLEEP.1132.
- 401 9. Schwartz J, Simon Jr RD. Sleep extension improves serving accuracy: A study with
402 college varsity tennis players. *Physiol Behav* 2015;151:541-
403 4.doi:10.1016/j.physbeh.2015.08.035.

- 404 10. Fullagar H, Skorski S, Duffield R, et al. The effect of an acute sleep hygiene strategy
405 following a late-night soccer match on recovery of players. *Chronobiol Int* 2016;33(5):490-
406 505.doi:10.3109/07420528.2016.1149190.
- 407 11. Bonnar D, Bartel K, Kakoschke N, et al. Sleep interventions designed to improve
408 athletic performance and recovery: a systematic review of current approaches. *Sports Med*
409 2018;48(3):683-703.doi:10.1007/s40279-017-0832-x.
- 410 12. Walsh NP, Halson SL, Sargent C, et al. Sleep and the athlete: narrative review and
411 2021 expert consensus recommendations. *Br J Sports Med* 2020;Nov 3(Online ahead of
412 print).doi:10.1136/bjsports-2020-102025.
- 413 13. Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews
414 and meta-analyses: the PRISMA statement. *PloS Medicine* 2009;6(7):1-6.doi:10.7326/0003-
415 4819-151-4-200908180-00135.
- 416 14. Cohen J. A coefficient of agreement for nominal scales. *Educ Psychol Meas*
417 1960;20(1):37-46.doi:10.1177/001316446002000104.
- 418 15. Sterne JAC, Savović J, Page MJ, et al. RoB 2: a revised tool for assessing risk of bias
419 in randomised trials. *BMJ* 2019;366:l4898.doi:10.1136/bmj.l4898.
- 420 16. Sterne JA, Hernán MA, Reeves BC, et al. ROBINS-I: a tool for assessing risk of bias
421 in non-randomised studies of interventions. *BMJ* 2016;355:i4919.doi:10.1136/bmj.i4919.
- 422 17. McGuinness LA, Higgins JPT. Risk-of-bias VISualization (robvis): An R package and
423 Shiny web app for visualizing risk-of-bias assessments. *Res Synth Methods* 2020;Online
424 ahead of print.doi:10.1002/jrsm.1411.
- 425 18. Campbell M, McKenzie JE, Sowden A, et al. Synthesis without meta-analysis
426 (SWiM) in systematic reviews: reporting guideline. *BMJ* 2020;368.doi:10.1136/bmj.l6890.
- 427 19. Cohen J. Statistical Power Analysis for the Behavioral Sciences—Second Edition. 12
428 Lawrence Erlbaum Associates Inc. *Hillsdale, New Jersey* 1988;13.

- 429 20. Roberts SSH, Teo W-P, Aisbett B, et al. Extended Sleep Maintains Endurance
430 Performance Better than Normal or Restricted Sleep. *Med Sci Sport Exer* 2019;51(12):2516-
431 23.doi:10.1249/MSS.0000000000002071.
- 432 21. Suppiah HT, Low CY, Choong GCW, et al. Restricted and unrestricted sleep
433 schedules of Asian adolescent, high-level student athletes: effects on sleep durations,
434 marksmanship and cognitive performance. *Biol Rhythm Res* 2016;47(4):505-
435 18.doi:10.1080/09291016.2016.1151102.
- 436 22. Petit E, Mougin F, Bourdin H, et al. A 20-min nap in athletes changes subsequent
437 sleep architecture but does not alter physical performances after normal sleep or 5-h phase-
438 advance conditions. *European Journal of Applied Physiology* 2014;114(2):305-
439 15.doi:10.1007/s00421-013-2776-7.
- 440 23. Boukhris O, Trabelsi K, Ammar A, et al. A 90 min Daytime Nap Opportunity Is
441 Better Than 40 min for Cognitive and Physical Performance. *Int J Environ Res Public Health*
442 2020;17(13).doi:10.3390/ijerph17134650.
- 443 24. Ancoli-Israel S, Martin JL, Blackwell T, et al. The SBSM guide to actigraphy
444 monitoring: clinical and research applications. *Behav Sleep Med* 2015;13(sup1):S4-
445 S38.doi:10.1080/15402002.2015.1046356.
- 446 25. Sargent C, Lastella M, Halson SL, et al. The validity of activity monitors for
447 measuring sleep in elite athletes. *J Sci Med Sport* 2016;19(10):848-
448 53.doi:10.1016/j.jsams.2015.12.007.
- 449 26. Knutson KL, Rathouz PJ, Yan LL, et al. Intra-individual daily and yearly variability in
450 actigraphically recorded sleep measures: the CARDIA study. *Sleep* 2007;30(6):793-
451 6.doi:10.1093/sleep/30.6.793.
- 452 27. Aili K, Åström-Paulsson S, Stoetzer U, et al. Reliability of actigraphy and subjective
453 sleep measurements in adults: the design of sleep assessments. *J Clin Sleep Med*
454 2017;13(1):39-47.doi:10.5664/jcsm.6384.

- 455 28. Chai Y, Fang Z, Yang FN, et al. Two nights of recovery sleep restores hippocampal
456 connectivity but not episodic memory after total sleep deprivation. *Sci Rep* 2020;10(1):1-
457 11.doi:10.1038/s41598-020-65086-x.
- 458 29. Pejovic S, Basta M, Vgontzas AN, et al. Effects of recovery sleep after one work week
459 of mild sleep restriction on interleukin-6 and cortisol secretion and daytime sleepiness and
460 performance. *Am J Physiol Endocrinol Metab* 2013;305(7):E890-
461 E6.doi:10.1152/ajpendo.00301.2013.
- 462 30. Reynolds AC, Banks S. Total sleep deprivation, chronic sleep restriction and sleep
463 disruption. *Progress in Brain Research*. 185: Elsevier; 2010. p. 91-103.
- 464 31. Aeschbach D, Sher L, Postolache TT, et al. A longer biological night in long sleepers
465 than in short sleepers. *J Clin Endocrinol Metab* 2003;88(1):26-30.doi:10.1210/jc.2002-
466 020827.
- 467 32. Webb WB. Are short and long sleepers different? *Psychol Rep* 1979;44(1):259-
468 64.doi:10.2466/pr0.1979.44.1.259.
- 469 33. Aeschbach D, Cajochen C, Landolt H, et al. Homeostatic sleep regulation in habitual
470 short sleepers and long sleepers. *AmJPhysiolC* 1996;270(1):R41-
471 R53.doi:10.1152/ajpregu.1996.270.1.R41.
- 472 34. Ferrara M, De Gennaro L. How much sleep do we need? *Sleep Med Rev*
473 2001;5(2):155-79.doi:10.1053/smr.2000.0138.
- 474 35. Violani C, Devoto A, Lucidi F. Individual traits of subjects able to extend their sleep
475 beyond 10 hours. *J Sleep Res* 1996;5:244.
- 476 36. Violani C, Lucidi F, Devoto A. Individual traits of subjects able to extend their sleep
477 beyond of of eleven hours. *J Sleep Res* 1998;7(Supp 2).
- 478 37. Queiroz S, Ruiz F, Prado J, et al. The consequences of partial sleep restriction for
479 habitual sleep duration, sleepiness and reaction time in healthy males. *Sleep Health* 2020;In
480 Press.doi:10.1016/j.sleh.2020.04.002.

- 481 38. Gagnon P, De Koninck J, Broughton R. Reappearance of electroencephalogram slow
482 waves in extended sleep with delayed bedtime. *Sleep* 1985;8(2):118-
483 28.doi:10.1093/sleep/8.2.118.
- 484 39. Hao Y-L, Zhang B, Jia F-J, et al. A three-phase epidemiological study of short and
485 long sleepers in a middle-aged Chinese population: prevalence and characteristics. *Braz J*
486 *Med Biol Res* 2014;47(2):157-65.doi:10.1590/1414-431X20133430.
- 487 40. Núñez P, Perillan C, Arguelles J, et al. Comparison of sleep and chronotype between
488 senior and undergraduate university students. *Chronobiol Int* 2019;36(12):1626-37.
- 489 41. Lastella M, Roach GD, Halson SL, et al. The chronotype of elite athletes. *J Hum Kinet*
490 2016;54(1):219-25.
- 491 42. Vitale JA, Weydahl A. Chronotype, physical activity, and sport performance: a
492 systematic review. *Sports Med* 2017;47(9):1859-68.
- 493 43. Brooks JL. Counterbalancing for serial order carryover effects in experimental
494 condition orders. *Psychol Methods* 2012;17(4):600-14.doi:10.1037/a0029310.
- 495 44. Green C, Strobach T, Schubert T. On methodological standards in training and
496 transfer experiments. *Psychol Res* 2014;78(6):756-72.doi:10.1007/s00426-013-0535-3.
- 497 45. Rupp TL, Wesensten NJ, Bliese PD, et al. Banking sleep: realization of benefits
498 during subsequent sleep restriction and recovery. *Sleep* 2009;32(3):311-21.
- 499 46. Arnal PJ, Sauvet F, Leger D, et al. Benefits of sleep extension on sustained attention
500 and sleep pressure before and during total sleep deprivation and recovery. *Sleep*
501 2015;38(12):1935-43.doi:10.5665/sleep.5244.
- 502 47. Malcata RM, Hopkins WG. Variability of competitive performance of elite athletes: a
503 systematic review. *Sports Med* 2014;44(12):1763-74.

504 48. Rylands LP, Roberts SJ, Hurst HT. Variability in laboratory vs. field testing of peak
505 power, torque, and time of peak power production among elite bicycle motocross cyclists. *J*
506 *Strength Cond Res* 2015;29(9):2635-40.

507 49. Morrison A, Polisena J, Husereau D, et al. The effect of English-language restriction
508 on systematic review-based meta-analyses: a systematic review of empirical studies.
509 *International Journal of Technology Assessment in Health Care*
510 2012;28(2):138.doi:10.1017/S0266462312000086.

511

512 Table 1. Characteristics of the included studies.

| Study | Study design | Sample | | | | Pre-intervention | | At-intervention | | Post-intervention |
|------------------------|--------------|----------|---------------------------|------------|-----------|-----------------------------------|-------------|-----------------|---|-----------------------------------|
| | | Gender | Age (years old) (M±SD) | Modality | Period | Sleep parameter (h:min) (M±SD) | Instrument | Period | Strategy | Sleep parameter (h:min) (M±SD) |
| Mah et al. (2011) | nRCT | 11 males | 19.4 ± 1.4 | Basketball | 2-4 weeks | TST: 06:40 ± 01:01 | Actigraphy | 5-7 weeks | > 10 hours in bed per night Including naps | TST: 08:27 ± 01:18 |
| Petit et al. (2014) | RCT | 16 males | 22.2 ± 1.7 | - | 30 days | TST: 08:11 ± 00:41 | Sleep diary | 1 day | 20 min of nap | - |

513

514

515

516

517

518 Table 1. Continued

| Study | Study design | Sample | Pre-intervention | | | At-intervention | | Post-intervention | | |
|---------------------------|--------------|----------------------|------------------|----------|--------|---|-------------|-------------------|--|---|
| | | Gender | Age (years old) | Modality | Period | Sleep parameter (h:min) (M±SD) | Instrument | Period | Strategy | Sleep parameter (h:min) (M±SD) |
| Schwartz and Simon (2015) | nRCT | 5 males 7 females | 20.2 | Tennis | 1 week | Sleep per night: 07:08 ± 00:48 | Sleep diary | 1 week | > 9 hours of sleep per night Including naps | sleep per night: 08:51 ± 00:36 |
| Fullagar et al. (2016) | RCT | 20 males | - | Soccer | 3 days | Sleep duration: Control: 06:38 ± 01:01 SHS: 06:54 ± 01:06 | Actigraphy | 1 day | Without specific target of hours No naps | Sleep duration: Control: 04:30 ± 0:27 SHS: 06:09 ± 0:43 |

| Study | Study design | Sample | | | | Pre-intervention | | At-intervention | | Post-intervention |
|-----------------------|--------------|------------------------|---------------------------|-----------------------------------|--------|---|------------|-----------------|--|---|
| | | Gender | Age (years old) (M±SD) | Modality | Period | Sleep parameter (h:min) (M±SD) | Instrument | Period | Strategy | Sleep parameter (h:min) (M±SD) |
| Suppiah et al (2016) | nRCT | 12 males 12 females | 14.1 ± 1.4 | Pistol shooters Rifle shooters | 2 days | TST: RC: 07:05 ± 01:05 URC: 06:54 ± 01:09 | Actigraphy | 5 days | > 9 hours of actual sleep No naps | TST: RC: 05:42 ± 00:44 URC: 06:08 ± 00:47 |
| Roberts et al. (2016) | nRCT | 9 males | 30 ± 6 | Cycling Triathlon | 4 days | TIB: D1: 07:06 ± 00:48 D2: 06:30 ± 01:00 D3: 06:54 ± 00:42 | Actigraphy | 4 days | + 30% of hTIB No naps | TIB: D1: 08:36 ± 01:00 D2: 08:18 ± 00:36 D3: 08:12 ± 00:36 |

521

522 Table 1. Continued.

| Study | Study design | Sample | Pre-intervention | | | At-intervention | | Post-intervention | |
|------------------------|--------------|-----------------|------------------|-------------------------------|--------------------------------|---|------------|-------------------|-------------------------------------|
| | Gender | Age (years old) | Modality | Period | Sleep parameter (h:min) (M±SD) | Instrument | Period | Strategy | Sleep parameter (h:min) (M±SD) |
| Boukhris et al. (2020) | RCT | 14 males | 20.3 ± 3.0 | Football, rugby, and handball | 1 day | TST N40: 08:35 ± 01:19 N90: 08:45 ± 01:51 | Actigraphy | 1 day | - 40-min of nap 90-min of nap |

523 Abbreviations: D1, day 1; D2, day 2; D3, day 3; hTIB, habitual time in bed; M, mean; min, minutes; nRCT, non-randomized controlled trial; RC, restricted
 524 condition; RCT, Randomized Controlled Trial; SD, standard deviation; SHS, sleep hygiene strategy; TIB, time in bed; TST, total sleep time, URC,
 525 unrestricted condition.

526

527 Table 2. Percentage of change, effect size and confidence interval of the included studies.

| Study | Performance parameters | Percentage of change (%) | Effect size | |
|--------------------------|-------------------------|--------------------------|-------------------------------------|----------------|
| | | | Cohen's d (95% Confidence Interval) | Classification |
| Mah et al. (2011) | 282 feet sprint | -4.3 | 1.215 (-2.07/-0.26) | Large |
| | Free throws | +11.4 | 0.918 (0.01/1.76) | Large |
| | Three-point field goals | +13.7 | 0.757 (-0.13/1.59) | Medium |
| Petit et al. (2014) | Wingate – peak power | +0.9 | 0.050 (-0.65/0.74) | Trivial |
| | Wingate – mean power | +0.6 | 0.049 (-0.65/0.74) | Trivial |
| Scwartz and Simon (2015) | Serving accuracy | +17.3 | -0.421 (-0.40/1.21) | Small |
| Suppiah et al. (2016) | Shooting performance | Not described | Not described | Not described |

528

Continued

529

530

531

532 Table 1. (continued). Percentage of change, effect size and confidence interval of the included studies.

| Study | Performance parameters | Percentage of change (%) | Effect size | |
|------------------------|--|--|-------------------------------------|----------------|
| | | | Cohen's d (95% Confidence Interval) | Classification |
| Fullagar et al. (2016) | Countermovement jump height | +2.8 | 0.25 (-0.38/0.86) | Small |
| | Countermovement jump force | +1.3 | 0.10 (-0.52/0.72) | Trivial |
| | Yoyo intermittent recovery test distance | -4.6 | -0.20 (-0.82/0.42) | Small |
| | Yoyo intermittent recovery test MHR | -0.5 | -0.13 (-0.75/0.49) | Trivial |
| Roberts et al. (2019) | Time trial | Normal sleep s vs sleep extension: -3.2 | 0.583 (-1.80/0.73) | Medium |
| Boukhris et al. (2020) | Maximal voluntary isometric contraction | 40-min: +5.6 | 0.44 (-0.32/1.18) | Small |
| | | 90-min: +9.6 | 0.75 (-0.03/1.50) | Moderate |
| | Shuttle run test- higher distance | 40-min: +7.8 | 1.13 (0.3/1.89) | Large |
| | | 90-min: +10.1 | 1.28 (0.44/2.06) | Large |
| | Shuttle run test- total distance | 40-min: +7.8 | 0.97 (0.16/1.72) | Large |
| | | 90-min: +12.6 | 1.70 (0.79/2.51) | Large |

533