



LJMU Research Online

Reguitti, A, Pumo, ML, Mazzali, PA, Pastorello, A, Pignata, G, Elias-Rosa, N, Prentice, SJ, Reynolds, T, Benetti, S, Rodriguez, O, Mattila, S and Kuncarayakti, H

Low-luminosity Type II supernovae - III. SN 2018hwm, a faint event with an unusually long plateau

<http://researchonline.ljmu.ac.uk/id/eprint/18015/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Reguitti, A, Pumo, ML, Mazzali, PA, Pastorello, A, Pignata, G, Elias-Rosa, N, Prentice, SJ, Reynolds, T, Benetti, S, Rodriguez, O, Mattila, S and Kuncarayakti, H (2020) Low-luminosity Type II supernovae - III. SN 2018hwm. a faint event with an unusually long plateau. Monthly Notices of

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

Erratum: Low luminosity Type II supernovae – III. SN 2018hwm, a faint event with an unusually long plateau

by A. Reguitti ^{1,2,3}★ M. L. Pumo, ^{1,4,5} P. A. Mazzali, ^{6,7} A. Pastorello, ¹ G. Pignata, ^{2,3} N. Elias-Rosa, ^{1,8} S. J. Prentice ⁹ T. Reynolds ¹⁰ S. Benetti ¹ S. Mattila, ¹⁰ H. Kuncarayakti ^{10,11} and O. Rodríguez ¹²

¹INAF - Osservatorio Astronomico di Padova, Vicoletto dell'Osservatorio 5, 35122 Padova, Italy

²Departamento de Ciencias Físicas - Universidad Andres Bello, Avda. República 252, Santiago, Chile

³Millennium Institute of Astrophysics, Nuncio Monsenor Sótero Sanz 100, Providencia, Santiago, Chile

⁴Università degli studi di Catania, Dipartimento di Fisica e Astronomia “E. Majorana”, Via S. Sofia 64, Catania, 95123, Italy

⁵Laboratori Nazionali del Sud-INFN, Via S. Sofia 64, Catania, 95123, Italy

⁶Astrophysics Research Institute, Liverpool John Moores University, ic2, 146 Brownlow Hill, Liverpool L3 5RF, UK

⁷Max-Planck Institut für Astrophysik, Karl-Schwarzschild-Str. 1, D-85741 Garching, Germany

⁸Institute of Space Sciences (ICE, CSIC), Campus UAB, Carrer de Can Magrans s/n, 08193 Barcelona, Spain

⁹School of Physics, Trinity College Dublin, The University of Dublin, Dublin 2, Ireland

¹⁰Tuorla Observatory, Department of Physics and Astronomy, FI-20014 University of Turku, Finland

¹¹Finnish Centre for Astronomy with ESO (FINCA), FI-20014 University of Turku, Finland

¹²School of Physics and Astronomy, Tel Aviv University, Tel Aviv 69978, Israel

Key words: errata – addenda – supernovae: general – supernovae: individual: SN 2018hwm.

The paper ‘Low luminosity Type II supernovae – III. SN 2018hwm, a faint event with an unusually long plateau’ was published in MNRAS, 501, 1059–1071 (2021). After the publication of the paper, we realised that there was an error in the bolometric light curve of SN 2018hwm reported in fig. 7 (top panel), which was one order of magnitude fainter than the luminosity reported correctly in fig. 6 (bottom panel). As a consequence, the average luminosity of the plateau, as shown in fig. 7, is $\sim 3 \times 10^{40}$ erg s $^{-1}$ instead of $\sim 3 \times 10^{41}$ erg s $^{-1}$. The input bolometric light curve used for the modelling was hence 10 times fainter than the correct luminosity, and the SN parameters inferred from the modelling need to be revised.

We re-run the hydrodynamical modelling code over the correct bolometric data adopting the same assumptions on the explosion epoch, distance modulus and reddening ($MJD = 58425.0 \pm 1.5$, $DM = 33.58 \pm 0.19$ mag, $A_V = 0.071$ mag, respectively). The best match between the hydrodynamical model and the observed data is shown in Fig. 1. This new figure is intended to replace fig. 7 of Reguitti et al. (2021). The new progenitor parameters we infer are: initial radius $R_0 = 5.9 \times 10^{13}$ cm ($845 R_\odot$), kinetic plus thermal energy $E = 0.075 \times 10^{51}$ erg, ejected mass $M_{ej} = 8.0 M_\odot$ and Nickel mass $M(^{56}\text{Ni}) = 0.0085 M_\odot$. These values replace those originally reported in Reguitti et al. (2021): $R_0 = 1.5 \times 10^{13}$ cm ($215 R_\odot$), $E = 0.055 \times 10^{51}$ erg, $M_{ej} = 8.0 M_\odot$ and $M(^{56}\text{Ni}) = 0.002 M_\odot$. In particular, with the correct bolometric light curve, we infer R_0 and $M(^{56}\text{Ni})$ values which are a factor of about four larger than estimated before. Although the above mentioned error required a revision of the previous SN parameters, the new estimates do not significantly affect the main conclusions of Reguitti et al. (2021).

The larger progenitor radius and the slightly higher ^{56}Ni mass makes SN 2018hwm similar to other low-luminosity SNe IIP studied in the past (e.g. Pastorello et al. 2004; Spiro et al. 2014). In particular, the more expanded progenitor for SN 2018hwm resembles the observed red supergiant (RSG) progenitors of SNe 2005cs (Maund, Smartt & Danziger 2005) and 2008bk (Mattila et al. 2008; Van Dyk et al. 2012; Maund et al. 2014b; O’Neill et al. 2021). Kozyreva et al. (2021) also pointed out that SN 2018hwm is probably a Fe-CC SN from a relatively low-mass RSG. However, on the basis of the new aforementioned parameters, the electron-capture (EC) SN explosion of a super-asymptotic giant branch star cannot be definitely ruled out.

The expansion velocities of SN 2018hwm during the photospheric phase are comparable to those of SN 2005cs (Pastorello et al. 2009) and the samples of faint SNe IIP (Pastorello et al. 2004; Spiro et al. 2014). For the energy versus ejected mass ratio, we now revise the value to $E/M_{ej} = 0.0094$ foe/ M_\odot (where 1 foe = 1×10^{51} erg), which is among the lowest calculated for a Type IIP SN (Pumo et al. 2017; Lisakov et al. 2018 and references therein), and is nearly half of what is derived for other low luminosity SNe IIP for which a 8–10 M_\odot RSG progenitor was directly observed in pre-explosion archive images.

How the revised parameters impact the progenitor and explosion scenarios will be presented in a forthcoming paper (Valerin et al. 2021, in preparation).

In conclusion, with the new parameters, both the ECSN and the Fe-CCSN scenarios are still plausible scenarios for SN 2018hwm.

We apologize for the inconvenience.

* E-mail: andreareguitti@gmail.com




Figure 1. Comparison of the evolution of the main observables of SN 2018hwm with the best-fit model computed with the general-relativistic, radiation-hydrodynamics code on the correct data. The new best-fit model parameters are: total energy 0.075×10^{51} erg, radius at explosion 5.9×10^{13} cm, $M(^{56}\text{Ni}) = 0.0085 M_{\odot}$ and ejected mass $8.0 M_{\odot}$. Top, middle, and bottom panels show the bolometric light curve, the photospheric velocity, and the black-body photospheric temperature as a function of time, respectively.

REFERENCES

- Kozyreva A., Baklanov P., Jones S., Stockinger G., Janka H.-T., 2021, *MNRAS*, 503, 797
 Lisakov S. M., Dessart L., John Hillier D., Waldman R., Livne E., 2018, *MNRAS*, 473, 3863
 Mattila S., Smartt S. J., Eldridge J. J., Maund J. R., Crockett R. M., Danziger I. J., 2008, *ApJ*, 688, 91
 Maund J. R., Smartt S. J., Danziger I. J., 2005, *MNRAS*, 364, 33
 Maund J. R., Mattila S., Ramirez-Ruiz E., Eldridge J. J., 2014, *MNRAS*, 438, 1577
 O'Neill D., Kotak R., Fraser M., Mattila S., Pietrzynski G., Prieto J. L., 2021, *A&A*, 645, 7
 Pastorello A. et al., 2004, *MNRAS*, 347, 74
 Pastorello A. et al., 2009, *MNRAS*, 394, 2266
 Pumo M. L., Zampieri L., Spiro S., Pastorello A., Benetti S., Cappellaro E., Manicò G., Turatto M., 2017, *MNRAS*, 464, 3013
 Reguitti A. et al., 2021, *MNRAS*, 501, 1059
 Spiro S. et al., 2014, *MNRAS*, 439, 2873
 Valerin G. et al., 2021, in preparation
 Van Dyk S. D. et al., 2012, *AJ*, 143, 19

This paper has been typeset from a TeX/LaTeX file prepared by the author.