



LJMU Research Online

Beigrezaei, S, Darabi, Z, Davies, IG, Mazidi, M, Ghayour-Mobarhan, M and Khayyatzadeh, SS

Higher global diet quality score is related to lower prevalence of depression and poor quality of life among adolescent girls

<http://researchonline.ljmu.ac.uk/id/eprint/24785/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Beigrezaei, S, Darabi, Z, Davies, IG, Mazidi, M, Ghayour-Mobarhan, M and Khayyatzadeh, SS (2023) Higher global diet quality score is related to lower prevalence of depression and poor quality of life among adolescent girls. BMC Psychiatry. 23 (1).

LJMU has developed [LJMU Research Online](#) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

RESEARCH

Open Access



Higher global diet quality score is related to lower prevalence of depression and poor quality of life among adolescent girls

Sara Beigrezaei^{1,2†}, Zahra Darabi^{1,2†}, Ian G. Davies³, Mohsen Mazidi^{4,5,6}, Majid Ghayour-Mobarhan⁷ and Sayyed Saeid Khayyat-zadeh^{1,2,8*}

Abstract

Background Adolescence is a key time for the development of depression symptoms and the diet quality may be associated with mental health conditions. The present study examined the association between depression and quality of life (QoL) and the global diet quality score (GDQS) as a simple and standardized metric diet quality in Iranian adolescents.

Methods This cross-sectional study was conducted on 733 adolescent girls recruited using a random cluster sampling method. A 147-item food frequency questionnaire (FFQ) was used for dietary intake assessment. The GDQS is gained by summing points of all the 25 food groups, ranged from 0 to 49. Depression symptoms were assessed using a Persian version of the Beck Depression Inventory (BDI). For assessment of health-related QoL, the Short Form 12 Survey–version 2 (SF-12v2) questionnaire was employed. Multivariable logistic regression examined the association of depression and QoL with GDQS in crude and adjusted models.

Results Adolescent girls in the highest tertile of GDQS score compared with the lowest tertile had a 41% lower odds of depressive symptoms (OR: 0.59; 95% CI: 0.39–0.90, $P = 0.01$). The participants in the third tertile of GDQS score had lower odds of poor QoL compared with the first tertile (OR: 0.56; 95% CI: 0.37–0.85, $P < 0.01$). These associations remained significant (both $P = 0.01$) after adjustment for age, energy intake, body mass index (BMI), physical activity, and menstruation (depressive symptoms: OR: 0.59; 95% CI: 0.38–0.92; QoL: OR: 0.59; 95% CI: 0.38–0.91, $P = 0.01$).

Conclusion We found that adolescent girls with a higher score of the GDQS had lower odds of depression and poor QoL. Prospective and interventional investigations are needed to reach a clear vision.

Keywords Depression, Quality of life, Global dietary quality score, GDQS, Cross-sectional, Adolescents

[†]Sara Beigrezaei and Zahra Darabi contributed equally to this work.

*Correspondence:

Sayyed Saeid Khayyat-zadeh
khayyat-zadeh@yahoo.com

¹Research Center for Food Hygiene and Safety, School of Public Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

²Department of Nutrition, School of Public Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

³Research Institute of Sport and Exercise Science, Liverpool John Moores University, Liverpool, UK

⁴Medical Research Council Population Health Research Unit, University of Oxford, Oxford, UK

⁵Nuffield Department of Population Health, Clinical Trial Service Unit and Epidemiological Studies Unit (CTSU), University of Oxford, Oxford, UK

⁶Department of Twin Research and Genetic Epidemiology, King's College London, South Wing St Thomas, London, UK

⁷International UNESCO center for Health-Related Basic Sciences and Human Nutrition, Mashhad University of Medical Sciences, Mashhad, Iran

⁸Nutrition and Food Security Research Center, Shahid Sadoughi University of Medical Sciences, Shohadaye gomnam BLD. ALEM square, Yazd, Iran



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

Depression is a common mental disorder worldwide [1], a major cause of disability, and is the main contributor to the global burden of disease [2]. Evidence has reported that women are more affected by depression than men [3]. Depression raises the risk of chronic diseases such as type 2 diabetes mellitus and cardiovascular diseases [2]; moreover, it is the main cause of death from suicide in the world [4]. Depression generally first emerges throughout adolescence and affects 8 to 20% of adolescents < 18 years worldwide [5]. In addition, depression can negatively affect the psychological and physical health of adolescents and result in some behavioral and social consequences including decreased quality of life (QoL) [6, 7]. QoL is a multi-dimensional concept of an individual's overall well-being situation in association with the value, cultural, social, and environmental circumstances in which they live [8].

As such, the identification and improvement of modifiable risk factors for adolescent depression can benefit their QoL [9]. Poor-quality diet is a major cause of unfavorable health outcomes [10, 11]. Depression has been linked to diet quality and a number of studies show associations between lower diet quality and higher depression risk in adolescents [12, 13]. For example, a cross-sectional study on adolescents aged 11–14 years reported an association between higher intake of unhealthy food items and symptoms of mental health [12]. Moreover, Jacka et al. [14] found that 10–14-year old adolescents with adherence to a less healthy diet had increased odds of depressive symptoms. Jacka et al. [15], in a separate study, also showed a dose response relationship between a higher healthy diet quality score and higher scores of health-related QoL in adolescents aged 11–18 years. However, McMartin et al. [16] did not reveal a link between diet quality and depression and anxiety in children aged 10–11 years.

Recently, the global dietary quality score (GDQS) was developed as a new tool to comprehensively evaluate the quality of diet [17].

The GDQS metric consists of 25 food groups that are based on current nutrition science and epidemiologic literature and that have a significant impact on nutrient intake and/or non-communicable disease (NCD) risk globally [17–19]. The GDQS sub metrics can be further subdivided to give more detailed information about the contribution of smaller sets of food groups or individual food groups to diet quality in populations [17]. Some of these food groups include whole/ refined grains, Nuts and seeds, deep orange tubers, deep orange vegetables and fruits, meats, dairies, etc. [17]. Though several metrics for diet quality have been developed worldwide, there is still not a validated, relatively simple, and widely used metric to assess the quality of diet in population-based

surveys [17]. The GDQS provides an appropriate simple and standardized metric for the population-based surveys of diet quality [17]. Evidence from prospective studies is consistent that adherence to this diet quality index is associated with a lower risk of diabetes and obesity and less weight gain in US women [20, 21].

To our knowledge, this is the first study to assess the association between mental health conditions and diet quality assessed by the GDQS, as a global score system. Therefore, in the present study, we aimed to evaluate diet quality in association with depression and QoL in adolescent girls.

Materials and methods

Study population

We used a random cluster sampling method to select 1026 adolescent girls (aged 12–18 years) from several schools in Mashhad and Sabzevar, northeastern Iran, for this cross-sectional study. In the first step, the 8 educational districts of Mashhad and Sabzevar were allocated to 4 clusters because of socioeconomic similarity. Then, one district from each cluster was randomly select. In the next step, two high schools will randomly select from each selected district. We excluded 38 subjects for various reasons and 255 subjects for under-reporting or over reporting their energy intake. Our final analysis included 733 adolescent girls.

Adolescent girls between the ages of 12 and 18 were included in this study. Participants with history of chronic diseases such as colitis, diabetes, cardiovascular disease (CVD), cancer, and hepatitis, individuals taking anti-inflammatory, antidepressant, anti-diabetic, or anti-obesity.

drugs as well as those on vitamin D, calcium supplementation, or hormone therapy within the last six months were not included. All adolescents and their parents provided complete written informed consent/assent before participating in the study. The study was approved by the ethic committee of Mashhad University of Medical Sciences (MUMS) (code number: 931,188).

Dietary assessment

A 147-item FFQ assessed dietary intakes by face-to-face interview, regarding daily, weekly, monthly and yearly intake, over the last year. Validity and reliability of this questionnaire has been previously reported [22, 23]. The reported portion sizes were converted to grams using household measures, and the energy and nutrient intakes were calculated using the Nutritionist IV software [10].

GDQS calculation

The GDQS metric is composed of 25 food groups (Supplementary Table 1). Points are attributed based on three or four categories of consumed amounts (defined in g/d)

specific to each group. Supplementary Table 1 described the scoring of 25 food groups. This was broken down into 16 healthy food groups, of which higher intake received higher points; seven unhealthy food groups, receiving more points for lower intake, and ultimately two food groups categorized as unhealthy when consumed in excessive amounts (more points for lower intake). The GDQS is scored by summing points of all 25 food groups, ranging from 0 to 49 [17].

Assessment of depression

Depression symptoms were assessed using a Persian version of the Beck Depression Inventory (BDI). The validity and reliability of this questionnaire was confirmed in previous studies [24, 25]. This questionnaire included 21 items that evaluated various symptoms of depression; feelings of guilt, hopelessness, and sadness; crying, sleep disturbance, fear and loss of appetite over the past 2 weeks. The range of scores for the BDI was between 0 and 63. If the BDI score was <16, the individuals were considered as not depressed, and if this score was ≥ 16 , they were categorized as depressed [26].

Quality of life assessment

For assessment of health-related QoL, the SF-12v2 questionnaire was used. This questionnaire was a short form of the SF-36 questionnaire and a modified version of the SF-12v1 [27]. The validity and reliability of the questionnaire were confirmed in a previous Iranian study [28]. The questionnaire includes 12 items that evaluated 8 domains of health including physical function, physical role (physical health problems), bodily pain, general health, vitality, social functioning, emotional role (emotional problems), and mental health. The QoL scores range from 0 to 100. We used a median cut-point for the definition of poor QoL; therefore, the subjects were categorized as having poor QoL if their score was <43.

Demographic data collection and anthropometric measurements

General demographic data were gathered by experienced interviewers. This data included age, smoking status, menstruation status, medical history, supplement use, psychological

treatment and chronic diseases. Anthropometric measurements were performed by trained investigators using standard protocols (including weight, height and waist circumferences (WC)).

Weight was measured using a digital scale with an accuracy of 100 g, while participants were without shoes in a minimal clothing state. Height (without shoes) was measured using a stadiometer with an accuracy of 0.1 cm. Body mass index (BMI) was calculated as weight (kg).

divided by height squared (m^2). In addition, WC was measured at the narrowest level between the costal margin and iliac crest during minimal respiration in a standing position. Physical activity level was assessed using the Persian-translated modifiable activity questionnaire (MAQ) with high reliability and moderate validity [29]. Data on the time and frequency of light, moderate, high, and very hard intensity activities were obtained according to the list of common activities of daily life over the past year, and these activity data were transformed into metabolic equivalent-hours/week (Met/hours/week; 1 MET = 3.5 mL $kg^{-1} min^{-1}$ of O_2 consumption).

Statistical analysis

The participants were classified into tertiles of GDQS. General characteristics, biochemical parameters and nutrient intake across tertiles of GDQS were expressed as means \pm standard deviations (SDs) for continuous variables, and as frequencies and percentages for categorical variables. To examine differences between tertiles, one-way-ANOVA and chi-square tests were used for continuous and categorical variables, respectively. To investigate the relationship between GDQS with depression and QoL, multivariate and linear regression were conducted with crude and adjusted models (Model 1: adjusted for age and energy intake; Model 2: additionally, adjusted for BMI percentile; Model 3: additionally, adjusted for physical activity, menstruation). All statistical analyses were conducted using SPSS version 21. P-values < 0.05 were considered statistically significant.

Results

General characteristics study participants

Current cross-sectional study indicated that higher adherence to GDQS inversely associated to poor QoL and depression prevalence.

The mean \pm SD age of the participants was 14.5 ± 1.53 years. The prevalence of depression and poor QoL were 24% and 49%, respectively. General characteristics and anthropometric variables of the participants across tertiles of GDQS score were presented in Table 1. The participants in the lowest tertile of GDQS score had significantly lower percentile BMI compared to participants in the highest tertile of GDQS score ($P < 0.01$). Age, waist circumference, waist-to-hip ratio, physical activity and menstruation were not different among tertiles of GDQS score.

Dietary intake of study participants

Dietary intake of study participants across tertiles of GDQS score are shown in Table 2. Participants in the highest tertile of GDQS score compared to the participants who were in the lowest tertile had higher intake of energy, percent of protein, carbohydrate and fat. Intake of

Table 1 General characteristics of study participants by tertile of GDQS

	T1	T2	T3	P value*
Age(year)	14.70±1.608	14.40±1.524	14.53±1.467	0.09
Percentile BMI (kg/m ²)	42.00±28.65310	49.36±28.48770	50.76±29.36819	<0.01
Waist circumference (cm)	69.47±8.121	70.47±9.460	71.62±9.221	0.07
Waist-to-hip ratio	0.76±0.05294	0.76±0.06753	0.76±0.05575	0.79
Physical activity (MET h/week)	45.05±2.85	45.39±3.63701	45.67±3.60948	0.21
Menstruation (yes) n (%)	170 (91.4)	322 (89.2)	170 (93.4)	0.26
Quality of life score	41.30±7.84	41.77±8.10	43.47±7.75	0.02
Depression score	11.56±8.88	11.48±9.50	9.16±9.03	0.01

BMI: Body mass index; GDQS: Global Diet Quality Score

Values are means±SD and number (prevalence percent) for quantitative and qualitative variables, respectively.

*Obtained from one way Anova for continuous variables and Chi-squared test for categorical variables

vitamins A, C, B6 and calcium were significantly higher in participants in the third tertile of the GDQS score compared to first tertile (per 1000 kcal). As expected, participants in the lowest tertile of the GDQS score had lower intake of citrus fruits, dark green leafy vegetables, cruciferous vegetables, deep orange vegetables, legumes, nuts and seeds, liquid oils, fish and shellfish, poultry and game meat, egg, and total dairy compared with the highest tertile.

Association between GDQS score and depression and poor QoL

Multi-variable adjusted odds ratios (ORs) for depression and poor QoL categories across tertiles of GDQS score are represented in Table 3. Adolescent girls in the highest tertile of GDQS score compared with the lowest tertile had a 41% lower probability of depressive symptoms (OR: 0.59; 95% CI: 0.39–0.90, $P=0.01$). The participants in the third tertile of GDQS score had lower odds of poor QoL compared with the subjects in the first tertile (OR: 0.56; 95% CI: 0.37–0.85, $P<0.01$). These associations remained significant after adjustment for age, energy intake, BMI percentile, physical activity and menstruation (Depression: OR: 0.59; 95% CI: 0.38–0.92; QoL: OR: 0.59; 95% CI: 0.38–0.91, $P\leq 0.01$ for both).

The association between Score of depression and quality of life with GDQS were presented in Table 4. There is a inverse relationship between GDQS with depression score ($\beta=-0.010$) and a positive relationship with quality of life score ($\beta=0.095$) in crude model.

Discussion

In the present cross-sectional study, we found that adolescent girls with a higher score of the GDQS had a lower risk of depression and poor QoL. This favorable association remained in multivariate models after controlling for some lifestyle-related variables. To our knowledge, this is the first study to investigate GDQS and risk of depression and poor QoL.

In line with our findings, other studies have reported significant relationships between depression and diet quality. Gibson-Smith et al. [30] showed the Mediterranean diet score and the Alternative healthy eating index (AHEI), as measures of diet quality, are lower in adults with depression. A prospective study in midaged females reported that long-term maintenance of good diet quality may be related to reduced depression risk (9). Moreover, a systematic review of observational studies showed a link between a diet rich in healthy foods and lower levels of depression in adolescents (31). Whereas, Winpenny et al. [32] reported no prospective or cross-sectionally associations between diet quality, fruit and vegetable intake, or fish intake with depressive symptoms in adolescence. Some studies were conducted in IRAN [33, 34]. Results of a cross-sectional study, showed that moderate adherence to the Dietary Approaches to Stop Hypertension (DASH) dietary pattern was inversely associated with depression in Iranian adults. The DASH-style diet is a healthy dietary pattern that consists of fruits, vegetables, low-fat dairy, and plant proteins from nuts and legumes, and that limits red meat, sweets, and sugar-sweetened beverages (33). Another study has suggested that Iranian adolescent females who consumed a pro-inflammatory diet, as measured by higher dietary inflammatory index (DII) scores, were more likely to have depressive symptoms (34).

In our study, GDQS was associated with adolescent QoL cross-sectionally. Jacka et al. [15] in both longitudinal and cross-sectional studies showed the diet quality of Australian adolescents was related to their mental health according to scores on the emotional subscale of the Pediatric Quality of Life Inventory. In addition, Esteban-Gonzalo et al. [35], from cross-sectional analysis, revealed that high-quality diet (using the Mediterranean diet) adherence was directly associated with better health-related –QoL.

There are many nutritional components of a diet with high GDQS that may have a direct effect on various mechanisms that underpin depression and QoL, including the immune system and oxidative stress [31].

Table 2 Energy and dietary intakes of study participants by tertile of GDSQ

	T1	T2	T3	P-value*
Citrus fruits (g)	24.12 ± 53.99	28.98 ± 32.95	38.77 ± 41.17	<0.01
Other fruits (g)	134.96 ± 132.69	179.16 ± 172.79	231.53 ± 168.67	<0.001
Dark green leafy vegetables (g)	16.91 ± 18.53	36.87 ± 40.79	58.44 ± 43.06	<0.001
Cruciferous vegetables (g)	6.94 ± 17.74	11.21 ± 18.10	17.06 ± 19.78	<0.001
Deep orange vegetables (g)	0.27 ± 1.11	0.54 ± 1.94	0.47 ± 1.68	0.15
Other vegetables (g)	103.35 ± 112.49	145.00 ± 156.23	179.14 ± 141.59	<0.001
Legumes (g)	49.88 ± 51.51	71.36 ± 50.51	92.76 ± 71.34	<0.001
Deep orange tubers (g)	40.32 ± 44.86	37.07 ± 30.91	44.69 ± 36.42	0.08
Nuts and seeds (g)	10.18 ± 26.69	15.43 ± 33.29	24.30 ± 27.63	<0.001
Whole grains (g)	60.31 ± 116.32	57.37 ± 106.03	51.15 ± 79.78	0.59
Liquid oils (g)	4.27 ± 7.41	5.75 ± 7.33	8.13 ± 8.81	<0.001
Fish and shellfish (g)	5.74 ± 8.97	7.57 ± 9.77	12.17 ± 21.22	<0.001
Poultry and game meat (g)	18.89 ± 22.05	23.93 ± 22.16	30.90 ± 27.05	<0.001
Low fat dairy (g)	180.17 ± 173.82	242.67 ± 195.04	309.57 ± 240.83	<0.001
Eggs (g)	15.31 ± 16.29	20.96 ± 20.99	25.43 ± 24.96	<0.001
High fat dairy (g)	143.65 ± 153.49	190.44 ± 180.66	222.21 ± 164.88	<0.001
Red meat (g)	11.45 ± 17.24	13.78 ± 17.02	15.30 ± 13.37	0.02
Processed meat (g)	7.31 ± 9.74	7.81 ± 10.11	7.68 ± 10.23	0.85
Refined grains and baked goods (g)	423.02 ± 236.07	413.33 ± 224.04	458.13 ± 228.43	0.08
Sweets and ice cream (g)	50.45 ± 37.63	49.28 ± 42.25	45.79 ± 49.08	0.45
Sugar-sweetened beverages (g)	80.76 ± 142.33	51.69 ± 89.52	39.67 ± 60.92	<0.001
Juice (g)	54.19 ± 97.95	50.85 ± 92.62	71.87 ± 194.00	0.19
White roots and tubers (g)	8.29 ± 15.712	9.29 ± 12.90	1.63 ± 26.17	0.13
Purchased deep fried foods (g)	30.65 ± 31.04	33.95 ± 32.11	33.43 ± 27.52	0.42
Energy intake (Kcal)	2453.92 ± 832.39	2660.46 ± 765.80	3033.56 ± 786.70	<0.001
Carbohydrate (% energy)	56.32 ± 8.08	54.17 ± 7.01	53.89 ± 6.43	<0.001
Protein (% energy)	12.87 ± 2.24	13.86 ± 2.13	14.22 ± 2.07	<0.001
Fat (% energy)	33.04 ± 8.66	34.29 ± 7.42	34.21 ± 6.77	0.12
Cholesterol (mg/1000 Kcal)	78.92 ± 47.03	93.88 ± 47.58	95.91 ± 39.43	<0.001
Saturated fatty acid (gr/1000 Kcal)	10.55 ± 3.36	11.66 ± 3.32	11.52 ± 3.28	<0.001
Monounsaturated fatty acid (gr/1000 Kcal)	12.05 ± 0.95	12.20 ± 3.26	12.13 ± 3.08	0.88
Polyunsaturated fatty acid (gr/1000 Kcal)	8.41 ± 3.61	8.41 ± 3.04	8.31 ± 2.76	0.92
Calcium (mg/1000 Kcal)	387.75 ± 125.18	436.79 ± 139.06	456.74 ± 130.93	<0.001
Iron (mg/1000 Kcal)	7.41 ± 1.62	7.29 ± 1.41	7.43 ± 1.33	0.55
Vitamin A (mcg/1000 Kcal)	192.01 ± 371.85	219.71 ± 114.60	262.46 ± 119.93	<0.01
Vitamin C (mg/1000 Kcal)	30.15 ± 20.39	35.83 ± 19.33	41.21 ± 20.41	<0.001
Folate (mcg/1000 Kcal)	226.80 ± 55.74	226.96 ± 45.36	231.20 ± 45.40	0.53
Vitamin B6 (mg/1000 Kcal)	0.67 ± 0.13	0.72 ± 0.12	0.73 ± 0.11	<0.001
Vitamin B12 (mcg/1000 Kcal)	1.55 ± 3.79	1.57 ± 0.75	1.65 ± 0.72	0.88

GDSQ: Global Diet Quality Score

Values are means ± SDs

*Obtained from One way Anova

Compelling evidence from recent meta-analyses has demonstrated that a more inflammatory diet increases the risk of depression [36].

Higher scores of GDQS are related to higher adherence to a healthy dietary pattern containing high amounts of fish, fruits, vegetables, whole grains, low-fat dairy, nuts, and legumes. Some of these healthy foods are of interest due to their role in reducing inflammation [37, 38]. Whole grains as a main source of fiber is associated with increasing adiponectin and decreasing inflammatory

markers [39]. Moreover, dietary phytochemicals like polyphenols have anti-inflammatory effects on neuro-psychiatric systems [37]. High concentrations of omega-3 polyunsaturated fatty acids in marine foods, due to their anti-inflammatory properties, delay the beginning of cytokine-induced depression and play a key role in the regulation of emotions [40, 41].

In addition, the high content of antioxidants such as carotenoids, vitamin E, and vitamin C in fruits and vegetables as part of the GDQS have protective impacts

Table 3 Multivariable-adjusted odds ratio (OR) of the associations between GDQS with poor quality of life and depression

	T1 Reference	T2 OR (95%CI)	T3 OR (95%CI)	P value ¹	P trend
Poor quality of life					
Crude	1.00	0.97 (0.68–1.39)	0.56(0.37–0.85)	<0.01	<0.01
Model1	1.00	0.98(0.68–1.41)	0.56(0.36–0.86)	<0.01	<0.01
Model2	1.00	1.01(0.70–1.45)	0.58(0.37–0.89)	0.01	0.01
Model3	1.00	1.04(0.72–1.51)	0.59(0.38–0.91)	0.01	0.01
Depression					
Crude	1.00	0.65(0.43–0.98)	0.59(0.39–0.90)	0.01	0.01
Model1	1.00	0.65(0.43–0.98)	0.59(0.39–0.91)	0.01	0.01
Model2	1.00	0.66(0.43–0.99)	0.60(0.39–0.93)	0.02	0.01
Model3	1.00	0.65 (0.42–0.98)	0.59(0.38–0.92)	0.01	0.01

GDQS: Global Diet Quality Score; OR: odds ratio; CI: confidence interval.

¹Last tertile compared to first tertile.

Model 1: Adjusted for age and energy intake.

Model 2: Additionally, adjusted for BMI percentile.

Model 3: Additionally, adjusted for physical activity, menstruation.

Table 4 The association between Score of depression and quality of life with GDQS

Food intake	Score of Quality of life		Score of Depression	
	B	P value	B	P value ¹
GDQS				
Crude	-0.1	<0.01	0.095	0.1
Model1	-0.11	0.004	0.106	<0.01
Model2	-0.11	0.005	0.106	<0.01
Model3	-0.11	0.004	0.101	0.01

GDQS: Global Diet Quality Score

¹Obtained from linear regression

Model 1: Adjusted for age and energy intake

Model 2: Additionally, adjusted for BMI percentile

Model 3: Additionally, adjusted for physical activity, menstruation

against depression risk [42, 43]. The protective effect of antioxidants can be attributed to mechanisms that repair neuronal damage, especially the hippocampus neurons, which are related to depression [43]. The hypothalamus-pituitary-adrenal (HPA) axis through regulating glucocorticoid production contributes to the pathophysiology of neuropsychiatric disorders [44]. Evidence suggests people living with depression have disturbances to the HPA axis and excessive cortisol production [44]. Clinical trials with nutrient interventions, such as vitamin C, have revealed a reduction in the reactivity of cortisol to acute physiological stress in healthy adults [45]. Additionally, interventions using polyphenol-rich foods have shown a remarkably decrease in cortisol levels [46, 47]. Some studies reported that omega-3 fatty acids could improve cortisol levels in healthy and depressive adults, as well [48].

On the other hand, high intake of processed foods is associated with inflammation, which can contribute in

the pathogenesis of depression [49, 50]. A cross-sectional study established that the Western-type diet, containing high amounts of processed meat, refined grains, high-fat and high-sugar foods was linked to greater levels of low-grade inflammation [51] and the subsequent atrophy of the brain, which was directly related to depression [52]. Moreover, evidence has demonstrated that low-grade inflammation is becoming growingly common in adolescents [15]. In addition, it is demonstrated that dietary patterns rich in sweets and soft drinks were associated with an increased depression risk due to the alteration of endorphin levels and oxidative stress [53].

Although it is thought that depression can cause adherence to an unhealthy diet, some longitudinal studies have proposed that this direction of causality is a less likely justification for long-term associations [54]. Previous evidence proposed that sugary and sweetened food items may alleviate the distress of people with mood disturbance in short term. This relationship can have calmativ impacts in the short term.

To the best of our knowledge, this is the first study to explore associations between GDQS and depression and QoL in adolescents. Our findings provide the basis for future investigation to identify any causal link between GDQS and depression and QoL. The large sample size and a wide range of covariates involved in the analyses add strength to our study. However, there are some limitations regarding our findings that should be considered. The direction of the associations cannot be determined in cross-sectional studies. In addition, evaluating dietary intakes with the FFQ is prone to misreporting (over/underestimation) and misclassification. This study was entirely conducted on adolescent girls, limiting the generalizability of our findings to populations beyond this sample.

Conclusion

In summary, our findings demonstrated that greater GDQS is associated with lower depression prevalence and poor QoL in adolescent girls. Children and their parents should be informed which dietary intakes of high sources of antioxidant and fiber and lower consumption of sugar and saturated fatty acids improve psychological health. To confirm these findings, further research needs to be conducted in different areas, including longitudinal studies with larger sample sizes.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12888-023-05313-7>.

Supplementary Material 1

Acknowledgements

The authors gratefully thank the participants of this study.

Authors' contributions

S.Kh designed and conducted the study. S.B. and Z.D. participated in the drafted and analysis the first version of the manuscript. S.Kh and M.Gh-M. supervised the study. All authors critically reviewed the final version of the manuscript.

Funding

Mashhad University of Medical Sciences (MUMS) supported this study.

Data Availability

The data and materials of the current study is available from the corresponding author on reasonable request.

Declarations

Competing interests

The authors declare no competing interests.

Ethics approval and consent to participate

The ethical committee of Mashhad University of Medical Sciences in Mashhad approved the study (code number: 931188). The written informed consent was signed by all participants before the beginning study. All methods were carried out in accordance with relevant guidelines and regulations.

Consent for publication

Not applicable.

Received: 12 June 2023 / Accepted: 26 October 2023

Published online: 28 November 2023

References

1. Institute of Health Metrics and Evaluation. Global Health Data Exchange (GHDx). <http://ghdx.healthdata.org/gbd-results-tool?params=gbd-api-2019-permalink/d780dffbe8a381b25e1416884959e88b> (Accessed 1 May 2021).
2. World Health Organization. Diabetes Fact sheet [updated September 2021]. Available from: <https://www.who.int/news-room/fact-sheets/detail/depression>.
3. World Health Organization. Depression and other common mental disorders: global health estimates (no. WHO/MSD/MER/2017.2). Geneva: World Health Organization; 2017.
4. Milloy M, Wood E. Withdrawal from Methadone in US prisons: cruel and unusual? *The Lancet*. 2015;386(9991):316–8.
5. Naicker K, Galambos NL, Zeng Y, Senthilselvan A, Colman I. Social, demographic, and health outcomes in the 10 years following adolescent depression. *J Adolesc Health: Official Publication Soc Adolesc Med*. 2013;52(5):533–8.
6. Arraras JJ, Ibañez B, Pereda N, Iribarren S, Basterra I. The association of clinical insight and depression with quality of life in schizophrenia. *Psychiatry Res*. 2019;279:350–2.
7. Li Y-N, Shapiro B, Kim JC, Zhang M, Porszasz J, Bross R, et al. Association between quality of life and anxiety, depression, physical activity and physical performance in maintenance hemodialysis patients. *Chronic Dis Translational Med*. 2016;2(02):110–9.
8. Kuyken W, Orley J, Power M, Herrman H, Schofield H, Murphy B, et al. The world-health-organization quality-of-life assessment (Whoqol)-position paper from the world-health-organization. *Soc Sci Med*. 1995;41(10):1403–9.
9. Lai JS, Hure AJ, Oldmeadow C, McEvoy M, Byles J, Attia J. Prospective study on the association between diet quality and depression in mid-aged women over 9 years. *Eur J Nutr*. 2017;56(1):273–81.
10. Olatona FA, Onabanjo OO, Ugbaja RN, Nnoaham KE, Adelekan DA. Dietary habits and metabolic risk factors for non-communicable Diseases in a university undergraduate population. *J Health Popul Nutr*. 2018;37(1):21.
11. Wu C-H, Lin C-Y, Hsieh Y-P, Strong C, Meshki C, Lin Y-C, et al. Dietary behaviors mediate the association between food insecurity and obesity among socio-economically disadvantaged youth. *Appetite*. 2019;132:275–81.
12. Jacka FN, Rothon C, Taylor S, Berk M, Stansfeld SA. Diet quality and mental health problems in adolescents from East London: a prospective study. *Soc Psychiatry Psychiatr Epidemiol*. 2013;48(8):1297–306.
13. Kohlboeck G, Sausenthaler S, Standl M, Koletzko S, Bauer C-P, Von Berg A, et al. Food intake, diet quality and behavioral problems in children: results from the GINI-plus/LISA-plus studies. *Annals of Nutrition and Metabolism*. 2012;60(4):247–56.
14. Jacka FN, Kremer PJ, Leslie ER, Berk M, Patton GC, Toumbourou JW, et al. Associations between diet quality and depressed mood in adolescents: results from the Australian healthy neighbourhoods study. *Aust N Z J Psychiatry*. 2010;44(5):435–42.
15. Jacka FN, Kremer PJ, Berk M, de Silva-Sanigorski AM, Moodie M, Leslie ER, et al. A prospective study of diet quality and mental health in adolescents. *PLoS ONE*. 2011;6(9):e24805.
16. McMartin SE, Kuhle S, Colman I, Kirk SF, Veugelers PJ. Diet quality and mental health in subsequent years among Canadian youth. *Public Health Nutr*. 2012;15(12):2253–8.
17. Bromage S, Batis C, Bhupathiraju SN, Fawzi WW, Fung TT, Li Y, et al. Development and validation of a novel food-based Global Diet Quality score (GDQS). *J Nutr*. 2021;151(Supplement2):755–925.
18. Willett W, Rockström J, Loken B, Springmann M, Lang T, Vermeulen S, et al. Food in the Anthropocene: the EAT–Lancet Commission on healthy diets from sustainable food systems. *The Lancet*. 2019;393(10170):447–92.
19. Micha R, Shulkin ML, Penalvo JL, Khatibzadeh S, Singh GM, Rao M, et al. Etiologic effects and optimal intakes of foods and nutrients for risk of Cardiovascular Diseases and Diabetes: systematic reviews and meta-analyses from the Nutrition and Chronic Diseases Expert Group (NutriCoDE). *PLoS ONE*. 2017;12(4):e0175149.
20. Fung TT, Li Y, Bromage S, Bhupathiraju SN, Batis C, Fawzi W, et al. Higher global Diet Quality score is associated with less 4-year weight gain in US women. *J Nutr*. 2021;151(Supplement2):1625–75.
21. Fung TT, Li Y, Bhupathiraju SN, Bromage S, Batis C, Holmes MD, et al. Higher global Diet Quality score is inversely associated with risk of type 2 Diabetes in US women. *J Nutr*. 2021;151(Supplement2):1685–755.
22. Esfahani FH, Asghari G, Mirmiran P, Azizi F. Reproducibility and relative validity of food group intake in a food frequency questionnaire developed for the Tehran lipid and glucose study. *J Epidemiol*. 2010;20(2):150–8.
23. Asghari G, Rezazadeh A, Hosseini-Esfahani F, Mehrabi Y, Mirmiran P, Azizi F. Reliability, comparative validity and stability of dietary patterns derived from an FFQ in the Tehran lipid and glucose study. *Br J Nutr*. 2012;108(6):1109–17.
24. Ghassemzadeh H, Mojtabei R, Karamghadiri N, Ebrahimkhani N. Psychometric properties of a persian-language version of the Beck Depression Inventory-Second edition: BDI-II-PERSIAN. *Depress Anxiety*. 2005;21(4):185–92.
25. Scogin F, Beutler L, Corbishley A, Hamblin D. Reliability and validity of the short form Beck Depression Inventory with older adults. *J Clin Psychol*. 1988;44(6):853–7.

26. Smarr KL, Keefer AL. Measures of depression and depressive symptoms: Beck depression Inventory-II (BDI-II), center for epidemiologic studies depression scale (CES-D), geriatric depression scale (GDS), hospital anxiety and depression scale (HADS), and patient health Questionnaire-9 (PHQ-9). *Arthritis Care Res.* 2011;63(S11):454–566.
27. Cheak-Zamora NC, Wyrwich KW, McBride TD. Reliability and validity of the SF-12v2 in the medical expenditure panel survey. *Qual Life Res.* 2009;18(6):727–35.
28. Montazeri A, Vahdaninia M, Mousavi SJ, Asadi-Lari M, Omidvari S, Tavousi M. The 12-item medical outcomes study short form health survey version 2.0 (SF-12v2): a population-based validation study from Tehran. *Iran Health Qual life Outcomes.* 2011;9(1):1–8.
29. Momenan AA, Delshad M, Sarbazi N, Rezaei_Ghaleh N, Ghanbarian A, Azizi F. Reliability and validity of the modifiable activity questionnaire (MAQ) in an Iranian urban adult population. *Arch Iran Med.* 2012;15(5):279–82.
30. Gibson-Smith D, Bot M, Brouwer IA, Visser M, Penninx BW. Diet quality in persons with and without depressive and anxiety disorders. *J Psychiatr Res.* 2018;106:1–7.
31. O'neil A, Quirk SE, Housden S, Brennan SL, Williams LJ, Pasco JA, et al. Relationship between diet and mental health in children and adolescents: a systematic review. *Am J Public Health.* 2014;104(10):e31–e42.
32. Winpenny EM, van Harmelen A-L, White M, van Sluijs EM, Goodyer IM. Diet quality and depressive symptoms in adolescence: no cross-sectional or prospective associations following adjustment for covariates. *Public Health Nutr.* 2018;21(13):2376–84.
33. Valipour G, Esmailzadeh A, Azadbakht L, Afshar H, Hassanzadeh A, Adibi P. Adherence to the DASH diet in relation to psychological profile of Iranian adults. *Eur J Nutr.* 2017;56:309–20.
34. Shivappa N, Hebert JR, Neshatbini Tehrani A, Bayzai B, Naja F, Rashidkhani B. A pro-inflammatory diet is associated with an increased odds of depression symptoms among Iranian female adolescents: a cross-sectional study. *Front Psychiatry.* 2018;9:400.
35. Esteban-Gonzalo L, Turner AI, Torres SJ, Esteban-Cornejo I, Castro-Piñero J, Delgado-Alfonso Á, et al. Diet quality and well-being in children and adolescents: the UP&DOWN longitudinal study. *Br J Nutr.* 2019;121(2):221–31.
36. Lassale C, Batty GD, Baghdadli A, Jacka F, Sánchez-Villegas A, Kivimäki M, et al. Healthy dietary indices and risk of depressive outcomes: a systematic review and meta-analysis of observational studies. *Mol Psychiatry.* 2019;24(7):965–86.
37. Yahfoufi N, Alsadi N, Jambi M, Matar C. The immunomodulatory and anti-inflammatory role of polyphenols. *Nutrients.* 2018;10(11):1618.
38. Marx W, Lane M, Hockey M, Aslam H, Berk M, Walder K, et al. Diet and depression: exploring the biological mechanisms of action. *Mol Psychiatry.* 2021;26(1):134–50.
39. Huang T, Xu M, Lee A, Cho S, Qi L. Consumption of whole grains and cereal fiber and total and cause-specific mortality: prospective analysis of 367,442 individuals. *BMC Med.* 2015;13(1):1–9.
40. Liao Y, Xie B, Zhang H, He Q, Guo L, Subramaniapillai M, et al. Correction: efficacy of omega-3 PUFAs in depression: a meta-analysis. *Translational Psychiatry.* 2021;11(1):465.
41. Su K-P, Matsuoka Y, Pae C-U. Omega-3 polyunsaturated fatty acids in Prevention of Mood and anxiety disorders. *Clin Psychopharmacol Neurosci.* 2015;13(2):129–37.
42. Akbaraly TN, Brunner EJ, Ferrie JE, Marmot MG, Kivimaki M, Singh-Manoux A. Dietary pattern and depressive symptoms in middle age. *Br J Psychiatry.* 2009;195(5):408–13.
43. Sarandol A, Sarandol E, Eker SS, Erdinc S, Vatansever E, Kirli S. Major depressive disorder is accompanied with oxidative stress: short-term antidepressant treatment does not alter oxidative–antioxidative systems. *Hum Psychopharmacology: Clin Experimental.* 2007;22(2):67–73.
44. Naughton M, Dinan TG, Scott LV. Corticotropin-releasing hormone and the hypothalamic–pituitary–adrenal axis in psychiatric Disease. *Handb Clin Neurol.* 2014;124:69–91.
45. Brody S, Preut R, Schommer K, Schürmeyer TH. A randomized controlled trial of high dose ascorbic acid for reduction of blood pressure, cortisol, and subjective responses to psychological stress. *Psychopharmacology.* 2002;159(3):319–24.
46. Tsang C, Hodgson L, Bussu A, Farhat G, Al-Dujaili E. Effect of polyphenol-rich dark chocolate on salivary cortisol and mood in adults. *Antioxidants.* 2019;8(6):149.
47. Tsang C, Smail NF, Almoosawi S, Davidson I, Al-Dujaili EA. Intake of polyphenol-rich pomegranate pure juice influences urinary glucocorticoids, blood pressure and homeostasis model assessment of insulin resistance in human volunteers. *J Nutritional Sci.* 2012;1.
48. Barbadoro P, Annino I, Ponzio E, Romanelli RM, D'Errico MM, Prospero E, et al. Fish oil supplementation reduces cortisol basal levels and perceived stress: a randomized, placebo-controlled trial in abstinent alcoholics. *Mol Nutr Food Res.* 2013;57(6):1110–4.
49. Lopez-Garcia E, Schulze MB, Fung TT, Meigs JB, Rifai N, Manson JE, et al. Major dietary patterns are related to plasma concentrations of markers of inflammation and endothelial dysfunction. *Am J Clin Nutr.* 2004;80(4):1029–35.
50. Tiemeier H, Hofman A, van Tuijl HR, Kiliaan AJ, Meijer J, Breteler MM. Inflammatory proteins and depression in the elderly. *Epidemiology.* 2003;14:103–7.
51. Liu S, Manson JE, Buring JE, Stampfer MJ, Willett WC, Ridker PM. Relation between a diet with a high glycemic load and plasma concentrations of high-sensitivity C-reactive protein in middle-aged women. *Am J Clin Nutr.* 2002;75(3):492–8.
52. Li Y, Lv MR, Wei YJ, Sun L, Zhang JX, Zhang HG, et al. Dietary patterns and depression risk: a meta-analysis. *Psychiatry Res.* 2017;253:373–82.
53. Westover AN, Marangell LB. A cross-national relationship between sugar consumption and major depression? *Depress Anxiety.* 2002;16(3):118–20.
54. Le Port A, Gueguen A, Kesse-Guyot E, Melchior M, Lemogne C, Nabi H, et al. Association between dietary patterns and depressive symptoms over time: a 10-year follow-up study of the GAZEL cohort. *PLoS ONE.* 2012;7(12):e51593.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.