



LJMU Research Online

Saini, J, McPhee, JS, Al-Dabbagh, S, Stewart, CE and Al-Shanti, N

Regenerative function of immune system: Modulation of muscle stem cells

<http://researchonline.ljmu.ac.uk/id/eprint/3497/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Saini, J, McPhee, JS, Al-Dabbagh, S, Stewart, CE and Al-Shanti, N (2016)
Regenerative function of immune system: Modulation of muscle stem cells.
Ageing Research Reviews, 27. pp. 67-76. ISSN 1568-1637**

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

1 **Regenerative function of immune system: Modulation of muscle stem cells**

2
3 Jasdeep Saini¹, Jamie S. McPhee¹, Sarah Al-Dabbagh¹, Claire E. Stewart² & Nasser Al-Shanti ^{1*}

4
5 ¹ Work was performed in: Healthcare Science Research Institute, School of Healthcare Science
6 Manchester Metropolitan University, John Dalton Building, Chester Street, M1 5GD, Manchester,
7 UK.

8
9 ² Present address: Research Institute for Sport & Exercise Sciences, School of Sport and Exercise
10 Sciences, Tom Reilly Building, Byrom Street Campus, Liverpool John Moores University, Liverpool,
11 L3 3AF.

12
13 * To whom correspondence should be addressed: Dr. NASSER AL-SHANTI. School of Healthcare
14 Science, John Dalton Building, Chester Street, M1 5GD, Manchester, UK. tel: 0044 161 2475712;
15 fax: 0044 0161 247 6831; email: n.al-shanti@mmu.ac.uk

16
17
18 Keywords: Muscle Treg, regeneration, skeletal muscle, immune system,

19
20
21
22
23 Running Head: The role of immune cells in muscle regeneration

30 **ABSTRACT**

31 Ageing is characterised by progressive deterioration of physiological systems and the loss of
32 skeletal muscle mass is one of the most recognisable, leading to muscle weakness and mobility
33 impairments. This review highlights interactions between the immune system and skeletal
34 muscle precursor cells (widely termed satellite cells or myoblasts) to influence satellite cell
35 behaviour during muscle regeneration after injury, and outlines deficits associated with ageing.
36 Resident neutrophils and macrophages in skeletal muscle become activated when muscle fibres
37 are damaged via stimuli (e.g. contusions, strains, avulsions, hyperextensions, ruptures) and
38 release high concentrations of cytokines, chemokines and growth factors into the
39 microenvironment. These localised responses serve to attract additional immune cells which can
40 reach in excess of 1×10^5 immune cell/mm³ of skeletal muscle in order to orchestrate the repair
41 process. T-cells have a delayed response, reaching peak activation roughly 4 days after the initial
42 damage. The cytokines and growth factors released by activated T-cells play a key role in muscle
43 satellite cell proliferation and migration, although the precise mechanisms of these interactions
44 remain unclear. T-cells in older people display limited ability to activate satellite cell proliferation
45 and migration which is likely to contribute to insufficient muscle repair and, consequently, muscle
46 wasting and weakness. If the factors released by T-cells to activate satellite cells can be identified,
47 it may be possible to develop therapeutic agents to enhance muscle regeneration and reduce the
48 impact of muscle wasting during ageing and disease.

49

50 **Highlights:**

- 51 • Immune cells infiltrate damaged skeletal muscles to release cytokines, chemokines and
52 growth factors into the localised area that alter the micro-environment to clear cellular
53 debris and activate muscle satellite cells.
- 54 • In young adults, the factors released by T-cells, in particular the regulatory T-cells, can
55 extend the period of satellite cell proliferation to enhance muscle repair.
- 56 • In old adults, the T-cells do not release appropriate factors into the micro-environment
57 and this may contribute to inadequate muscle recovery and consequently, to age-related
58 deficits in muscle size and function.
- 59 • Identification of the factors released by young immune cells to regulate muscle
60 regeneration could lead to the development of novel therapeutic agents to treat muscle
61 wasting disorders and ageing.

62

63

64

65

66

67

68

69

70

71

72

73

74

75	CONTENTS
76	
77	1. INTRODUCTION
78	1.1 Myogenesis and satellite cell activation
79	1.2 Young myogenesis
80	1.3 Aged myogenesis
81	
82	2. INNATE IMMUNITY & MUSCLE REGENERATION
83	2.1 Innate immune response to acute damage and repair
84	2.2 Regulation of skeletal muscle regeneration via innate immune cell signalling
85	
86	3. ADAPTIVE IMMUNITY & MUSCLE REGENERATION
87	3.1 Adaptive immune response to muscle damage
88	3.2 Regulation of skeletal muscle regeneration via Regulatory T-cells
89	
90	4. CONCLUSIONS
91	
92	5. REFERENCES
93	
94	
95	
96	
97	
98	
99	
100	
101	
102	
103	

104 **1. INTRODUCTION**

105 In older age, skeletal muscle atrophies considerably (Maden-Wilkinson et al., 2014; Lexell et al.,
106 1988; Janssen et al., 2002; Lexell, 1995; Morley et al., 2001), which contributes to weakness and
107 mobility impairments inherent to sarcopenia (Janssen, 2011; Cruz-Jentoft et al., 2010) and frailty
108 (Fried et al., 2001). Loss of skeletal muscle mass and function with ageing are associated with
109 altered immune, hormonal and metabolic factors directly impacting on muscle (Narici & Maffulli,
110 2010) and resulting in motor unit remodelling (Piasecki, Ireland, Jones, et al., 2015; Piasecki,
111 Ireland, Stashuk, et al., 2015). This review will first outline the role of the immune system in
112 myogenesis that occurs after injury and then discuss how changes in immune cells may
113 contribute to ageing-related muscle impairments. Identification of the signalling molecules
114 exchanged between immune and satellite cells may lead to novel therapeutic strategies to
115 preserve muscle with advancing old age and muscle wasting conditions.

116

117 **1.1 Myogenesis and satellite cell activation**

118 Skeletal muscle is the most abundant tissue type in healthy humans (Yin et al., 2013). It powers
119 movements and contributes to metabolism by storing amino acids, glucose and fatty acids as well
120 as oxidising substrates to replenish adenosine triphosphate stores (Leto & Saltiel, 2012). Muscles
121 also release cytokines and growth factors into the extra-cellular compartments to act locally or
122 systemically (Pedersen, 2011). The production of skeletal muscle cells occurs during embryonic
123 myogenesis and thereafter myofibres themselves are incapable of proliferation (Bentzinger et
124 al., 2012). Hence, the number of skeletal muscle fibres is largely determined before birth.
125 Postnatal muscle growth arises by adapting and remodelling pre-existing fibres and through
126 recruitment of resident, non-fused, self-renewing satellite cells (Tedesco et al., 2010). Satellite
127 cells reside beneath the basal lamina of mature fibres in a quiescent state, they neither undergo
128 cell division nor differentiation unless they are specifically activated to do so (Kuang et al., 2007).
129 Damage to the muscle through injury or very intense prolonged unaccustomed exercise training
130 are examples of principal activators of quiescent satellite cells.

131

132

133 **1.2 Young myogenesis**

134 Increases in muscle mass (hypertrophy) and adapted metabolism after exercise training in adults
135 improves athletic performance and health (Egan & Zierath, 2013). The training-induced
136 hypertrophy can depend on satellite cell proliferation and differentiation (Joanisse et al., 2013;
137 Yin et al., 2013). However, hypertrophy may not necessarily require activation of satellite cells,
138 since a satellite cell deficient mouse model showed normal training-induced hypertrophy
139 (McCarthy et al., 2011; D. J. Glass, 2003). Satellite cells are, however, centrally involved in muscle
140 regeneration after damage (Lepper et al., 2011; Yin et al., 2013). Some minor muscle damage can
141 be a feature of everyday living that goes largely unnoticed by the individual due to minimal
142 muscle tenderness and no apparent effect on function. More painful and functionally impairing
143 damage can occur after repeated intense or rapid muscular activations, especially following
144 unaccustomed high-load eccentric contractions (lengthening under strain) performed across a
145 large range of motion (Paulsen et al., 2012) or electrical stimulation protocols (Crameri et al.,
146 2007; Mackey et al., 2008; Nosaka et al., 2011). External stressors such as heavy impact causing
147 contusion, traumatic puncture wounds or pathogen invasion can also damage otherwise healthy
148 muscle, and in animal models, damage can be induced through injection of substances such as
149 cardiotoxin (Ctx) (Sousa-Victor et al., 2014). Once activated, satellite cells migrate to the
150 damaged site and re-enter into the cell cycle (Tedesco et al., 2010; Siegel et al., 2009) to generate
151 the required concentration of myoblasts through several cycles of proliferation to regenerate
152 damaged fibres. Although the majority of activated satellite cells differentiate into myotubes, a
153 population of satellite cells return to a quiescent state (self-renewal) to maintain their numbers
154 for the next incidence of muscle injury (Relaix & Zammit, 2012; Yin et al., 2013). The
155 differentiated myotubes either fuse with pre-existing damaged myofibers to provide additional
156 myonuclei during muscle regeneration, or fuse with each other forming de novo myofibers to
157 replace the damaged myofibres during muscle regeneration (Adams, 2006; Siegel et al., 2011).

158

159 Satellite cells do not function in an isolated environment, a number of non-myogenic cells also
160 populate muscle and influence the regenerative actions of satellite cells (Cerletti et al., 2008). For
161 example, mesenchymal interstitial cells (Farup et al., 2015; Uezumi et al., 2014) and infiltrating

162 immune cells secrete numerous cytokines and growth factors into the localised
163 microenvironment that orchestrate muscle regenerative mechanisms by clearing cellular debris
164 and facilitating repair (Tedesco et al., 2010). These cytokines are not necessarily released into the
165 general circulation to act systemically (Steensberg et al., 2002). Thus, effective muscle repair and
166 regeneration relies not only on muscle satellite cells (known as the intrinsic niche) but also on
167 other distinct cell types and their locally secreted cytokines (termed the extrinsic niche).

168

169 **1.3 Aged Myogenesis**

170 Muscle from aged mice was estimated to contain around 65% fewer functioning satellite cells
171 than muscle from young mice (Cosgrove et al., 2014) and the overall number of satellite cells was
172 also lower in aged mouse muscle (Chakkalakal et al., 2012). However, this was not the main cause
173 of sarcopenia, at least not in mice, where induced depletion of satellite cells in young adults had
174 little impact on the rate of muscle ageing (Fry et al., 2015). It is interesting to note that healthy
175 older people do maintain the ability to activate satellite cells after intense exercise (Verdijk et al.,
176 2009). However, if the activation of satellite cells cannot keep pace with damage, then muscle
177 wasting or atrophy will inevitably occur. The loss of muscle mass with ageing has been linked to
178 the reduced regenerative actions of older satellite cells and altered immune response to damage
179 (Peake et al., 2010; Degens, 2010). There are reports of intrinsic deficiencies within satellite cells
180 that reduce their activity. For instance, two-thirds of satellite cells in older mice showed reduced
181 capacity for muscle regeneration due to elevated activity of p38 α and p38 β mitogen-activated
182 kinase signalling which was not overcome by transplantation into a young recipient (Cosgrove et
183 al., 2014). However, the debate continues as to whether or not satellite cell intrinsic deficits can
184 be overcome by exposure to a 'young' microenvironment (reviewed elsewhere: (Brack & Munoz-
185 Canoves, 2015)). There is strong evidence implicating the aged microenvironment with reduced
186 satellite cell responses (Chakkalakal et al., 2012; Barberi et al., 2013). Transplanted muscle from
187 young into old mice fails to regenerate, but transplanted muscle from old into young regenerate
188 (B. M. Carlson & Faulkner, 1989), but might have a delayed regenerative response (Smythe et al.,
189 2008). Moreover, 'rejuvenating' the microenvironment in older mice enhanced activation of

190 satellite cells through increased Notch signalling, as shown in heterochronic parabiosis models
191 (Conboy et al., 2005; Morgan E. Carlson et al., 2008).

192

193 Lower satellite cell function with ageing was linked to increased activity of the transforming
194 growth factor beta (TGF- β) family of molecules within satellite cells that are negative regulators
195 of growth and restrict the proliferative responses (M. E. Carlson et al., 2009; Sousa-Victor et al.,
196 2014; Yousef et al., 2015). Elevated fibroblast growth factor (FGF) signalling from the aged
197 microenvironment was associated with depletion of the stem cell population and impaired
198 regenerative capacity, but was countered in the aged satellite cells that had higher levels of
199 Sprouty1 (Spry1) to inhibit FGF signalling (Chakkalakal et al., 2012). By altering satellite cell
200 signalling through Notch, Wnt and receptor tyrosine kinases/extracellular signal-regulated kinase
201 (RTK/ERK) it has been possible to overcome deficits in aged satellite cell function (Brack & Rando,
202 2007; Morgan E. Carlson et al., 2008; Naito et al., 2012). Circulating soluble factors, such as
203 hormones, or other molecules released locally into the microenvironment may influence the
204 intracellular satellite cell signalling to regulate proliferative and differentiation responses. For
205 example, elevating the circulating oxytocin had rejuvenating effects for satellite cells (Elabd et
206 al., 2014); increasing circulating levels of growth differentiation factor 11 (GDF-11) also
207 rejuvenated satellite cells (Sinha et al., 2014). However, alternative research investigating the
208 effect of GDF-11 on myogenesis observed a significant inhibition of skeletal muscle regeneration
209 (Brun & Rudnicki, 2015). Additionally, elevated levels of osteopontin in aged mice was associated
210 with impaired satellite cell responses to damage and this was overcome by reducing osteopontin
211 *in vitro* and *in vivo* (Paliwal et al., 2012). Thus, a key detail, which has not yet been fully
212 understood, is how the satellite cells respond to the rapidly changing microenvironment
213 occurring soon after muscle damage, which is heavily influenced by the infiltrating immune cells.

214

215 **2. INNATE IMMUNITY & MUSCLE REGENERATION**

216 Changes in immune cells with ageing have been well characterised and the observations of
217 elevated systemic inflammation led to the term 'inflamm-ageing' (Franceschi et al., 2000).
218 Human immunity is subdivided into two main areas, often described as *innate* and *adaptive*

219 immunity. Innate immunity describes the primary capacity of the immune system to respond to
220 pathophysiological triggers such as injury or pathogens and is mediated mainly through the
221 myeloid progenitor cells (e.g. neutrophils, macrophages, dendritic cells, natural killer cells, mast
222 cells, eosinophils, basophils) (Plackett et al., 2004). During normal physiological conditions,
223 immune cells circulate within the blood and the lymphatic system, with considerable
224 accumulations in lymphoid organs and most tissues of the body. Peripheral tissues also contain
225 a population of resident immune cells, primarily consisting of macrophages and dendritic cells.
226 However, during pathophysiological conditions supplementary leukocytes rapidly permeate
227 tissues. During muscle regeneration, there can be in excess of 1×10^5 immune cell/ mm^3 of skeletal
228 muscle (Wehling et al., 2001). When activated, these immune cells secrete cytokines and growth
229 factors which regulate the damaged muscle microenvironment (Merly et al., 1999; Warren et al.,
230 2004; Smith et al., 2008).

231

232 **2.1 Innate Immune response to acute damage and repair**

233 The regulation of infiltrating inflammatory cells is a dynamic process which varies depending on
234 the extent of muscle damage and the time required to repair (Paulsen et al., 2012). Minor muscle
235 damage, such as that which occurs after exercise, causes only a modest inflammatory response
236 and may not cause substantial leukocyte cell infiltration, while more severe muscle damage
237 occurring after very intense, unaccustomed exercise with high eccentric loads causes a
238 considerably greater muscle tenderness, immune cell (e.g. neutrophil, macrophage and muscle
239 T reg) infiltration (Fig. 1) of the damaged area and inflammatory responses consistent with
240 rhabdomyolysis (reviewed in (Paulsen et al., 2012)).

241

242 The innate immune response to damage involves infiltration of inflammatory cells, but studies in
243 aged mice have revealed a delayed inflammatory response (Shavlakadze et al., 2010). In healthy
244 muscle, neutrophils show a transient response, infiltrating the extracellular space around the
245 damaged fibres within 2 hours before concentrations decline to negligible levels within 3 or 4
246 days. The mechanisms of neutrophil infiltration remain unclear, but the resulting perpetuation
247 of inflammatory damage is believed to be important for initiating the reparative process (Dumont

248 et al., 2008). Neutrophils release interleukin 1 (IL-1) and interleukin 8 (IL-8) which act as
249 chemoattractants for macrophages, inducing the initial macrophage infiltration to the injury site
250 (Fujishima et al., 1993; Cassatella, 1999). Resident macrophages within the endomysium and
251 perimysium are also involved in phagocytosis and secrete enzymes, growth factors and
252 cytokines/chemokines aiding the recruitment of additional immune cells (Wang et al., 2014).

253
254 Macrophages go through various stages of activation. *Classic activation* of macrophages is
255 denoted as the M1 phenotype, where the increase in numbers and expression of
256 proinflammatory mediators, cytokines and chemokines are observed from 24 hrs and reach peak
257 activation around 2 or 3 days after damage (Rodriguez-Prados et al., 2010; Saclier et al., 2013)
258 (Villalta et al., 2009). The M1 phenotype macrophages originating from the blood as monocytes
259 are distinguishable by their expression of the glycoprotein lymphocyte antigen 6C (Ly6C) as well
260 as receptors for the CX3C chemokine receptor 1 (CX3CR1) and C-C chemokine receptor type 2
261 (CCR2) (Geissmann et al., 2003). The chemokine CCR2 and its ligand CCL2 (or MCP-1) which are
262 mainly produced by monocytes/macrophages coordinate the recruitment of macrophage Ly6C⁺
263 to the site of injury supporting the proinflammatory response. Ly6C⁺ monocytes differentiate into
264 M1 macrophages in tissue and produce proinflammatory cytokines (Jetten et al., 2014). Ly6C⁻
265 cells are recruited to the area by CX3CR1 and CCR2 chemokine receptor signalling and
266 differentiate into M2 macrophages to perform anti-inflammatory and pro-myogenic functions
267 that contribute to the later stages of regeneration (Forbes & Rosenthal, 2014). The M2
268 phenotype is known as *alternative activation* and peaks between 4 and 6 days (see Fig. 1) during
269 the reparative process, where expression of anti-inflammatory mediators, cytokines and
270 chemokines supports the regeneration through satellite cell activation (J. G. Tidball, 2005; Arnold
271 et al., 2007). In cases of severe muscle damage causing fibre necrosis, macrophages can be found
272 infiltrating the intracellular areas of fibres several days post-injury, and elevated macrophage
273 concentrations are evident in muscle tissue up to 3 weeks later (Paulsen et al., 2010).

274

275 **Fig. 1:** Timeline of inflammatory responses and immune cells during regeneration.

276

277 CCR2^{-/-} mice show impaired monocyte recruitment to the site of injury, while neutrophil and
278 other T-cells remain unaffected (Abbadie et al., 2003). The CCR2^{-/-} mice also show impaired
279 muscle regeneration, arrested angiogenesis along with increased fibrosis and excess adipocyte
280 accumulation at the injury site (Martinez et al., 2010). Bone marrow transplants from wild-type
281 mice into CCR2^{-/-} mice recovered the regenerative capacity of skeletal muscle of the CCR2^{-/-} mice.
282 These results show that CCR2, released by proliferating myocytes and resident immune cells,
283 recruits bone marrow derived monocytes (Sun et al., 2009). However, the same results are not
284 observed in studies involving CCL2^{-/-} mice. The CCL2^{-/-} mice have only a mild deficiency in
285 regeneration, which may indicate that alternative chemokine (C-C motif) ligands can bind with
286 the CCR2 receptor and support the recruitment of monocytes and ultimately improve
287 regenerative capacity (Lu, Huang, Ransohoff, et al., 2011).

288

289 **2.2 Regulation of skeletal muscle regeneration via innate immune cell signalling**

290 In response to muscle injury, the innate immune system is activated, to enhance repair damaged
291 tissue by secreting several cytokines (summarised in Fig. 2) (Madaro & Bouche, 2014). The
292 cytokine interleukin 6 (IL-6) is involved in the initial infiltration of monocytes and macrophages
293 during the inflammatory response shortly after muscle damage. Studies involving IL-6^{-/-} mice
294 revealed a significant decrease in the early infiltration of monocytes and macrophages to the
295 injury site, resulting in diminished myofibre mass and more fibrosis of the muscle (Zhang et al.,
296 2013). In the wild-type mice, much of the IL-6 produced soon after injury comes from the early
297 monocyte and macrophage infiltration (Zhang et al., 2013). IL-6 also stimulates macrophage
298 expression of another important molecule, granulocyte colony-stimulating factor (G-CSF), which
299 is involved in normal myoblast proliferation and myofibre differentiation throughout the muscle
300 regeneration process (Zhang et al., 2013; Wright et al., 2015). IL-6^{-/-} mice show slower rates of
301 hypertrophic muscle growth than wild-type animals (Serrano et al., 2008). This study also found
302 that IL-6^{-/-} animals have considerably lower levels of myogenin expression, but MyoD expression
303 was unaffected, which helps to explain why myofibre differentiation was lower in IL-6^{-/-} animals
304 compared with wild-type.

305

306 Supplementary to IL-6 a rapid expression of tumour necrosis factor alpha (TNF α) after injury
307 serves to intensify inflammation in the early stages following muscle damage and is linked to the
308 innate immune response (Warren et al., 2002). TNF α is released by the resident neutrophils,
309 along with interferon gamma (IFN γ) and Interleukin-1 beta (IL-1 β), which can promote monocyte
310 differentiation to M1 phenotype macrophages (Arango Duque & Descoteaux, 2014).
311 Interestingly, as neutrophils and TNF α concentrations peak after 2 days post-injury, the quick
312 tapering of neutrophils (3-4 days) is not paralleled by reductions in TNF α levels, which remain
313 elevated for approximately 14 days after injury (Novak et al., 2014). This indicates that TNF α is
314 not only involved with the early inflammatory process, but potentially has functions throughout
315 muscle regeneration. Together, IL-6 and TNF α can enhance the proliferation of myoblasts,
316 function as chemo-attractants aimed at myoblasts and immune cells, hinder the fusion of
317 myocytes and affect development of stimulated satellite cells to the early phases of
318 differentiation.

319

320 As mentioned earlier macrophages undergo various phases of activation. Specific cytokines such
321 as the ones described (i.e. CCL2, IL-6, and TNF α) are observed to be critically linked with
322 classically activated M1 macrophage infiltration to the site of muscle damage through the initial
323 inflammatory response. However, the differentiation of M2 phenotype macrophages is more
324 complex than that of M1 (Mantovani et al., 2004). The Sub-phenotype M2a macrophages emerge
325 from the exposure to cytokines secreted by adaptive immune responses, including interleukin 4
326 (IL-4) and interleukin 13 (IL-13), which stimulate the complex phases of tissue restoration and
327 injury healing. The arrival of M2b macrophages are believed to begin with the provocation of Toll-
328 like receptor immune complexes, leading to the release of anti-inflammatory chemokines such
329 as IL-10 and the inflammatory cytokines TNF α and IFN γ (J. G. Tidball et al., 2014). TNF α can
330 activate NF- κ B within macrophages, which then induce the production and upregulation of
331 additional proinflammatory mediators, including TNF α , which are then subsequently secreted by
332 the macrophages into the microenvironment of the regenerating muscle. Research using TNF α :
333 $^{-/-}$ mouse models showed a reduction in myogenic differentiation when compared to wild-type
334 mice, this suggests that TNF α signalling within the immuno-muscular microenvironment

335 performs a regulatory role in muscle regeneration. (Chen et al., 2005). Alternatively, *in vitro*
336 models using C2C12 murine myoblasts indicated that elevated TNF α hindered the myoblast
337 capability to exit the cell cycle, indicating that TNF α prolonged myoblast proliferation while
338 inhibiting myogenic differentiation.

339
340 TNF α can also activate NF- κ B within myoblasts, resulting in myoblast proliferation through the
341 up-regulation of cyclin D1 while suppressing differentiation, as well as inhibiting MyoD
342 expression, further suppressing differentiation (Langen et al., 2004). Along with TNF α increasing
343 proliferation and inhibiting differentiation through the NF- κ B signalling pathway, NF- κ B
344 activation in myoblasts promotes the activation of p38 kinase. Animal studies have demonstrated
345 that suppressing p38 leads to reductions in myotube formation along with lower levels of
346 myogenin (Liu et al., 2012). When NF- κ B signalling is activated within myoblasts via stimulation
347 from TNF α secreted within the immuno-muscular microenvironment, an increase of IL-6 is also
348 observed, delivering a supplementary route to enhancing the effects that NF- κ B has on
349 proliferation and increasing its suppression of differentiation. *In vitro* cell culture experiments
350 where mouse myoblasts were treated with IL-6 displayed increases in myoblast proliferation, but
351 not cell fusion (Pelosi et al., 2014). Likewise, *In vitro* cell culture experiments have shown that
352 TNF α increase the migration capacity of myoblasts, demonstrating its role as a chemoattractant
353 (Torrente et al., 2003). Providing further evidence that TNF α production by neutrophils and M1
354 phenotype macrophages following muscle damage promotes muscle regeneration via the
355 attraction of satellite cells to the site of damage. Fig.2 shows the interaction of TNF α and the NF-
356 κ B signalling pathway and its influence on skeletal muscle regeneration.

357
358 The release of IL-10 by M2b macrophages also supports the recruitment of M2c macrophages,
359 which release cytokines that are essential for the cessation of M1 macrophage infiltration and
360 activity (J. G. Tidball et al., 2014). The IL-10 released by both M2b and M2c macrophages
361 stimulates the proliferation of myoblasts needed for muscle growth and regeneration (Deng et
362 al., 2012). Sub-phenotypes of M2b macrophages are observed throughout the repair process
363 since their production of IL-10 is needed to promote anti-inflammatory actions during muscle

364 regeneration (Bosurgi et al., 2012). IL-10^{-/-} mice show impaired transition of macrophages from
365 the M1 to M2 phenotypes, resulting in a corresponding impairment to muscle regeneration. It is
366 interesting to note that IL-10^{-/-} mice are also used as an animal model of early-onset frailty with
367 poor muscle mass and function in older age (Walston et al., 2008). Furthermore, mouse myoblast
368 cell cultures supplemented with IL-10 and M2 macrophages resulted in enhanced myoblast
369 proliferation (Deng et al., 2012). Therefore, IL-10 can mediate the transition of M1 to M2
370 macrophages after muscle damage occurs and encourages the proliferation of myoblasts and
371 maturation of myofibers.

372

373 As myoblasts switch from proliferation to differentiation, a shift from M1 macrophages and
374 proinflammatory cytokines to M2 macrophages and anti-inflammatory cytokines occurs
375 concurrently. This cytokine transference diminishes the proinflammatory response and supports
376 the differentiation of myofibres (Deng et al., 2012), thereby positively influencing the
377 regenerative process (J. G. Tidball, 2005; Arnold et al., 2007). This is in part linked to insulin-like
378 growth factor I (IGF-I), a protein known for its growth-promoting properties and anabolic-
379 inducing effects through the up-regulation of myogenic regulatory factors (MRFs) (Chakravarthy
380 et al., 2000; Mourkioti & Rosenthal, 2005; Xu & Wu, 2000). Importantly, IGF-I is also secreted by
381 M2 macrophages during muscle regeneration (Tonkin et al., 2015; James G. Tidball & Welc, 2015).
382 When observing the infiltration of monocytes and macrophages into the muscle injury site of
383 CCR2^{-/-} mice a considerable reduction of infiltrating cells is observed when compared to the
384 controls. Interestingly, a reduction of circulating IGF-1 is also observed in conjunction with the
385 reduced number of infiltrating immune cells. (Lu, Huang, Saederup, et al., 2011). This fascinating
386 discovery indicates that macrophages provide growth factors that aid in the repair of muscle
387 tissue damage by encouraging IGF-I stimulated satellite cell proliferation.

388

389 Overall, the regeneration of healthy young muscle occurs by rapid recruitment of immune cells
390 to the damaged site in order to orchestrate the regenerative process by removing necrotic
391 cellular debris, coordinating pro/anti-inflammatory events and activating satellite cells through
392 strictly regulated signalling and chemo-attractant molecules (i.e. cytokines, chemokines).

393 Although damaged muscle fibres secrete a number of cytokines, chemokines and growth factors,
394 it is the resident and infiltrating immune cells that are the main producers of these regenerating
395 mediators. Consequently, any alterations to the numbers or types of cytokines, chemokines, or
396 growth factors as a result of age related immune dysfunction has a considerable potential to
397 disrupt the ability of satellite cells in elderly muscle to become activated, migrate to the site of
398 injury, proliferate in adequate quantities and/or differentiate appropriately, resulting in an age
399 linked decline of muscle size and function. Investigations regarding age associated changes to
400 innate immune cell signalling molecules have discovered substantial difference when compared
401 to young counterparts. Specifically, an increase in proinflammatory cytokines (i.e. IL-6, TNF α , IL-
402 1 β) is observed, leading to the chronic inflammatory state often observed in the elderly
403 (Bruunsgaard et al., 2003; Ershler & Keller, 2000; O'Mahony et al., 1998). Increases in
404 proinflammatory cytokines have been identified in the advancement of many geriatric disorders
405 (Franceschi & Campisi, 2014). Thus, it can be appreciated that inflamm-ageing is also having a
406 detrimental effect on the innate immune cells ability to properly coordinate the precisely
407 programmed stages of muscle regeneration, due to their inability to appropriately regulate the
408 signalling molecules circulating within the immuno-muscular microenvironment during skeletal
409 muscle regeneration.

410

411 **Fig. 2:** Innate immune signalling pathways in skeletal muscle regeneration.

412

413 **3. ADAPTIVE IMMUNITY & MUSCLE REGENERATION**

414 Adaptive immunity is observed as a secondary onset response to a pathophysiological incident,
415 which is primarily mediated through lymphoid stem cells such as the T-cells and B-cells (Kim et
416 al., 2007). There has been a remarkable increase in the number of descriptive studies detailing
417 the interactions between innate immune responses and muscle regeneration. However,
418 understanding of the role of the adaptive immune system in muscle regeneration is limited. Just
419 as macrophages and cells involved with innate immunity are detected during acute muscle injury,
420 adaptive immune cells such as T-cells are also present during the regeneration process (Cheng et
421 al., 2008).

422

423

424 **3.1 Adaptive immune response to muscle damage**

425 T-cell infiltration to the site of injury is apparent approximately 3 days after injury and remains
426 elevated for at least 10 days (Cheng et al., 2008). The satellite cells begin to migrate in damaged
427 muscle in the initial 24 hours and begin to proliferate rapidly thereafter. These initial activities
428 are likely regulated via cytokines secreted by innate immune cells (e.g. macrophages). However,
429 adaptive immune responses to damaged muscle via the delayed release of cytokines by T-cells
430 will promote continued satellite cell proliferation. The sustained T-cell presence throughout the
431 regenerative process suggests that T-cells are fundamentally involved with skeletal muscle repair,
432 but the mechanisms of these interactions are not well understood.

433

434 Experiments conducted with T-cell deficient mice resulted in a significant reduction in the early
435 growth and development of muscle (Morrison et al., 2005). Cell culture investigations observing
436 the impact of activated murine splenic T-cell cytokine secretions (secretome) on satellite cell
437 function presented a ~24% increase in the proliferation of satellite cells isolated from young (3
438 months old) mouse muscle, compared to non-secretome treated satellite cell cultures (Dumke &
439 Lees, 2011). Conversely, there was no significant effect on the proliferation of aged (32 months
440 old) mouse muscle satellite cells when exposed to the same T-cell secretome. Furthermore, T-
441 cells signalling (i.e. chemokines) also increased the rate of migration of young satellite cells but
442 not old. However, T-cell secretome significantly reduced the ability of aged satellite cells to
443 differentiate when compared to young satellite cells (Dumke & Lees, 2011). Additionally, recent
444 research employing a mouse model observed that adding the secretome from human T-cells onto
445 a punch-biopsy muscle wound accelerated healing (Mildner et al., 2013).

446 These findings reveal T-cell regulation of muscle repair, as well as the possibility that ageing may
447 diminish T-cell regulated satellite cell function. Further research has explored the impact of T-cell
448 secretome from activated and non-activated T-cells isolated from young (20-25 years old) human
449 blood on immortalized murine satellite cells. The young activated-T-cell secretome enhanced
450 proliferation of the satellite cells and reduced differentiation (Al-Shanti et al., 2014).

451 Demonstrating that regenerating muscle is influenced by, and responds to, a typical ‘young’
452 adaptive immune response. Follow-on work showed that the secretome from young (18-25 years
453 old) activated T-cells enhances both proliferation and migration in immortalized murine satellite
454 cell, however, the secretome from old (78-85 years old) activated T-cells induced premature
455 differentiation similar to control conditions, with no effects on proliferation or migration of the
456 satellite cells (Al-Dabbagh et al., 2015). This outcome implies that proteins secreted by the
457 adaptive immune cells in young people enhance satellite cell proliferation and migration,
458 whereas secreted proteins by the adaptive immune cells of old people attenuates satellite cell
459 proliferation and migration by prematurely stimulating differentiation. These studies indicate
460 that impairments in the ability of satellite cells in elderly people to appropriately proliferate and
461 migrate to the site of muscle injury are related to age-associated T-cell deficiencies, promoting
462 age-related reductions in skeletal muscle size and function.

463

464 Various studies have established that T-cells secrete growth factors and cytokines, some of which
465 can influence satellite cell function (e.g. FGF2, IFN γ , TGF β , TNF α , and IL4) (Blotnick et al., 1994;
466 De Rosa et al., 2004; Levings et al., 2002). The challenge for future studies will be to determine
467 how advanced ageing alters the specific types and concentrations of proteins secreted by old T-
468 cells when compared to young T-cells. This will identify the up- and/or down- regulated immune
469 factors responsible for altering satellite cell function during muscular regeneration in elderly
470 people. Conceivably, these discoveries could lead to the manipulation of immune factors in the
471 immuno-muscular microenvironment of elderly people, possibly replicating a young immuno-
472 muscular microenvironment and overcoming the age associated defects in aged satellite cell
473 function.

474

475 **3.2 Regulation of skeletal muscle regeneration via Regulatory T-cells**

476 Much of the early research beginning to expose the role of adaptive immunity on muscle
477 regeneration has focused on investigating all T-cells as a single component of immunity
478 interacting with satellite cells (Fig. 3). However, there are several different sub-phenotypes of T-
479 cells and distinguishing between them during regeneration may be crucial for identifying which

480 T-cell sub- phenotypes are up and/or down regulating cytokines and growth factors that
481 influence satellite cell function. Attention has been drawn to a specific population of immune
482 response regulatory T-cells (Treg), denoted as the CD4⁺Foxp3⁺ sub-phenotype. Not only are these
483 Treg cells involved with immune response regulation (Josefowicz et al., 2012), they have also
484 been detected at concentrations of $1.05 \pm 0.38 \times 10^4$ cells/g of muscle 28 days after injury.
485 However, alternate T-cell sub-phenotype populations decrease to pre-injury levels of 0.13 ± 0.06
486 $\times 10^4$ cells/g of muscle by the same time point of the repair process (Dalia Burzyn et al., 2013).
487 This finding indicates that Treg cells may be a vital immune cell type influencing muscle
488 regeneration.

489
490 Using mouse models with muscular injury induced via Ctx, it was shown that the Treg cell
491 concentrations increased within the injured muscle as the innate immune cells shifted from a
492 pro- inflammatory to anti-inflammatory phenotype (i.e. M1 to M2) (Dalia Burzyn et al., 2013). It
493 was also discovered that Treg cells found specifically in muscle (mTreg) produce distinctive
494 proteins from their counterparts found in other tissues. These proteins include the anti-
495 inflammatory cytokine IL-10 and the growth factors amphiregulin and platelet-derived growth
496 factor (PDGF), all of which have been shown to influence typical muscle regeneration (Dalia
497 Burzyn et al., 2013; Huey et al., 2008; Yablonkareuveni et al., 1990). Furthermore, experiments
498 where Treg cells were prevented from entering the Ctx injured mouse muscle resulted in innate
499 immune cells failing to switch from pro-inflammatory M1 phenotype to the anti-inflammatory
500 M2 phenotype. Treg ablation from damaged muscle also caused an abnormal inflamed
501 morphology of the regenerating muscle fibres with fibrosis (Castiglioni et al., 2015). Treg-
502 stimulated satellite cells showed sustained proliferation and delayed differentiation (Castiglioni
503 et al., 2015).

504
505 Although evidence has been presented outlining the role Treg cells perform during muscle
506 regeneration, further research is required to fully understand how Treg cells are recruited and
507 expanded within damaged muscle. It is also interesting to consider that Treg cells are able to
508 influence muscle repair via interaction with innate immune cells (i.e. macrophages) as well as

509 activating satellite cells (D. Burzyn et al., 2013). These observations may help to serve as a
510 foundation for future studies looking at the impact ageing has on Treg cells and whether ageing
511 causes a reduction or increase in the number of Treg cells infiltrating the site of muscle damage.
512 These studies may also help to determine if ageing impacts Treg cells' ability to produce the
513 appropriate types and concentrations of cytokines and growth factors needed for normal muscle
514 repair and regeneration.

515

516 **4. CONCLUSIONS**

517 Immune cell infiltration into the site of muscle damage and subsequent release of signalling
518 molecules (i.e. cytokines and growth factors) into the microenvironment regulate muscle repair
519 and regeneration through direct interaction with satellite cells (see Fig. 3). Immune factors
520 released within an aged immuno-muscular microenvironment differ from those of young.
521 Investigating specific populations and sub-phenotypes of both innate and adaptive immune cells,
522 in both young and elderly people, will provide insight into the mechanisms of age-associated
523 muscle wasting. Developing novel therapies to treat sarcopenia by manipulating the aged
524 immuno-muscular microenvironment during regeneration may enhanced muscle size and
525 restore muscle function in the elderly. Current strategies to promote muscle regeneration and
526 maintenance in elderly people are primarily focused on nutrition and physical activity (English &
527 Paddon-Jones, 2010; Moore, 2014). These approaches may alleviate the progression and
528 trajectory of sarcopenia, but only to a relatively minor degree. These therapies are only able to
529 delay the inevitable loss of skeletal muscle mass, function and regenerative capacity associated
530 with progressive ageing. A number of pharmacological strategies to tackle muscle wasting have
531 been proposed, although no treatments are currently in clinical use that block or reverse the loss
532 of muscle in the elderly (D. Glass & Roubenoff, 2010). Therefore, developing a novel approach to
533 prevent sarcopenia is essential and elucidating the role of the immune system in muscle
534 regeneration will help to identify regulatory processes that are candidates for intervention.

535

536 **5. REFERENCES**

537

538 Abbadie, C., Lindia, J. A., Cumiskey, A. M., Peterson, L. B., Mudgett, J. S., Bayne, E. K., DeMartino, J. A.,
539 MacIntyre, D. E. and Forrest, M. J. (2003) 'Impaired neuropathic pain responses in mice lacking the
540 chemokine receptor CCR2.' *Proc Natl Acad Sci U S A*, 100(13), Jun 24, pp. 7947-7952.
541

542 Adams, G. R. (2006) 'Satellite cell proliferation and skeletal muscle hypertrophy.' *Appl Physiol Nutr*
543 *Metab*, 31(6), Dec, pp. 782-790.
544

545 Al-Dabbagh, S., McPhee, J. S., Murgatroyd, C., Butler-Browne, G., Stewart, C. E. and Al-Shanti, N. (2015)
546 'The lymphocyte secretome from young adults enhances skeletal muscle proliferation and migration,
547 but effects are attenuated in the secretome of older adults.' *Physiol Rep*, 3(11), Nov,
548

549 Al-Shanti, N., Durcan, P., Al-Dabbagh, S., Dimchev, G. A. and Stewart, C. E. (2014) 'Activated
550 Lymphocytes Secretome Inhibits Differentiation and Induces Proliferation of C2C12 Myoblasts.' *Cellular*
551 *Physiology and Biochemistry*, 33(1) pp. 117-128.
552

553 Arango Duque, G. and Descoteaux, A. (2014) 'Macrophage cytokines: involvement in immunity and
554 infectious diseases.' *Front Immunol*, 5 p. 491.
555

556 Arnold, L., Henry, A., Poron, F., Baba-Amer, Y., van Rooijen, N., Plonquet, A., Gherardi, R. K. and
557 Chazaud, B. (2007) 'Inflammatory monocytes recruited after skeletal muscle injury switch into
558 antiinflammatory macrophages to support myogenesis.' *J Exp Med*, 204(5), May 14, pp. 1057-1069.
559

560 Barberi, L., Scicchitano, B. M., De Rossi, M., Bigot, A., Duguez, S., Wielgosik, A., Stewart, C., McPhee, J.,
561 Conte, M., Narici, M., Franceschi, C., Mouly, V., Butler-Browne, G. and Musarò, A. (2013) 'Age-
562 dependent alteration in muscle regeneration: the critical role of tissue niche.' *Biogerontology*, 14(3), Jun,
563 pp. 273-292.
564

565 Bentzinger, C. F., Wang, Y. X. and Rudnicki, M. A. (2012) 'Building muscle: molecular regulation of
566 myogenesis.' *Cold Spring Harb Perspect Biol*, 4(2), Feb,
567

568 Blotnick, S., Peoples, G. E., Freeman, M. R., Eberlein, T. J. and Klagsbrun, M. (1994) 'T lymphocytes
569 synthesize and export heparin-binding epidermal growth factor-like growth factor and basic fibroblast
570 growth factor, mitogens for vascular cells and fibroblasts: differential production and release by CD4+
571 and CD8+ T cells.' *Proc Natl Acad Sci U S A*, 91(8), Apr 12, pp. 2890-2894.
572

573 Bosurgi, L., Corna, G., Vezzoli, M., Touvier, T., Cossu, G., Manfredi, A. A., Brunelli, S. and Rovere-Querini,
574 P. (2012) 'Transplanted Mesoangioblasts Require Macrophage IL-10 for Survival in a Mouse Model of
575 Muscle Injury.' *Journal of Immunology*, 188(12), Jun, pp. 6267-6277.
576

577 Brack, A. S. and Rando, T. A. (2007) 'Intrinsic changes and extrinsic influences of myogenic stem cell
578 function during aging.' *Stem Cell Rev*, 3(3), Fall, pp. 226-237.
579

580 Brack, A. S. and Munoz-Canoves, P. (2015) 'The ins and outs of muscle stem cell aging.' *Skelet Muscle*, 6
581 p. 1.
582

583 Brun, Caroline E. and Rudnicki, Michael A. (2015) 'GDF11 and the Mythical Fountain of Youth.' *Cell*
584 *Metabolism*, 22(1) pp. 54-56.
585

586 Bruunsgaard, H., Andersen-Ranberg, K., Hjelmberg, J., Pedersen, B. K. and Jeune, B. (2003) 'Elevated
587 levels of tumor necrosis factor alpha and mortality in centenarians.' *Am J Med*, 115(4), Sep, pp. 278-283.
588

589 Burzyn, D., Benoist, C. and Mathis, D. (2013) 'Regulatory T cells in nonlymphoid tissues.' *Nat Immunol*,
590 14(10), Oct, pp. 1007-1013.
591

592 Burzyn, D., Kuswanto, W., Kolodin, D., Shadrach, Jennifer L., Cerletti, M., Jang, Y., Sefik, E., Tan, Tze G.,
593 Wagers, Amy J., Benoist, C. and Mathis, D. (2013) 'A Special Population of Regulatory T Cells Potentiates
594 Muscle Repair.' *Cell*, 155(6), 12/5/, pp. 1282-1295.
595

596 Carlson, B. M. and Faulkner, J. A. (1989) 'Muscle transplantation between young and old rats: age of
597 host determines recovery.' *Am J Physiol*, 256(6 Pt 1), Jun, pp. C1262-1266.
598

599 Carlson, M. E., Hsu, M. and Conboy, I. M. (2008) 'Imbalance between pSmad3 and Notch induces CDK
600 inhibitors in old muscle stem cells.' *Nature*, 454(7203), 07/24/print, pp. 528-532.
601

602 Carlson, M. E., Suetta, C., Conboy, M. J., Aagaard, P., Mackey, A., Kjaer, M. and Conboy, I. (2009)
603 'Molecular aging and rejuvenation of human muscle stem cells.' *EMBO Mol Med*, 1(8-9), Nov, pp. 381-
604 391.
605

606 Cassatella, M. A. (1999) 'Neutrophil-derived proteins: selling cytokines by the pound.' *Adv Immunol*, 73
607 pp. 369-509.
608

609 Castiglioni, A., Corna, G., Rigamonti, E., Basso, V., Vezzoli, M., Monno, A., Almada, A. E., Mondino, A.,
610 Wagers, A. J., Manfredi, A. A. and Rovere-Querini, P. (2015) 'FOXP3+ T Cells Recruited to Sites of Sterile
611 Skeletal Muscle Injury Regulate the Fate of Satellite Cells and Guide Effective Tissue Regeneration.' *PLoS*
612 *One*, 10(6) p. e0128094.
613

614 Cerletti, M., Shadrach, J. L., Jurga, S., Sherwood, R. and Wagers, A. J. (2008) 'Regulation and function of
615 skeletal muscle stem cells.' *Cold Spring Harb Symp Quant Biol*, 73 pp. 317-322.
616

617 Chakkalakal, J. V., Jones, K. M., Basson, M. A. and Brack, A. S. (2012) 'The aged niche disrupts muscle
618 stem cell quiescence.' *Nature*, 490(7420), Oct 18, pp. 355-360.
619

620 Chakravarthy, M. V., Abraha, T. W., Schwartz, R. J., Fiorotto, M. L. and Booth, F. W. (2000) 'Insulin-like
621 growth factor-I extends in vitro replicative life span of skeletal muscle satellite cells by enhancing G1/S
622 cell cycle progression via the activation of phosphatidylinositol 3'-kinase/Akt signaling pathway.' *J Biol*
623 *Chem*, 275(46), Nov 17, pp. 35942-35952.
624

625 Chen, S. E., Gerken, E., Zhang, Y., Zhan, M., Mohan, R. K., Li, A. S., Reid, M. B. and Li, Y. P. (2005) 'Role of
626 TNF- α signaling in regeneration of cardiotoxin-injured muscle.' *Am J Physiol Cell Physiol*, 289(5),
627 Nov, pp. C1179-1187.
628

629 Cheng, M., Nguyen, M. H., Fantuzzi, G. and Koh, T. J. (2008) 'Endogenous interferon-gamma is required
630 for efficient skeletal muscle regeneration.' *Am J Physiol Cell Physiol*, 294(5), May, pp. C1183-1191.
631

632 Conboy, I. M., Conboy, M. J., Wagers, A. J., Girma, E. R., Weissman, I. L. and Rando, T. A. (2005)
633 'Rejuvenation of aged progenitor cells by exposure to a young systemic environment.' *Nature*,
634 433(7027), 02/17/print, pp. 760-764.
635
636 Cosgrove, B. D., Gilbert, P. M., Porpiglia, E., Mourkioti, F., Lee, S. P., Corbel, S. Y., Llewellyn, M. E., Delp,
637 S. L. and Blau, H. M. (2014) 'Rejuvenation of the muscle stem cell population restores strength to injured
638 aged muscles.' *Nat Med*, 20(3), 03//print, pp. 255-264.
639
640 Cramer, R. M., Aagaard, P., Qvortrup, K., Langberg, H., Olesen, J. and Kjaer, M. (2007) 'Myofibre damage
641 in human skeletal muscle: effects of electrical stimulation versus voluntary contraction.' *J Physiol*, 583(Pt
642 1), Aug 15, pp. 365-380.
643
644 Cruz-Jentoft, A. J., Baeyens, J. P., Bauer, J. M., Boirie, Y., Cederholm, T., Landi, F., Martin, F. C., Michel, J.
645 P., Rolland, Y., Schneider, S. M., Topinkova, E., Vandewoude, M. and Zamboni, M. (2010) 'Sarcopenia:
646 European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia
647 in Older People.' *Age Ageing*, 39(4), Jul, pp. 412-423.
648
649 De Rosa, S. C., Lu, F. X., Yu, J., Perfetto, S. P., Falloon, J., Moser, S., Evans, T. G., Koup, R., Miller, C. J. and
650 Roederer, M. (2004) 'Vaccination in humans generates broad T cell cytokine responses.' *J Immunol*,
651 173(9), Nov 1, pp. 5372-5380.
652
653 Degens, H. (2010) 'The role of systemic inflammation in age-related muscle weakness and wasting.'
654 *Scand J Med Sci Sports*, 20(1), Feb, pp. 28-38.
655
656 Deng, B., Wehling-Henricks, M., Villalta, S. A., Wang, Y. and Tidball, J. G. (2012) 'IL-10 Triggers Changes in
657 Macrophage Phenotype That Promote Muscle Growth and Regeneration.' *Journal of Immunology*,
658 189(7), Oct, pp. 3669-3680.
659
660 Dumke, B. R. and Lees, S. J. (2011) 'Age-related impairment of T cell-induced skeletal muscle precursor
661 cell function.' *Am J Physiol Cell Physiol*, 300(6), Jun, pp. C1226-1233.
662
663 Dumont, N., Bouchard, P. and Frenette, J. (2008) 'Neutrophil-induced skeletal muscle damage: a
664 calculated and controlled response following hindlimb unloading and reloading.' *American Journal of
665 Physiology-Regulatory Integrative and Comparative Physiology*, 295(6), Dec, pp. R1831-R1838.
666
667 Egan, B. and Zierath, J. R. (2013) 'Exercise metabolism and the molecular regulation of skeletal muscle
668 adaptation.' *Cell Metab*, 17(2), Feb 5, pp. 162-184.
669
670 Elabd, C., Cousin, W., Upadhyayula, P., Chen, R. Y., Chooljian, M. S., Li, J., Kung, S., Jiang, K. P. and
671 Conboy, I. M. (2014) 'Oxytocin is an age-specific circulating hormone that is necessary for muscle
672 maintenance and regeneration.' *Nat Commun*, 5, 06/10/online,
673
674 English, K. L. and Paddon-Jones, D. (2010) 'Protecting muscle mass and function in older adults during
675 bed rest.' *Curr Opin Clin Nutr Metab Care*, 13(1), Jan, pp. 34-39.
676
677 Ershler, W. B. and Keller, E. T. (2000) 'Age-associated increased interleukin-6 gene expression, late-life
678 diseases, and frailty.' *Annu Rev Med*, 51 pp. 245-270.
679

680 Farup, J., Madaro, L., Puri, P. L. and Mikkelsen, U. R. (2015) 'Interactions between muscle stem cells,
681 mesenchymal-derived cells and immune cells in muscle homeostasis, regeneration and disease.' *Cell*
682 *Death Dis*, 6 p. e1830.

683

684 Forbes, S. J. and Rosenthal, N. (2014) 'Preparing the ground for tissue regeneration: from mechanism to
685 therapy.' *Nat Med*, 20(8), Aug, pp. 857-869.

686

687 Franceschi, C. and Campisi, J. (2014) 'Chronic inflammation (inflammaging) and its potential contribution
688 to age-associated diseases.' *J Gerontol A Biol Sci Med Sci*, 69 Suppl 1, Jun, pp. S4-9.

689

690 Franceschi, C., Bonafe, M., Valensin, S., Olivieri, F., De Luca, M., Ottaviani, E. and De Benedictis, G.
691 (2000) 'Inflamm-aging. An evolutionary perspective on immunosenescence.' *Ann N Y Acad Sci*, 908, Jun,
692 pp. 244-254.

693

694 Fried, L. P., Tangen, C. M., Walston, J., Newman, A. B., Hirsch, C., Gottdiener, J., Seeman, T., Tracy, R.,
695 Kop, W. J., Burke, G. and McBurnie, M. A. (2001) 'Frailty in older adults: evidence for a phenotype.' *J*
696 *Gerontol A Biol Sci Med Sci*, 56(3), Mar, pp. M146-156.

697

698 Fry, C. S., Lee, J. D., Mula, J., Kirby, T. J., Jackson, J. R., Liu, F., Yang, L., Mendias, C. L., Dupont-
699 Versteegden, E. E., McCarthy, J. J. and Peterson, C. A. (2015) 'Inducible depletion of satellite cells in
700 adult, sedentary mice impairs muscle regenerative capacity without affecting sarcopenia.' *Nat Med*,
701 21(1), 01//print, pp. 76-80.

702

703 Fujishima, S., Hoffman, A. R., Vu, T., Kim, K. J., Zheng, H., Daniel, D., Kim, Y., Wallace, E. F., Larrick, J. W.
704 and Raffin, T. A. (1993) 'Regulation of neutrophil interleukin 8 gene expression and protein secretion by
705 LPS, TNF-alpha, and IL-1 beta.' *J Cell Physiol*, 154(3), Mar, pp. 478-485.

706

707 Geissmann, F., Jung, S. and Littman, D. R. (2003) 'Blood monocytes consist of two principal subsets with
708 distinct migratory properties.' *Immunity*, 19(1), Jul, pp. 71-82.

709

710 Glass, D. and Roubenoff, R. (2010) 'Recent advances in the biology and therapy of muscle wasting.' *Ann*
711 *N Y Acad Sci*, 1211, Nov, pp. 25-36.

712

713 Glass, D. J. (2003) 'Molecular mechanisms modulating muscle mass.' *Trends Mol Med*, 9(8), Aug, pp. 344-
714 350.

715

716 Huey, K. A., Meador, B. M., Krzyzstton, C. P. and Johnson, R. W. (2008) 'Effects of IL-10 and age on IL-6, IL-
717 1 beta, and TNF alpha responses to an acute inflammatory insult in mouse skeletal muscle.' *Faseb*
718 *Journal*, 22, Apr, p. 1.

719

720 Janssen, I. (2011) 'The Epidemiology of Sarcopenia.' *Clinics in Geriatric Medicine*, 27(3), Aug, pp. 355-+.

721

722 Janssen, I., Heymsfield, S. B. and Ross, R. (2002) 'Low relative skeletal muscle mass (sarcopenia) in older
723 persons is associated with functional impairment and physical disability.' *Journal of the American*
724 *Geriatrics Society*, 50(5), May, pp. 889-896.

725

726 Jetten, N., Verbruggen, S., Gijbels, M. J., Post, M. J., De Winther, M. P. J. and Donners, M. (2014) 'Anti-
727 inflammatory M2, but not pro-inflammatory M1 macrophages promote angiogenesis in vivo.'
728 *Angiogenesis*, 17(1), Jan, pp. 109-118.
729
730 Joannis, S., Gillen, J. B., Bellamy, L. M., McKay, B. R., Tarnopolsky, M. A., Gibala, M. J. and Parise, G.
731 (2013) 'Evidence for the contribution of muscle stem cells to nonhypertrophic skeletal muscle
732 remodeling in humans.' *Faseb j*, 27(11), Nov, pp. 4596-4605.
733
734 Josefowicz, S. Z., Lu, L. F. and Rudensky, A. Y. (2012) 'Regulatory T cells: mechanisms of differentiation
735 and function.' *Annu Rev Immunol*, 30 pp. 531-564.
736
737 Kim, K. D., Zhao, J., Auh, S., Yang, X. M., Du, P. S., Tang, H. and Fu, Y. X. (2007) 'Adaptive immune cells
738 temper initial innate responses.' *Nature Medicine*, 13(10), Oct, pp. 1248-1252.
739
740 Kuang, S., Kuroda, K., Le Grand, F. and Rudnicki, M. A. (2007) 'Asymmetric self-renewal and commitment
741 of satellite stem cells in muscle.' *Cell*, 129(5), Jun 1, pp. 999-1010.
742
743 Langen, R. C., Van Der Velden, J. L., Schols, A. M., Kelders, M. C., Wouters, E. F. and Janssen-Heininger, Y.
744 M. (2004) 'Tumor necrosis factor-alpha inhibits myogenic differentiation through MyoD protein
745 destabilization.' *Faseb j*, 18(2), Feb, pp. 227-237.
746
747 Lepper, C., Partridge, T. A. and Fan, C. M. (2011) 'An absolute requirement for Pax7-positive satellite
748 cells in acute injury-induced skeletal muscle regeneration.' *Development*, 138(17), Sep, pp. 3639-3646.
749
750 Leto, D. and Saltiel, A. R. (2012) 'Regulation of glucose transport by insulin: traffic control of GLUT4.' *Nat*
751 *Rev Mol Cell Biol*, 13(6), Jun, pp. 383-396.
752
753 Levings, M. K., Bacchetta, R., Schulz, U. and Roncarolo, M. G. (2002) 'The role of IL-10 and TGF-beta in
754 the differentiation and effector function of T regulatory cells.' *Int Arch Allergy Immunol*, 129(4), Dec, pp.
755 263-276.
756
757 Lexell, J. (1995) 'HUMAN AGING, MUSCLE MASS, AND FIBER-TYPE COMPOSITION.' *Journals of*
758 *Gerontology Series a-Biological Sciences and Medical Sciences*, 50, Nov, pp. 11-16.
759
760 Lexell, J., Taylor, C. C. and Sjostrom, M. (1988) 'WHAT IS THE CAUSE OF THE AGING ATROPHY - TOTAL
761 NUMBER, SIZE AND PROPORTION OF DIFFERENT FIBER TYPES STUDIED IN WHOLE VASTUS LATERALIS
762 MUSCLE FROM 15-YEAR-OLD TO 83-YEAR-OLD MEN.' *Journal of the Neurological Sciences*, 84(2-3), Apr,
763 pp. 275-294.
764
765 Liu, Q. C., Zha, X. H., Faralli, H., Yin, H., Louis-Jeune, C., Perdiguero, E., Pranckeviciene, E., Munoz-
766 Canoves, P., Rudnicki, M. A., Brand, M., Perez-Iratxeta, C. and Dilworth, F. J. (2012) 'Comparative
767 expression profiling identifies differential roles for Myogenin and p38 alpha MAPK signaling in
768 myogenesis.' *Journal of Molecular Cell Biology*, 4(6), Dec, pp. 386-397.
769
770 Lu, H., Huang, D., Ransohoff, R. M. and Zhou, L. (2011) 'Acute skeletal muscle injury: CCL2 expression by
771 both monocytes and injured muscle is required for repair.' *Faseb j*, 25(10), Oct, pp. 3344-3355.
772

773 Lu, H., Huang, D., Saederup, N., Charo, I. F., Ransohoff, R. M. and Zhou, L. (2011) 'Macrophages recruited
774 via CCR2 produce insulin-like growth factor-1 to repair acute skeletal muscle injury.' *Faseb j*, 25(1), Jan,
775 pp. 358-369.
776
777 Mackey, A. L., Bojsen-Moller, J., Qvortrup, K., Langberg, H., Suetta, C., Kalliokoski, K. K., Kjaer, M. and
778 Magnusson, S. P. (2008) 'Evidence of skeletal muscle damage following electrically stimulated isometric
779 muscle contractions in humans.' *J Appl Physiol* (1985), 105(5), Nov, pp. 1620-1627.
780
781 Madaro, L. and Bouche, M. (2014) 'From innate to adaptive immune response in muscular dystrophies
782 and skeletal muscle regeneration: the role of lymphocytes.' *Biomed Res Int*, 2014 p. 438675.
783
784 Maden-Wilkinson, T. M., McPhee, J. S., Rittweger, J., Jones, D. A. and Degens, H. (2014) 'Thigh muscle
785 volume in relation to age, sex and femur volume.' *Age (Dordr)*, 36(1), Feb, pp. 383-393.
786
787 Mantovani, A., Sica, A., Sozzani, S., Allavena, P., Vecchi, A. and Locati, M. (2004) 'The chemokine system
788 in diverse forms of macrophage activation and polarization.' *Trends Immunol*, 25(12), Dec, pp. 677-686.
789
790 Martinez, C. O., McHale, M. J., Wells, J. T., Ochoa, O., Michalek, J. E., McManus, L. M. and Shireman, P. K.
791 (2010) 'Regulation of skeletal muscle regeneration by CCR2-activating chemokines is directly related to
792 macrophage recruitment.' *Am J Physiol Regul Integr Comp Physiol*, 299(3), Sep, pp. R832-842.
793
794 McCarthy, J. J., Mula, J., Miyazaki, M., Erfani, R., Garrison, K., Farooqui, A. B., Srikuea, R., Lawson, B. A.,
795 Grimes, B., Keller, C., Van Zant, G., Campbell, K. S., Esser, K. A., Dupont-Versteegden, E. E. and Peterson,
796 C. A. (2011) 'Effective fiber hypertrophy in satellite cell-depleted skeletal muscle.' *Development*, 138(17),
797 Sep, pp. 3657-3666.
798
799 Merly, F., Lescaudron, L., Rouaud, T., Crossin, F. and Gardahaut, M. F. (1999) 'Macrophages enhance
800 muscle satellite cell proliferation and delay their differentiation.' *Muscle Nerve*, 22(6), Jun, pp. 724-732.
801
802 Mildner, M., Hacker, S., Haider, T., Gschwandtner, M., Werba, G., Barresi, C., Zimmermann, M., Golabi,
803 B., Tschachler, E. and Ankersmit, H. J. (2013) 'Secretome of peripheral blood mononuclear cells
804 enhances wound healing.' *PLoS One*, 8(3) p. e60103.
805
806 Moore, D. R. (2014) 'Keeping Older Muscle "Young" through Dietary Protein and Physical Activity12.' *Adv*
807 *Nutr*, 5(5), Sep, pp. 599s-607s.
808
809 Morley, J. E., Baumgartner, R. N., Roubenoff, R., Mayer, J. and Nair, K. S. (2001) 'Sarcopenia.' *Journal of*
810 *Laboratory and Clinical Medicine*, 137(4), Apr, pp. 231-243.
811
812 Morrison, J., Partridge, T. and Bou-Gharios, G. (2005) 'Nude mutation influences limb skeletal muscle
813 development.' *Matrix Biol*, 23(8), Jan, pp. 535-542.
814
815 Mourkioti, F. and Rosenthal, N. (2005) 'IGF-1, inflammation and stem cells: interactions during muscle
816 regeneration.' *Trends Immunol*, 26(10), Oct, pp. 535-542.
817
818 Naito, A. T., Sumida, T., Nomura, S., Liu, M. L., Higo, T., Nakagawa, A., Okada, K., Sakai, T., Hashimoto, A.,
819 Hara, Y., Shimizu, I., Zhu, W., Toko, H., Katada, A., Akazawa, H., Oka, T., Lee, J. K., Minamino, T., Nagai, T.,
820 Walsh, K., Kikuchi, A., Matsumoto, M., Botto, M., Shiojima, I. and Komuro, I. (2012) 'Complement C1q

821 activates canonical Wnt signaling and promotes aging-related phenotypes.' *Cell*, 149(6), Jun 8, pp. 1298-
822 1313.

823

824 Narici, M. V. and Maffulli, N. (2010) 'Sarcopenia: characteristics, mechanisms and functional
825 significance.' *Br Med Bull*, 95 pp. 139-159.

826

827 Nosaka, K., Aldayel, A., Jubeau, M. and Chen, T. C. (2011) 'Muscle damage induced by electrical
828 stimulation.' *Eur J Appl Physiol*, 111(10), Oct, pp. 2427-2437.

829

830 Novak, M. L., Weinheimer-Haus, E. M. and Koh, T. J. (2014) 'Macrophage Activation and Skeletal Muscle
831 Healing Following Traumatic Injury.' *J Pathol*, 232(3), Feb, pp. 344-355.

832

833 O'Mahony, L., Holland, J., Jackson, J., Feighery, C., Hennessy, T. P. and Mealy, K. (1998) 'Quantitative
834 intracellular cytokine measurement: age-related changes in proinflammatory cytokine production.' *Clin
835 Exp Immunol*, 113(2), Aug, pp. 213-219.

836

837 Paliwal, P., Pishesha, N., Wijaya, D. and Conboy, I. M. (2012) 'Age dependent increase in the levels of
838 osteopontin inhibits skeletal muscle regeneration.' *Aging (Albany NY)*, 4(8), Aug, pp. 553-566.

839

840 Paulsen, G., Mikkelsen, U. R., Raastad, T. and Peake, J. M. (2012) 'Leucocytes, cytokines and satellite
841 cells: what role do they play in muscle damage and regeneration following eccentric exercise?' *Exerc
842 Immunol Rev*, 18 pp. 42-97.

843

844 Paulsen, G., Cramer, R., Benestad, H. B., Fjeld, J. G., Morkrid, L., Hallen, J. and Raastad, T. (2010) 'Time
845 course of leukocyte accumulation in human muscle after eccentric exercise.' *Med Sci Sports Exerc*, 42(1),
846 Jan, pp. 75-85.

847

848 Peake, J., Della Gatta, P. and Cameron-Smith, D. (2010) 'Aging and its effects on inflammation in skeletal
849 muscle at rest and following exercise-induced muscle injury.' *Am J Physiol Regul Integr Comp Physiol*,
850 298(6), Jun, pp. R1485-1495.

851

852 Pedersen, B. K. (2011) 'Muscles and their myokines.' *J Exp Biol*, 214(Pt 2), Jan 15, pp. 337-346.

853

854 Pelosi, M., De Rossi, M., Barberi, L., Musar, #xf2 and , A. (2014) 'IL-6 Impairs Myogenic Differentiation by
855 Downmodulation of p90RSK/eEF2 and mTOR/p70S6K Axes, without Affecting AKT Activity.' *BioMed
856 Research International*, 2014 p. 12.

857

858 Piasecki, M., Ireland, A., Jones, D. A. and McPhee, J. S. (2015) 'Age-dependent motor unit remodelling in
859 human limb muscles.' *Biogerontology*, Dec 14,

860

861 Piasecki, M., Ireland, A., Stashuk, D., Hamilton-Wright, A., Jones, D. A. and McPhee, J. S. (2015) 'Age-
862 related neuromuscular changes affecting human vastus lateralis.' *The Journal of Physiology*, pp. n/a-n/a.

863

864 Plackett, T. P., Boehmer, E. D., Faunce, D. E. and Kovacs, E. J. (2004) 'Aging and innate immune cells.'
865 *Journal of Leukocyte Biology*, 76(2), Aug, pp. 291-299.

866

867 Relaix, F. and Zammit, P. S. (2012) 'Satellite cells are essential for skeletal muscle regeneration: the cell
868 on the edge returns centre stage.' *Development*, 139(16), Aug, pp. 2845-2856.

869
870 Rodriguez-Prados, J. C., Traves, P. G., Cuenca, J., Rico, D., Aragonés, J., Martín-Sanz, P., Cascante, M. and
871 Bosca, L. (2010) 'Substrate Fate in Activated Macrophages: A Comparison between Innate, Classic, and
872 Alternative Activation.' *Journal of Immunology*, 185(1), Jul, pp. 605-614.
873
874 Saclier, M., Cuvellier, S., Magnan, M., Mounier, R. and Chazaud, B. (2013) 'Monocyte/macrophage
875 interactions with myogenic precursor cells during skeletal muscle regeneration.' *Febs Journal*, 280(17),
876 Sep, pp. 4118-4130.
877
878 Serrano, A. L., Baeza-Raja, B., Perdiguero, E., Jardí, M. and Muñoz-Canoves, P. (2008) 'Interleukin-6 is an
879 essential regulator of satellite cell-mediated skeletal muscle hypertrophy.' *Cell Metab*, 7(1), Jan, pp. 33-
880 44.
881
882 Shavlakadze, T., McGeachie, J. and Grounds, M. D. (2010) 'Delayed but excellent myogenic stem cell
883 response of regenerating geriatric skeletal muscles in mice.' *Biogerontology*, 11(3), Jun, pp. 363-376.
884
885 Siegel, A. L., Kuhlmann, P. K. and Cornelison, D. D. (2011) 'Muscle satellite cell proliferation and
886 association: new insights from myofiber time-lapse imaging.' *Skelet Muscle*, 1(1) p. 7.
887
888 Siegel, A. L., Atchison, K., Fisher, K. E., Davis, G. E. and Cornelison, D. D. (2009) '3D timelapse analysis of
889 muscle satellite cell motility.' *Stem Cells*, 27(10), Oct, pp. 2527-2538.
890
891 Sinha, M., Jang, Y. C., Oh, J., Khong, D., Wu, E. Y., Manohar, R., Miller, C., Regalado, S. G., Loffredo, F. S.,
892 Pancoast, J. R., Hirshman, M. F., Lebowitz, J., Shadrach, J. L., Cerletti, M., Kim, M. J., Serwold, T.,
893 Goodyear, L. J., Rosner, B., Lee, R. T. and Wagers, A. J. (2014) 'Restoring systemic GDF11 levels reverses
894 age-related dysfunction in mouse skeletal muscle.' *Science*, 344(6184), May 9, pp. 649-652.
895
896 Smith, C., Kruger, M. J., Smith, R. M. and Myburgh, K. H. (2008) 'The inflammatory response to skeletal
897 muscle injury: illuminating complexities.' *Sports Med*, 38(11) pp. 947-969.
898
899 Smythe, G. M., Shavlakadze, T., Roberts, P., Davies, M. J., McGeachie, J. K. and Grounds, M. D. (2008)
900 'Age influences the early events of skeletal muscle regeneration: studies of whole muscle grafts
901 transplanted between young (8 weeks) and old (13-21 months) mice.' *Exp Gerontol*, 43(6), Jun, pp. 550-
902 562.
903
904 Sousa-Victor, P., Gutarra, S., Garcia-Prat, L., Rodriguez-Ubreva, J., Ortet, L., Ruiz-Bonilla, V., Jardí, M.,
905 Ballestar, E., Gonzalez, S., Serrano, A. L., Perdiguero, E. and Muñoz-Canoves, P. (2014) 'Geriatric muscle
906 stem cells switch reversible quiescence into senescence.' *Nature*, 506(7488), Feb 20, pp. 316-321.
907
908 Steensberg, A., Keller, C., Starkie, R. L., Osada, T., Febbraio, M. A. and Pedersen, B. K. (2002) 'IL-6 and
909 TNF-alpha expression in, and release from, contracting human skeletal muscle.' *Am J Physiol Endocrinol*
910 *Metab*, 283(6), Dec, pp. E1272-1278.
911
912 Sun, D., Martínez, C. O., Ochoa, O., Ruiz-Willhite, L., Bonilla, J. R., Centonze, V. E., Waite, L. L., Michalek,
913 J. E., McManus, L. M. and Shireman, P. K. (2009) 'Bone marrow-derived cell regulation of skeletal muscle
914 regeneration.' *Faseb j*, 23(2), Feb, pp. 382-395.
915

916 Tedesco, F. S., Dellavalle, A., Diaz-Manera, J., Messina, G. and Cossu, G. (2010) 'Repairing skeletal
917 muscle: regenerative potential of skeletal muscle stem cells.' *J Clin Invest*, 120(1), Jan, pp. 11-19.
918

919 Tidball, J. G. (2005) 'Inflammatory processes in muscle injury and repair.' *Am J Physiol Regul Integr Comp*
920 *Physiol*, 288(2), Feb, pp. R345-353.
921

922 Tidball, J. G. and Welc, S. S. (2015) 'Macrophage-Derived IGF-1 Is a Potent Coordinator of Myogenesis
923 and Inflammation in Regenerating Muscle.' *Mol Ther*, 23(7), 07//print, pp. 1134-1135.
924

925 Tidball, J. G., Dorshkind, K. and Wehling-Henricks, M. (2014) 'Shared signaling systems in myeloid cell-
926 mediated muscle regeneration.' *Development*, 141(6), Mar, pp. 1184-1196.
927

928 Tonkin, J., Temmerman, L., Sampson, R. D., Gallego-Colon, E., Barberi, L., Bilbao, D., Schneider, M. D.,
929 Musaro, A. and Rosenthal, N. (2015) 'Monocyte/Macrophage-derived IGF-1 Orchestrates Murine
930 Skeletal Muscle Regeneration and Modulates Autocrine Polarization.' *Mol Ther*, 23(7), Jul, pp. 1189-
931 1200.
932

933 Torrente, Y., El Fahime, E., Caron, N. J., Del Bo, R., Belicchi, M., Pisati, F., Tremblay, J. P. and Bresolin, N.
934 (2003) 'Tumor necrosis factor-alpha (TNF-alpha) stimulates chemotactic response in mouse myogenic
935 cells.' *Cell Transplant*, 12(1) pp. 91-100.
936

937 Uezumi, A., Ikemoto-Uezumi, M. and Tsuchida, K. (2014) 'Roles of nonmyogenic mesenchymal
938 progenitors in pathogenesis and regeneration of skeletal muscle.' *Front Physiol*, 5
939

940 Verdijk, L. B., Gleeson, B. G., Jonkers, R. A., Meijer, K., Savelberg, H. H., Dendale, P. and van Loon, L. J.
941 (2009) 'Skeletal muscle hypertrophy following resistance training is accompanied by a fiber type-specific
942 increase in satellite cell content in elderly men.' *J Gerontol A Biol Sci Med Sci*, 64(3), Mar, pp. 332-339.
943

944 Villalta, S. A., Nguyen, H. X., Deng, B., Gotoh, T. and Tidball, J. G. (2009) 'Shifts in macrophage
945 phenotypes and macrophage competition for arginine metabolism affect the severity of muscle
946 pathology in muscular dystrophy.' *Hum Mol Genet*, 18(3), Feb 1, pp. 482-496.
947

948 Walston, J., Fedarko, N., Yang, H., Leng, S., Beamer, B., Espinoza, S., Lipton, A., Zheng, H. and Becker, K.
949 (2008) 'The physical and biological characterization of a frail mouse model.' *J Gerontol A Biol Sci Med Sci*,
950 63(4), Apr, pp. 391-398.
951

952 Wang, H., Melton, D. W., Porter, L., Sarwar, Z. U., McManus, L. M. and Shireman, P. K. (2014) 'Altered
953 Macrophage Phenotype Transition Impairs Skeletal Muscle Regeneration.' *The American journal of*
954 *pathology*, 184(4) pp. 1167-1184.
955

956 Warren, G. L., Hulderman, T., Jensen, N., McKinstry, M., Mishra, M., Luster, M. I. and Simeonova, P. P.
957 (2002) 'Physiological role of tumor necrosis factor alpha in traumatic muscle injury.' *Faseb j*, 16(12), Oct,
958 pp. 1630-1632.
959

960 Warren, G. L., O'Farrell, L., Summan, M., Hulderman, T., Mishra, D., Luster, M. I., Kuziel, W. A. and
961 Simeonova, P. P. (2004) 'Role of CC chemokines in skeletal muscle functional restoration after injury.'
962 *Am J Physiol Cell Physiol*, 286(5), May, pp. C1031-1036.
963

964 Wehling, M., Spencer, M. J. and Tidball, J. G. (2001) 'A nitric oxide synthase transgene ameliorates
965 muscular dystrophy in mdx mice.' *J Cell Biol*, 155(1), Oct 1, pp. 123-131.
966

967 Wright, C. R., Brown, E. L., Della Gatta, P. A., Fatouros, I. G., Karagounis, L. G., Terzis, G., Mastorakos, G.,
968 Michailidis, Y., Mandalidis, D., Spengos, K., Chatzinikolaou, A., Methenitis, S., Draganidis, D., Jamurtas, A.
969 Z. and Russell, A. P. (2015) 'Regulation of Granulocyte Colony-Stimulating Factor and Its Receptor in
970 Skeletal Muscle is Dependent Upon the Type of Inflammatory Stimulus.' *J Interferon Cytokine Res*, 35(9),
971 Sep, pp. 710-719.
972

973 Xu, Q. and Wu, Z. (2000) 'The insulin-like growth factor-phosphatidylinositol 3-kinase-Akt signaling
974 pathway regulates myogenin expression in normal myogenic cells but not in rhabdomyosarcoma-
975 derived RD cells.' *J Biol Chem*, 275(47), Nov 24, pp. 36750-36757.
976

977 Yablonkareuveni, Z., Balestreri, T. M. and Bowenpope, D. F. (1990) 'REGULATION OF PROLIFERATION
978 AND DIFFERENTIATION OF MYOBLASTS DERIVED FROM ADULT-MOUSE SKELETAL-MUSCLE BY SPECIFIC
979 ISOFORMS OF PDGF.' *Journal of Cell Biology*, 111(4), Oct, pp. 1623-1629.
980

981 Yin, H., Price, F. and Rudnicki, M. A. (2013) 'Satellite cells and the muscle stem cell niche.' *Physiol Rev*,
982 93(1), Jan, pp. 23-67.
983

984 Yousef, H., Conboy, M. J., Morgenthaler, A., Schlesinger, C., Bugaj, L., Paliwal, P., Greer, C., Conboy, I. M.
985 and Schaffer, D. (2015) *Systemic attenuation of the TGF- β pathway by a single drug simultaneously*
986 *rejuvenates hippocampal neurogenesis and myogenesis in the same old mammal.*
987

988 Zhang, C., Li, Y., Wu, Y., Wang, L., Wang, X. and Du, J. (2013) 'Interleukin-6/signal transducer and
989 activator of transcription 3 (STAT3) pathway is essential for macrophage infiltration and myoblast
990 proliferation during muscle regeneration.' *J Biol Chem*, 288(3), Jan 18, pp. 1489-1499.
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009





Fig. 1. Timeline of inflammatory responses and immune cell during muscle

The immune system responds to muscle damage by recruiting a precise sequence of pro and anti-inflammatory immune cells to the site of injury. Immune cells are observed from the initial pro-inflammatory phase required for removal of cellular debris through to the final repair of the damaged muscle fibres. Neutrophils rapidly infiltrate the extracellular space around the damaged fibres within 2 hours and peak in number between 6 and 24 hours followed by rapid decline of neutrophils to negligible levels within 72 to 96 hours. This initial infiltration of neutrophils further contributes to the inflammatory damage to the injured muscle fibres. Subsequently, M1 macrophage concentrations rapidly begin to increase at the site of injury and initiate the pro-inflammatory functions of muscle repair through secretion of several cytokines and mediators. The number of M1 macrophages will continue to increase until peak concentrations at 72 to 96 hours after injury and then begin to decline sharply. This is followed by the increase in numbers of anti-inflammatory and pro-myogenic M2 macrophages, which reach peak concentrations in the regenerating muscle at roughly 120 to 144 hours post injury, remaining significantly elevated for several days following. Finally activated T-cells are recruited to the regenerating muscle damage site, with concentrations beginning to peak as M2 macrophage number begin to decline. Populations of T cells, specifically mTreg remain significantly elevated for 30 days following the initial injury causing muscle damage. (Modified from **Tidball & Villalta, 2010; Forbes & Rosenthal, 2014**).


1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037



1038
 1039
 1040
 1041
 1042
 1043
 1044
 1045
 1046
 1047
 1048
 1049
 1050
 1051
 1052
 1053
 1054
 1055
 1056
 1057
 1058
 1059

Fig. 2. Innate immune signalling pathways in skeletal muscle regeneration

The activation of NF-κB in either muscle cells or macrophages can affect muscle cell proliferation and differentiation. Cytokines (IL1β, TNFα and IFNγ) can increase NF-κB activation in both muscle and/or macrophages. The cytokines contribute to further activation of NF-κB in macrophages and muscle cells or they can act on the muscle cells themselves to affect their proliferation or differentiation. TNFα can activate NF-κB within macrophages, which then induces the production of additional proinflammatory mediators. NF-κB activation can promote proliferation of muscle cells through the up regulation of transcripts needed for cell cycle progression (cyclin D1), while suppressing differentiation by decreasing the expression or destabilizing transcripts needed for muscle to experience early and terminal differentiation (MyoD and myogenin). Along with TNFα promoting proliferation and inhibiting differentiation through the NF-κB signalling pathway, it can also promote later stages of differentiation through the activation of p38 kinase. Nuclear factor-kappa B (NF- κB), interferon gamma (IFNγ), Interleukin-1 beta (IL-1β), Tumour necrosis factor (TNFα), (Modified from **Tidball & Villalta, 2010, Pillon et al., 2013; Forbes & Rosenthal, 2014**).



1060
 1061
 1062
 1063
 1064
 1065
 1066
 1067
 1068
 1069
 1070
 1071
 1072
 1073
 1074
 1075
 1076
 1077

Fig. 3. Summary of general interactions between immune and muscle cells following acute muscle injury. Following muscle damage quiescent satellite cells become activated and begin to migrate to the site of injury. The satellite cells re-enter the cell cycle and begin to proliferate until a proliferative threshold is met. A required quantity of the proliferating satellite cells will self-renew to replenish the pool of quiescent cells while the remaining proliferating cells will continue to differentiate to repair the damaged muscle fibres. Importantly, the phases of satellite cell activation, migration, proliferation and differentiation are regulated by immune cells. The immune system responds to muscle damage with a complex sequence of reactions, which ultimately lead to inflammation followed by muscle regeneration. The initial infiltration of transient neutrophils contain and localize the damage in the muscle and clean up cellular debris. M1 macrophages secrete cytokines that induce satellite cell activation and proliferation. M2 macrophages that then promote muscle repair, differentiation and recruit T-cells to the injured muscle site. T-cells such as mTreg cells secreting numerous growth factors (e.g. IGF-I, amphiregulin) and cytokines, which may contribute to facilitating muscle regeneration. Insulin-like growth factor I (IGF-I), Interferon gamma (IFN γ), Interleukin-1 beta (IL-1 β), Tumour necrosis factor (TNF α), fibroblast growth factor (FGF-2), muscle regulatory factors 4 (MRF4), Interleukin-4 (IL-4), Interleukin-10 (IL-10) (Modified from Siegel, A. et al, 2009; Tidball & Villalta, 2010, Pillon et al., 2013; Forbes & Rosenthal, 2014).