



LJMU Research Online

Maessen, MFH, Schalkwijk, CG, Verheggen, RJHM, Aengevaeren, VL, Hopman, MTE and Eijsvogels, TMH

A comparison of dicarbonyl stress and advanced glycation endproducts in lifelong endurance athletes vs. sedentary controls

<http://researchonline.ljmu.ac.uk/id/eprint/6320/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Maessen, MFH, Schalkwijk, CG, Verheggen, RJHM, Aengevaeren, VL, Hopman, MTE and Eijsvogels, TMH (2017) A comparison of dicarbonyl stress and advanced glycation endproducts in lifelong endurance athletes vs. sedentary controls. Journal of Science and Medicine in Sport. ISSN

LJMU has developed [LJMU Research Online](http://researchonline.ljmu.ac.uk/) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

A comparison of dicarbonyl stress and advanced glycation endproducts in lifelong endurance athletes *versus* sedentary controls

Martijn F.H. Maessen MSc ^a

Casper G. Schalkwijk PhD ^b

Rebecca J.H.M. Verheggen MSc, MD ^a

Vincent L. Aengevaeren MSc, MD ^a

Maria T.E. Hopman MD, PhD ^a

Thijs M.H. Eijsvogels PhD ^{a,c}

Affiliations:

^a Department of Physiology, Radboud university medical center, Nijmegen, The Netherlands.

^b Department of Internal Medicine, CARIM School for Cardiovascular Diseases, Maastricht University Medical Centre, The Netherlands.

^c Research Institute for Sports and Exercise Sciences, Liverpool John Moores University, Liverpool, United Kingdom.

Total word count: **3159**

Word count abstract: **250**

Total number of Figures: **1**

Total number of Tables: **2**

Reprints and correspondence:

Dr. Thijs Eijsvogels PhD, Dept. of Physiology (392), Radboud university medical center, P.O. Box 9101, 6500 HB Nijmegen, The Netherlands. E-mail: Thijs.Eijsvogels@radboudumc.nl.

Tel. (+31) (0)24 36 14200 Fax. (+31) (0)24 36 68340

4 **ABSTRACT**

5 **Objectives.** Dicarbonyl stress and high concentrations of advanced glycation endproducts (AGEs)
6 relate to an elevated risk for cardiovascular diseases (CVD). Exercise training lowers the risk for
7 future CVD. We tested the hypothesis that lifelong endurance athletes have lower dicarbonyl stress
8 and AGEs compared to sedentary controls and that these differences relate to a better cardiovascular
9 health profile. **Design.** Cross-sectional study

10 **Methods.** We included 18 lifelong endurance athletes (ATH, 61±7 years) and 18 sedentary controls
11 (SED, 58±7 years) and measured circulating glyoxal (GO), methylglyoxal (MGO) and 3-
12 deoxyglucosone (3DG) as markers of dicarbonyl stress. Furthermore, we measured serum levels of
13 protein-bound AGEs N^ε-(carboxymethyl)lysine (CML), N^ε-(carboxyethyl)lysine (CEL),
14 methylglyoxal-derived hydroimidazolone-1 (MG-H1), and pentosidine. Additionally, we measured
15 cardiorespiratory fitness (VO₂peak) and cardiovascular health markers.

16 **Results.** ATH had lower concentrations of MGO (196 [180-246] vs. 242 [207-292] nmol/mmol lysine,
17 P=0.043) and 3DG (927 [868-972] vs. 1061 [982-1114] nmol/mmol lysine, P<0.01), but no GO
18 compared to SED. ATH demonstrated higher concentrations CML and CEL compared to SED.
19 Pentosidine did not differ across groups and MG-H1 was significantly lower in ATH compared to
20 SED. Concentrations of MGO en 3DG were inversely correlated with cardiovascular health markers,
21 whereas CML and CEL were positively correlated with VO₂peak and cardiovascular health markers.

22 **Conclusion.** Lifelong exercise training relates to lower dicarbonyl stress (MGO and 3DG) and the
23 AGE MG-H1. The underlying mechanism and (clinical) relevance of higher CML and CEL
24 concentrations among lifelong athletes warrants future research, since it conflicts with the idea that
25 higher AGE concentrations relate to poor cardiovascular health outcomes.

26

27 **Key words:** oxidative stress; cardiovascular disease; physical activity; exercise physiology

28 **Introduction**

29 Advanced glycation endproducts (AGEs) are a complex group of modified proteins or lipids that are
30 formed by a process of non-enzymatically glycation and oxidation. AGEs formation is a slow process
31 (*i.e.*, weeks to months) and depends on the extent of oxidative stress, degree of hyperglycemia, and
32 turnover rate of proteins.^{1, 2} The formation of AGEs is irreversible and AGEs accumulate with
33 increasing age. Highly reactive dicarbonyls (α -oxoaldehydes) are involved in the fast formation of
34 AGEs and accumulation of dicarbonyls is known as dicarbonyl stress.^{1, 2} Dicarbonyls are precursors
35 for AGEs³ and the most important dicarbonyl marker is the highly reactive methylglyoxal (MGO).⁴
36 Dicarbonyl stress and a high concentration of AGEs are linked to the development of cardiovascular
37 diseases.⁴⁻⁶

38

39 Higher levels of circulating AGEs are also related to higher vascular stiffness.⁷⁻⁹ There are several
40 mechanisms proposed how AGEs may affect the vascular wall properties, such as binding to receptor
41 AGEs (RAGEs) and cross-linking matrix proteins in the vessel wall.^{2, 10} AGE-binding to RAGEs
42 leads to an upregulation of inflammation and production of reactive oxygen species.^{11, 12} These
43 processes augment vascular dysfunction and may promote vascular stiffness.^{11, 12} Alternatively, AGEs
44 can also bind to collagen and elastin to form crosslinks with matrix proteins, which promotes vascular
45 stiffness.¹² Strategies to lower the burden of high levels of AGEs may improve cardiovascular health
46 and need to be explored.

47

48 Regular exercise training is part of a healthy lifestyle and is an effective strategy to reduce the risk for
49 cardiovascular morbidity and mortality.^{13, 14} Exercise training attenuates the age-associated decline in
50 cardiovascular function,^{15, 16} and improves glucose¹⁷ and lipid metabolism.¹⁸ Findings from animal
51 studies suggest that these health benefits of exercise training may relate to a reduction of dicarbonyl
52 stress and AGEs concentrations.^{19, 20} Clinical studies linking exercise training with dicarbonyl stress
53 or AGEs are, however, sparse and conflicting.²¹⁻²³ A previous study demonstrated that 12 months of
54 tai chi training for 2 sessions/week significantly reduced serum AGEs concentrations in asymptomatic
55 middle-aged adults.²³ However, another study found no effect on serum AGEs concentrations in

56 middle-aged overweight or obese men after a 3-month aerobic moderate intensity exercise training
57 program ²¹. Variation in study outcomes may partially relate to the training duration (3 vs. 12 months),
58 exercise intensity (light vs. moderate), or study population (asymptomatic vs. overweight/obese).
59 Lifelong endurance athletes may provide better insight to what extent exercise is related to attenuated
60 AGEs formation.

61

62 Therefore, we tested the hypothesis that lifelong endurance athletes have lower dicarbonyl stress and a
63 lower concentration of AGEs compared to sedentary controls. Additionally, we explored whether
64 lower dicarbonyl stress and lower concentration of AGEs relate to a better cardiovascular health
65 profile.

66 **Methods**

67 Thirty-six male participants aged >45 years were included and stratified into 2 groups based on their
68 lifelong exercise patterns: 1) lifelong endurance athletes (ATH, n=18), 2) sedentary controls (SED,
69 n=18). ATH had to perform ≥ 20 years of endurance exercise training (e.g., running or cycling) for ≥ 4
70 hours/week, whereas SED had to report ≥ 20 years of habitual physical activity <2 hours/week. Current
71 smokers, participants with a history of diabetes mellitus or cardiovascular disease, or participants not
72 able to perform an incremental maximal cycling test were not included in the study. The Local
73 Committee on Research Involving Human Subjects of the region Arnhem and Nijmegen approved the
74 study. All participants gave their written informed consent prior to study participation.

75
76 During this cross-sectional study, participants visited our laboratory on 2 separate days. On day 1,
77 participants were medically screened for eligibility, followed by an incremental maximal cycling test
78 to determine their physical fitness. On day 2, pulse wave velocity was measured as an index of
79 vascular stiffness and blood samples were obtained under fasting conditions. Both testing days were
80 scheduled within a 14-day time-frame, with at least 1 recovery day between measurement day 1 and 2.

81
82 A physician medically screened the participants by taking a detailed medical history, physical
83 examination, and 12-lead electrocardiogram. After screening, participants performed an incremental
84 maximal cycling test to determine the cardiorespiratory fitness and peak oxygen uptake (VO_{2peak} ,
85 mLO_2/min). The test took place in a temperature-controlled room (18-19°C) and under the supervision
86 of a physician. Participants cycled with 60-80 rotations per minute while the workload increased with
87 20 Watt/min for ATH and 10 Watt/min for CON. Heart rate was continuously measured via a 12 lead-
88 electrocardiogram. Oxygen uptake (VO_2 [mL/min]), carbon dioxide output (VCO_2 [mL/min]), and
89 respiratory exchange ratio (RER) were continuously measured via a gas analyser (CPET, Cosmed
90 v9.1b, Rome, Italy). Lactate concentration (mmol/L) was measured (Lactate Pro™ 2, Arkray, type LT-
91 1730, Kyoto, Japan) via a capillary blood sample taken 1.5 minute after cessation of the exercise test.
92 The incremental maximal cycling test was considered successful when 2 of the 4 criteria were met: 1)

93 RER \geq 1.05, II) achievement of at least 85% of age-predicted maximal heart rate (220 – age), III)
94 blood lactate \geq 6.00 mmol/L, or IV) flattening of VO₂ uptake curve (\leq 150 mL increase during the last
95 minute).^{24, 25}

96
97 Lifelong exercise patterns were queried via an exercise history questionnaire, distinguishing 5 age-
98 periods: I) 20-29 years, II) 30-39 years, III) 40-49 years, IV) 50-59 years and V) >60 years. Each
99 category consisted of 2 queries: 1) type of activity (*e.g.*, running, cycling, etc., or nothing) and 2)
100 exercise time (hours) per activity per week. Based on the Compendium of Physical Activities²⁶, the
101 corresponding metabolic equivalent of task (MET) score per exercise activity was determined.
102 Vigorous exercise activities were defined as a MET score >6. Subsequently, exercise volume (MET-
103 hours/week) was calculated by multiplying exercise time with accompanying MET score. The average
104 exercise time and dose were calculated over the last 2 decades.

105
106 Before the second testing day, participants were asked to abstain from I) (vigorous) physical activities
107 for 24 hours, II) caffeine, alcohol, or vitamin supplement intake for at least 18 hours, and III) food
108 intake for \geq 6 hours. Central and peripheral pulse wave velocity was assessed with a three-lead
109 electrocardiogram and an echo-Doppler ultrasound machine (WakiLoki Doppler, 4 MHz, Atys) at the
110 left carotid artery, right common femoral artery, and radial artery. The distances between sternal notch
111 and site of measurement for the carotid artery and between radial artery and common femoral artery
112 via the umbilicus were measured.²⁷ At least 10 cardiac cycles were recorded for analyses. Based on
113 the R-R interval and onset of the Doppler waveform, central and peripheral pulse wave velocities were
114 calculated in Matlab R2014 (The MathWorks Inc., United States).

115
116 Following vascular measurements, a fasting blood sample (8 mL) was obtained from an antecubital
117 vein for the assessment of concentrations of dicarbonyl stress and AGEs. Additionally, lysine and
118 traditional cardiovascular risk factors (total-, high-density lipoproteins [HDL]-, low-density
119 lipoproteins [LDL]-cholesterol, triglycerides, glycated hemoglobin [HbA1C], and glucose) were
120 determined. Homeostasis model assessment of insulin resistance (HOMA-IR) was calculated based on

121 glucose and insulin concentrations ($IR = (\text{fasting insulin [mU/L]} \times \text{fasting glucose [mmol/L]})/22.5$).²⁸
122 To gain insight in the cardiovascular (risk) profile of ATH and SED, the 10-year CVD risk was
123 calculated via the Framingham Risk Score (FRS).²⁹
124
125 For measurement of serum levels of diarbonyl components and AGEs, we used ultra-performance
126 liquid chromatography tandem mass spectrometry (UPLC-MS/MS, Waters, Milford Massachusetts,
127 USA). UPLC-MS/MS combines liquid chromatography for separation and tandem mass spectrometry
128 for specific detection.
129
130 Whole blood samples in serum-separating tubes were centrifuged after collection (10 min, 4°C, 3,000
131 g) and supernatant was stored at -80°C until analysis. Serum levels of dicarbonyl compounds glyoxal
132 (GO), MGO, and 3-deoxyglucosone (3DG) were analysed following a previously described protocol.³
133 Briefly, serum samples were deproteinized using perchloric acid and subsequently derivatized with o-
134 phenylenediamine. GO, MGO, and 3DG concentrations were measured using stable isotope-dilution
135 UPLC-MS/MS (Waters, Milford Massachusetts, USA) with a run-to-run time of 8 min. Intra-run and
136 inter-run variations were 4.3% and 14.3% for GO, 2.9% and 7.3% for MGO, and 2.4% and 12.0% for
137 3DG, respectively.³
138
139 Protein-bound serum AGEs N^ε-(carboxymethyl)lysine (CML), N^ε-(carboxyethyl)lysine (CEL),
140 methylglyoxal-derived hydroimidazolone-1 (MG-H1), and lysine were measured with UPLC-MS/MS
141 (Waters, Milford Massachusetts, USA), as previously described.^{30,31} Pentosidine was measured with
142 high-performance liquid chromatography and fluorescent detection.³¹ Intra-run and inter-run
143 variations were 2.8% and 7.1% for CML, 3.7% and 6.4% for CEL, 3.7% and 5.1% for MG-H1, and
144 2.0% and 3.1% for pentosidine.^{30,31} All serum AGEs were adjusted for lysine concentrations as a
145 marker of total protein concentration.
146
147 Participant characteristics were summarized with means and standard deviations or median and
148 interquartile range (IQR), when appropriate. Categorical data were analysed using the *Fisher's exact*

149 test. Parameters were checked for normality using a *Shapiro-Wilk* test and Q-Q plots. Skewed
150 variables were log_e-transformed before statistical analyses were conducted. Differences in participant
151 characteristics, lifelong exercise patterns, and cardiovascular health markers between ATH and SED
152 were analysed using an independent *Student's t* test. As an overall measure of pulse wave velocity, z-
153 scores of central and peripheral pulse wave velocities were averaged. Correlations between markers
154 for dicarbonyl stress or AGEs and markers for cardiovascular health (BMI, pulse wave velocity,
155 cardiorespiratory fitness, Framingham risk score, and glucose metabolism) were evaluated using
156 *Spearman's rank* test. All statistical analyses were performed using SPSS 21.0 software (IBM Corp.
157 Released 2012. IBM SPSS Statistics for Windows, Version 21.0. Armonk, NY: IBM Corp.). Statistical
158 significance was assumed at $p < 0.05$ (two-sided).

159 **Results**

160 Age, height, mean arterial pressure, and smoking history did not differ between groups, but ATH
161 demonstrated a lower body weight and Body Mass Index compared to SED (Table 1). HbA1c, total
162 cholesterol, and glucose concentrations did not differ between groups, but ATH demonstrated a higher
163 HDL cholesterol concentration and lower LDL cholesterol, triglycerides, and HOMA-IR compared to
164 SED (Table 1). The median time between smoking cessation and study participation was 28 years
165 (Q_{25} : 12 to Q_{75} : 40) in ATH *versus* 25 years (Q_{25} : 15 to Q_{75} : 37) in SED ($P=0.78$).

166
167 ATH showed a significantly higher weekly exercise time and dose compared to SED (Table 1). ATH
168 mostly performed vigorous-intensity exercise activities (e.g. running or road cycling). We observed a
169 higher VO_{2peak} in ATH (3544 ± 651 mL/min) compared to SED (2843 ± 519 mL/min, $p<0.01$).
170 Likewise, ATH reached a higher power output during the incremental exercise test compared to SED
171 ($p<0.01$, Table 1).

172
173 Central pulse wave velocity was significantly lower in ATH (7.0 ± 2.2 m/s) compared to SED (9.2 ± 2.3
174 m/s, $P<0.01$). Peripheral pulse wave velocity was significantly lower in ATH (8.1 ± 1.5 m/s) compared
175 to SED (9.4 ± 1.6 m/s, $p=0.017$).

176
177 MGO (196 [180-246] *vs.* 242 [207-292] nmol/mmol lysine, $P=0.043$) and 3DG (927 [868-972] *vs.*
178 1061 [982-1114] nmol/mmol lysine, $p<0.01$) concentrations were lower in ATH compared to SED
179 (Figure 1). Glyoxal concentrations did not differ between ATH *vs.* SED (314 [202-451] *vs.* 342 [266-
180 388] nmol/mmol lysine, $p=0.86$, Figure 1).

181
182 CML was significantly higher in ATH (80 [73-89] nmol/mmol lysine) *vs.* SED (68 [56-76]
183 nmol/mmol lysine, $p<0.01$, Figure 2). Similarly, CEL was significantly higher in ATH (35 [28-41]
184 nmol/mmol lysine) compared to SED (28 [24-34] nmol/mmol lysine, $p=0.035$). Pentosidine (0.63
185 [0.59-0.86] *vs.* 0.56 [0.48-0.67] nmol/mmol lysine, $p=0.11$) did not differ between groups (Figure 2).

186 MG-H1 concentration was significantly lower in ATH (363 [288-468] nmol/mmol lysine) compared to
187 SED (460 [340-536] nmol/mmol lysine, $p=0.043$, Figure 2).

188

189 MGO was positively correlated with BMI, central PWV, and FRS. (Table 2). 3DG was negatively
190 correlated with VO_2 peak, but positively correlated with BMI, central and peripheral PWV, FRS, and
191 glucose (Table 2). GO did not correlate with cardiovascular health parameters (Table 2).

192

193 CML was negatively correlated with BMI and peripheral PWV, but positively correlated with
194 VO_2 peak. MG-H1 was negatively correlated with VO_2 peak (Table 2). Pentosidine was negatively
195 correlated with peripheral PWV and glucose (Table 2). CEL did not correlate with cardiovascular
196 health parameters (Table 2).

197 **Discussion**

198 This study aimed to compare markers of dicarbonyl stress and circulating AGEs between lifelong
199 endurance athletes and sedentary controls. MGO and 3DG were significantly lower in ATH compared
200 to SED, and were related to a better cardiovascular health profile. However, we also found that CML
201 and CEL were significantly higher in ATH compared to SED.

202

203 The benefits of exercise training on cardiovascular health are indisputable ¹⁴⁻¹⁶, but underlying
204 mechanisms explaining the lower risk for cardiovascular events in physically active individuals are not
205 fully understood ¹⁶. Our results suggest that benefits of exercise training relate to a lower concentration
206 of MGO and 3DG. These findings are in line with a recent study in rats, which demonstrated that
207 running exercise was associated with a reduction in dicarbonyl stress ¹⁹. In general, we found that
208 markers of dicarbonyl stress showed a moderate, yet significant correlation with cardiovascular health
209 or metabolic markers. For example, lower concentration MGO and 3DG were correlated to low
210 Framingham risk score, lower insulin concentration, and better HOMA-IR. Reducing hyperglycaemia
211 and improving insulin sensitivity may be a first step to reduce accumulation of MGO ^{4,32} and 3DG ³².
212 High levels of dicarbonyl stress, and especially MGO, increase morbidity risk ⁴⁻⁶. MGO is highly
213 reactive and is mainly catabolized via glyoxalase I of the glyoxalase system. The activity of the
214 glyoxalase system depends on concentrations of reduced glutathione (GSH) ^{4,33}. Biosynthesis of GSH
215 is heavily dependent of the antioxidant response element-nuclear respiratory factor (ARE-Nrf)
216 pathway. Animal and human studies demonstrated that an acute bout of swimming or moderate
217 intensity endurance exercise training upregulate the ARE-Nrf pathway and GSH biosynthesis. This led
218 to the hypothesis that exercise training enhances the glyoxalase system and may lower MGO and MG-
219 H1 concentrations. ³⁴ Based on our data, it can be speculated that exercise training possibly lowers the
220 levels of MGO and MG-H1 via an upregulation of the glyoxalase system. Further research is
221 warranted to explore these pathways. Taken together, our data demonstrated that exercise training is
222 related to lower levels of MGO, 3DG, and MG-H1.

223

224 In contrast to our hypothesis, we found that 2 of the 4 AGEs (CML and CEL) were significantly
225 higher in ATH, whereas MG-H1 was significantly lower in ATH compared to SED. Although MG-H1
226 is a AGE, it is produced in a much shorter timeframe and is less stable than CML, CEL or pentosidine.
227 ³⁵ MG-H1 may, therefore, better relate to abnormal accumulation of dicarbonyl stress. ³⁵ This could
228 explain why MG-H1 showed opposite results compared to the other AGEs, since dicarbonyl stress was
229 lower in ATH compared to SED.

230
231 Previous studies indicated that an increase in AGEs concentration relates to poor health outcomes. ⁴⁻⁶
232 Our findings are contradictory to this concept, as we found an inverse relation between circulating
233 CML and pulse wave velocity, BMI, and cardiorespiratory fitness. A potential explanation for this
234 finding could be that exercise enhances collagen turnover rate, which breaks and prevents AGE cross-
235 links in the vessel wall. ^{12, 36, 37} This may contribute to higher levels of circulating AGEs, but this
236 hypothesis needs to be reinforced with future studies. Alternatively, a recent animal study
237 demonstrated that a 12-week running exercise training leads to suppressed RAGEs activation in the
238 aorta of aged rats. ³⁸ It could be speculated that attenuated RAGEs activity limits the uptake of AGEs
239 from the circulation to the surrounding tissue, ³⁹ leading to increased levels of circulating AGEs. Thus,
240 the observation of higher AGEs in lifelong endurance athletes may relate to a higher collagen turnover
241 and/or suppression of RAGEs due to long-term exercise training.

242
243 Another possible explanation for the higher AGEs concentrations in ATH vs. SED may relate to the
244 (vigorous) exercise training regimes of our lifelong endurance athletes. Acute exercise induces a
245 transient increase in oxidative stress, ⁴⁰ which upregulates the formation of AGEs. ^{1, 2} Mice deficient in
246 NADPH (nicotinamide adenine dinucleotide phosphate) oxidase, a pathway involved in the generation
247 of reactive oxygen species, showed an impaired CML generation, which suggests that oxidative stress
248 is a potential stimulus to generate CML. ⁴¹ Although the sudden increase in oxidative stress is a
249 necessary stimulus to enhance the anti-oxidative defence mechanism (*i.e.*, glyoxalase system), ⁴² it is
250 possible that the formation of AGEs is simultaneously upregulated. The positive relation between
251 exercise dose / time and CML concentrations found in the present study (Table 2) may relate to the

252 effects of sustained exposure to vigorous exercise training. Hence, lifelong and repetitive exposure to
253 vigorous exercise increases oxidative stress and may boost the accumulation of circulating AGEs in
254 the blood. Future research is warranted to elucidate the underlying mechanisms and (clinical) impact
255 of higher AGEs (CML and CEL) concentrations in athletes, as this observation contradicts with the
256 general believe that high concentrations of circulating AGEs relate to CVD.

257
258 This cross-sectional study is inherent to some limitations. First, the comparison between athletes and
259 sedentary individuals does not prove that exercise can attenuate the formation of dicarbonyl stress. A
260 randomized clinical trial would be needed to confirm causation. However, our results indicate that
261 exercise training is related to lower dicarbonyl stress. Unfortunately, we do not have information about
262 the dietary habits of the participants. The absorption, bioavailability, and effects of dietary AGEs are
263 poorly understood in vivo,⁴³ and it could be that diet patterns may contribute to the differences in
264 AGEs between ATH and SED. AGE-rich food intake has been associated with higher levels of serum
265 AGEs, whereas an AGE-restricted diet has been associated with lower serum AGEs.²¹ However,
266 whether food AGEs influence protein bound AGEs, as measured in this study, is not clear. Free AGEs
267 may be relatively quickly absorbed, biotransformed, and excreted. On the other hand, high molecule
268 weight AGEs, such as protein bound AGEs, may not be very extensively absorbed due to insufficient
269 degradation by gastrointestinal enzymes.⁴³ Further research is warranted to establish a direct relation
270 between dietary AGEs and protein-bound AGEs. Finally, all the participants of the study were men
271 and the lifelong athletes performed endurance exercise activities only, which limits the generalizability
272 of the present study.

273

274 **Conclusion**

275 Findings of the present study indicate that lifelong exercise training is associated with lower
276 dicarbonyl stress (MGO and 3DG), which is related to improved cardiovascular health. Although MG-
277 H1 was lower in lifelong endurance athletes compared to sedentary controls, AGEs concentrations of
278 CML and CEL were significantly higher in athletes compared to sedentary controls. The underlying
279 mechanism and (clinical) relevance of higher CML and CEL concentrations among lifelong athletes

280 warrants future research, since it conflicts with the idea that higher AGEs concentrations relate to poor
281 cardiovascular health.
282

283 **Practical Implications**

- 284 • Results of our study support the cardiovascular health benefits of lifelong exercise training, as
285 lifelong endurance athletes demonstrated a better cardiovascular risk profile compared to
286 sedentary controls.
- 287 • Lifelong exercise training is related to lower dicarbonyl stress, as veteran athletes had lower
288 concentrations of methylglyoxal and 3-deoxyglucosone compared to sedentary controls.
- 289 • Lifelong exercise training is related to higher concentrations of advanced glycation
290 endproducts (N^ε-(carboxymethyl)lysine and N^ε-(carboxyethyl)lysine). Although previous
291 studies indicated that higher concentrations of advanced glycation endproducts were
292 associated with adverse outcomes, the clinical significance of our findings in a highly active
293 population is unknown.

294

295 **Acknowledgements**

296 Dr. TE is financially supported by a European Commission Horizon 2020 grant [Marie Skłodowska-
297 Curie Fellowship]. The remaining authors report no conflicts of interest.

298 **References**

- 299 1. Goldin A, Beckman JA, Schmidt AM, et al. Advanced glycation end products:
300 sparking the development of diabetic vascular injury. *Circulation*. 2006; 114(6):597-
301 605.
- 302 2. Brownlee M. Biochemistry and molecular cell biology of diabetic complications.
303 *Nature*. 2001; 414(6865):813-820.
- 304 3. Scheijen JL, Schalkwijk CG. Quantification of glyoxal, methylglyoxal and 3-
305 deoxyglucosone in blood and plasma by ultra performance liquid chromatography
306 tandem mass spectrometry: evaluation of blood specimen. *Clin Chem Lab Med*. 2014;
307 52(1):85-91.
- 308 4. Maessen DE, Stehouwer CD, Schalkwijk CG. The role of methylglyoxal and the
309 glyoxalase system in diabetes and other age-related diseases. *Clin Sci (Lond)*. 2015;
310 128(12):839-861.
- 311 5. Hanssen NM, Beulens JW, van Dieren S, et al. Plasma advanced glycation end
312 products are associated with incident cardiovascular events in individuals with type 2
313 diabetes: a case-cohort study with a median follow-up of 10 years (EPIC-NL).
314 *Diabetes*. 2015; 64(1):257-265.
- 315 6. Hanssen NM, Wouters K, Huijberts MS, et al. Higher levels of advanced glycation
316 endproducts in human carotid atherosclerotic plaques are associated with a rupture-
317 prone phenotype. *Eur Heart J*. 2014; 35(17):1137-1146.
- 318 7. McNulty M, Mahmud A, Feely J. Advanced glycation end-products and arterial
319 stiffness in hypertension. *Am J Hypertens*. 2007; 20(3):242-247.
- 320 8. Semba RD, Najjar SS, Sun K, et al. Serum carboxymethyl-lysine, an advanced
321 glycation end product, is associated with increased aortic pulse wave velocity in
322 adults. *Am J Hypertens*. 2009; 22(1):74-79.

- 323 9. van Eupen MG, Schram MT, van Sloten TT, et al. Skin Autofluorescence and
324 Pentosidine Are Associated With Aortic Stiffening: The Maastricht Study.
325 *Hypertension*. 2016; 68(4):956-963.
- 326 10. Smit AJ, Lutgers HL. The clinical relevance of advanced glycation endproducts
327 (AGE) and recent developments in pharmaceuticals to reduce AGE accumulation. *Curr*
328 *Med Chem*. 2004; 11(20):2767-2784.
- 329 11. Basta G. Receptor for advanced glycation endproducts and atherosclerosis: From basic
330 mechanisms to clinical implications. *Atherosclerosis*. 2008; 196(1):9-21.
- 331 12. Sell DR, Monnier VM. Molecular basis of arterial stiffening: role of glycation - a
332 mini-review. *Gerontology*. 2012; 58(3):227-237.
- 333 13. Blair SN, Kampert JB, Kohl HW, 3rd, et al. Influences of cardiorespiratory fitness and
334 other precursors on cardiovascular disease and all-cause mortality in men and women.
335 *JAMA*. 1996; 276(3):205-210.
- 336 14. Maessen MF, Verbeek AL, Bakker EA, et al. Lifelong Exercise Patterns and
337 Cardiovascular Health. *Mayo Clin Proc*. 2016; 91(6):745-754.
- 338 15. Eijssvogels TM, Molossi S, Lee DC, et al. Exercise at the Extremes: The Amount of
339 Exercise to Reduce Cardiovascular Events. *J Am Coll Cardiol*. 2016; 67(3):316-329.
- 340 16. Mora S, Cook N, Buring JE, et al. Physical activity and reduced risk of cardiovascular
341 events: potential mediating mechanisms. *Circulation*. 2007; 116(19):2110-2118.
- 342 17. Thomas DE, Elliott EJ, Naughton GA. Exercise for type 2 diabetes mellitus. *Cochrane*
343 *Database Syst Rev*. 2006(3):CD002968.
- 344 18. Kraus WE, Houmard JA, Duscha BD, et al. Effects of the amount and intensity of
345 exercise on plasma lipoproteins. *N Engl J Med*. 2002; 347(19):1483-1492.

- 346 19. Ito D, Cao P, Kakihana T, et al. Chronic Running Exercise Alleviates Early
347 Progression of Nephropathy with Upregulation of Nitric Oxide Synthases and
348 Suppression of Glycation in Zucker Diabetic Rats. *PLoS One*. 2015; 10(9):e0138037.
- 349 20. Boor P, Celec P, Behuliak M, et al. Regular moderate exercise reduces advanced
350 glycation and ameliorates early diabetic nephropathy in obese Zucker rats.
351 *Metabolism*. 2009; 58(11):1669-1677.
- 352 21. Macias-Cervantes MH, Rodriguez-Soto JM, Uribarri J, et al. Effect of an advanced
353 glycation end product-restricted diet and exercise on metabolic parameters in adult
354 overweight men. *Nutrition*. 2015; 31(3):446-451.
- 355 22. Oudegeest-Sander MH, Olde Rikkert MG, Smits P, et al. The effect of an advanced
356 glycation end-product crosslink breaker and exercise training on vascular function in
357 older individuals: a randomized factorial design trial. *Exp Gerontol*. 2013;
358 48(12):1509-1517.
- 359 23. Goon JA, Aini AH, Musalmah M, et al. Effect of Tai Chi exercise on DNA damage,
360 antioxidant enzymes, and oxidative stress in middle-age adults. *J Phys Act Health*.
361 2009; 6(1):43-54.
- 362 24. Balady GJ, Arena R, Sietsema K, et al. Clinician's Guide to cardiopulmonary exercise
363 testing in adults: a scientific statement from the American Heart Association.
364 *Circulation*. 2010; 122(2):191-225.
- 365 25. Edvardsen E, Hem E, Anderssen SA. End criteria for reaching maximal oxygen
366 uptake must be strict and adjusted to sex and age: a cross-sectional study. *PLoS One*.
367 2014; 9(1):e85276.
- 368 26. Ainsworth BE, Haskell WL, Herrmann SD, et al. 2011 Compendium of Physical
369 Activities: a second update of codes and MET values. *Med Sci Sports Exerc*. 2011;
370 43(8):1575-1581.

- 371 27. Laurent S, Cockcroft J, Van Bortel L, et al. Expert consensus document on arterial
372 stiffness: methodological issues and clinical applications. *Eur Heart J.* 2006;
373 27(21):2588-2605.
- 374 28. Matthews DR, Hosker JP, Rudenski AS, et al. Homeostasis model assessment: insulin
375 resistance and beta-cell function from fasting plasma glucose and insulin
376 concentrations in man. *Diabetologia.* 1985; 28(7):412-419.
- 377 29. D'Agostino RB, Sr., Vasan RS, Pencina MJ, et al. General cardiovascular risk profile
378 for use in primary care: the Framingham Heart Study. *Circulation.* 2008; 117(6):743-
379 753.
- 380 30. Hanssen NM, Engelen L, Ferreira I, et al. Plasma levels of advanced glycation
381 endproducts Nepsilon-(carboxymethyl)lysine, Nepsilon-(carboxyethyl)lysine, and
382 pentosidine are not independently associated with cardiovascular disease in
383 individuals with or without type 2 diabetes: the Hoorn and CODAM studies. *J Clin*
384 *Endocrinol Metab.* 2013; 98(8):E1369-1373.
- 385 31. Scheijen JL, van de Waarenburg MP, Stehouwer CD, et al. Measurement of
386 pentosidine in human plasma protein by a single-column high-performance liquid
387 chromatography method with fluorescence detection. *J Chromatogr B Analyt Technol*
388 *Biomed Life Sci.* 2009; 877(7):610-614.
- 389 32. Thornalley PJ, Langborg A, Minhas HS. Formation of glyoxal, methylglyoxal and 3-
390 deoxyglucosone in the glycation of proteins by glucose. *Biochem J.* 1999; 344 Pt
391 1:109-116.
- 392 33. Thornalley PJ. Pharmacology of methylglyoxal: formation, modification of proteins
393 and nucleic acids, and enzymatic detoxification-a role in pathogenesis and
394 antiproliferative chemotherapy. *Gen Pharmacol.* 1996; 27(4):565-573.

- 395 34. Dieter BP, Vella CA. A proposed mechanism for exercise attenuated methylglyoxal
396 accumulation: activation of the ARE-Nrf pathway and increased glutathione
397 biosynthesis. *Med Hypotheses*. 2013; 81(5):813-815.
- 398 35. Ahmed N, Argirov OK, Minhas HS, et al. Assay of advanced glycation endproducts
399 (AGEs): surveying AGEs by chromatographic assay with derivatization by 6-
400 aminoquinolyl-N-hydroxysuccinimidyl-carbamate and application to Nepsilon-
401 carboxymethyl-lysine- and Nepsilon-(1-carboxyethyl)lysine-modified albumin.
402 *Biochem J*. 2002; 364(Pt 1):1-14.
- 403 36. Coupe C, Svensson RB, Grosset JF, et al. Life-long endurance running is associated
404 with reduced glycation and mechanical stress in connective tissue. *Age (Dordr)*. 2014;
405 36(4):9665.
- 406 37. Ziemann SJ, Melenovsky V, Kass DA. Mechanisms, pathophysiology, and therapy of
407 arterial stiffness. *Arterioscler Thromb Vasc Biol*. 2005; 25(5):932-943.
- 408 38. Gu Q, Wang B, Zhang XF, et al. Contribution of receptor for advanced glycation end
409 products to vasculature-protecting effects of exercise training in aged rats. *Eur J*
410 *Pharmacol*. 2014; 741:186-194.
- 411 39. Gaens KH, Goossens GH, Niessen PM, et al. Nepsilon-(carboxymethyl)lysine-
412 receptor for advanced glycation end product axis is a key modulator of obesity-
413 induced dysregulation of adipokine expression and insulin resistance. *Arterioscler*
414 *Thromb Vasc Biol*. 2014; 34(6):1199-1208.
- 415 40. Powers SK, Jackson MJ. Exercise-induced oxidative stress: cellular mechanisms and
416 impact on muscle force production. *Physiol Rev*. 2008; 88(4):1243-1276.
- 417 41. Anderson MM, Heinecke JW. Production of N(epsilon)-(carboxymethyl)lysine is
418 impaired in mice deficient in NADPH oxidase: a role for phagocyte-derived oxidants

- 419 in the formation of advanced glycation end products during inflammation. *Diabetes*.
420 2003; 52(8):2137-2143.
- 421 42. Fisher-Wellman K, Bloomer RJ. Acute exercise and oxidative stress: a 30 year
422 history. *Dyn Med*. 2009; 8:1.
- 423 43. Poulsen MW, Hedegaard RV, Andersen JM, et al. Advanced glycation endproducts in
424 food and their effects on health. *Food Chem Toxicol*. 2013; 60:10-37.
- 425

426 **Figure legend**

Figure 1. Individual and average values of markers for (A) dicarbonyl stress and (B) advanced glycation endproducts in lifelong athletes (circles) and sedentary controls (squares). For dicarbonyl markers, GO concentrations did not differ between groups, whereas MGO and 3DG were significantly lower in athletes compared to controls. For advanced glycation endproducts, CML and CEL concentrations were higher in athletes compared to controls. Concentrations of pentosidine did not differ between groups. Concentrations of MG-H1 were lower in athletes compared to controls. P-value refers to an *independent Student's t* or (¥) *Mann-Whitney U* test. Group averages are presented as median and interquartile range.

427

Table 1. Participants' characteristics of lifelong endurance athletes (ATH, $n=18$) and sedentary controls (SED, $n=18$). Data is presented as mean and standard deviation or median and interquartile range (IQR). P-value refers to an *independent Student's t* test or *Mann-Whitney U* (*) test.

<i>n</i>	ATH	SED	<i>p</i> -value
CHARACTERISTICS			
Age (years)	61±7	58±7	0.29
Height (m)	179±8	181±6	0.31
Weight (kg)	74±8	87±10	<0.01
Body Mass Index (kg/m ²) ‡	23.6 (21.1-24.9)	26.7 (25.0-27.4)	<0.01
Mean arterial pressure (mmHg) *	98 (90-106)	103 (93-107)	0.70
Systolic blood pressure (mmHg)	134±17	137±16	0.53
Diastolic blood pressure (mmHg)	84±10	84±10	0.92
Smoking history (%yes [n])	10 (56)	15 (83)	0.15
CARDIOVASCULAR HEALTH PARAMETERS			
Pulse Wave Velocity			
Central PWV (m/s)	7.0±2.2	9.2±2.3	<0.01
Peripheral PWV (m/s)	8.1±1.5	9.4±1.6	0.017
Framingham Risk Score (%) *	10.1 (7.5-20.3)	16.5 (10.1-19.5)	0.12
VO ₂ peak (mL/min)	3544±651	2843±519	<0.01
Fasting blood levels			
HbA1c (mmol/mol) ‡	35.5 (34.4-38.3)	35.5 (35.5-38.3)	0.53
Cholesterol (mmol/L)	5.4±0.8	5.9±0.9	0.07
LDL (mmol/L)	3.3±0.8	4.0±0.8	0.012
HDL (mmol/L)	1.8±0.3	1.4±0.3	<0.01
Triglycerides (mmol/L) *	0.8 (0.7-1.2)	1.3 (1.0-2.4)	<0.01
Glucose (mmol/L) *	4.6 (4.4-5.0)	4.7 (4.4-4.9)	0.66
Insulin (mU/L)	2.8±1.8	6.8±2.9	<0.01
HOMA-IR *	0.5 (0.3-0.9)	1.3 (0.8-2.2)	<0.01
LIFELONG EXERCISE PATTERNS			
Exercise time (hours/week) ‡	7.1 (5.8-11.9)	0.5 (0.0-1.4)	<0.01
Exercise dose (MET-hours/week) ‡	60 (47-110)	4 (0-12)	<0.01
INCREMENTAL EXERCISE TEST			
Maximal heart rate (beats/min)	165±13	171±15	0.29
RER (ratio: VCO ₂ / VO ₂) *	1.13 (1.06-1.17)	1.08 (1.05-1.14)	0.029
Lactate (mmol/L) *	11.6 (8.9-12.3)	11.1 (9.4-12.8)	0.77
Power Output (W)	319±58	209±46	<0.01

HbA1c: Glycated haemoglobin; HDL: High-density lipoprotein; HOMA-IR: homeostasis model assessment of insulin resistance; LDL: low-density lipoprotein; MET: Metabolic Equivalent of Task; PWV: pulse wave velocity; RER: respiratory exchange ratio; VO₂peak: peak oxygen uptake;

* Data were log_e-transformed before statistical analysis

‡ non-parametrically tested via Mann-Whitney U

Table 2. Spearman's Rank (ρ) correlations between dicarbonyl stress, advanced glycation endproducts, and cardiovascular health parameters

	Dicarbonyl stress			Advanced glycation endproducts			
	GO	MGO	3DG	CML	CEL	Pentosidine	MG-H1
CARDIOVASCULAR HEALTH MARKERS							
BMI	0.02	0.35*	0.40*	-0.53**	-0.13	-0.31	0.19
Average PWV	0.12	0.35*	0.55**	-0.54**	0.10	-0.31	0.04
Central PWV	0.24	0.51**	0.46**	-0.30	0.07	-0.10	0.03
Peripheral PWV	-0.05	0.10	0.44**	-0.58**	0.11	-0.43*	0.02
VO ₂ peak (mL/min)	0.01	-0.32	-0.47**	0.34*	0.33	0.18	-0.55**
FRS	0.24	0.52**	0.43**	-0.23	-0.06	-0.09	0.11
Glucose	-0.19	0.15	0.46**	-0.13	0.13	-0.41*	-0.09
Insulin	0.04	0.36*	0.44**	-0.36*	-0.24	-0.12	0.35*
HOMA-IR	-0.01	0.34	0.49**	-0.36*	-0.21	-0.16	0.33
LIFELONG EXERCISE PATTERNS							
Exercise time	-0.04	-0.34*	-0.53**	0.46**	0.32	0.28	-0.36*
Exercise dose	-0.04	-0.34*	-0.53**	0.45**	0.36*	0.30	-0.37*


3DG: 3-deoxyglucosone; CEL: N_ε-(carboxyethyl)lysine; CML: N_ε-(carboxymethyl)lysine; FRS: Framingham risk score; GO: glyoxal; HOMA-IR: homeostasis model assessment of insulin resistance; MG-H1: Methylglyoxal-derived hydroimidazolone-1; MGO: methylglyoxal; VO₂peak: peak oxygen uptake (cardiorespiratory fitness); Average PWV: average pulse wave velocity, the average of the z-scores of central and peripheral PWV;

Correlation is significant at *0.05 or **0.01 level (two-sided).


Figure_1

A. Markers for dicarbonyl stress


I. Glyoxal (GO)



II. Methylglyoxal (MGO)




III. 3-deoxyglucosone (3-DG)




B. Markers for advanced glycation endproducts


I. N^ε-(carboxymethyl)lysine (CML)



II. N^ε-(carboxyethyl)lysine (CEL)



III. Pentosidine



IV. Methylglyoxal-derived hydroimidazolone-1 (MG-H1)

