



## LJMU Research Online

**Lomax, DR, Falkingham, PL, Schweigert, G and Jiménez, AP**

**An 8.5 m long ammonite drag mark from the Upper Jurassic Solnhofen Lithographic Limestones, Germany**

<http://researchonline.ljmu.ac.uk/id/eprint/6408/>

### Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Lomax, DR, Falkingham, PL, Schweigert, G and Jiménez, AP (2017) An 8.5 m long ammonite drag mark from the Upper Jurassic Solnhofen Lithographic Limestones, Germany. PLoS ONE, 12 (5). ISSN 1932-6203**

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)

<http://researchonline.ljmu.ac.uk/>

RESEARCH ARTICLE

# An 8.5 m long ammonite drag mark from the Upper Jurassic Solnhofen Lithographic Limestones, Germany

Dean R. Lomax<sup>1\*</sup>, Peter L. Falkingham<sup>2</sup>, Günter Schweigert<sup>3</sup>, Alejandro P. Jiménez<sup>4</sup>

**1** School of Earth and Environmental Sciences, The University of Manchester, Manchester, United Kingdom, **2** Liverpool John Moores University, School of Natural Sciences and Psychology, Liverpool, United Kingdom, **3** Staatliches Museum für Naturkunde, Stuttgart, Germany, **4** CosmoCaixa Museum, Barcelona, Spain

\* [dean.lomax@manchester.ac.uk](mailto:dean.lomax@manchester.ac.uk)



**OPEN ACCESS**

**Citation:** Lomax DR, Falkingham PL, Schweigert G, Jiménez AP (2017) An 8.5 m long ammonite drag mark from the Upper Jurassic Solnhofen Lithographic Limestones, Germany. PLoS ONE 12 (5): e0175426. <https://doi.org/10.1371/journal.pone.0175426>

**Editor:** Steffen Kiel, Naturhistoriska riksmuseet, SWEDEN

**Received:** December 23, 2016

**Accepted:** March 24, 2017

**Published:** May 10, 2017

**Copyright:** © 2017 Lomax et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** 3D model generated via photogrammetry, including source photos taken by DRL and APJ. Model consists of 5,000,000 faces, downsampled from an original model of 20,206,944 faces. Texture resolution is 8000x8000. Available from: <https://doi.org/10.6084/M9.figshare.4479734>. Digital model of the complete drag mark, generated using photogrammetry of 645 photographs, showing close-up of the drag mark and the producer. Available from: <https://doi.org/10.6084/m9.figshare.4502807>.

## Abstract

Trackways and tracemakers preserved together in the fossil record are rare. However, the co-occurrence of a drag mark, together with the dead animal that produced it, is exceptional. Here, we describe an 8.5 m long ammonite drag mark complete with the preserved ammonite shell (*Subplanites rueppellianus*) at its end. Previously recorded examples preserve ammonites with drag marks of < 1 m. The specimen was recovered from a quarry near Solnhofen, southern Germany. The drag mark consists of continuous parallel ridges and furrows produced by the ribs of the ammonite shell as it drifted just above the sediment surface, and does not reflect behaviour of the living animal.

## Introduction

The Upper Jurassic Lithographic Limestones of Solnhofen and surrounding areas in southern Germany are renowned for exceptionally well preserved fossils. Various types of ichnofossil have been described from these Solnhofen-type limestones, and some document the trace and tracemaker together. Recorded examples include arthropods such as limulids and crustaceans [1], bivalves and snails [2,3], and fish [4], which have all been found at the end of their traces. These traces are known as mortichnia and record the last movements of an animal before death [1,5]. The longest mortichnion described to date is a 9.7 m long horseshoe crab trackway [6].

Other examples where an animal and its trace are preserved together include arthropod moulting traces [7], arm crawling crinoids [8], animals inside burrows (e.g. [9,10]), and animals found atop nests (e.g. [11,12]). Such fossils capture a specific moment in time, which is important in understanding and interpreting different behaviours in the fossil record [13].

Dead animals may also leave surface structures behind, but it is rare to find them preserved together. These structures, although in association with a body do not represent behaviour and therefore are better considered as sole marks, drag marks, or other non-biogenic structures, rather than as trace fossils (see [14]). Simple drag marks have been reported from the Solnhofen limestones, including those caused by jellyfish, driftwood, and some ammonite drag and

**Funding:** The authors received no funding for this work.

**Competing interests:** The authors have declared that no competing interests exist.


roll marks [1–3]. Here, we describe an 8.5 m long drag mark created by the shell of a dead ammonite that is preserved at the end of its mark (MCFO 0492, in the permanent collections of the CosmoCaixa Museum, Barcelona, Spain) (Fig 1). The studied specimen was collected in the late 1990s, probably from a quarry in the Langenthalheim Haardt district, near the village of Solnhofen, Bavaria, Germany and prepared in 1998 (Fig 2). The counterpart of the studied specimen exists but it is held in a private collection. The ammonite is identified as *Subplanites rueppellianus* (Quenstedt, 1888), the index fossil of the *rueppellianus* biohorizon of the Upper Jurassic (Tithonian) Hybonotum ammonite Zone [15].

No permits were required for the described study, which complied with all relevant regulations.

### Ammonite tool marks


Ammonite shells are among the most common fossils in the Upper Jurassic Solnhofen Lithographic Limestones. However, they have been rarely in the focus of palaeontological research due to their strongly compressed, unattractive preservation [16]. Recently, it became evident that ammonites provide an important tool for high-resolution biostratigraphy when comparing the ages of the various Solnhofen limestone localities [15,17].

Several ammonite tool marks have been described from the Solnhofen limestones. As reviewed by Maeda and Seilacher [18], the first ammonite touch marks (made by the ammonite shell briefly touching the sediment surface) were initially misidentified as trace fossils produced by vertebrates; such as claw scratches, or ripple effects caused by fish swimming just above the sediment surface. Other ammonite touch marks have been reported from the Jurassic of the Champagnole region, France [19] and from the Upper Cretaceous of the Western Interior Seaway, USA [20]. Marks produced by rolling ammonite shells have also been described. The first description of several ammonite roll marks was by Abel [21], however,



**Fig 1. MCFO 0492, the entire drag mark created by the drifting shell of a dead ammonite (*Subplanites rueppellianus*), with close-up of several portions.** A. The first portion of the drag mark clearly showing two prominent ridges. B. Drag mark showing two prominent ridges with additional faint ridges. C. Drag mark showing four prominent ridges and a gentle curve. D. Drag mark showing numerous prominent ridges, along with the ammonite. Large scale measures 1 m. Small scales measure 10 cm.

<https://doi.org/10.1371/journal.pone.0175426.g001>



**Fig 2. Various Plattenkalk localities of the Franconian and Swabian Alb.** Note the location of Langenaltheim, near Solnhofen, the probable location of MCFO 0492. Reprinted from [35] under a CC BY license, with permission from Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen, original copyright 2007.

<https://doi.org/10.1371/journal.pone.0175426.g002>

Abel misidentified these as trackways produced by turtles and coelacanths. Similar roll marks have been described from Slovenia [22]. Some marks have been produced by bouncing ammonites [1,23], where a rolling mark is punctuated by periodic gaps where the shell has hit the aperture while rolling.



The first report of an ammonite body fossil and drag mark was given by Rothpletz [24], who described a specimen from the Solnhofen limestone consisting of a short mark and fragmentary shell. Since then, similar drag marks from the Solnhofen limestones have been documented by Trusheim [25], Kolb [26], Seilacher [23], Barthel [27], Viohl [28], and Keupp and Schweigert [29]. The latter three are represented by relatively complete ammonites, but with drag marks all less than 1 m in length. Similarly, another specimen that has not been described in the literature (SMF XXX 838a+b, Senckenberg Museum Frankfurt, Germany) comprises a drag mark of less than 2 m, with the ammonite preserved. However, for the entire length of this specimen, the drifting ammonite shell has been dragged through a mass of algal mats.

## Description

The perisphinctid ammonite *Subplanites* is common from the early Tithonian Solnhofen limestones [29]. Perisphinctids are well-known for roll and drag marks [1]. The preserved ammonite of MCFO 0492 measures 114 x 101 mm, although it may have been slightly larger (Fig 3). It is somewhat poorly preserved and a crack runs anteroposteriorly through the ammonite, which has been restored. Another fragment of the lateroposterior portion has also been restored. The damage is probably due to extraction of the specimen when collected. Due to the small size and lack of a lappeted aperture, the ammonite is interpreted as a sub-adult male specimen. Since there is no aptychus (lower Jaw) in the body-chamber, the shell must be from a dead ammonite, where the soft parts including the calcified aptychus have been lost during decay.

The well-preserved drag mark of MCFO 0492 was extracted in numerous pieces, prepared, and pieced together (Fig 1). The total length, measured along the central axis of the preserved drag mark, is 8.5 m. A digital model (<https://doi.org/10.6084/m9.figshare.4479734>) and movie (<https://doi.org/10.6084/m9.figshare.4502807>) of the complete drag mark was generated using photogrammetry [30] from 645 photographs (12mp, Canon PowerShot SX40 HS). The point at which the ammonite first contacted the substrate is not preserved, and thus the drag is incomplete with no indication of how much is missing. We interpret MCFO 0492 as being a drag mark on the primary surface of the ancient sea floor because the ammonite shell, which was probably buoyant with the aid of decaying gases, could only have affected the uppermost layer of the carbonate mud. This is in accordance with the ecdysichnia described by Vallon et al. [7], but in contrast to most trace fossils described by Seilacher [5] who regarded, especially mortichnia, as undertracks.

The drag mark is comprised of a series of ridges and furrows made by the ribs of the ammonite being dragged along the substrate (Fig 3). The drag mark is largely straight, but with minor changes in lateral direction. The preserved start begins with two prominent ridges, with a single furrow. Here, the mark width measures 5.7 mm. From this point, the drag mark width was measured at approximately every 50 cm (Table 1). At one metre, additional ridges created by the ribs of the ammonite appear in the substrate, but they are faint and poorly preserved. Noticeably, at 1.7 m, an additional three ridges are present but disappear again.

Four ridges appear consistently from around 2 m (Fig 1), until about 6.5 m, where five prominent ridges appear. At approximately 7.5 m, only four prominent ridges can be seen, but beyond this point the drag mark preserves five very prominent ridges. It is not until the drag mark is nearly terminating, at 30 cm anterior to the ammonite, where six ridges are present and prominent. At 3 cm from the ammonite, the number of ridges increases to 11, showing that more of the ammonite is clearly in contact with the substrate (Fig 3). Here, the orientation of the ridges turns from being parallel to the long axis of the specimen to almost perpendicular



**Fig 3. The ammonite *Subplanites rueppellianus*, the producer of the drag mark (MCFO 0492).** Note the touch down mark which changes the orientation (and number) of the ridges in the substrate, anteroventral to the ammonite. Scale measures 5 cm.

<https://doi.org/10.1371/journal.pone.0175426.g003>

to it, and increase in number to 18. Here, the ridges and furrows in the substrate mirror the spacing of the ammonite ribs that are well preserved, indicative of a touch down mark (Fig 3).

### Interpretation and review

Seilacher [1] stated that non-living objects, such as the shells of dead ammonites, may be transported along the substrate by waves, winds and currents, which would result in tool marks that can be easily mistaken for animal traces. The first interpretation of an ammonite drag mark was given by Trusheim [25], although he regarded the mark as a trace fossil created by an ammonite crawling along the seafloor. Kolb [26,31] was the first person to interpret an ammonite drag mark as the result of an ammonite swaying along the seafloor, leaving the rib



**Table 1. Measurements of the drag mark width of MCFO 0492, including the number of ridges present.** Measurements marked with an \* are estimated. Estimates were made when the total number of ridges were unclear.

Measurement number	Number of ridges present	Comments	Measurement along drag mark length (cm)	Width of drag mark (in mm)
1	2	Preserved start of drag mark	0	5.7
2	2		50	5.5
3	2	See 3a	100	5.5
3a	?4	Second measurement including faint ridges	100	*11.4
4	2	See 4a	150	5.6
4a	?5/6	Second measurement including faint ridges	150	*17.5
5	5		200	14.8
6	5	Includes very faint ridges	250	13
7	4	Four prominent ridges	300	11.9
8	4		350	12.2
9	4	Some of the ridges are faint	400	12.7
10	4		450	11.8
11	4	Four ridges present, but the mark is wider	500	14.5
12	4		550	13
13	4		600	11.7
14	5		650	16
15	5		700	14.3
16	4	Very prominent ridges	750	12.6
17	5	"	800	16
18	6	"	820	19
19	11	Last measurement taken anterior to the ammonite	845	34.8
20	18	Measurement of ridges in touch down mark lying anteroventral to the ammonite	850	63.2

<https://doi.org/10.1371/journal.pone.0175426.t001>

impressions of the ammonite shell. He also noted lateral structures and interpreted them as imprints left by the tentacles of the dead ammonite. Seilacher [23] provided an interpretation of how different ammonite marks, including drag, roll, bounce and swaying marks, came to existence by waterlogged ammonite shells. Some modes of preservation were illustrated by Seilacher ([1] plate 57). Such scenarios have been simulated by rolling replica ammonites on carbonate mud and clay [1,3,27,32].

Seilacher [1] noted that ammonite roll marks were probably driven by turbidity currents and that some marks were likely caused by ammonites that retained some buoyancy. Kolb [26] suggested that these marks were probably produced in very shallow water. Although MCFO 0492 is exceptionally long but still incomplete, the preserved total length suggests it must have been created by a very calm but constant current, otherwise the ammonite shell would have started rolling. The Solnhofen limestones are considered to have formed in relatively shallow water (20 to 60 m) in a subtropical, probably semi-arid zone [3,33,34]. The seafloor itself was not as hostile as previously thought since there are numerous examples of life such as feeding traces around fish carcasses or moulting traces of lobsters [7]. It is also conceivable that the drag mark formed on a gentle palaeo-slope and the ammonite was moved via gravity (sinking), however, a lack of precise locality data leaves this assertion unsupported.

The number of ridges, and thus the width of the drag mark, differs throughout its length. When more ridges are present, we infer that more of the ammonite must have been in contact with the substrate (Table 1). This suggests that the orientation of the ammonite shell changed subtly as it drifted, with more or fewer ribs of the shell in contact with the substrate as it rotated.

Positioned anteroventral to the ammonite are 18 ridges. The transition of the ridges in the substrate from parallel to almost perpendicular to the longitudinal axis of the drag mark suggests that, before the ammonite came to rest on the substrate, the shell touched the substrate and rotated slightly (Fig 3).

## Conclusions

This exceptionally long fossil was produced by an ammonite shell post-mortem. The shell must have been partially buoyant, firstly because only a small portion of the shell contacts the substrate over the length of the mark, and secondly to be moved by a current that was gentle enough not to disturb the surrounding sediment. It is likely that the ammonite was losing buoyancy over the length of the drag mark, which resulted in eventual loss of all buoyancy and the ammonite falling on its side.

The drag mark of the studied specimen does not represent a mortichnion because it was not created by the animal when alive. Rather, this structure should more correctly be considered a tool mark. As such, behaviour must not be inferred from the drag mark of specimens such as MCFO 0492, and they have to be interpreted as non-biogenic structures produced by physical means [14]. MCFO 0492 represents the hitherto longest fossil drag mark created by a dead animal, complete with the animal preserved at the end.

## Acknowledgments

We thank the staff of the CosmoCaixa Museum for allowing DRL access to study the specimen described in this paper. We greatly appreciate the input from R. Albersdörfer, F. Bachhia and G. Rockers regarding the history of the studied specimen. DRL would also like to thank A. Lord, C. Franz and R. Brocke for providing photographs and information for the specimen held in the collections of the Senckenberg Museum Frankfurt, Germany. Thanks are also due to J. Adams for assisting DRL with photographs and measurements, and J. Nudds and W. Wahl for discussing the studied specimen with DRL. Finally, thanks to L. Vallon and an anonymous reviewer for helpful comments in order to progress the manuscript.

## Author Contributions

**Conceptualization:** DRL.

**Data curation:** DRL PLF.

**Formal analysis:** DRL PLF.

**Funding acquisition:** DRL.

**Investigation:** DRL.

**Methodology:** DRL PLF.

**Project administration:** DRL.

**Software:** PLF.

**Supervision:** DRL APJ.

**Validation:** DRL PLF GS APJ.

**Visualization:** DRL PLF GS APJ.

**Writing – original draft:** DRL.

Writing – review & editing: DRL PLF GS APJ.

## References

1. Seilacher A. Trace Fossil Analysis. Springer Verlag, Heidelberg; 2007.
2. Barthel KW, Solnhofen. Ein Blick in die Erdgeschichte. Ott-Verlag, Thun; 1978.
3. Barthel KW, Swinburne NHM, Conway Morris S. Solnhofen—A Study in Mesozoic Palaeontology. Cambridge University Press, Cambridge; 1990.
4. Schweigert G, Maxwell E, Dietl G. First record of a true mortichnium produced by a fish. *Ichnos*. 2016; 23: 71–76.
5. Seilacher A. Biomats, biofilms, and bioglime as preservational agents for arthropod trackways. *Palaeogeography, Palaeoclimatology, Palaeoecology*. 2008; 270: 252–257.
6. Lomax DR, Racay CA. A long mortichnial trackway of *Mesolimulus walchi* from the Upper Jurassic Solnhofen Lithographic Limestone near Wintershof, Germany. *Ichnos*. 2012; 19: 189–197.
7. Vallon LH, Schweigert G, Bromley RG, Röper M, Ebert M. Ecdysichnia—A new ethological category for trace fossils produced by moulting. *Annales Societatis Geologorum Poloniae*. 2015; 85: 433–444.
8. Neto de Carvalho C, Pereira B, Klompmaker A, Baucon A, Moita JA, Pereira P, et al. Running crabs, walking crinoids, grazing gastropods: behavioral diversity and evolutionary implications of the Cabeço da Ladeira Lagerstätte (Middle Jurassic, Portugal). *Comunicações Geológicas*. 2015; 103: 39–54.
9. Varricchio DJ, Martin AJ, Katsura Y. First trace and body fossil evidence of a burrowing, denning dinosaur. *Proceedings of the Royal Society B: Biological Sciences*. 2007; 274: 1361–1368. <https://doi.org/10.1098/rspb.2006.0443> PMID: 17374596
10. Fernandez V, Abdala F, Carlson KJ, Cook DC, Rubidge BS, Yates A, et al. Synchrotron reveals Early Triassic odd couple: injured amphibian and aestivating therapsid share burrow. *PLoS ONE*. 2013; 8: e64978. <https://doi.org/10.1371/journal.pone.0064978> PMID: 23805181
11. Norell MA, Clark JM, Chiappe LM, Dashzeveg D. A nesting dinosaur. *Nature*. 1995; 378: 774–776.
12. Hastings AK, Hellmund M. Rare *in situ* preservation of adult crocodylian with eggs from the Middle Eocene of Geiseltal, Germany. *Palaios*. 2015; 30: 446–461.
13. Boucot AJ, Poinar GO Jr. Fossil behaviour compendium. CRC Press; 2010.
14. Vallon LH, Rindsberg AK, Martin AJ. The use of the terms trace, mark and structure. *Annales Societatis Geologorum Poloniae*. 2015; 85: 527–528.
15. Schweigert G. Biostratigraphie der Plattenkalke der Südlichen Frankenalb. In: Arratia G, Schultze H-P, Tischlinger H, Viohl G, editors. Solnhofen—Ein Fenster in die Jurazeit. München, Verlag Dr. Friedrich Pfeil; 2015. pp. 57–60
16. Seilacher A, Andalib F, Dietl G, Gocht H. Preservational history of compressed Jurassic ammonites from Southern Germany. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*. 1976; 152: 307–356.
17. Schweigert G. Ammonite biostratigraphy as a tool for dating Upper Jurassic lithographic limestones from South Germany—first results and open questions. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*. 2007; 245: 117–125.
18. Maeda H, Seilacher A. Ammonoid taphonomy. In: Landman NH, Tanabe K, Davis RA, editors. Ammonoid Paleobiology. New York, Plenum Press, 1996.
19. Gaillard C. Cannelures d'érosion et figures d'impact dues à des coquilles d'ammonites à épines (Oxfordien supérieur du Jura français). *Eclogae geologicae Helveticae*. 1977; 70: 701–715.
20. Landman NH, Cobban WA. Ammonite touch marks in Upper Cretaceous (Cenomanian-Santonian) deposits of the Western Interior Seaway. In: Landman NH, Davis RA, Mapes R H, editors. *Cephalopods Present and Past: New insights and fresh perspectives*. Springer; 2007. pp. 396–422.
21. Abel O. Fahrtenstudien. I. Über Schwimmfahrten von Fischen und Schildkröten aus den lithographischen Schiefen Bayerns. *Palaeobiologica*. 1930; 3: 371–412.
22. Summesberger H, Jurkovšek B, Kolar-Jurkovšek T. Rollmarks of soft parts and a possible crop content of Late Cretaceous ammonites from the Slovenian Karst. In: Olóriz F, Rodríguez-Tovar FJ, editors. *Advancing Research on Living and Fossil Cephalopods*. New York, Kluwer Academic/Plenum; 1999. pp. 335–344.
23. Seilacher A. Umlagerung und Rolltransport von Cephalopoden-Gehäusen. *Neues Jahrbuch für Geologie und Paläontologie Monatshefte*. 1963; 1963: 593–615.



24. Rothpletz A. Ueber die Einbettung der Ammoniten in die Solnhofener Schichten. *Abhandlungen der Mathematisch-Physikalischen Klasse der Königlich Bayerischen Akademie der Wissenschaften*. 1909; 24, 311–337.
25. Trusheim F. Eine neue Lebensspur aus den lithographischen Schiefen Süddeutschlands. *Paläontologische Zeitschrift*. 1934; 16: 1–14.
26. Kolb A. Die Ammoniten als Dibranchiata: Untersuchung einer Schleifmarke aus den Solnhofener Schiefen. *Geologische Blätter für Nordost-Bayern und angrenzende Gebiete*. 1961; 11: 1–28.
27. Barthel KW. Zur Entstehung der Solnhofener Plattenkalke (unteres Unterjura). *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und historische Geologie*. 1964; 4: 37–69.
28. Viohl G. Die lithographischen Plattenkalke im engeren Sinne. In: Arratia G, Schultze H-P, Tischlinger H, Viohl G, editors. *Solnhofen—Ein Fenster in die Jurazeit*. München, Verlag Dr. Friedrich Pfeil; 2015. pp. 78–100.
29. Keupp H, Schweigert G. Kopffüßer (Cephalopoda). Ectocochlia: Perlboote (Nautiloidea) und Ammonshörner (Ammonoidea). In: Arratia G, Schultze H-P, Tischlinger H, Viohl G, editors. *Solnhofen—Ein Fenster in die Jurazeit*. München, Verlag Dr. Friedrich Pfeil; 2015. pp. 218–228.
30. Falkingham PL. Acquisition of high resolution three-dimensional models using free, open-source, photogrammetric software. *Palaeontologia Electronica*. 2012; 15: 1–15.
31. Kolb A. Ammoniten-Marken aus den Solnhofener Schiefen bei Eichstätt: ein weiterer Beweis für die Oktopoden-Organisation der Ammoniten. *Geologische Blätter für Nordost-Bayern und angrenzende Gebiete*. 1967; 17: 21–37.
32. Dzulynski S, Sanders JE. Current marks on firm mud bottoms. *Transactions of the Connecticut Academy of Arts and Sciences*. 1962; 42: 57–96.
33. Keupp H, Koch R, Schweigert G, Viohl G. Geological history of the Southern Franconian Alb—the area of the Solnhofen Lithographic Limestone. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*. 2007; 245: 3–21.
34. Selden P, Nudds J. *Evolution of fossil ecosystems*. Manson Publishing Ltd; 2012.
35. Fürsich FT, Mäuser M, Schneider S, Werner W. The Wattendorf Plattenkalk (Upper Kimmeridgian)—a new conservation lagerstätte from the northern Franconian Alb, southern Germany. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*. 2007; 245: 45–58.