



## LJMU Research Online

**Bennett, SJ, Uji, M and Baures, R**

**Asymmetrical time-to-contact error with two moving objects persists across different vertical locations**

<http://researchonline.ljmu.ac.uk/id/eprint/7970/>

### Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Bennett, SJ, Uji, M and Baures, R (2018) Asymmetrical time-to-contact error with two moving objects persists across different vertical locations. Acta Psychologica, 185. pp. 146-154. ISSN 0001-6918**

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)

<http://researchonline.ljmu.ac.uk/>

**Asymmetrical time-to-contact error with two moving objects persists across different vertical separations**

Simon J Bennett<sup>1</sup>, Makoto Uji<sup>1</sup> & Robin Baurès<sup>2</sup>

<sup>1</sup> Research Institute for Exercise & Sport Sciences, Faculty of Science, Liverpool John Moores University, Liverpool, UK

<sup>2</sup> CerCo, Université de Toulouse, CNRS, UPS, France

Address of the corresponding author:

Simon Bennett

Professor of Sensorimotor Neuroscience

School of Sport and Exercise Sciences

Tom Reilly Building, Byrom Street, Liverpool, L3 3AF

0151 904 6257

[s.j.bennett@ljmu.ac.uk](mailto:s.j.bennett@ljmu.ac.uk)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15

### Abstract

When human observers estimate the time-to-contact (TTC) of more than one object there is an asymmetric pattern of error consistent with prioritizing the lead object at the expense of the trail object. Here, we examined TTC estimation in a prediction motion task where two objects moved along horizontal trajectories (5 or 7.5 deg/s) that had different vertical separation, and thus placed specific demands on visuospatial attention. Results showed that participants were able to accurately judge arrival order, irrespective of vertical separation, in all but two conditions where the object trajectories crossed close to the arrival location. Constant error was significantly higher for the object that trailed, as opposed to led, by 250 or 500 ms. Asymmetry in constant error between the lead and trail object was not influenced by vertical separation, and was also evident across a range of arrival times. However, while the lag between the two consecutive TTC estimations was scaled to the actual difference in object arrival times, lag did increase with vertical separation. Taken together, our results confirm that TTC estimation of two moving objects in the prediction motion task suffers from an asymmetrical interference, which is likely related to factors that influence attentional allocation.

**Keywords:** Time-to-Contact; Multiple Objects; Attentional Allocation; Asymmetrical Estimation Error

## 1 **Asymmetrical time-to-contact error with two moving objects persists across different vertical** 2 **locations**

3 An individual's capacity to estimate the arrival time of a single moving object at a specific  
4 location, which is also known as time-to-contact (TTC), has often been assessed with the prediction  
5 motion (PM) task. Having seen the initial part of an object's trajectory prior to occlusion, the participant  
6 is required to make a response (e.g., button press) that coincides with arrival time of the now unseen  
7 object at a specified location. Typically, there is a linear relationship between estimated and actual  
8 TTC, with a slope that is less than one (Caird & Hancock, 1994; Yakimoff, Bocheva, & Mitrani, 1987;  
9 Yakimoff, Mateeff, Ehrenstein, & Hohnsbein, 1993), and a transition from overestimation to  
10 underestimation of TTC around 800-900 ms (Benguigui, Ripoll & Broderik, 2003; Manser & Hancock,  
11 1996; Schiff & Detwiler, 1979; Schiff & Oldak, 1990). The implication is that participants misperceive  
12 the object's actual TTC, and are thus delayed (overestimation) or premature (underestimation) in  
13 pressing the response key. Importantly, however, this linear relationship between estimated and actual  
14 estimated TTC does not hold when the PM task involves two moving objects approaching the same  
15 location (Baurès, Oberfeld & Hecht, 2010; 2011). This situation requires the participant to make two  
16 concurrent TTC estimations and results in an asymmetrical pattern of error. Participants exhibit the  
17 expected level of accuracy for the object that arrives first (i.e., lead object) but significantly  
18 overestimate TTC of the second object when it trails (the lead object) by a short temporal delay  
19 (Baurès, DeLucia, & Olson, 2017).

20 The asymmetrical pattern of error when estimating the arrival time of two objects has been  
21 described with reference to the Psychological Refractory Period (e.g., Pashler, 1994), according to  
22 which the realization of a primary task (i.e., TTC estimation of the lead object) disrupts the completion  
23 of a second task using the same central resource (i.e., TTC estimation of the trail object). As explained  
24 by Baurès et al. (2011), TTC estimation in the PM task requires 4 steps: (1) sensory registration of the  
25 TTC-relevant optical variables, (2) computation of an absolute TTC estimate on the basis of the  
26 information about the objects' motion extracted at step 1, (3) preparation/timing of the motor response  
27 to coincide with the estimated TTC, and (4) initiation and execution of the button press indicating the  
28 estimated TTC. Using a Sperling-like (Sperling, 1960) variation of the PM task where a cue indicated  
29 in advance which object's TTC had to be estimated, Baurès et al. (2011) ruled out the involvement of  
30 steps 3 and 4 in the occurrence of the PRP-like effect (i.e., there was only one motor response and

1 thus attention sharing was not required). It was concluded that when two TTC estimations compete for  
2 the same limited resource during steps 1 or 2, priority is given to the lead object at the expense of the  
3 trail object. In this respect, it is feasible that the asymmetric pattern of error in the PM task is  
4 consistent with over-allocation of attention to the lead object rather than a capacity limitation (Arend,  
5 Johnston & Shapiro, 2006; Martens & Wyble, 2010). By focusing attention on the lead object,  
6 participants are able to extract the necessary information (i.e., position and velocity) for accurate TTC  
7 estimation of that object alone.

8 Unlike the rapid serial visual presentation (RSVP) task typically used to examine the PRP, in  
9 the PM studies described above the two objects were both present, separated by 2 degrees in the  
10 vertical axis, during the initial visible period leading up to occlusion. Therefore, it could be reasoned  
11 that sufficient information regarding the motion of the two objects should have been available for  
12 estimating TTC. However, it is worth noting that the two objects had identical size, shape and color  
13 (e.g., black circles that subtended 1 degree), which when combined with the vertical separation, could  
14 have impacted upon the ability to disambiguate the motion paths and thus estimate TTC. For instance,  
15 it is known that motion perception and pursuit eye movements both initially involve a process that  
16 averages spatially separate inputs (see Heinen & Watamaniuk, 1998), with the weighting influenced by  
17 spatial (Lisberger & Fererra, 1997) and temporal (Marinovic & Wallis, 2011) proximity. This averaging  
18 process is subsequently surpassed by a winner-takes-all response once the decision has been made  
19 to overtly attend to a particular (e.g., lead) object (for the locus of attention during smooth pursuit see  
20 Khan, Lefèvre, Heinen & Blohm, 2010; Van Donkelaar & Drew, 2002). From this point onwards,  
21 pursuit of a moving object places specific demands on visuospatial attention, which can influence  
22 processing of other objects depending on their relative location (Kerzel & Ziegler, 2005; Müller,  
23 Mollenhauer, Rösler, & Kleinschmidt, 2005).

24 In the current study, therefore, we conducted two experiments that examined the influence of  
25 vertical separation between two moving objects on accuracy of TTC estimation. In Experiment 1, we  
26 replicated the object features used in previous work (Baurès et al., 2010; 2011; 2017), whereas in  
27 Experiment 2 we modified the shape of one object in order to facilitate disambiguation. Importantly, the  
28 evolving horizontal separation between the two objects was dependent on their respective velocity and  
29 actual TTC, and thus would not independently account for any differences as a function of vertical  
30 separation. In addition, we ensured that the motion paths (horizontal axis) of the two objects did not

1 cross prior to occlusion, thus minimizing this potential cue regarding arrival order and TTC. Based on  
2 our previous work, we expected that participants would accurately judge arrival order. In addition, we  
3 expected that TTC estimation error would be significantly greater for the object that trailed, as opposed  
4 to led, by a short temporal delay. Given the somewhat mixed findings regarding the effect of relative  
5 location on processing of multiple objects, we did not have a clear expectation regarding the effect of  
6 vertical separation. Shim, Alvarez and Jiang (2009) reported that participants exhibit an impaired  
7 ability to track objects that move in near proximity (i.e.,  $\leq 2$  deg) because of limitations in spatial  
8 resolution of attention. On the other hand, it has been shown that when overt attention is focused on a  
9 moving object, participants are less able to remember the location of stationary targets presented in  
10 the periphery than the fovea (Kerzel & Ziegler, 2005). In the PM task where participants are required  
11 to perform two concurrent TTC estimations, it follows that vertical separation between the two objects  
12 could influence the allocation of attention and thus impact upon TTC estimation error.

13

14

## Experiment 1

### 15 Participants

16 Sixteen male volunteers ( $M_{age} = 21$  years) completed the experiment having provided written  
17 consent. They reported having normal or corrected-to-normal vision, were healthy and without any  
18 known oculomotor abnormalities. Participants were familiarized to the task and procedure, which was  
19 carried out in accordance with the Declaration of Helsinki and approved by the host University local  
20 ethics committee.

### 21 Materials and Procedure

22 Participants were sat in a purpose-built dark room, facing a 22" CRT monitor (Iiyama Vision  
23 Master 505) located on a workbench at a viewing distance of 0.9 m. The head was supported with a  
24 height-adjustable chin rest. Experimental stimuli were generated on a host PC (Dell Precision 670)  
25 using the COGENT toolbox (developed by John Romaya at the Laboratory of Neurobiology at the  
26 Wellcome Department of Imaging Neuroscience) implemented in MATLAB (Mathworks Inc). The  
27 stimuli were presented with a spatial resolution of 1280x1024 pixels and a refresh rate of 85 Hz.  
28 Estimation of TTC was determined from the moment the Y and B keys were pressed on a Razer  
29 Arcosa keyboard (1000 Hz Ultrapolling) with a QWERTY key layout.



1 control for potential effects of condition order, half of the participants completed the three blocks with  
2 the two objects separated by 0.5 deg in the vertical axis followed three blocks with the two objects  
3 separated by 3 deg. The condition order was reversed for the other participants. To control for  
4 potential effects of object position on the vertical axis, the reference object was presented at the lower  
5 or upper vertical position for an equal number of trials. This had the additional advantage of minimizing  
6 any potential influence of hand preference on participants' manual response.

7  
8 Insert Figure 2 About Here

## 8 **Data Analysis**

9 For each participant, the number of correct responses was calculated for each combination of  
10 independent variables: 2 (vertical separation) x 4 ( $\Delta$ TTC) x 2 (reference object velocity) x 2 (distractor  
11 object velocity). The data was then analysed in RStudio (Version 0.99.902) using a generalized linear  
12 mixed model (R Core Team), with a binomial distribution and logistic link function (i.e., binomial logistic  
13 regression). Starting with the full model, we followed an iterative process in order to find the simplest  
14 model that accounted for the highest proportion of variance in the data. We included only those terms  
15 involved with significant main and/or interaction effects, determined by the Wald Chi Square tests (i.e.,  
16  $p > 0.05$ ).

17 We used two approaches for quantifying error in TTC estimation. Similar to Baurès et al.  
18 (2010, 2011, 2017), we first calculated constant error (CE) for each of the two objects relative to their  
19 respective arrival time (i.e., 1900ms for the reference, 1400, 1650, 2150 or 2400 ms for the distractor).  
20 Positive CE indicates an overestimation of the objects arrival time, whereas negative CE indicates an  
21 underestimation of the objects arrival time. Second, the TTC estimation of the lead object was  
22 subtracted from the TTC estimation of the trail object, thus giving a positive measure of lag. Analysis  
23 of lag was important because it permitted us to determine if participants were simply responding to the  
24 trail object at a fixed time after the lead object, or if they were modulating the second response with  
25 respect to the actual difference in arrival times between the two objects. To minimize the influence of  
26 errors in perceiving arrival order on the effects of interest, such trials were excluded from the  
27 calculation of intra-participant mean data. CE and lag were analysed using a linear mixed model (lme4  
28 v1.1-7; Bates, Maechler, Bolker, Walker, Christensen, Singmann & Dai, 2014), following the same  
29 iterative process described above in order to determine the most parsimonious model. Participants  
30 were included as a random effect (i.e., intercept) and the combination of independent variables input

1 as fixed effects: 2 (vertical separation) x 4 ( $\Delta$ TTC) x 2 (reference object velocity) x 2 (distractor object  
2 velocity). The inclusion of random intercepts for each participant was important in order to account for  
3 inter-participant variability in the magnitude of TTC estimation error.

## 4 **Results**

### 5 **Arrival Order**

6 Arrival order was incorrectly perceived in 247 trials of a total 3072 trials (9%), with 1 participant  
7 exhibiting no correct trials in two of the conditions. As shown in Figure 3, participants judged arrival  
8 order of the two objects with similar accuracy irrespective of vertical separation. Mean number of  
9 correct responses was 5.5 (CI.95% = 4.2 : 6.8) in the 0.5 deg condition and 5.5 (CI.95% = 4.2 : 6.8) in  
10 the 3 deg condition. The lack of moderation by vertical separation on the number of correct responses  
11 was confirmed by binomial logistic regression, which indicated no significant contribution from this  
12 factor when it was included as a main or interaction effect. The removal of vertical separation  
13 produced a reduced model that fit the data better than the null model ( $\chi^2_{(15)} = 339.69, p < .001$ ), and  
14 accounted for 47% of the overall variance (conditional R-square). A further reduction to a main effects  
15 only model produced a significantly worse fit of the data ( $\chi^2_{(10)} = 213.9, p < .001$ ) that accounted for  
16 only 29% of the overall variance. Therefore, the reduced model including main and interaction effects  
17 was accepted. As shown in Table 1, Wald Chi Square tests indicated the number of correct responses  
18 was significantly affected by the interaction between  $\Delta$ TTC, reference object velocity and distractor  
19 object velocity. Tukey pairwise comparisons indicated that participants made more errors in judging  
20 arrival order when the lead object moved at 7.5 deg/s and the trail object moved at 5 deg/s with a  
21 delay of 250 ms (reference: M = 3.9; CI.95% = 1.6 : 6.2; distractor: M = 4.4; CI.95% = 2.5 : 6.6).

22 Insert Table 1 About Here

23 Insert Figure 3 About Here

### 24 **CE Reference Object**

25 A full factorial model indicated that vertical separation did not moderate accuracy of estimated  
26 arrival time of the reference object. Mean CE was 529 ms (CI.95% = 283 : 775) in the 0.5 deg vertical  
27 separation condition and 500 ms (CI.95% = 254 : 747) in the 3 deg vertical separation condition. In a  
28 subsequent reduced factorial model, Wald Chi Square tests indicated there were main and interaction  
29 effects for  $\Delta$ TTC, reference object velocity and distractor object velocity (see Table 2 upper rows). The  
30 reduced model produced an equally good fit as the full factorial model ( $\chi^2_{(16)} = 5.73, p > .1$ ) and a

1 significantly better fit of the data than the intercept-only model ( $\chi^2_{(15)} = 336.78, p < .001$ ). The reduced  
2 model accounted for 76% of the overall variance (conditional R-square). Tukey pairwise comparisons  
3 indicated that CE was greater ( $p < .0001$ ) when both the reference and distractor object moved at 7.5  
4 deg/s compared to all other combinations of object velocity. Independent of object velocity, there was  
5 also a significant effect of  $\Delta\text{TTC}$  ( $p < .0001$ ). As can be seen in Figure 4, CE was significantly lower  
6 when the reference object arrived before ( $\Delta\text{TTC} -250$ : M = 257 ms; CI.95% = 9 : 506,  $\Delta\text{TTC} -500$ : M =  
7 242 ms; CI.95% = -7 : 490) compared to after ( $\Delta\text{TTC} 250$ : M = 835 ms; CI.95% = 586 : 1083,  $\Delta\text{TTC}$   
8 500: M = 725 ms; CI.95% = 477 : 974) the distractor object.

9  
10 Insert Table 2 About Here

### 10 **CE Distractor Object**

11 Although the reference and distractor objects had identical visual features and an equal  
12 probability of moving at 5 or 7.5 deg/s in the upper or lower vertical location, TTC of the reference  
13 object was fixed at 1900 ms, whereas TTC of the distractor varied by  $\pm 250$  ms or  $\pm 500$  ms. Therefore,  
14 the pattern of CE described above and reported in previous work (Baurès et al., 2010; 2011; 2017)  
15 could be specific to TTC of the reference object, which was constant across all trial types. To examine  
16 this issue, we repeated the same analysis on CE of the distractor object. The findings for the distractor  
17 object mirrored those of the reference object, thus indicating the effects were not specific to a single  
18 TTC (i.e., 1900ms). Once again we found no significant effect of vertical separation on CE for the  
19 distractor object. Mean CE was 534 ms (CI.95% = 292 : 777) in the 0.5 deg vertical separation  
20 condition and 509 ms (CI.95% = 267 : 752) in the 3 deg vertical separation condition. In a subsequent  
21 reduced factorial model, there were main and interaction effects for  $\Delta\text{TTC}$ , reference object velocity  
22 and distractor object velocity (see Table 2 lower rows). The reduced model produced a significantly  
23 better fit of the data than the intercept-only model ( $\chi^2_{(15)} = 289.57, p < .001$ ) and accounted for 74% of  
24 the overall variance (conditional R-square). CE was greater ( $p < .0001$ ) when the reference and  
25 distractor object both moved at 7.5 deg/s (M = 674 ms; CI.95% = 429 : 919) compared to all other  
26 combinations of object velocity. As can be seen in Figure 4, CE was significantly lower when the  
27 distractor object arrived before ( $\Delta\text{TTC} -250$ : M = 286 ms; CI.95% = 41 : 531,  $\Delta\text{TTC} -500$ : M = 272 ms;  
28 CI.95% = 28 : 517) compared to after ( $\Delta\text{TTC} 250$ : M = 808 ms; CI.95% = 564 : 1053,  $\Delta\text{TTC} 500$ : M =  
29 722 ms; CI.95% = 477 : 966) the reference object.

30  
31 Insert Figure 4 About Here

## 1 Lag between TTC estimations

2 A full factorial model indicated significant main effects for all factors, and an interaction between  
3 reference and distractor object velocities. A subsequent main-effects only model produced a better fit  
4 than the full factorial model ( $\chi^2_{(25)} = 45.29$ ,  $p < .01$ ), as well as the intercept-only model ( $\chi^2_{(6)} = 184.61$ ,  
5  $p < .001$ ). The accepted main-effects model accounted for 63% of the overall variance (conditional R-  
6 square). Tukey pairwise comparisons indicated that lag was shorter ( $p < 0.01$ ) when the objects were  
7 located closer ( $M = 856$  ms;  $CI.95\% = 733 : 986$ ) rather than further ( $M = 903$  ms;  $CI.95\% = 777 : 1030$ )  
8 in the vertical axis. Also, lag was significantly shorter when the temporal separation between the  
9 reference and distractor objects ( $\Delta TTC$ ) was -250 ms ( $M = 798$  ms;  $CI.95\% = 669 : 926$ ) and 250 ms ( $M$   
10  $= 795$  ms;  $CI.95\% = 666 : 924$ ) compared to -500 ms ( $M = 980$  ms;  $CI.95\% = 852 : 1108$ ) and 500 ms  
11 ( $M = 953$  ms;  $CI.95\% = 825 : 1081$ ). Therefore, while participants did not make their second TTC  
12 estimation at a fixed time after the first TTC estimation, perceived lag between the two objects was  
13 modulated by vertical separation (see Figure 4).

## 14 Discussion

15 While temporal proximity is undoubtedly a key factor in the asymmetric pattern of error found  
16 when making two concurrent TTC estimations in the PM task, here we examined if there was also an  
17 influence of vertical separation between the two objects. Consistent with Baurès et al. (2010, 2011,  
18 2017), we found that temporal estimation was significantly more accurate for the lead object than the  
19 trail object. Extending upon previous work, we also found that the overestimation in CE for the trail  
20 object compared to the lead object was similar across a range of TTCs. Analysis of the lag between  
21 the two successive TTC estimations ruled out the possibility that participants gave their second TTC  
22 estimation at a fixed interval after the first estimation. Despite being overestimated per se, lag  
23 increased in accord with the actual difference between the arrival times (i.e., 250 and 500ms).  
24 Interestingly, however, we did find that lag was shorter when the objects were located closer together  
25 in the vertical axis. It is not obvious from the CE data why this effect occurred. For instance, there was  
26 no interaction between  $\Delta TTC$  and vertical separation, whereby participants consistently  
27 underestimated TTC of the lead object and/or overestimated TTC of the trail object. The finding that  
28 vertical separation mediated participants' overestimation of the interval between arrival of successive  
29 objects warrants further investigation.



1 effects of condition order, half of the participants completed the three blocks with the two objects  
2 separated in the vertical axis followed three blocks with the two objects aligned in vertical axis. The  
3 condition order was reversed for the other participants.

## 4 **Results**

### 5 **Arrival Order**

6 Arrival order was incorrectly perceived in 369 trials of a total 3456 trials (approximately 11%),  
7 with 5 participants exhibiting no correct trials in some of the conditions. Analysis of the full model  
8 indicated that arrival order was judged with similar accuracy irrespective of vertical separation. Mean  
9 number of correct responses was 5.3 (CI.95% = 3.8 : 6.8) in the aligned condition and 5.4 (CI.95% =  
10 3.9 : 6.8) in the 3° vertical separation condition. The removal of vertical separation produced a  
11 reduced model that fit the data better than the null model ( $\chi^2_{(15)} = 530.99, p < .001$ ), and accounted for  
12 47% of the overall variance (conditional R-square). A main effects only model was rejected as it  
13 produced a significantly worse fit of the data than the reduced model ( $\chi^2_{(10)} = 333.83, p < .001$ ), and  
14 accounted for only 28% of the overall variance. Wald Chi Square tests on the reduced model indicated  
15 the number of correct responses was significantly affected by  $\Delta$ TTC, as well as the interaction  
16 between reference object velocity and distractor object velocity. Tukey pairwise comparisons indicated  
17 that participants made more errors in judging arrival order when the reference and distractor moved at  
18 a different compared to same velocity. Although not quite reaching conventional levels of significance,  
19 it can be seen in Figure 5 that participants again tended to make more errors in estimating arrival  
20 order when the lead object moved at 7.5 deg/s and the trail object moved at 5 deg/s with a delay of  
21 250 ms (reference: M = 3.5; CI.95% = 1.1 : 5.8; distractor: M = 3.7; CI.95% = 1.4 : 6.1).

22

23 Insert Table 3 About Here

24 Insert Figure 5 About Here

25

### 26 **Reference Object**

27 As can be seen in Figure 6, the results were very similar to those of Experiment 1, with  
28 accuracy of estimated arrival time of both objects being unaffected by vertical separation. Mean CE  
29 was 420 ms (CI.95% = 34 : 806) in the aligned condition and 402 ms (CI.95% = 16 : 788) in the 3 deg  
30 vertical separation condition. A reduced model (see Table 4) not including vertical separation

1 produced a significantly better fit of the data than the intercept-only model ( $\chi^2_{(15)} = 324.33$ ,  $p < .001$ )  
2 and accounted for 91% of the overall variance (conditional R-square). Observation of the group mean  
3 data (see Figure 6), and the outcome of Tukey pairwise comparisons, indicated that CE was greatest  
4 ( $p < .0001$ ) when the reference and distractor object both moved at 7.5 deg/s ( $M = 570$  ms;  $CI.95\% =$   
5  $183 : 957$ ). Independent of object velocity, there was also a significant effect of  $\Delta TTC$  ( $p < .0001$ ). As  
6 can be seen in Figure 5, CE was significantly lower when the reference object arrived before ( $\Delta TTC -$   
7  $250$ :  $M = 206$  ms;  $CI.95\% = -181 : 593$ ,  $\Delta TTC -500$ :  $M = 212$  ms;  $CI.95\% = -175 : 599$ ) compared to  
8 after ( $\Delta TTC 250$ :  $M = 650$  ms;  $CI.95\% = 263 : 1037$ ,  $\Delta TTC 500$ :  $M = 577$  ms;  $CI.95\% = 190 : 964$ ) the  
9 distractor object.

10 Insert Table 4 About Here

### 11 **Distractor Object**

12 The findings for the distractor object mirrored those of the reference object. There were no  
13 significant main or interaction effects involving vertical separation. Mean CE was 424 ms ( $CI.95\% = 45$   
14  $: 804$ ) in the aligned condition and 412 ms ( $CI.95\% = 32 : 791$ ) in the 3 deg vertical separation  
15 condition. In a subsequent reduced factorial model, there were main and interaction effects for  $\Delta TTC$ ,  
16 reference object velocity and distractor object velocity (see Table 4). The reduced model produced a  
17 significantly better fit of the data than the intercept-only model ( $\chi^2_{(15)} = 243.43$ ,  $p < .001$ ) and  
18 accounted for 88% of the overall variance (conditional R-square). CE was greater ( $p < .0001$ ) when  
19 the reference and distractor object both moved at 7.5 deg/s ( $M = 576$  ms;  $CI.95\% = 195 : 956$ ). CE  
20 was significantly lower when the distractor object arrived before ( $\Delta TTC -250$ :  $M = 238$  ms;  $CI.95\% = -$   
21  $143 : 619$ ,  $\Delta TTC -500$ :  $M = 220$  ms;  $CI.95\% = -161 : 601$ ) compared to after ( $\Delta TTC 250$ :  $M = 631$  ms;  
22  $CI.95\% = 250 : 1012$ ,  $\Delta TTC 500$ :  $M = 583$  ms;  $CI.95\% = 202 : 963$ ) the reference object (see Figure  
23 6).

24 Insert Figure 6 About Here

### 25 **Lag between TTC estimations**

26 A full factorial model indicated significant main effects for all factors, but no interactions. A  
27 main-effects only model produced an equal fit as the full factorial model ( $\chi^2_{(25)} = 12.49$ ,  $p > .1$ ), and a  
28 significantly better fit than the intercept-only model ( $\chi^2_{(6)} = 123.84$ ,  $p < .001$ ). The reduced model  
29 accounted for 60% of the overall variance (conditional R-square). Tukey pairwise comparisons  
30 indicated that lag was shorter ( $p < .01$ ) when the objects were aligned ( $M = 729$  ms;  $CI.95\% = 595 :$

1 862) rather than separated ( $M = 801$  ms;  $CI.95\% = 668 : 934$ ) in the vertical axis. Also, lag was  
2 significantly shorter when the temporal separation between the reference and distractor objects  
3 ( $\Delta TTC$ ) was  $-250$  ms ( $M = 672$  ms;  $CI.95\% = 536 : 807$ ) and  $250$  ms ( $M = 662$  ms;  $CI.95\% = 526 :$   
4  $797$ ) compared to  $-500$  ms ( $M = 869$  ms;  $CI.95\% = 733 : 1004$ ) and  $500$  ms ( $M = 857$  ms;  $CI.95\% =$   
5  $721 : 992$ ). Again, while participants did not make their second TTC estimation at a fixed time after the  
6 first TTC estimation, perceived lag between the two objects was modulated by vertical separation (see  
7 Figure 6).

## 8 **Discussion**

9 We compared TTC estimations when two objects with different features (i.e., circle and  
10 square) moved on horizontal trajectories that were aligned or separated in the vertical axis. Our results  
11 confirmed the presence of an asymmetric pattern of error (i.e., PRP-like effect), with more accurate  
12 TTC estimation for the lead object than the trail object. This was evident across a range of absolute  
13 arrival times and occurred irrespective of vertical separation. Analysis of the lag between the two  
14 successive TTC estimations confirmed that participants moderated their response in accord with the  
15 difference between the object arrival times. However, while participants waited on average and extra  
16  $170$  ms between their two responses when  $\Delta TTC$  was  $500$  compared to  $250$  ms, lag per se was  
17 largely overestimated. As can be seen in the CE data, this was predominantly due to overestimating  
18 TTC of the trail object. We also found that vertical separation moderated lag such that it was shorter  
19 when the objects were aligned. Observation of the CE data indicated that this was not due to a  
20 systematic misestimation in TTC of either the lead or trail object. It would seem, therefore, that vertical  
21 separation between two moving objects does exert a small but significant on the delay between  
22 successive TTC estimations.

## 23 **General Discussion**

24 During our daily interactions within our normal surrounds, it is not unusual to make TTC  
25 estimations regarding the approach of more than one object. For instance, while cycling in a town or  
26 city one might follow the motion of other road users as they approach a junction or several pedestrians  
27 while walking along a busy street (Gould, Poulter, Helman, & Wann, 2012; Baurès, Oberfeld, Tournier,  
28 Hecht, & Cavallo, 2014). Such behaviours require attention to be allocated to multiple objects that can  
29 have different spatiotemporal properties and physical features (for a commentary on different  
30 attentional models see Tombu & Seiffert, 2008). Notably, while individuals are able to keep track of the

1 spatial evolution of multiple objects with reasonable accuracy (Cavanagh & Alvarez, 2005; Pylyshyn &  
2 Storm, 1988), there is a systematic pattern of error when estimating TTC of two objects at a known  
3 location. Specifically, it has been shown using a prediction motion (PM) task that TTC estimation error  
4 of the lead object is similar to single-object conditions, whereas TTC estimation error of the trail object  
5 is significantly increased when it arrives after a short delay (Baurès et al., 2010, 2011, 2017). This  
6 pattern of error is akin to the well-known Psychological Refractory Period (PRP), which is thought to  
7 be a result of attentional allocation rather than a capacity limitation (Arend et al., 2006; Martens &  
8 Wyble, 2010). In the PM task, for example, it is possible that participants increase overt attentional  
9 focus on the lead object, to the detriment of the trail object, because the former demands the more  
10 behaviorally urgent response (Lin, Franconeri, & Enns 2008).

11         The current study compared TTC estimation in two experiments where the two moving objects  
12 had different vertical separation. The logic was that vertical separation might modulate allocation of  
13 attention between the lead and trail object (He et al., 1997; Shim et al., 2008), thereby influencing the  
14 pattern of TTC estimation error. In both experiments, each with different groups of participants, we  
15 found the expected asymmetrical error in TTC estimation (Baurès et al., 2010, 2011). Participants  
16 exhibited much larger error in estimating TTC of the trail object compared to the lead object when they  
17 had close temporal proximity (i.e., <750 ms; Baurès et al., 2017). In addition, we showed here for the  
18 first time within a single study that this effect was not specific to a single TTC. However, and  
19 somewhat contrary to our initial expectations, we found no effect of vertical separation between the  
20 two objects on their respective constant error. The next part of our analysis examined if participants  
21 made their second response at a constant delay after the first response, such as might be a strategy if  
22 they were only able to determine arrival order. We ruled out this explanation by showing that  
23 participants modulated the lag between successive responses in accord with the difference between  
24 the object arrival times (i.e., 250 or 500 ms). In other words, participants showed evidence of  
25 estimating TTC of the two objects and not TTC of the lead object only. That said, lag per se was  
26 overestimated by approximately 300-600 ms, predominantly due to greater error in response to the  
27 trail object. Moreover, overestimation was reduced when the two objects were close together (0.5 deg  
28 in Experiment 1) or aligned (Experiment 2) in the vertical axis. Despite being of small amplitude (i.e.,  
29 approximately 60 ms), the effect of vertical separation on lag was present in both experiments (with  
30 different participants) and was not due to a systematic misestimation of either the lead or trail object.

1           How, then, do we interpret the combined findings for constant error and lag between  
2 successive TTC estimations? To answer this question, we start from the position that TTC estimation  
3 in the PM task involves several stages that are influenced by attention. As described above, we  
4 suggest that the asymmetrical error in TTC estimation is consistent with participants increasing  
5 attention on the lead object because it demanded the more behaviorally urgent response (Lin et al.,  
6 2008). An increase in attention on the lead object likely coincides with gaze location. For instance, we  
7 have previously shown that TTC estimation is more accurate when participants are permitted to  
8 pursue the moving object (Bennett, Baurès, Hecht & Benguigui, 2010), and that having been cued to  
9 overtly pursue the trail object during the initial visible period, participants shift gaze to the lead object  
10 during occlusion (Baurès, Bennett & Causer, 2016). Although yet to be confirmed, we suspect that  
11 having made their first response (TTC estimation) with gaze located on the lead object, participants in  
12 the current study shifted overt attention to the trail object, which added a small but significant delay  
13 when the two objects were located further apart in the vertical axis. A shift of overt attention could add  
14 delay through a combination of saccadic programming and interrupted processing of the trail object  
15 due to saccadic suppression. The implication is that the effect of vertical separation was a  
16 consequence of attentional allocation that occurred at a later stage than the perception of information  
17 required for accurate TTC estimation. It is important to recognize, however, that we were careful to  
18 ensure the horizontal trajectories did not cross during the initial visible period, thereby eliminating this  
19 simple cue to arrival order. Had this not been the case, estimation of TTC may have been mediated by  
20 vertical separation. For example, the crossing of horizontal paths during the initial visible period might  
21 exert a stronger influence on motion processing (e.g., distraction, vector averaging, assimilation) that  
22 underpins perception of TTC if the two objects are located close together or aligned in the vertical axis.

23           When modifying velocity and TTC of two objects in the PM task, there will be a unique change  
24 in horizontal separation between the evolving trajectories (see Figure 2). Although this spatial variable  
25 would not independently account for any differences as a function of vertical separation in the current  
26 study, the influence of horizontal separation on accuracy of arrival order, and TTC estimation error,  
27 was indirectly considered in our regression modelling. For estimation of arrival order in experiment 1,  
28 the significant three-way interaction between velocity of the two objects and  $\Delta$ TTC provided some  
29 indication that a spatial variable could have been involved for specific combinations of our parameters.  
30 For instance, participants made more errors in judging arrival order in trials where the lead object (i.e.,

1 reference or distractor) moved at 7.5 deg/s and the trail object moved at 5 deg/s with a 250 ms delay.  
2 A similar effect was evident in experiment 2, although the three-way interaction did not quite reach the  
3 conventional level of significance. Notably, however, 3 of the 18 participants did in fact exhibit no  
4 correct trials in these two conditions. It is possible, therefore, that participants failed to perceive that  
5 the horizontal motion paths of the two objects crossed late during the occlusion interval (see Figure 2),  
6 and thus at a time when the ability to extrapolate object motion has begun to deteriorate (Bennett &  
7 Benguigui, 2016; Tanaka, Worringham & Kerr, 2009; Wexler & Klam, 2001). Consequently, they may  
8 have incorrectly estimated that the formerly closer object (in space) also had the shorter TTC. As often  
9 found in children (Benguigui, Broderick, Baurès, & Amorim, 2008; Keshavarz, Landwehr, Baurès,  
10 Oberfeld, Hecht, & Benguigui, 2010), one explanation is that on some trials adult participants used a  
11 heuristic (e.g., distance) that did not provide reliable information to accurately estimate TTC (DeLucia,  
12 2004). Intermittent use of either a temporal or spatial variable is supported by the finding that there  
13 was no effect of this particular combination of parameters on TTC estimation error or lag between TTC  
14 estimations (i.e., errorful trials omitted). It will be interesting in future work to compare a wider range  
15 of conditions in which the motion paths cross at different times during the occlusion period.

16 Together with the results of our recent series of studies, here we confirmed that participants  
17 are unable to perform two concurrent TTC estimations with similarly high accuracy. Consistent with  
18 over-allocation of attention on the most salient object, participants systematically overestimated TTC  
19 of the trail object. Although recent work has indicated that this asymmetric pattern of error is not  
20 identical to the PRP effect exhibited in the RSVP task (for a detailed discussion see Baurès et al.,  
21 2017), these findings could have some important practical consequences. For instance, there could be  
22 some value in making participants aware that there is a tendency to over-allocate attention to the lead  
23 of two approaching objects, and then provide training or stimulus conditions that encourage a more  
24 even allocation of attention. This might be important in numerous ball-sport situations, where the  
25 player has to estimate TTC of the ball while concurrently estimating TTC between themselves and  
26 several opponents. Novice players are known to “ball watch” and are thus less aware of their  
27 surroundings. If the novice player does not correctly estimate the closing gap (and thus TTC) between  
28 themselves and surrounding players, this could result in a collision or give an advantage to the  
29 opposition. A similar situation could occur for the novice driver, who has to decide whether or not there  
30 is enough time to exit a junction when there are two cars approaching from the opposite direction. By

1 recognizing and then over-allocating attention to the car that will pass the junction first, the driver might  
2 not update their TTC estimate of the second car, resulting in an inappropriately timed behaviour.  
3 Future studies with stimuli that are more representative of real world settings are required to confirm  
4 whether over-allocating attention on the lead object does indeed occur outside of the laboratory PM  
5 task in situations where asymmetrical estimation of TTC could have serious consequences.

6

7

### References

8 Arend, I., Johnston, S., & Shapiro, K. (2006). Task-irrelevant visual motion and flicker attenuate the  
9 attentional blink. *Psychonomic Bulletin & Review*, *13*(4), 600-607.

10 Bates, D., Maechler, M., Bolker, B., Walker, S., Christensen, R. H. B., Singmann, H., & Bolker, M. B.  
11 (2012). Package 'lme4'. CRAN. *R Foundation for Statistical Computing, Vienna, Austria*.

12 Baurès, R., Bennett, S.J., & Causer, J. (2015). Temporal estimation with two moving objects: overt  
13 and covert pursuit. *Experimental Brain Research*, *233*(1), 253-261.

14 Baurès, R., DeLucia, P.R., & Olson, M. (2017). Asymmetric interference in concurrent time-to-contact  
15 estimation: Cousin or twin of the psychological refractory period effect? *Attention, Perception  
16 & Psychophysics*, *79*(2), 698-711. doi:10.3758/s13414-016-1244-y

17 Baurès, R., Oberfeld, D., & Hecht, H. (2010). Judging the contact-times of multiple objects: Evidence  
18 for asymmetric interference. *Acta Psychologica*, *134*(3), 363-371.

19 doi:10.1016/j.actpsy.2010.03.009

20 Baurès, R., Oberfeld, D., & Hecht, H. (2011). Temporal-range estimation of multiple objects: Evidence  
21 for an early bottleneck. *Acta Psychologica*, *137*(1), 76-82. doi:10.1016/j.actpsy.2011.03.002.

22 Baurès, R., Oberfeld, D., Tournier, I., Hecht, H., & Cavallo, V. (2014). Arrival-time judgments on  
23 multiple-lane streets: The failure to ignore irrelevant traffic. *Accident Analysis & Prevention*,  
24 *65*, 72-84.

25 Benguigui, N., & Bennett, S. J. (2010). Ocular pursuit and the estimation of time-to-contact with  
26 accelerating objects in prediction motion are controlled independently based on first-order  
27 estimates. *Experimental Brain Research*, *202*(2), 327-339.

28 Benguigui, N., Broderick, M. P., Baurès, R. and Amorim, M.-A. (2008). Motion prediction and the  
29 velocity effect in children. *British Journal of Developmental Psychology*, *26*, 389–407.

30 doi:10.1348/026151008X295146

- 1 Bennett, S. J., Baurès, R., Hecht, H., & Benguigui, N. (2010). Eye movements influence estimation of  
2 time-to-contact in prediction motion. *Experimental Brain Research*, 206(4), 399-407.
- 3 Bennett, S. J., & Benguigui, N. (2016). Spatial estimation of accelerated stimuli is based on a linear  
4 extrapolation of first-order information. *Experimental Psychology*, 63(2), 98-106. doi:  
5 10.1027/1618-3169/a000318
- 6 Blaser, E., Pylyshyn, Z. W., & Holcombe, A. O. (2000). Tracking an object through feature space.  
7 *Nature*, 408(6809), 196-199.
- 8 Caird, J. K., & Hancock, P. A. (1994). The perception of arrival time for different oncoming vehicles at  
9 an intersection. *Ecological Psychology*, 6, 83-109.
- 10 Cavanagh, P., & Alvarez, G. A. (2005). Tracking multiple targets with multifocal attention. *Trends in*  
11 *Cognitive Sciences*, 9(7), 349-354.
- 12 DeLucia, P. R. (2004). Multiple sources of information influence time-to-contact judgments: Do  
13 heuristics accommodate limits in sensory and cognitive processes? *Advances in Psychology*,  
14 135, 243-285.
- 15 Gould, M., Poulter, D. R., Helman, S., & Wann, J. P. (2012). Errors in judging the approach rate of  
16 motorcycles in nighttime conditions and the effect of an improved lighting configuration,  
17 *Accident Analysis & Prevention*, 45, 432-437. doi: 10.1016/j.aap.2011.08.012.
- 18 He, S., Cavanagh, P., & Intriligator, J. (1997). Attentional resolution. *Trends in Cognitive Sciences*,  
19 1(3), 115-121.
- 20 Heinen, S. J., & Watamaniuk, S. N. (1998). Spatial integration in human smooth pursuit. *Vision*  
21 *Research*, 38(23), 3785-3794.
- 22 Johnson, C.A., Keltner, J. L., & Balestrery, F. (1978). Effects of target size and eccentricity on visual  
23 detection and resolution. *Vision Research*, 18(9),1217-1222.
- 24 Kerzel, D., & Ziegler, N. E. (2005). Visual short-term memory during smooth pursuit eye movements.  
25 *Journal of Experimental Psychology: Human Perception and Performance*, 31(2), 354.
- 26 Keshavarz, B., Landwehr, K., Baurès, R., Oberfeld, D., Hecht, H., & Benguigui, N. (2010). Age-  
27 correlated incremental consideration of velocity information in relative time-to-arrival  
28 judgments. *Ecological Psychology*, 22(3), 212-221.
- 29 Khan, A. Z., Lefèvre, P., Heinen, S. J., & Blohm, G. (2010). The default allocation of attention is  
30 broadly ahead of smooth pursuit. *Journal of Vision*, 10(13), 1-17.

- 1 Lin, J.Y., Franconeri, S., & Enns, J.T. (2008) Objects on a collision path with the observer demand  
2 attention. *Psychological Science*, *19*, 686–692.
- 3 Lisberger, S. G., & Ferrera, V. P. (1997). Vector averaging for smooth pursuit eye movements initiated  
4 by two moving targets in monkeys. *Journal of Neuroscience*, *17*(19), 7490-7502.
- 5 Manser, M.P., & Hancock, P.A. (1996). Influence of approach angle on estimates of time-to-contact.  
6 *Ecological Psychology*, *8*, 71–99.
- 7 Marinovic, W., & Wallis, G. (2011). Visual attention affects temporal estimation in anticipatory motor  
8 actions. *Experimental Brain Research*, *212*, 613–621.
- 9 Martens, S., & Wyble, B. (2010). The attentional blink: Past, present, and future of a blind spot in  
10 perceptual awareness. *Neuroscience and Biobehavioural Reviews*, *34*, 947-957.
- 11 McKee, S.P., & Nakayama, K. (1984). The detection of motion in the peripheral field. *Vision Research*,  
12 *24*(1), 25-32.
- 13 Müller, N. G., Mollenhauer, M., Rösler, A., & Kleinschmidt, A. (2005). The attentional field has a  
14 Mexican hat distribution. *Vision Research*, *45*(9), 1129-1137.
- 15 Pashler, H. (1994). Dual-task interference in simple tasks: Data and theory. *Psychological Bulletin*,  
16 *116*(2), 220–244. doi:10.1037/0033-2909.116.2.220.
- 17 Pylyshyn, Z. W., & Storm, R. W. (1988). Tracking multiple independent targets: Evidence for a parallel  
18 tracking mechanism. *Spatial Vision*, *3*(3), 179-197.
- 19 Schiff, W., & Oldak, R. (1990). Accuracy of judging time to arrival: Effects of modularity, trajectory, and  
20 gender. *Journal of Experimental Psychology: Human Perception and Performance*, *16*, 303–  
21 316.
- 22 Schiff, W., & Detwiler, M. L. (1979). Information used in judging impending collision. *Perception*, *8*(6),  
23 647-658. doi:10.1068/p080647.
- 24 Shim, W. M., Alvarez, G. A., & Jiang, Y. V. (2008). Spatial separation between targets constrains  
25 maintenance of attention on multiple objects. *Psychonomic Bulletin & Review*, *15*(2), 390-397.
- 26 Sperling, G. (1960). The information available in brief visual presentations. *Psychological Monographs:*  
27 *General and Applied*, *74*(11), 1–30.
- 28 Tanaka, H., Worringham, C., & Kerr, G. (2009). Contributions of vision–proprioception interactions to  
29 the estimation of time-varying hand and target locations. *Experimental Brain Research*, *195*,  
30 371-382.

- 1 Tombu, M., & Seiffert, A. E. (2008). Attentional costs in multiple-object tracking. *Cognition*, 108(1), 1-  
2 25.
- 3 Van Donkelaar, P., & Drew, A. S. (2002). The allocation of attention during smooth pursuit eye  
4 movements. *Progress in Brain Research*, 140, 267-277.
- 5 Wexler, M., & Klam, F. (2001). Movement prediction and movement production. *Journal of*  
6 *Experimental Psychology: Human Perception and Performance*, 27, 48-64.
- 7 Yakimoff, N., Bocheva, N., & Mitrani, L. (1987). A linear model for the response time in motion  
8 prediction. *Acta Neurobiologiae Experimentalis*, 47, 55-62.
- 9 Yakimoff, N., Mateeff, S., Ehrenstein, W. H., & Hohnsbein, J. (1993). Motion extrapolation  
10 performance: a linear model approach. *Human Factors*, 35, 501-510.

**Table 1:** Type II Wald Chi-Square tests for the fixed effects included in the binomial logistic regression on number of correct responses in experiment 1. The accepted reduced model is shown. Factors included were: Delta.TTC ( $\Delta$ TTC); Vref (reference object velocity); Vdis (distractor object velocity).

	Chisq	df	p value	
Delta.TTC	52.85	3	0.000	***
Vref	0.78	1	0.378	
Vdis	6.48	1	0.011	*
Delta.TTC:Vref	3.24	3	0.356	
Delta.TTC:Vdis	9.58	3	0.023	*
Vref:Vdis	75.73	1	0.000	***
Delta.TTC:Vref:Vdis	16.06	3	0.001	**

**Table 2:** Type II Wald Chi-Square tests for the fixed effects included in linear mixed model regression on constant error of the reference (upper rows) and distractor (lower rows) object in experiment 1. The accepted reduced model is shown. Factors included were: Delta.TTC ( $\Delta$ TTC); Vref (reference object velocity); Vdis (distractor object velocity).

	Chisq	df	p value	
<u>Reference</u>				
Delta.TTC	395.19	3	0.000	***
Vref	19.83	1	0.000	***
Vdis	17.12	1	0.000	***
Delta.TTC:Vref	10.59	3	0.014	*
Delta.TTC:Vdis	6.45	3	0.092	.
Vref:Vdis	7.96	1	0.005	**
Delta.TTC:Vref:Vdis	10.82	3	0.013	*
<u>Distractor</u>				
Delta.TTC	320.06	3	0.000	***
Vref	19.79	1	0.000	***
Vdis	17.23	1	0.000	***
Delta.TTC:Vref	7.48	3	0.058	
Delta.TTC:Vdis	10.38	3	0.016	*
Vref:Vdis	6.08	1	0.014	*
Delta.TTC:Vref:Vdis	0.73	3	0.865	

**Table 3:** Type II Wald Chi-Square tests for the fixed effects included in the binomial logistic regression on number of correct responses in experiment 2. The accepted reduced model is shown. Factors included were: Delta.TTC ( $\Delta$ TTC); Vref (reference object velocity); Vdis (distractor object velocity).

	Chisq	df	p value	
Delta.TTC	89.81	3	0.000	***
Vref	18.90	1	0.000	***
Vdis	2.00	1	0.158	
Delta.TTC:Vref	1.05	3	0.790	
Delta.TTC:Vdis	3.73	3	0.292	
Vref:Vdis	151.61	1	0.000	***
Delta.TTC:Vref:Vdis	7.22	3	0.065	.

**Table 4:** Type II Wald Chi-Square tests for the fixed effects included in linear mixed model regression on constant error of the reference (upper rows) and distractor (lower rows) object in experiment 2. The accepted reduced model is shown. Factors included were: Delta.TTC ( $\Delta$ TTC); Vref (reference object velocity); Vdis (distractor object velocity).

	Chisq	df	p value	
<i>Reference</i>				
Delta.TTC	346.06	3	0.000	***
Vref	10.69	1	0.001	**
Vdis	39.81	1	0.000	***
Delta.TTC:Vref	6.81	3	0.078	.
Delta.TTC:Vdis	2.40	3	0.494	
Vref:Vdis	23.06	1	0.000	***
Delta.TTC:Vref:Vdis	1.81	3	0.614	
<i>Distractor</i>				
Delta.TTC	230.64	3	0.000	***
Vref	8.92	1	0.003	**
Vdis	22.47	1	0.000	***
Delta.TTC:Vref	0.72	3	0.868	
Delta.TTC:Vdis	6.31	3	0.098	.
Vref:Vdis	22.29	1	0.000	***
Delta.TTC:Vref:Vdis	4.47	3	0.215	

## 1 **Figure Legends**

2

3 **Figure 1.** Representation of the visual stimulus in experiment 1 (left panel) and experiment 2 (right  
4 panel). A. The visual scene initially contains two stationary objects and an arrival line (full black  
5 rectangle). The two dashed rectangles represent the forthcoming occlusion of the objects, but were  
6 not visible to the participants during the experiment. The two objects then move rightwards for 600 ms  
7 toward the arrival line with a velocity of either 5 or 7.5 deg/s. B. Both objects are occluded at the same  
8 time, with the reference object reaching the arrival line after 1900 ms and the distractor object arriving  
9 either earlier or later by 250 or 500 ms. C. Participants press a key with the right and left index finger  
10 to coincide with the moment each object would have made contact with the arrival line. In experiment  
11 1, the two objects are separated in the vertical axis by 0.5 or 3 deg. In experiment 2, the two objects  
12 are aligned or separated in the vertical axis by 3 deg (NB. not shown to avoid replication). To avoid  
13 feature assimilation in experiment 2, the two objects are either a circle or square.

14

15 **Figure 2.** Horizontal object position as a function of time. The solid black and red lines depict the  
16 reference object, which has a TTC of 1900 ms and was presented in every trial. The broken black  
17 lines depict the distractor object, which has  $\Delta$ TTC of  $\pm$  250 or 500 ms. Panel A shows all position  
18 trajectories that included the 5 deg/s object. Panel B shows all position trajectories that included the  
19 7.5 deg/s object. The light grey bar in each panel represents the onset of occlusion (600 ms) and  
20 arrival time of the reference object. NB. None of the objects became visible after they reached the  
21 arrival line. The double horizontal lines represent the location of the arrival line, which was constant at  
22 11 deg from screen centre.

23

24 **Figure 3.** Group mean number of correct responses in experiment 1 as a function of  $\Delta$ TTC and  
25 velocity of the two objects (reference, distractor). NB. Negative  $\Delta$ TTC indicates that the reference was  
26 the lead object, whereas positive  $\Delta$ TTC indicates that the reference was the trail object.

27

28 **Figure 4.** Group mean CE ( $\pm$  95% CI) as a function of Delta TTC, Vertical Separation (Close = 0.5 deg;  
29 Far = 3 deg) for the reference object (squares on black and grey lines) and distractor object (triangles  
30 on black and grey lines). Delta TTC is expressed relative to the reference object. Accordingly, -500 and  
31 -250 ms indicate the reference was the lead object and the distractor was the trail object. Conversely,  
32 500 and 250 ms indicate the reference was the trail object and the distractor was the lead object. NB.  
33 To aid interpretation of the factor, Vertical Separation, an offset has been introduced on the horizontal  
34 axis.

35

36 **Figure 5.** Group mean number of correct responses in experiment 2 as a function of  $\Delta$ TTC and  
37 velocity of the two objects (reference, distractor). NB. Negative  $\Delta$ TTC indicates that the reference was  
38 the lead object, whereas positive  $\Delta$ TTC indicates that the reference was the trail object.

39

1 **Figure 6.** Group mean CE ( $\pm$  95% CI) as a function of Delta TTC, Vertical Separation (Close = 0.5 deg;  
2 Far = 3 deg) for the reference object (squares on black and grey lines) and distractor object (triangles  
3 on black and grey lines). Delta TTC is expressed relative to the reference object. Accordingly, -500 and  
4 -250 ms indicate the reference was the lead object and the distractor was the trail object. Conversely,  
5 500 and 250 ms indicate the reference was the trail object and the distractor was the lead object. NB.  
6 To aid interpretation of the factor, Vertical Separation, an offset has been introduced on the horizontal  
7 axis.




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6