



LJMU Research Online

Pomeroy, E, Wells, JCK, Stanojevic, S, Jaime Miranda, J, Moore, LG, Cole, TJ and Stock, JT

Surname-Inferred Andean Ancestry Is Associated with Child Stature and Limb Lengths at High Altitude in Peru, but not at Sea Level

<http://researchonline.ljmu.ac.uk/id/eprint/8450/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Pomeroy, E, Wells, JCK, Stanojevic, S, Jaime Miranda, J, Moore, LG, Cole, TJ and Stock, JT (2015) Surname-Inferred Andean Ancestry Is Associated with Child Stature and Limb Lengths at High Altitude in Peru, but not at Sea Level. American Journal of Human Biology. 27 (6). pp. 798-806. ISSN 1042-

LJMU has developed [LJMU Research Online](#) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

Original Research Article

Surname-Inferred Andean Ancestry Is Associated with Child Stature and Limb Lengths at High Altitude in Peru, but not at Sea Level

EMMA POMEROY,^{1,2*} JONATHAN C.K. WELLS,^{2,3} SANJA STANOJEVIC,⁴ J. JAIME MIRANDA,⁵ LORNA G. MOORE,⁶ TIM J. COLE,⁷ AND JAY T. STOCK²¹Newnham College, University of Cambridge, Cambridge, United Kingdom²Division of Biological Anthropology, Department of Archaeology and Anthropology, University of Cambridge, United Kingdom³Childhood Nutrition Research Centre, UCL Institute of Child Health, University College London, London, United Kingdom⁴Division of Respiratory Medicine, The Hospital for Sick Children, Toronto, Ontario, Canada⁵CRONICAS Center of Excellence in Chronic Diseases and Department of Medicine, School of Medicine, Universidad Peruana Cayetano Heredia, Lima, Peru⁶Department of Obstetrics / Gynecology, University of Colorado Denver, Aurora, Colorado⁷Population, Policy and Practice Programme, UCL Institute of Child Health, University College London, United Kingdom

Objectives: Native Andean ancestry gives partial protection from reduced birthweight at high altitude in the Andes compared with European ancestry. Whether Andean ancestry is also associated with body proportions and greater post-natal body size at altitude is unknown. Therefore, we tested whether a greater proportion of Andean ancestry is associated with stature and body proportions among Peruvian children at high and low altitude.

Methods: Height, head circumference, head-trunk height, upper and lower limb lengths, and tibia, ulna, hand and foot lengths, were measured in 133 highland and 169 lowland children aged 6 months to 8.5 years. For highland and lowland groups separately, age-sex-adjusted anthropometry *z* scores were regressed on the number of indigenous parental surnames as a proxy for Andean ancestry, adjusting for potential confounders (maternal age and education, parity, altitude [highlands only]).

Results: Among highland children, greater Andean ancestry was negatively associated with stature and tibia, ulna, and lower limb lengths, independent of negative associations with greater altitude for these measurements. Relationships were strongest for tibia length: each additional Andean surname or 1,000 m increase at altitude among highland children was associated with 0.18 and 0.65 *z* score decreases in tibia length, respectively. Anthropometry was not significantly associated with ancestry among lowland children.

Conclusions: Greater Andean ancestry is associated with shorter stature and limb measurements at high but not low altitude. Gene-environment interactions between high altitude and Andean ancestry may exacerbate the trade-off between chest dimensions and stature that was proposed previously, though we could not test this directly. *Am. J. Hum. Biol.* 27:798–806, 2015. © 2015 The Authors American Journal of Human Biology Published by Wiley Periodicals, Inc.

Studies of responses to hypoxia among populations resident at high altitude for many generations provide a classic means of investigating both genetic adaptation and plasticity in development and growth (Baker and Little 1976; Beall, 2013; Frisancho, 1993). Native high-altitude populations, including Andeans and Tibetans, are partially protected from the negative effects of hypoxia on birth weight compared with more recent migrants to altitude (Julian et al., 2007; Krampfl et al., 2000; Zamudio et al., 1993). However, birth weight still continues to decrease with altitude among residents of Peru, the focus of this study, such that mean birth weight falls by ~130 g per 1000 m increase in altitude (Mortola et al., 2000).

Despite their generally lower socioeconomic status (SES) compared with Andean residents of European ancestry, and this dose-response “altitude-penalty” on birth weight, neonates of Andean descent have higher birth weight than those of European ancestry at equivalent altitudes above ~2500m. For example, Julian et al. reported that infants born in Bolivia to parents of Andean ancestry weighed on average 200–250 g more than those of European ancestry at the same altitude, controlling for gestational age and other known influences on birth weight (Julian et al., 2009; Vargas et al., 2007). The degree of protection from decreased birth weight correlates positively with the proportion of Andean ancestry

(Julian et al., 2007; Soria et al., 2013), and paternal Andean ancestry exerts a stronger effect than maternal Andean ancestry by ~80 g (Bennett et al., 2008).

Given that most studies investigating the relationship between ancestry and growth at altitude have been restricted to birth weight, there is a particular need to extend such research postnatally and to consider whether outcomes such as growth are affected. Understanding the relationship between Andean ancestry and growth, as

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Contract grant sponsor: Arts and Humanities Research Council (UK); Contract grant sponsor: University of Cambridge Centre for Latin American Studies Abbey-Santander Travel Grants; Contract grant sponsor: Newnham College, Cambridge (Henry Sidgwick Research Fellowship to EP); Contract grant sponsor: National Institutes of Health; Contract grant number: HHSN268200900033C, HLBI-079647, and TW #001188; Contract grant sponsor: UK Medical Research Council; Contract grant number: MR/J004839/1.

Additional Supporting Information may be found in the online version of this article.

*Correspondence to: Dr Emma Pomeroy, Newnham College, Cambridge, CB3 9DF, UK. E-mail: emma.pomeroy@cantab.net

Received 12 November 2014; Revision received 1 February 2015; Accepted 9 March 2015

DOI: 10.1002/ajhb.22725

Published online 11 May 2015 in Wiley Online Library (wileyonlinelibrary.com).

measured by body size and proportions, at altitude can offer new insight into the developmental etiology of adaptations to hypoxia among native high-altitude residents. Highland Andean children and adults are generally shorter than low-altitude populations, with larger chests and shorter legs relative to the trunk (although see Frisancho et al., 1975a; Stinson and Frisancho, 1978), but some of these differences may be largely attributable to contrasts in healthcare, SES, and temperature which correlate negatively with altitude, rather than hypoxia per se (Frisancho et al., 1970; Greksa, 2006; Greksa et al., 1984; Niermeyer et al., 2009; Pawson and Huicho, 2010; Pawson et al., 2001). With decreasing temperature (increasing latitude), limbs are shorter relative to the trunk and distal limb segments become similarly shorter relative to proximal segments among modern and ancient human populations (Katzmarzyk and Leonard, 1998; Roberts, 1978; Trinkaus, 1981). However, the extent to which such variation in body proportions is genetically determined (Cowgill et al., 2012; Holliday, 1997) or is a direct effect of ambient temperature on growth (Serrat et al., 2008) remains uncertain.

Several hypotheses are proposed concerning the relationship between Andean ancestry and postnatal body size and proportions. Hypothesis 1 is that, other factors being equal, individuals with greater Andean ancestry grow taller and are heavier postnatally as well as prenatally at altitude as a result of genetic adaptations to hypoxia. Recent studies have detected signatures of natural selection among Andeans in genes associated with the hypoxia-inducible factor (HIF) pathway, which is involved in tissue oxygen homeostasis (Bigham et al., 2009, 2010). However, the phenotypic effects of these genotypes are uncertain, although progress is being made in this regard (Bigham et al., 2014). Andeans do not have major genetic variants that elevate arterial oxygen saturation, unlike Tibetans (Beall et al., 1994, 1999), and variants of the genes *EGLN1* and *EPAS1*, which show evidence of positive selection in highland Andeans, are not associated with blood hemoglobin concentration (Bigham et al., 2013). Alternatively, numerous studies show that highland Andean children have larger chests and lung volumes for their age and/or height compared with non-Andeans at high altitude (Beall, 1982; Brutsaert et al., 1999; Frisancho et al., 1975a; Hoff, 1973; Mueller et al., 1978b; Palomino et al., 1978; Stinson, 1980, 1982), although Tibetans and Europeans are also reported to have increased chest and/or lung dimensions at high altitude compared with their low-altitude counterparts (Beall 1982; DeGraff et al. 1970; Droma et al. 1991; Greksa 1986; Greksa 1988; Greksa and Haas 1982; Greksa et al. 1988).

While larger chest/lungs could increase arterial oxygenation and thus reduce hypoxic constraints on growth (consistent with Hypothesis 1), others have proposed that the greater acceleration of chest growth among highland Andeans is compensated for by reduced height (Frisancho, 2013; Frisancho et al., 1975b; Mueller et al., 1978a; Pawson and Huicho, 2010). Hypothesis 2 is therefore that Andean ancestry is associated with altered postnatal growth patterns, including shorter stature. Various studies suggest that under conditions of environmental stress (e.g., hypoxia, malnutrition) growth is prioritized in some parts of the body, typically the brain (Barbiro-Michaely et al., 2007; Barker, 1998; Giussani, 2011; Wells, 2013) at the particular expense of lower limb and distal limb segment (tibia) length and, to a lesser extent, trunk length

(Bailey and Hu, 2002; Bailey et al., 2007; Bogin et al., 2002; Gunnell et al., 1998; Lampl et al., 2003; Pomeroy et al., 2012; Pomeroy et al., 2013; Whitley et al., 2008). Thus, if brain and chest size are prioritized over stature among children of Andean ancestry, we might also anticipate that lower limb and especially tibia length experience the greatest reductions in growth, altering relative trunk and limb proportions.

Two variants of Hypothesis 2 may also be proposed. Hypothesis 2a is that patterns of chest, stature, and limb proportions associated with Andean ancestry are genetic adaptations to hypoxia and expressed regardless of altitude. Hypothesis 2b is that the Andean growth pattern results from gene-environment interactions whereby greater chest size has a genetic basis, but is facultatively expressed during development at high altitude. Previous studies suggest that accelerated Andean chest development is conditional upon high-altitude exposure and ancestry: children of European ancestry also develop relatively larger chests at high altitude (Greksa, 1986, 1988; Greksa and Haas, 1982; Greksa et al., 1988), but to a lesser extent than Andeans (Brutsaert et al., 1999; Greksa, 1986, 1988; Greksa et al., 1988; Palomino et al., 1978; Stinson, 1982). If the highland Andean body plan is contingent on hypoxia exposure, Andean ancestry would be expected to be associated with stature and limb proportions among highland but not lowland children. At low altitude, no increase in chest size would be favored; hence, limb growth would not be sacrificed.

The purpose of this study was to examine the relationship between native Andean ancestry (inferred from surname analysis) and the size and proportions of the head, trunk, limbs, and limb segments among highland and lowland Peruvian children. We aimed to investigate which of the hypotheses outlined best accounts for any variation in postnatal body size and proportions associated with Andean ancestry by evaluating the following predictions:

Hypothesis 1. If genetic adaptation to hypoxia benefits overall growth at altitude, highland children with a greater proportion of Andean ancestry will have a taller stature and longer limbs;

Hypothesis 2a. If genetic adaptation to hypoxia among native Andeans favors a larger chest at the expense of stature and limb length, a higher proportion of Andean ancestry will be associated with shorter stature and limbs regardless of altitude;

Hypothesis 2b. If genetic adaptation to hypoxia among native Andeans favors a larger chest at the expense of stature and limb length only on exposure to hypoxia, a greater degree of Andean ancestry will be associated with shorter stature and limbs among highland children only.

Methods

The sample is part of that from a larger study of body size and proportions among Peruvian children (Pomeroy et al., 2012). A convenience sample of 447 children was recruited from two populations. The first population was the “*pueblo joven*” (shanty town) of Pampas de San Juan de Miraflores on the south side of Peru’s capital, Lima (latitude -12.0 , longitude -77.0 , altitude 140 m). The settlement was unplanned and comprised of migrants predominantly from highland Peru. Although it remains a relatively low SES community, many houses are now more substantially constructed and are connected to water and

sewage systems (Checkley et al., 2004; Masterson Creber et al., 2010; Miranda et al., 2009; Sterling et al., 2012). There are also schools and health posts in the settlement. The highland sample came from small, rural communities in the Vinchos and Santillana districts of Peru's Ayacucho Region (latitude -13.2 , longitude -74.2 for Ayacucho city). Communities were located at 3,100–4,400 m altitude, and comprised mainly of subsistence agropastoralists (Masterson Creber et al., 2010; Miranda et al., 2009).

Participants of the original study were aged 6 months to 14 years, falling within specific target age groups (Pomeroy et al., 2012). Date of birth was confirmed using national identity documents, birth record cards, or school records. Participants had been born and raised in their study region, and did not have any medical conditions, which might compromise growth aside from general undernutrition. A maximum of one child per household participated in the study on a voluntary basis. A parent or guardian provided written informed consent, and participants aged 6 years or over gave verbal or written assent. The study was approved by the Institutional Ethics Committee at the Universidad Peruana Cayetano Heredia, Lima, and the Health Directorate for Ayacucho Region (Dirección Regional de Salud Ayacucho, DIRESA).

Anthropometry was measured by one trained observer (EP) following standard techniques (Cameron, 2004; Lohman et al., 1988). Standing and head-trunk (sitting) heights were measured to the nearest mm with a Seca Leicester Height Measure, or as crown-heel and crown-rump length in children aged less than 2 years using a Rollametre (Dunmow, UK). Total lower limb length was calculated by subtracting head-trunk from standing height (or crown-rump from crown-heel length). Tibia, ulna, upper arm, hand, and foot lengths were measured to the nearest mm using sliding callipers, and total upper limb length calculated by summing ulna and upper arm length. Head circumference was measured using a 15 mm-wide nonstretch fiberglass tape (Hoechtmass, Germany). Intraobserver error on adult subjects gave a technical error of measurement below 1% and a coefficient of reliability (R) of at least 0.97 for all measurements, which were judged to be within acceptable limits (Ulijaszek and Kerr, 1999).

Parental surnames were used as a proxy for ancestry as in various studies in the Andes and elsewhere (Azcorra et al., 2013; Chakraborty et al., 1989; Colantonio et al., 2003; Relethford, 1995). Peruvians inherit the paternal surname of their mother and father, giving each individual two surnames, and they do not change their surnames at marriage. Therefore, the linguistic origin of a participant's parents' surnames (giving 4 surnames for analysis) can be used to infer the proportion of indigenous ancestry of that individual. While an imperfect marker (Brutsaert, 2001; Greksa, 1992), this approach has been used successfully in numerous studies of Andean populations that have aimed to relate birth characteristics to ancestry (Bennett et al., 2008; Gonzales et al., 1996; Julian et al., 2007; Mueller et al., 1978a; Soria et al., 2013) and the association between native ancestry and surname origin in the Andes has been previously validated (Chakraborty et al., 1989). The four parental surnames of each individual were classified as "indigenous," "mestizo," (mixed) or "foreign" based on their linguistic origin using a surname dictionary compiled by one of the authors (LGM).

Sex-age-specific internal z scores were calculated for anthropometry in the combined lowland and highland

sample after fitting centiles using the LMS method (Cole and Green, 1992), and analyses used these z scores throughout. For the highland and lowland groups, anthropometry z scores were regressed separately on the number of indigenous surnames (entered as a continuous variable, range 0–4) and potential confounding variables using a backwards stepwise model where potential confounders with $P < 0.1$ were retained. Potential confounders were altitude (highland sample only), maternal education (an aggregate marker of SES, e.g., Jansen et al., 2009; Kramer et al., 2000; Raum et al., 2001; van den Berg et al., 2013; which was used as such in previous study of Andean surnames and growth: Bennett et al., 2008), maternal age and birth order. These variables were included as they have previously been shown to relate to child growth and/or body proportions (Dasgupta et al., 2008; Hatton and Martin, 2010; Lawson and Mace, 2008; Li et al., 2007; Li and Power, 2004; Semba et al., 2008; Whitley et al., 2008), and because they are correlated with the number of indigenous parental surnames in our sample. Indigenous surnames increase in frequency with altitude in the Andes because European migrants arriving from the 16th century onwards generally settled at low to mid altitudes, and perhaps because indigenous Andeans tend to adopt more Spanish-sounding surnames as they acculturate (Schull and Rothhammer, 1977), while SES and access to education decrease at higher altitude (Niermeyer et al., 2009; Rivera-Ch et al., 2008).

As patterns of growth are complex and transient during puberty, analyses were restricted to children aged 6 months to 8.5 years (lowland $n = 169$, highland $n = 133$). Although pubertal onset was not assessed directly, the vast majority of children aged below 8.5 years were prepubertal in a similar low SES population in the Americas (Wilson et al., 2011). Analyses were restricted to individuals who had all four parental surnames recorded, as well as all potential confounding variables for the regression models (exclusions are detailed in Supporting Information Table S1). Within each sample, t -tests or chi-square tests as appropriate confirmed that there were no significant differences between those with and without full surname data in terms of altitude, maternal age or education, or anthropometry z scores (height, head circumference, sitting height, and tibia length: $P > 0.05$, results not shown). Analyses of relative total limb, ulna, tibia hand and foot lengths were performed by adjusting for head-trunk height z score in the regression model. Analyses were conducted in SPSS v.21, with P values below 0.05 considered significant.

Results

Tables 1–3 summarize the characteristics and anthropometry of the highland and lowland samples. Samples were approximately evenly divided between males and females. Most children had a mix of indigenous and mestizo or foreign surnames, and foreign surnames were more common in the lowland sample (Fig. 1). Seven percent of highland children had four indigenous surnames, while 9% had none, compared with 2 and 34% of lowland children respectively (highland-lowland difference by χ^2 : $P < 0.001$). Only 6% of highland mothers had secondary education or higher, while 19% had not attended school. In contrast all lowland mothers had some level of education and 77% had complete secondary education or higher (χ^2 : $P < 0.001$). Half of lowland children were first-born,

while only 30% of highland children were first-born and one-third were the fourth child or later born (χ^2 : $P < 0.001$). Highland children had lower mean anthropometry z scores for all measurements than lowland children (Table 3: highland-lowland differences have $P < 0.001$ for all measurements using t-test). Further details of these samples and differences in anthropometry were reported previously (Pomeroy et al., 2012, 2013, 2014).

Among highland children, more indigenous parental surnames were associated with shorter tibia length, relative tibia length, stature, lower limb length, ulna length and relative lower limb length (in order of decreasing regression coefficients for number of indigenous parental surnames: Table 4). Increased altitude was similarly associated with shorter measurement z scores except for head-trunk height, head circumference, and foot length. An additional indigenous surname was associated with a 0.18 z -score decrease in tibia length, while 1,000 m increase in altitude was associated with a 0.65 z -score decrease in

tibia length. This compares with 0.14 and 0.62 z -score decreases in stature with the same increases in indigenous surnames and altitude respectively, clearly very similar. Full details of the regression models including potential confounders included in each model are provided in Supporting Information Table S2.

Among lowland children, anthropometry z scores were not associated with the number of indigenous parental surnames (Table 4, full model details in Supporting Information Table S3).

TABLE 3. Summary statistics for child anthropometry, giving mean measurements at age 4 years by sex, and mean internal age-sex adjusted z scores by site

TABLE 1. Age and sex distribution of the sample

Age group (years)	Lowland			Highland		
	Female (n)	Male (n)	Total (n)	Female (n)	Male (n)	Total (n)
0.5–1.49	14	15	29	13	13	26
1.5–2.49	16	14	30	7	10	17
2.5–3.49	13	14	27	15	11	26
3.5–4.49	13	12	25	13	14	27
5.5–6.49	16	15	31	9	13	22
7.5–8.49	14	13	27	9	6	15
Total	86	83	169	66	67	133

Variable	Males		Females		Sex-age specific z scores					
	Mean	SD ^b	Mean	SD	Highland	Lowland	n	Mean	SD	n
Head circumference	49.5	1.3	48.9	1.4	-0.4	0.8	125	0.3	1.0	165
Stature	98.1	5.0	97.2	5.2	-0.8	0.7	133	0.7	0.7	167
Head-trunk height	57.9	2.5	57.3	2.7	-0.6	0.8	131	0.5	0.9	167
Total upper limb length	33.7	2.3	33.4	2.5	-0.9	0.7	106	0.6	0.7	154
Ulna length	14.7	1.0	14.5	1.1	-0.9	0.7	108	0.6	0.7	163
Hand length	11.3	0.7	11.3	0.7	-0.8	0.7	103	0.5	0.8	156
Total lower limb length	40.3	3.0	39.9	3.0	-0.8	0.8	131	0.7	0.7	167
Tibia length	19.4	1.7	19.3	1.6	-0.9	0.7	116	0.7	0.6	162
Foot length	15.5	1.0	15.3	1.0	-0.7	0.7	116	0.6	0.8	163

^aDerived from LMS model (Cole and Green, 1992) for the full dataset.

^bSD: standard deviation.

TABLE 2. Summary statistics on parental surnames, altitude, birth order, maternal education, maternal age, and offspring anthropometry z scores for the highland and lowland study samples

Variables	Highland			Lowland		
	n	%	Regression category ^a	n	%	Regression category
Categorical						
No. indigenous parental surnames						
0	12	9		56	33	
1	39	29		68	41	
2	40	30		35	21	
3	32	24		5	3	
4	10	8		3	2	
Birth order						
1	40	30		84	50	
2	29	22		51	31	
3	19	14		20	12	
4+	45	34		12	7	
Maternal education						
None, illiterate	23	17	1	0		1
None, literate	2	2	1	0		1
Primary, incomplete	70	53	2	3	2	1
Primary, complete	30	22	3	6	4	1
Secondary, incomplete	4	3	3	29	17	1
Secondary, complete	3	2	3	68	41	2
Postsecondary, incomplete	0	0	3	25	15	3
Postsecondary, complete	1	1	3	36	22	4
Continuous						
	Mean	SD		Mean	SD	
Maternal age at birth of child (yrs)	25.1	7.6		26.4	6.3	
	Median	IQR				
Altitude measured (m)	3,730	3,320–3,830		–	–	

^aRegression Category^a indicates how original categories were recoded for the purpose of the regression model: categories with small numbers of cases were combined.

Discussion

Our results indicate that a greater number of indigenous parental surnames was associated with reduced stature and the lengths of the tibia, lower limb and ulna, but not head circumference or hand and foot lengths, among highland children after adjusting for relevant factors including altitude. The strongest associations were observed with tibia length. In contrast, there were no such associations among lowland children. Among highland children, anthropometry was negatively associated with higher altitude independent of the number of indigenous surnames. We considered that these results were most consistent with Hypothesis 2b; namely, that acceler-

ated chest growth at altitude, dependent both on hypoxia exposure and augmented by Andean ancestry, results in reduced growth elsewhere in the body (see Introduction). While this is consistent with what others have proposed (Frisancho, 2013; Frisancho et al., 1975b; Mueller et al., 1978a; Pawson and Huicho, 2010), we lacked chest size data to test this directly. The strongest associations with ancestry among highland children were for tibia length, consistent with previous studies that suggest that tibia length is particularly sensitive to environmental conditions (see Introduction). This pattern of results supports the interpretation that ancestry-related differences in body dimensions result from a growth trade-off at altitude. The results were not consistent with an adaptation among native Andeans that promotes greater height and longer limbs at altitude (Hypothesis 1). The lack of association between ancestry and height or body proportions among lowland children does not suggest that reduced stature and limb lengths is a general characteristic of Andeans regardless of altitude (Hypothesis 2a).

If accelerated chest growth among high-altitude Andeans represents an adaptation to reduce tissue hypoxia (Frisancho, 2013; Frisancho et al., 1975b; Mueller et al., 1978a; Pawson and Huicho, 2010), it does not seem to lead to increased growth and stature. Any advantage in terms of body size may only be achieved once growth is complete, since growth velocity has been noted to be slower at high altitude (Frisancho, 1976). Some studies indicate that increased chest size relative to stature is more marked in children than adults (Hoff, 1973; Mueller et al., 1978a, 1978b; Palomino et al., 1978), suggesting that height “catches up” with chest growth later in development. Alternatively, the larger chest dimensions of highland Andeans are associated with improved lung function (Frisancho, 1969; Mueller et al., 1978b; Whitaker, 1992; but see Brutsaert et al., 1999; Tarazona-Santos et al., 2000) and so may improve work efficiency rather than impacting growth. Finally, others have suggested that only the most nutritionally-stressed Andean populations show increased chest growth (Pawson and




Fig. 1. Numbers of indigenous parental surnames among highland and lowland Peruvian children in the sample.

TABLE 4. Results of regression models of anthropometry z scores on the number of indigenous parental surnames among highland and lowland Peruvian children, adjusting for potential confounding variables

Measurement z score	Highland					Lowland	
	Altitude (km)		Number of indigenous parental surnames			Number of indigenous parental surnames	
	B	P	B	Standard error (B)	P	B	P
Tibia length	-0.65	<0.001	-0.18	0.05	0.002	0.03	0.6
Relative tibia length	-0.49	0.001	-0.13	0.05	0.005	-0.004	0.9
Lower limb length	-0.66	0.001	-0.16	0.06	0.01	-0.02	0.7
Stature	-0.62	0.001	-0.14	0.06	0.01	0.04	0.5
Ulna length	-0.72	<0.001	-0.12	0.05	0.03	0.01	0.9
Relative lower limb length	-0.57	0.002	-0.12	0.06	0.04	-0.04	0.5
Head-trunk height		ns	-0.12	0.06	0.06	0.08	0.3
Head circumference		ns	-0.11	0.07	0.08	-0.03	0.7
Relative ulna length	-0.49	0.001	-0.08	0.04	0.09	-0.02	0.7
Upper limb length	-0.61	0.001	-0.09	0.06	0.1	0.02	0.7
Hand length		ns	-0.09	0.06	0.2	-0.03	0.7
Foot length	-0.37	0.07	-0.09	0.06	0.2	-0.02	0.7
Relative upper limb length	-0.40	0.003	-0.05	0.05	0.5	-0.01	0.8

Bold indicates $P < 0.05$. Blank cells denote variable excluded from model as $P < 0.1$.

See Supporting Information Tables S2 and S3 for details of confounders for highland and lowland samples respectively. Variables are ordered in the table by P values for number of indigenous surnames among highland children.

Huicho, 2010; Pawson et al., 2001), but the relationship between nutritional status, chest size and linear growth remains to be directly tested.

Alternative explanations for our findings include the possibility that differences in limb and body proportions are due entirely to genetic factors. This would seem unlikely given that, while worldwide population differences in body size and proportions are thought to include a genetic component (Eveleth and Tanner, 1990), it probably accounts for only a small proportion of global variation in relative sitting height (head-trunk height relative to stature: $\sim 3.6\%$; Bogin et al., 2001). Furthermore, differences in relative limb proportions are greater between populations of European vs. African or Australian ancestry than between those of European vs. Asian (including native American) ancestry (Bogin et al., 2001; Feldesman and Fountain, 1996; Stinson, 2009), and most migrants to Peru have come from Europe (Bigham et al., 2008; Sanchez et al., 2010).

Whether Andean ancestry is associated with genetic differences in body proportions has not been extensively studied. Adaptation to cold climates (e.g., Trinkaus, 1981) or moving over steep terrain (e.g., Higgins and Ruff, 2011) has been proposed to result in relatively shorter limbs and especially distal limb segments (tibia), but evidence for such relationships among highland Andeans is lacking. The Multinational Andean Genetic and Health Program studies of north Chilean Aymara-speakers gave contradictory results regarding the relationship between ancestry, altitude, and body size and proportions (Mueller et al., 1978a; Palomino et al., 1978). Stinson (2009) investigated stature and relative sitting height among poorer rural and wealthier urban Bolivian children aged 8–13. Although ancestry, nutrition, SES and time spent at altitude were related and difficult to separate, she suggested that Bolivian ancestry (parents born in Bolivia) and time at altitude might both be associated with relatively shorter legs within the wealthier urban sample, although the relative influences of multiple factors could not be assessed. Variation with ancestry was not investigated within her rural sample. The relationships between these different environmental characteristics, ancestry, and body size and proportions thus also remain to be elucidated.

The lack of any relationship between growth measures and parental surnames among lowland children in our study suggests that the variation in body size and proportions with inferred ancestry among highland children may result from gene-environment interactions, although this interpretation has to be cautious. Acculturation, associated with the tendency to Hispanicize indigenous surnames (Schull and Rothhammer, 1977), is greater in lowland communities and the lowland sample likely represents a greater mix of different indigenous populations than the highland sample, since the settlement comprises migrants from various rural highland regions of Peru (Checkley et al., 2002), as well as a small contingent from the coast and Amazon. Although the majority of children in the lowland sample were likely to have been of largely highland Andean ancestry, we can neither confirm this objectively nor exclude a genetic difference in body proportions between the highland and lowland samples that is not environment-dependent. Another potential explanation for our results among highland children is that parental surnames are related not only to ancestry but also to other environmental variables like SES, which the available

variables may not have controlled for adequately. This possibility also requires further investigation.

Whether high-altitude neonates show any relationship between body size and proportions and Andean ancestry is unclear. While the effects of Andean ancestry on birth weight at altitude are consistent across studies, few consider differences in other neonatal anthropometry. Birth length, head circumference, limb lengths, skinfold thickness and weight are generally reduced at altitude (Giusani et al., 2001; Haas, 1976; Haas et al., 1977, 1980; McClung, 1969; Mortola et al., 2000; Soria et al., 2013). Haas et al. (1980) suggested that the decrease in crown-heel length with altitude was greater among neonates of European compared with Andean ancestry, though Soria et al. (2013) did not replicate this finding.

Other studies demonstrate a positive association between birth weight and adult height and weight (Eide et al., 2005; Sørensen et al., 1999; Yliharsila et al., 2007), so the apparent loss of this early growth advantage among children of Andean ancestry at altitude requires explanation. It may reflect the fact that higher birth weight at altitude among Andeans results from maternal physiological adaptations, including greater maternal uterine artery blood flow, which is augmented in the third trimester by $\sim 36\%$ (Julian et al., 2008; but see Postigo et al., 2009; Wilson et al., 2007; Zamudio et al., 2007), and increased resting respiratory rate (Vargas et al., 2007) compared with non-Andean mothers.

Higher birth weight is also positively associated with neonatal survival (except at extremely high birthweight: Karn and Penrose, 1951; Wilcox and Russell, 1983), therefore the greater birth weight associated with Andean ancestry may confer benefits in early postnatal survival at altitude rather than growth. Such a survival advantage has been shown for Tibetan (high-altitude native) vs. Han Chinese (recent migrants) at high altitude (Moore et al., 2001). Greater fat mass rather than lean mass may be key to early infant survival at altitude (Wiley, 1994), and newborns at altitude in Denver, Colorado have reduced subcutaneous fat but not lean mass, head circumference or femur length (Galan et al., 2001). This may suggest that fat mass is disproportionately affected by high-altitude gestation. In a Bolivian high-altitude sample, surname-inferred Andean ancestry was associated with greater birth weight but no difference in length compared with European ancestry, which would be consistent with greater fat accrual among Andean neonates at altitude, although direct data on body composition were unavailable and differences in ponderal index insignificant (Soria et al., 2013). If the greater birth weight of highland Andean infants reflects greater fat rather than lean mass, this could explain the positive effects of Andean ancestry on birth weight but not postnatal height and lower limb length in the highlands.

Strengths of our study include the detailed anthropometry, including limb segment lengths, which allowed us to investigate in greater detail the relationship between body proportions and ancestry than is possible in many studies. Furthermore, while a large number of studies have considered ancestry in relation to neonatal anthropometry, few have considered the relationship between ancestry and body size or proportions in infants and children. Unfortunately, we were unable to explore whether the associations between anthropometry and ancestry might reflect trade-offs with chest growth, or whether

they persist into adulthood. Investigating these problems in the future would help to shed light on the nature and cause of differences in body size and proportions by ancestry in the Andean highlands. Furthermore, our data were cross-sectional, and therefore we were unable to assess longitudinal differences in body size, proportions, and growth patterns according to ancestry in our samples.

The use of genetic data rather than surnames (which are less accurate markers of ancestry) would also serve to elucidate the nature of links between ancestry and body size and proportions. While Chakraborty et al. (1989) showed that the number of indigenous surnames correlated with genetic markers of Andean ancestry, their study considered north Chilean and Bolivian populations only, reported a limited range of markers (as expected for a study of its date), and showed that even those with no indigenous parental surnames had a significant proportion of Andean admixture. Those with no indigenous surnames had on average 64% “native American” loci, compared with 85% for those with 1–2 indigenous names, and 89% for those with 3–4. A more recent study of highland and lowland Bolivian women similarly showed a good correspondence between surnames and genetic indigenous or European ancestry using over 100 Ancestry Informative Markers (AIMs) (Julian et al., 2009). While the relationship has not been directly tested in our study samples, it is likely that surnames provide a reasonable proxy for ancestry in our analyses.

In conclusion, greater Andean ancestry (inferred from numbers of indigenous parental surnames) was negatively associated with anthropometry among highland, but not lowland, Peruvian children. In particular, highland children with more indigenous parental surnames had significantly shorter tibia length (absolutely and relative to trunk length), lower limb length, ulna length, and stature. Within the highland sample, altitude also showed a negative relationship with anthropometry z scores, and the ancestry and altitude associations with anthropometry in the highland sample were independent. The results suggested that the effect of indigenous Andean ancestry on increasing birth weight at altitude as demonstrated in other studies does not translate into larger postnatal body size. Previously proposed trade-offs between stature and chest size, contingent on Andean ancestry and exposure to hypoxia, may explain the pattern of results, but we were unable to test this hypothesis directly.

ACKNOWLEDGMENTS

The authors thank the participants and their families for kindly taking part in the study, Lilia Cabrera and Angela Huamán Gómez of PRISMA, and all the field staff for their assistance in participant recruitment and data collection. They also thank Dr. Antonio Bernabe Ortiz for helping to establish the project; the Ayacucho Department Health Directorate (DIRESA) for supporting the study; Dr Enrique Vargas (Instituto Boliviano de Biología de Altura, La Paz, Bolivia) for his assistance in classifying additional surnames; and Daniel Antiporta and Melina Paola Zevallos Guarda for administrative assistance.

LITERATURE CITED

Azcorra H, Varela-Silva MI, Rodriguez L, Bogin B, Dickinson F. 2013. Nutritional status of Maya children, their mothers, and their grand-

- mothers residing in the City of Merida, Mexico: Revisiting the leg-length hypothesis. *Am J Hum Biol* 25:659–665.
- Bailey SM, Hu XM. 2002. High-altitude growth differences among Chinese and Tibetan children. In: Gilli G, Schell LM, Benso L, editors. *Human Growth from Conception to Maturity*. London: Smith-Gordon. p 237–247.
- Bailey SM, Xu J, Feng JH, Hu X, Zhang C, Qui S. 2007. Tradeoffs between oxygen and energy in tibial growth at high altitude. *Am J Hum Biol* 19:662–668.
- Baker PT, Little MA. 1976. *Man in the Andes: A Multidisciplinary Study of High Altitude Quechua*. Stroudsburg, Pennsylvania: Dowden, Hutchinson and Ross, Inc.
- Barbiro-Michaely E, Tolmasov M, Rinkevich-Shop S, Sonn J, Mayevsky A. 2007. Can the “brain-sparing effect” be detected in a small-animal model? *Med Sci Monit* 13:BR211–219.
- Barker DJP. 1998. *Mothers, Babies, and Disease in Later Life*. Edinburgh: Churchill Livingstone.
- Beall CM. 1982. A comparison of chest morphology in high altitude Asian and Andean populations. *Hum Biol* 54:145–163.
- Beall CM. 2013. Human adaptability studies at high altitude: Research designs and major concepts during fifty years of discovery. *Am J Hum Biol* 25:141–147.
- Beall CM, Almsay LA, Blangero J, Williams-Blangero S, Brittenham GM, Strohl KP, Decker MJ, Vargas E, Villena M, Soria R, Alarcon AM, Gonzales C. 1999. Percent of oxygen saturation of arterial hemoglobin among Bolivian Aymara at 3,900–4,000 m. *Am J Phys Anthropol* 108:41–51.
- Beall CM, Blangero J, Williams-Blangero S, Goldstein MC. 1994. Major gene for percent of oxygen saturation of arterial hemoglobin in Tibetan highlanders. *Am J Phys Anthropol* 95:271–276.
- Bennett A, Sain SR, Vargas E, Moore LG. 2008. Evidence that parent-of-origin affects birth-weight reductions at high altitude. *Am J Hum Biol* 20:592–597.
- Bigham A, Bauchet M, Pinto D, Mao X, Akey JM, Mei R, Scherer SW, Julian CG, Wilson MJ, López Herráez D, Brutsaert T, Parra EJ, Moore LG, Shriver MD. 2010. Identifying signatures of natural selection in Tibetan and Andean populations using dense genome scan data. *PLoS Genet* 6:e1001116.
- Bigham AW, Julian CG, Wilson MJ, Vargas E, Browne VA, Shriver MD, Moore LG. 2014. Maternal PRKAA1 and EDNRA genotypes are associated with birth weight, and PRKAA1 with uterine artery diameter and metabolic homeostasis at high altitude. *Physiol Genomics* 46:687–697.
- Bigham AW, Kiyamu M, León-Velarde F, Parra EJ, Rivera-Ch M, Shriver MD, Brutsaert TD. 2008. Angiotensin-converting enzyme genotype and arterial oxygen saturation at high altitude in Peruvian Quechua. *High Alt Med Biol* 9:167–178.
- Bigham AW, Mao X, Mei R, Brutsaert T, Wilson MJ, Julian CG, Parra EJ, Akey JM, Moore LG, Shriver MD. 2009. Identifying positive selection candidate loci for high-altitude adaptation in Andean populations. *Hum Genomics* 4:79–90.
- Bigham AW, Wilson MJ, Julian CG, Kiyamu M, Vargas E, León-Velarde F, Rivera-Chira M, Rodríguez C, Browne VA, Parra E, Brutsaert T, Moore LG, Shriver MD. 2013. Andean and Tibetan patterns of adaptation to high altitude. *Am J Hum Biol* 25:190–197.
- Bogin B, Kapell M, Silva MIV, Orden B, Smith PK, Loucky J. 2001. How genetic are human body proportions? In: Dasgupta P, Hauspie R, editors. *Perspectives in Human Growth, Development and Maturation: Global Tributes to S.R. Das, the Pioneering Indian Auxologist*. New York: Kluwer Academic Publishers. p 205–221.
- Bogin B, Smith P, Orden AB, Varela Silva MI, Loucky J. 2002. Rapid change in height and body proportions of Maya American children. *Am J Hum Biol* 14:753–761.
- Brutsaert TD. 2001. Limits on inferring genetic adaptation to high altitude in Himalayan and Andean populations. *High Alt Med Biol* 2:211–225.
- Brutsaert TD, Soria R, Caceres E, Spielvogel H, Haas JD. 1999. Effect of developmental and ancestral high altitude exposure on chest morphology and pulmonary function in Andean and European/North American natives. *Am J Hum Biol* 11:383–395.
- Cameron N. 2004. Measuring growth. In: Hauspie RC, Cameron N, Molinari L, editors. *Methods in Human Growth Research*. Cambridge: Cambridge University Press. p 68–107.
- Chakraborty R, Barton SA, Ferrell RE, Schull WJ. 1989. Ethnicity determination by names among the Aymara of Chile and Bolivia. *Hum Biol* 61:159–177.
- Checkley W, Gilman RH, Black RE, Epstein LD, Cabrera L, Sterling CR, Moulton LH. 2004. Effect of water and sanitation on childhood health in a poor Peruvian peri-urban community. *Lancet* 363:112–118.
- Checkley W, Gilman RH, Black RE, Lescano AG, Cabrera L, Taylor DN, Moulton LH. 2002. Effects of nutritional status on diarrhea in Peruvian children. *J Pediatr* 140:210–218.

- Colantonio SE, Lasker GW, Kaplan BA, Fuster V. 2003. Use of surname models in human population biology: A review of recent developments. *Hum Biol* 75:785–807.
- Cole TJ, Green PJ. 1992. Smoothing reference centile curves: The LMS method and penalized likelihood. *Stat Med* 11:1305–1319.
- Cowgill LW, Eleazer CD, Auerbach BM, Temple DH, Okazaki K. 2012. Developmental variation in ecogeographic body proportions. *Am J Phys Anthropol* 148:557–570.
- Dasgupta P, Saha R, Nubé M. 2008. Changes in body size, shape and nutritional status of Middle-Class Bengali boys of Kolkata, India, 1982–2002. *Econ Hum Biol* 6:75–94.
- DeGraff AC, Grover RF, Johnson RL, Hammond JW, Miller JM. 1970. Diffusing capacity of the lung in Caucasians native to 3,100 m. *J Appl Physiol* 29:71–76.
- Droma T, McCullough RG, McCullough RE, Zhuang J, Cymerman A, Sun S, Sutton JR, Moore LG. 1991. Increased vital and total lung capacities in Tibetans compared to Han residents of Lhasa (3,658 m). *Am J Phys Anthropol* 86:341–351.
- Eide MG, Øyen N, Skjøerven R, Nilsen ST, Bjerkedal T, Grethe ST. 2005. Size at birth and gestational age as predictors of adult height and weight. *Epidemiol* 16:175–181.
- Eveleth PB, Tanner JM. 1990. *Worldwide Variation in Human Growth*. Cambridge: Cambridge University Press.
- Feldesman MR, Fountain RL. 1996. "Race" specificity and the femur/stature ratio. *Am J Phys Anthropol* 100:207–224.
- Frisancho AR. 1969. Human growth and pulmonary function of a high altitude Peruvian Quechua population. *Hum Biol* 41:365–379.
- Frisancho AR. 1976. Growth and morphology at high altitude. In: Baker PT, Little MA, editors. *Man in the Andes: A Multidisciplinary Study of High Altitude Quechua*. Stroudsburg, Pennsylvania: Dowden, Hutchinson and Ross, Inc. p 180–207.
- Frisancho AR. 1993. Human adaptation and accommodation to environmental stress. Ann Arbor, MI: University of Michigan Press.
- Frisancho AR. 2013. Developmental functional adaptation to high altitude: Review. *Am J Hum Biol* 25:151–168.
- Frisancho AR, Borkan GA, Klayman JE. 1975a. Pattern of growth of lowland and highland Peruvian Quechua of similar genetic composition. *Hum Biol* 47:233–243.
- Frisancho AR, Newman MT, Baker P. 1970. Differences in stature and cortical thickness among highland Quechua Indian boys. *Am J Clin Nutr* 23:382–385.
- Frisancho AR, Velasquez T, Sanchez J. 1975b. Possible adaptive significance of small body size in the attainment of aerobic capacity among high-altitude Quechua natives. In: Watts ES, Johnston FE, Lasker GW, editors. *Biosocial Interrelations in Population Adaptation*. The Hague: Moulton. p 55–64.
- Galan HL, Rigano S, Radaelli T, Cetin I, Bozzo M, Chyu J, Hobbins JC, Ferrazzi E. 2001. Reduction of subcutaneous mass, but not lean mass, in normal fetuses in Denver, Colorado. *Am J Obstet Gynecol* 185:839–844.
- Giussani DA. 2011. The vulnerable developing brain. *Proc Natl Acad Sci USA* 108:2641–2642.
- Giussani DA, Phillips PS, Anstee S, Barker DJ. 2001. Effects of altitude versus economic status on birth weight and body shape at birth. *Pediatr Res* 49:490–494.
- Gonzales GF, Villena A, Ubilluz M. 1996. Age at menarche in Peruvian girls at sea level and at high altitude: Effect of ethnic background and socioeconomic status. *Am J Hum Biol* 8:457–463.
- Greksa LP. 1986. Chest morphology of young Bolivian high-altitude residents of European ancestry. *Hum Biol* 58:427–443.
- Greksa LP. 1988. Effect of altitude on the stature, chest depth and forced vital capacity of low-to-high altitude migrant children of European ancestry. *Hum Biol* 60:23–32.
- Greksa LP. 1992. Surnames as indicators of European admixture in Andean Indians. *Int J Anthropol* 7:41–49.
- Greksa LP. 2006. Growth and development of Andean high altitude residents. *High Alt Med Biol* 7:116–124.
- Greksa LP, Haas JD. 1982. Physical growth and maximal work capacity in preadolescent boys at high altitude. *Hum Biol* 54:677–695.
- Greksa LP, Spielvogel H, Paredes-Fernandez L, Paz-Zamora M, Caceres E. 1984. The physical growth of urban children at high altitude. *Am J Phys Anthropol* 65:315–322.
- Greksa LP, Spielvogel H, Paz-Zamora M, Caceres E, Paredes-Fernandez L. 1988. Effect of altitude on the lung function of high altitude residents of European ancestry. *Am J Phys Anthropol* 75:77–85.
- Gunnell DJ, Smith GD, Frankel SJ, Kemp M, Peters TJ. 1998. Socio-economic and dietary influences on leg length and trunk length in childhood: A reanalysis of the Carnegie (Boyd Orr) survey of diet and health in prewar Britain (1937–39). *Paediatr Perinat Epidemiol* 12(Suppl1):96–113.
- Haas JD. 1976. Prenatal and infant growth and development. In: Baker PT, Little MA, editors. *Man in the Andes: A Multidisciplinary Study of High Altitude Quechua*. Stroudsburg, Pennsylvania: Dowden, Hutchinson and Ross. p 161–179.
- Haas JD, Baker PT, Hunt EE Jr. 1977. The effects of high altitude on body size and composition of the newborn infant in Southern Peru. *Hum Biol* 49:611–628.
- Haas JD, Frongillo EA Jr., Stepick CD, Beard JL, Hurtado L. 1980. Altitude, ethnic and sex differences in birth weight and length in Bolivia. *Hum Biol* 52:459–477.
- Hatton TJ, Martin RM. 2010. The effects on stature of poverty, family size, and birth order: British children in the 1930s. *Oxf Econ Pap* 62:157–184.
- Higgins RW, Ruff CB. 2011. The effects of distal limb segment shortening on locomotor efficiency in sloped terrain: Implications for Neandertal locomotor behavior. *Am J Phys Anthropol* 146:336–345.
- Hoff C. 1973. Altitudinal variations in the physical growth and development of Peruvian Quechua. *Homo* 24:87–99.
- Holliday TW. 1997. Body proportions in Late Pleistocene Europe and modern human origins. *J Hum Evol* 32:423–448.
- Jansen PW, Tiemeier H, Looman CWN, Jaddoe VVW, Hofman A, Moll HA, Steegers EAP, Verhulst FC, Mackenbach JP, Raat H. 2009. Explaining educational inequalities in birthweight: The Generation R Study. *Paediatr Perinat Epidemiol* 23:216–228.
- Julian CG, Galan HL, Wilson MJ, Desilva W, Cioffi-Ragan D, Schwartz J, Moore LG. 2008. Lower uterine artery blood flow and higher endothelin relative to nitric oxide metabolite levels are associated with reductions in birth weight at high altitude. *Am J Physiol Regul Integr Comp Physiol* 295:R906–915.
- Julian CG, Vargas E, Armaza JF, Wilson MJ, Niermeyer S, Moore LG. 2007. High-altitude ancestry protects against hypoxia-associated reductions in fetal growth. *Arch Dis Child Fetal Neonatal Ed* 92: F372–377.
- Julian CG, Wilson MJ, Lopez M, Yamashiro H, Tellez W, Rodriguez A, Bigham AW, Shriver MD, Rodriguez C, Vargas E, Moore LG. 2009. Augmented uterine artery blood flow and oxygen delivery protect Andeans from altitude-associated reductions in fetal growth. *Am J Physiol Regul Integr Comp Physiol* 296:R1564–1575.
- Karn MN, Penrose LS. 1951. Birth weight and gestation time in relation to maternal age, parity and infant survival. *Ann Eugen* 16:147–164.
- Katzmarzyk PT, Leonard WR. 1998. Climatic influences on human body size and proportions: Ecological adaptations and secular trends. *Am J Phys Anthropol* 106:483–503.
- Kramer MS, Séguin L, Lydon J, Goulet L. 2000. Socio-economic disparities in pregnancy outcome: Why do the poor fare so poorly? *Paediatr Perinat Epidemiol* 14:194–210.
- Krampl E, Lees C, Bland JM, Espinoza Dorado J, Moscoso G, Campbell S. 2000. Fetal biometry at 4300 m compared to sea level in Peru. *Ultrasound Obstet Gynecol* 16:9–18.
- Lampl M, Kuzawa CW, Jeanty P. 2003. Prenatal smoke exposure alters growth in limb proportions and head shape in the midgestation human fetus. *Am J Hum Biol* 15:533–546.
- Lawson DW, Mace R. 2008. Sibling configuration and childhood growth in contemporary British families. *Int J Epidemiol* 37:1408–1421.
- Li L, Dangour AD, Power C. 2007. Early life influences on adult leg and trunk length in the 1958 British birth cohort. *Am J Hum Biol* 19:836–843.
- Li L, Power C. 2004. Influences on childhood height: Comparing two generations in the 1958 British birth cohort. *Int J Epidemiol* 33:1320–1328.
- Lohman TG, Roche AF, Martorell R, editors. 1988. *Anthropometric Standardization Reference Manual*. Champaign, Illinois: Human Kinetics Books.
- Masterson Creber RM, Smeeth L, Gilman RH, Miranda JJ. 2010. Physical activity and cardiovascular risk factors among rural and urban groups and rural-to-urban migrants in Peru: A cross-sectional study. *Rev Panam Salud Publica* 28:1–8.
- McClung J. 1969. *Effects of High Altitude on Human Birth: Observations on Mothers, Placentas, and the Newborn in Two Peruvian Populations*. Cambridge, MA: Harvard University Press.
- Miranda JJ, Gilman R, Garcia H, Smeeth L. 2009. The effect on cardiovascular risk factors of migration from rural to urban areas in Peru: PERU MIGRANT Study. *BMC Cardiovasc Disord* 9:23.
- Moore LG, Young D, McCullough RE, Droma T, Zamudio S. 2001. Tibetan protection from intrauterine growth restriction (IUGR) and reproductive loss at high altitude. *Am J Hum Biol* 13:635–644.
- Mortola JP, Frappell PB, Aguero L, Armstrong K. 2000. Birth weight and altitude: A study in Peruvian communities. *J Pediatr* 136:324–329.
- Mueller WH, Schull VN, Schull WJ, Soto P, Rothhammer F. 1978a. A multinational Andean genetic and health program: Growth and development in an hypoxic environment. *Ann Hum Biol* 5:329–352.
- Mueller WH, Yen F, Rothhammer F, Schull WJ. 1978b. Multinational Andean genetic and health program: VII. Lung function and physical growth - multivariate analyses in high- and low-altitude populations. *Aviat Space Environ Med* 49:1188–1196.

- Niermeyer S, Andrade Mollinedo P, Huicho L. 2009. Child health and living at high altitude. *Arch Dis Child* 94:806–811.
- Palomino H, Mueller WH, Schull WJ. 1978. Altitude, heredity and body proportions in northern Chile. *Am J Phys Anthropol* 50:39–50.
- Pawson IG, Huicho L. 2010. Persistence of growth stunting in a Peruvian high altitude community, 1964–1999. *Am J Hum Biol* 22:367–374.
- Pawson IG, Huicho L, Muro M, Pacheco A. 2001. Growth of children in two economically diverse Peruvian high-altitude communities. *Am J Hum Biol* 13:323–340.
- Pomeroy E, Stock JT, Stanojevic S, Miranda JJ, Cole TJ, Wells JCK. 2012. Trade-offs in relative limb length among Peruvian children: Extending the thrifty phenotype hypothesis to limb proportions. *PLoS One* 7: e51795.
- Pomeroy E, Stock JT, Stanojevic S, Miranda JJ, Cole TJ, Wells JCK. 2013. Associations between arterial oxygen saturation, body size and limb measurements among high-altitude Andean children. *Am J Hum Biol* 25:629–636.
- Pomeroy E, Wells JCK, Stanojevic S, Jaime Miranda J, Cole TJ, Stock JT. 2014. Birth month associations with height, head circumference, and limb lengths among Peruvian children. *Am J Phys Anthropol* 154:115–124.
- Postigo L, Heredia G, Illsley NP, Torricos T, Dolan C, Echalar L, Tellez W, Maldonado I, Brimacombe M, Balanza E, Vargas E, Zamudio S. 2009. Where the O₂ goes to: preservation of human fetal oxygen delivery and consumption at high altitude. *J Physiol* 587:693–708.
- Raum E, Arabin B, Schlaud M, Walter U, Schwartz FW. 2001. The impact of maternal education on intrauterine growth: A comparison of former West and East Germany. *Int J Epidemiol* 30:81–87.
- Relethford J. 1995. The use of surnames in the study of human variation and plasticity. In: Mascie-Taylor G, Bogin B, editors. *Human variability and plasticity*. Cambridge: Cambridge University Press. p 146–158.
- Rivera-Ch M, Castillo A, Huicho L. 2008. Hypoxia and other environmental factors at high altitude. *Int J Environ Health* 2:92–106.
- Roberts DF. 1978. *Climate and Human Variability*. Menlo Park, CA: Cummings.
- Sanchez E, Webb RD, Rasmussen A, Kelly JA, Riba L, Kaufman KM, Garcia-de la Torre I, Moctezuma JF, Maradiaga-Ceceña MA, Cardiel-Rios MH, Acevedo E, Cucho-Venegas M, Garcia MA, Gamron S, Pons-Estel BA, Vasconcelos C, Martin J, Tusié-Luna T, Harley JB, Richardson B, Sawalha AH, Alarcón-Riquelme ME. 2010. Genetically determined amerindian ancestry correlates with increased frequency of risk alleles for systemic lupus erythematosus. *Arthritis Rheum* 62:3722–3729.
- Schull WJ, Rothhammer F. 1977. A Multinational Andean Genetic and Health Programme: Rationale and design for a study of adaptation to the hypoxia of altitude. In: Weiner JS, editor. *Physiological Variation and its Genetic Basis* London: Taylor and Francis.
- Semba RD, de Pee S, Sun K, Sari M, Akhter N, Bloem MW. 2008. Effect of parental formal education on risk of child stunting in Indonesia and Bangladesh: A cross-sectional study. *Lancet* 371:322–328.
- Serrat MA, King D, Lovejoy CO. 2008. Temperature regulates limb length in homeotherms by directly modulating cartilage growth. *Proc Natl Acad Sci USA* 105:19348–19353.
- Sørensen HT, Sabroe S, Rothman KJ, Gillman M, Steffensen FH, Fischer P, Serensen TIA. 1999. Birth Weight and length as predictors for adult height. *Am J Epidemiol* 149:726–729.
- Soria R, Julian CG, Vargas E, Moore LG, Giussani DA. 2013. Graduated effects of high-altitude hypoxia and highland ancestry on birth size. *Pediatr Res* 74:633–638.
- Sterling R, Miranda JJ, Gilman RH, Cabrera L, Sterling CR, Bern C, Checkley W. 2012. Early anthropometric indices predict short stature and overweight status in a cohort of Peruvians in early adolescence. *Am J Phys Anthropol* 148:451–461.
- Stinson S. 1980. The physical growth of high altitude Bolivian Aymara children. *Am J Phys Anthropol* 52:377–385.
- Stinson S. 1982. The effect of high altitude on the growth of children of high socioeconomic status in Bolivia. *Am J Phys Anthropol* 59:61–71.
- Stinson S. 2009. Nutritional, developmental, and genetic influences on relative sitting height at high altitude. *Am J Hum Biol* 21:606–613.
- Stinson S, Frisancho AR. 1978. Body proportions of highland and lowland Peruvian Quechua children. *Hum Biol* 50:57–68.
- Tarazona-Santos E, Lavine M, Pastor S, Fiori G, Pettener D. 2000. Hematological and pulmonary responses to high altitude in Quechuas: A multivariate approach. *Am J Phys Anthropol* 111:165–176.
- Trinkaus E. 1981. Neanderthal limb proportions and cold adaptation. In: Stringer CB, editor. *Aspects of Human Evolution*. London: Taylor and Francis. p 187–224.
- Ulijaszek SJ, Kerr DA. 1999. Anthropometric measurement error and the assessment of nutritional status. *Br J Nutr* 82:165–177.
- van den Berg G, van Eijsden M, Galindo-Garre F, Vrijkotte TGM, Gemke RBB. 2013. Smoking overrules many other risk factors for small for gestational age birth in less educated mothers. *Early Hum Dev* 89:497–501.
- Vargas M, Vargas E, Julian CG, Armaza JF, Rodriguez A, Tellez W, Niermeyer S, Wilson M, Parra E, Shriver M, Moore LG. 2007. Determinants of blood oxygenation during pregnancy in Andean and European residents of high altitude. *Am J Physiol Regul Integr Comp Physiol* 293: R1303–1312.
- Wells JC. 2013. The thrifty phenotype and the hierarchical preservation of tissues under stress. *Int J Epidemiol* 42:1223–1227.
- Whitley E, Gunnell D, Davey Smith G, Holly JMP, Martin RM. 2008. Childhood circumstances and anthropometry: The Boyd Orr cohort. *Ann Hum Biol* 35:518–534.
- Whittaker KL. 1992. The relationship of forced vital capacity to morphology and fat-free mass in high-altitude adult Aymara men. In: Eckhardt RB, Melton TW, editors. *Population Studies on Human Adaptation and Evolution in the Peruvian Andes*: Matson Museum of Anthropology, Pennsylvania State University. p 50–62.
- Wilcox AJ, Russell IT. 1983. Birthweight and Perinatal Mortality: II. On Weight-Specific Mortality. *Int J Epidemiol* 12:319–325.
- Wiley AS. 1994. Neonatal size and infant mortality at high altitude in the western Himalaya. *Am J Phys Anthropol* 94:289–305.
- Wilson HJ, Dickinson F, Griffiths PL, Azcorra H, Bogin B, Varela-Silva MI. 2011. How useful is BMI in predicting adiposity indicators in a sample of Maya children and women with high levels of stunting? *Am J Hum Biol* 23:780–789.
- Wilson MJ, Lopez M, Vargas M, Julian C, Tellez W, Rodriguez A, Bigham A, Armaza JF, Niermeyer S, Shriver M, Vargas E, Moore LG. 2007. Greater uterine artery blood flow during pregnancy in multigenerational (Andean) than shorter-term (European) high-altitude residents. *Am J Physiol Regul Integr Comp Physiol* 293:R1313–1324.
- Yliharsila H, Kajantie E, Osmond C, Forsen T, Barker DJP, Eriksson JG. 2007. Birth size, adult body composition and muscle strength in later life. *Int J Obes* 31:1392–1399.
- Zamudio S, Droma T, Norkyel KY, Acharya G, Zamudio JA, Niermeyer SN, Moore LG. 1993. Protection from intrauterine growth retardation in Tibetans at high altitude. *Am J Phys Anthropol* 91:215–224.
- Zamudio S, Postigo L, Illsley NP, Rodriguez C, Heredia G, Brimacombe M, Echalar L, Torricos T, Tellez W, Maldonado I, Balanza E, Alvarez T, Ameller J, Vargas E. 2007. Maternal oxygen delivery is not related to altitude- and ancestry-associated differences in human fetal growth. *J Physiol* 582:883–895.