

Li, C, Tian, Q, Yu, R, Zhou, B, Xia, J, Burke, C, Dong, B, Tett, SFB, Freychet, N, Lott, F and Ciavarella, A

Attribution of extreme precipitation in the lower reaches of the Yangtze River during May 2016

<http://researchonline.ljmu.ac.uk/id/eprint/8928/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Li, C, Tian, Q, Yu, R, Zhou, B, Xia, J, Burke, C, Dong, B, Tett, SFB, Freychet, N, Lott, F and Ciavarella, A (2018) Attribution of extreme precipitation in the lower reaches of the Yangtze River during May 2016. Environmental Research Letters. 13 (1). ISSN 1748-9326

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

LETTER • OPEN ACCESS

Attribution of extreme precipitation in the lower reaches of the Yangtze River during May 2016

To cite this article: Chunxiang Li *et al* 2018 *Environ. Res. Lett.* **13** 014015

View the [article online](#) for updates and enhancements.

Related content

- [Attribution of the July–August 2013 heat event in Central and Eastern China to anthropogenic greenhouse gas emissions](#)
Shuangmei Ma, Tianjun Zhou, Dáithí A Stone *et al.*
- [Multi-method attribution analysis of extreme precipitation in Boulder, Colorado](#)
Jonathan M Eden, Klaus Wolter, Friederike E L Otto *et al.*
- [A comparison of model ensembles for attributing 2012 West African rainfall](#)
Hannah R Parker, Fraser C Lott, Rosalind J Cornforth *et al.*

Environmental Research Letters



LETTER

OPEN ACCESS

RECEIVED

24 June 2017

REVISED

31 August 2017

ACCEPTED FOR PUBLICATION

27 October 2017

PUBLISHED

11 January 2018

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Attribution of extreme precipitation in the lower reaches of the Yangtze River during May 2016

Chunxiang Li¹, Qinhuo Tian², Rong Yu³, Baiquan Zhou^{4,8}, Jiangjiang Xia¹, Claire Burke⁵, Buwen Dong⁶, Simon F B Tett⁷, Nicolas Freychet⁷, Fraser Lott⁵ and Andrew Ciavarella⁵

¹ CAS Key Laboratory of Regional Climate-Environment for Temperate East Asia, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, People's Republic of China

² National Climate Center, China Meteorological Administration, Beijing, People's Republic of China

³ College of Atmospheric Science, Nanjing University of Information Science & Technology, Nanjing 210044

⁴ State Key Laboratory of Severe Weather, Chinese Academy of Meteorological Sciences, CMA, Beijing, People's Republic of China

⁵ Met Office Hadley Centre, Met Office, United Kingdom

⁶ National Centre for Atmospheric Science, Department of Meteorology, University of Reading, United Kingdom

⁷ School of Geosciences, University of Edinburgh, Edinburgh, United Kingdom

⁸ Author to whom any correspondence should be addressed.

E-mail: quan20080141@126.com

Keywords: extreme rainfall, extreme event attribution, El Niño, risk ratio, anthropogenic influence

Supplementary material for this article is available [online](#)

Abstract

May 2016 was the third wettest May on record since 1961 over central eastern China based on station observations, with total monthly rainfall 40% more than the climatological mean for 1961–2013. Accompanying disasters such as waterlogging, landslides and debris flow struck part of the lower reaches of the Yangtze River. Causal influence of anthropogenic forcings on this event is investigated using the newly updated Met Office Hadley Centre system for attribution of extreme weather and climate events. Results indicate that there is a significant increase in May 2016 rainfall in model simulations relative to the climatological period, but this increase is largely attributable to natural variability. El Niño years have been found to be correlated with extreme rainfall in the Yangtze River region in previous studies—the strong El Niño of 2015–2016 may account for the extreme precipitation event in 2016. However, on smaller spatial scales we find that anthropogenic forcing has likely played a role in increasing the risk of extreme rainfall to the north of the Yangtze and decreasing it to the south.

1. Introduction

In the context of global warming, more intense and more frequent extreme precipitation events have long been projected (Hartmann *et al* 2013, Hirsch and Archfield 2015). Faced with increasing climate change risks, society around the world needs a better understanding of such risks to prevent climate-related hazards, especially for regions with high exposure and vulnerability. China, occupying a region as large as Europe, has experienced more intense and frequent precipitation extremes in recent several decades (Zhai and Sun 1999, Yan and Yang 2000, Zhai *et al* 2005, Qian *et al* 2007). The lower reaches of the Yangtze River Valley—one of the most important industrial and agricultural zones, and densely populated regions in

China—experienced extreme rainfall in May 2016 during which there was one day when the daily precipitation broke the 56-year May daily maximum records at 25 stations (CMA 2016). This heavy and sustained precipitation event led to waterlogging, landslides, debris flow and some other disasters over part of central eastern China, resulting in severe damage to crops and disrupting agricultural production (CMA 2016). Due to the damage-causing capacity and general increasing trend of persistent heavy rainfall (Zolina *et al* 2010, Chen and Zhai 2013, IPCC AR5 2013), changes in the risk of such precipitation extremes are of great concern to policy-makers and the general public.

Min *et al* (2011) showed that anthropogenic forcings have contributed to the observed intensification of heavy precipitation events over large parts of the

Northern Hemisphere's land area. Zhang *et al* (2013) found consistency between the multimodel simulated response to the effects of anthropogenic forcing and observed changes in extreme precipitation on average over Northern Hemisphere land. There also exists evidence indicating that changes in heavy rainfall in some regions of southeast China is attributable to anthropogenic forcings (e.g. Burke *et al* 2016). Though there are numerous studies focusing on this issue, geographical coverage of events remains patchy (Stott *et al* 2016). In addition, there is a growing demand to do attribution analysis of extreme events timely to inform adaptation planning for flood defences and to support climate risk management in China. In this study, we intend to address how the probability of anomalous wet conditions, similar to those in May 2016 in the lower reaches of the Yangtze River Valley, has changed due to human-induced climate change.

2. Data

The observed daily precipitation data are obtained from Climate Data Center of China National Meteorological Information Center (NMIC), which span the period of 1961–2016. After conducting rigorous quality control as described in Chen and Zhai (2014), 1277 stations are retained around China. Two model ensembles generated by the newly updated Met Office Hadley Centre system (HadGEM3-GA6, Walters *et al* 2017) for Attribution of extreme weather and Climate Events (updated from Christidis *et al* 2013, see Ciavarella *et al* 2017) at a N216 resolution ($0.56^\circ \times \sim 0.83^\circ$ horizontal resolution) are used in our attribution analysis. We use two ensembles. One is forced with a combination of anthropogenic and natural forcings (ALL) and the other is forced by natural forcing only (NAT). The ALL experiment employs observed sea surface temperatures (SSTs) and sea-ice data (HadISST; Rayner *et al* 2003) as boundary conditions. The boundary conditions for the NAT experiment are provided from the HadISST observational dataset minus an estimate of the anthropogenic contribution derived from the Coupled Model Intercomparison Project Phase 5 (CMIP5) coupled model simulations. More detailed information about the system and forcings used can be found in Christidis *et al* (2013). During the period 1961–2013, each ensemble comprises 15 members, subsequently expanded to 105 and 525 members for 2014–2015 and 2016 respectively. Since 2014–2015 is used for test cases and ensemble size is not consistent with that for 1961–2013, the reference period is chosen as 1961–2013.

We also adopt the ALL experiments of 37 models (supplementary table S1, available at stacks.iop.org/ERL/13/014015/mmedia) from CMIP5 (Taylor *et al* 2012) to further assess whether the attribution results with HadGEM3-GA6 are robust. Since the ALL experiments end at 2005, we take 1986–2005 as our present day (PD) and 1901–1920 as an early period (EP). By

comparing the changes between two periods, we assess impact of changes in forcing, presumably dominated by changes in anthropogenic forcing, on probability of extreme event from EP to PD.

3. Methodology

For observed monthly mean precipitation in May, the average is a simple average of the station data. For data from model ensembles, the average is calculated from the area-weighted model grid box which fall within the mesh containing some station observations. The mesh grid is the same as the model grid. We use the 'risk ratio' (RR), a probabilistic extreme event attribution approach (Allen 2003, National Academies of Sciences, Engineering, and Medicine 2016), to characterize the anthropogenic contribution to the occurrence of the extreme event, by considering occurrence probabilities $P(\text{ALL})$ and $P(\text{NAT})$. These quantities represent the probability that the risk change of an event in the 'real' world with all forcings, $P(\text{ALL})$, to that in the 'natural' world with only natural forcings, $P(\text{NAT})$. From these estimated probabilities, the RR can be calculated, where $\text{RR} = P(\text{ALL})/P(\text{NAT})$. Bootstrapping with replacement from 1000 samples is used to estimate the error on the probability and risk ratio for the occurrence of an event for at least 90% confidence interval. The 90% confidence interval extends from 5%–95% percentiles of the probabilities or risk ratios achieved from the 1000 probability density functions (PDFs) of the 1000 bootstrapped samples. If the resample distribution is slightly asymmetric, the wider side of the distribution is used to define the width of the confidence interval.

3.1. Region selection

We examine the region (117°E – 121°E and 26°N – 34°N), containing 108 stations, which have continuously recorded since 1961, as depicted in figure 1(b) in the lower reaches of the Yangtze River Valley, where persistent heavy precipitation in May 2016 was concentrated. As indicated in figure 1(a) and table 1, the area-averaged monthly mean precipitation in May 2016 over this region is 7.9 mm day^{-1} , 40% more than the climatology (5.6 mm day^{-1}) over the period of 1961–2013, and is the third highest during the period of 1961–2016.

Rainfall tends to be a very localized phenomenon, evidenced by a dipole pattern of anomalous rainfall with opposite anomalies presented in Wang *et al* (2011) between the areas north and south of about 30°N in the lower reaches of Yangtze River Valley over the Meiyu period at interannual time scales. In addition, past research has revealed differences of attributable anthropogenic influence in spatially adjacent regions (e.g. Min *et al* 2011, Burke *et al* 2016, Burke and Stott 2017). Therefore, we divided the flood-affected




Figure 1. (a) Standardized anomaly of monthly mean precipitation, average of available station data within the study region for May over central eastern China. Red dashed line indicates the 1.5σ threshold, equivalent to the magnitude of 2016 anomaly. (b) Distribution of stations (black dots) in the study region.

Table 1. Areal mean May precipitation and precipitation inter-annual variability for the climatological period (1961–2013) and 2016.

Statistic\Year	Observations		ALL Model	
	1961–2013	2016	1961–2013	2016
Mean total rainfall (mm day^{-1})	5.6	7.9	6.8	8.0
Standard deviation for $R(i)$ time series (mm day^{-1})	1.5	/	2.0	/

area into two sub-regions, the Northern (30°N – 34°N) and Southern parts (26°N – 30°N), to examine anthropogenic influences in these two regions separately.

3.2. Threshold

Normalized departures from climatology are widely used to objectively rank synoptic-scale events (e.g. Hart and Grumm 2001, Junker *et al* 2009, Graham and Grumm 2010). We can objectively compare precipitation variability and extremes across space and time by normalizing climate data relative to a reference period (Duan *et al* 2015, Jiang *et al* 2016). In this study, in May of each year i , the standardized anomaly, $RA(i)$, is derived by subtracting the local mean, μ , from the area-average monthly mean precipitation of the available station data within the study region, $R(i)$, and divided by the corresponding standard deviation of inter-annual variability, σ , estimated from the reference period (1961–2013). It can be written as;

$$RA(i) = \frac{R(i) - \mu}{\sigma}, \quad (1)$$

$RA(i)$ is used to rank events based on departures from the local climatology in units of standard deviation (σ).

The observed areal averaged precipitation anomaly for May 2016 exceeds 1.5σ from the 1961–2013 mean, as shown in figure 1(a). The PDF of RA for all years between 1961–2013 indicates that the probability of precipitation anomaly greater than that for May 2016 is about 4% (figure 3(a)), corresponding to a return time of at least 13 years at 90% confidence level. The anomaly of 1.5σ is selected as the threshold for extreme precipitation for our attribution analysis. Using the same method, we calculated anomalies and thresholds for the two sub-regions for both observations and model simulations.

The thresholds are 1.66σ and 1.86σ for the Northern part and Southern part respectively (see table 2).

3.3. Model assessment

RA for both the ALL and NAT model simulations were calculated in the same way as in observations, but precipitation means and standard deviations were computed using all ensemble members of the ALL experiment in the period 1961–2013 using all data in the regions. Then for each ensemble member, all simulations were normalized using these values. By doing this, any systematic bias in area-averaged monthly mean precipitation and in inter-annual variability in model simulations are removed in the normalized precipitation time series.

Evaluation of the model simulations was carried out to see if the model could accurately reproduce the characteristics of precipitation in this region. Comparison of the normalized time series between the model ensemble and the observations indicates that the model is generally skillful in reproducing characteristics of May precipitation for the whole target region (figure 2(b)), southern (figure 2(d)) and northern sub-regions (figure 2(f)). The inter-annual variability of the observed time series is included in the range of the model ensemble for these three study regions (with the exception of 1963 and 1973). On average over the entire region, the model overestimates the absolute amount of monthly rainfall by about 21% (see table 1) and its interannual variability by about 33%.

The PDFs of the standardized precipitation anomaly for the observations and model simulations during our climatological period 1961–2013 are visually similar to each other (figure 2(a)). A Kolmogorov–Smirnov two-sample test (Wilks 2006), confirms that there is no significant difference between the





Table 2. The probability of the extreme precipitation with intensity as high as that in May 2016 for various forcings over three different regions.

	Whole region	Northern region	Southern region
Observed 1961–2013	0.057 ± 0.05	0.038 ± 0.03	0.057 ± 0.05
ALL 1961–2013	0.108 ± 0.02	0.053 ± 0.01	0.053 ± 0.01
NAT 2016	0.259 ± 0.03	0.095 ± 0.02	0.133 ± 0.03
ALL 2016	0.261 ± 0.03	0.156 ± 0.03	0.091 ± 0.02

distributions constructed with ALL experiments and observations over period 1961–2013 (P -value of 0.67). Low P -values are needed to reject the null hypothesis that two data are from the same continuous distribution. For the northern and southern sub-regions, ALL experiments and observations over period 1961–2013 also have similar PDF distributions with P -values of 0.93 and 0.99 respectively. Whilst the model overestimates the absolute amount of precipitation during May and its interannual variability, the standardized precipitation anomaly corrects these systematic biases. Using this metric, the corrected model precipitation performs well in reproducing May precipitation variability and is suitable for attribution studies of May precipitation in the lower reaches of the Yangtze River Valley.

4. Results

For the whole target region, the probability of areal averaged precipitation anomaly greater than the threshold defined above (1.5σ) is 11% for the ALL experiments over the period 1961–2013 (see figure 3(a)). This probability increases to 26% in 2016. For 2016, the risk of an extreme event with total rainfall as high as that in May 2016 is 2.4 times as large as the climatology. From figure 3, comparing the PDFs of both the ALL and NAT experiments for 2016 with those for climatology, it appears that natural forcing dominates in increasing the probability of the 2016 extreme event. As shown in table 2, the probability of exceeding the threshold of 1.5σ in ALL experiment for 2016 is $P(\text{ALL}) = 0.261 \pm 0.03$, while that in NAT



experiment is $P(\text{NAT}) = 0.259 \pm 0.03$. Correspondingly, the estimated risk ratio is $\text{RR} = 1.01 \pm 0.17$, indicating little change in the probability of such event due to human-induced climate change. The strong El Niño that occurred in 2015/2016 may account for the significant enhancement of May precipitation in the Yangtze River Valley in China.

The link between the El Niño Southern Oscillation (ENSO) signal and the spring precipitation over the Yangtze river has been demonstrated in several studies (Zhang and Sumi 2002, Ying *et al* 2015, Zhai *et al* 2016). As it is a main control of the precipitation inter-annual variability, it is important to verify that the model can reproduce this link. To do so, the correlation between May precipitation and May sea level pressure is computed. The correlation with the previous winter surface temperature is also computed (when ENSO signal is the strongest). Results are displayed in figure 4 for ERA Interim reanalysis and for the ensemble mean of 15 members of the N216-ALL simulations. This analysis was also conducted with the NCEP-NCAR reanalysis. Results were found to be very close to ERA Interim and are not displayed. The sea level pressure correlation during May (figures 4(a) and (b)) shows a clear positive pattern over the western North Pacific, associated with an anticyclonic anomaly and an increased transport of moisture from the tropics along the East China coast. Moreover, the ENSO signal is clear in the winter temperature (figures 4(c) and (d)) for both model simulations and reanalysis. Although ENSO

may be climate change influenced, ENSO variability, even in the 21st century seems to be within the bounds of the natural variation range based on model simulations (Chen *et al* 2017). The correlations in the model and in reanalysis show consistent patterns which are closely related to May precipitation over the Yangtze river. Along with greenhouse warming, the extreme rainfall events associated with ENSO events are expected to increase as frequency of extreme El Niño events increases (Cai *et al* 2014, 2015). A stronger correlation in the model simulations may be due to a too stronger impact of the ocean on the atmosphere, due to the SST forcing, or to other processes affecting the May precipitation not represented in the model. Nevertheless, it appears that the HadGEM3-GA6 N216 model is able to reproduce the main natural variability and is consistent with previous studies on interannual variability in the Yangtze river basin precipitation (Zhang and Sumi 2002, Ying *et al* 2015, Zhai *et al* 2016). Therefore, the shift in PDFs of both ALL and NAT experiments for May 2016 compared to climatology, as seen in figure 3, suggests that El Niño had a significant effect on rainfall for this region in 2016. Furthermore, the impact of El Niño seems to be greater than that of anthropogenic climate change in the lower Yangtze valley.

Interestingly, an effect from anthropogenic climate change on the changing risk of extreme precipitation is discovered for the two sub-regions. From the PDFs in figure 3(b), we find that the probability of the event occurring in the northern sub-region is larger in




Figure 4. Correlation between the May precipitation over the Yangtze river and May sea level pressure for ERA Interim (a) and N216-ALL ensemble mean (b). Correlation between the May precipitation over the Yangtze river and the 2 m temperatures in the preceding DJF for ERA Interim (c) and the near-surface air temperature for N216 ensemble mean (d). The precipitation time series used for the correlation with ERA Interim variables is that used in figure 1 (observations). For the N216 ensemble, each member is correlated with its precipitation signal before averaging the ensemble correlations.




Figure 5. The probability density functions (PDFs) of May monthly precipitation anomaly estimated from CMIP5 models for 1901–1920 and 1986–2005 respectively in the three parts of the study region: (a) the whole target region, (b) the northern part of the target region, (c) the southern part of the target region. Blue lines correspond to the CMIP5 multi-model mean for 1901–1920; red for 1986–2005; corresponding shadings denote the full range of 37 CMIP5 models. The vertical lines are the standardized observed precipitation anomalies of May 2016, which are chosen as the thresholds.

the ALL experiment for 2016 compared to the NAT experiment, with a RR of 1.64 ± 0.52 . In contrast, risk of extreme precipitation in the southern sub-region shows a decline due to human-induced climate change ($RR = 0.68 \pm 0.22$). To check the robustness of this result, 37 CMIP5 models are used to examine whether there exists same risk change of extreme May precipitation in ALL experiments. Change of risk in extreme May precipitation are estimated between the period 1986–2005 and the early period 1901–1920.

For the whole region and southern region, there seems to be no significant anthropogenic impact on risk of regional extreme rainfall with the RR around 1. However, the RR rises to 1.17 ± 0.43 in the northern region. The risk change in extreme May precipitation is generally consistent with those achieved using HadGEM3-GA6, which demonstrates the results we get is not sensitive to the model selected. Notably, the extent of anthropogenic influence on these two sub-regions differs considerably. This highlights the spatial




Figure 6. The probability density functions (PDFs) for the standardized anomaly of May mean precipitation difference between the northern sub-region and the southern sub-region. Red lines correspond to HadGEM3-GA6 all forcings run (ALL); blue is natural forcings only run (NAT); dashed lines indicate historical (1961–2013) PDF and solid lines indicate 2016; black indicates observations. The black vertical line is the value for 2016.

variability of rainfall and the sensitivity of extreme precipitation attribution to the selection of study region. More importantly, in the condition of nearly negligible human-induced risk change of extreme precipitation for the whole study region, anthropogenic influences played a role in increasing the probability of extreme precipitation events as intense as that in May 2016 in the northern sub-region and decreasing that in the southern sub-region. Thus, we have reason to believe that human influences have shifted the precipitation northwards in the study region. Similar detection was conducted on mean precipitation difference between the northern sub-region and the southern sub-region. The threshold for risk estimation is also selected as the standardized precipitation difference between the northern sub-region and the southern sub-region in May 2016. The result shown in figure 6 implies that human influences have increased the probability of positive anomaly of difference between the two regions, with a risk ratio of 1.73 ± 0.43 at 90% confidence level based on the threshold of May 2016. It evidences that human influences contribute to the shift of precipitation from the southern sub-region to the northern, thus causing the corresponding increase and decrease of occurring risk of extreme precipitation in the northern and southern sub-region respectively. Based on the urban map of East Asia (Schneider *et al* 2009, Zhang *et al* 2010), several megacities like Shanghai, Nanjing, and Hangzhou are clustered in the northern sub-region. This region of high urbanization which is exposed to higher risks of extreme precipitation due to anthropogenic influence has implication for city planning and mitigation of extremes.

5. Conclusion

In May 2016, persistent extreme precipitation was observed over the region located in the lower reaches of the Yangtze valley. We find that the increasing likelihood of an extreme rainfall event as intense as that observed in May 2016 is mainly attributable to natural variability, presumably the strong El Niño of

2015–2016, with a risk 2.4 times as large as the climatology. The attributable contribution from anthropogenic climate change is negligible. Interestingly, for the southern sub-region, human influences appear to have reduced the risk of such event with precipitation greater than or equal to that observed in May 2016. Conversely, anthropogenically forced climate change has increased the probability of this kind of intense rainfall in the northern sub-region by a factor of 1.64. Through it is difficult to estimate the role of ENSO on the risk of precipitation extremes using CMIP5 simulations, results show the change in risk of extreme precipitation due to all forcing changes between two periods, presumably dominated by changes in anthropogenic forcing, in CMIP5 models is consistent with HadGEM3-GA6 model. This evidences that the results we found is not sensitive to the model used. Further attribution on precipitation difference between the two regions evidences that anthropogenic influences have shifted precipitation from the southern sub-region to the northern, thus causing the corresponding increase and decrease of occurring risk of extreme precipitation in the two sub-regions respectively.

Acknowledgments

We thank Dr Siyan Dong for helpful comments and discussions. LC and JX were funded by the National Key R&D Program of China (2016YFA0600400) and the National Natural Science Foundation of China (NFSC, 41605066), QT was funded by the NFSC (41675074) and Climate Change Project (CCSF201704), RY was funded by the Postgraduate Research and Practice Innovation Program of Jiangsu Province (KYLX16_0941), and BZ was funded by the NFSC (41575094). This work was largely carried out during a workshop on Operational Attribution at the University of Edinburgh supported by the UK-China Research & Innovation Partnership Fund through the Met Office Climate Science for Service Partnership (CSSP) China as part of the Newton Fund, China.

ORCID iDs

Baiquan Zhou  <https://orcid.org/0000-0002-3709-5336>

Buwen Dong  <https://orcid.org/0000-0003-0809-7911>

Simon F B Tett  <https://orcid.org/0000-0001-7526-560X>

Nicolas Freychet  <https://orcid.org/0000-0003-2207-4425>

References

- Allen M 2003 Liability for climate change *Nature* **421** 891–2
- Burke C, Stott P A, Sun Y and Ciavarella A 2016 Attribution of extreme rainfall in Southeast China during may 2015 *Bull. Am. Meteorol. Soc.* **97** 92–6
- Burke C and Stott P A 2017 Impact of anthropogenic climate change on the East Asian summer monsoon *J. Clim.* **30** 5205–20
- Cai W *et al* 2014 Increasing frequency of extreme El Niño events due to greenhouse warming *Nat. Clim. Change* **4** 111–6
- Cai W *et al* 2015 ENSO and greenhouse warming *Nat. Clim. Change* **5** 849–59
- Chen C *et al* 2017 ENSO in the CMIP5 simulations: life cycles, diversity, and responses to climate change *J. Clim.* **30** 775–801
- Christidis N, Stott P, Scaife A, Arribas A, Jones G, Copset D, Knight J and Tennant W 2013 A new HadGEM3-A-based system for attribution of weather- and climate-related extreme events *J. Clim.* **26** 2756–83
- CMA 2016 (www.scio.gov.cn/xwfbh/gbwxxwfbh/xwfbh/qxj/Document/1479487/1479487.htm) (in Chinese)
- Chen Y and Zhai P 2013 Persistent extreme precipitation events in China during 1951–2010 *Clim. Res.* **57** 143–55
- Chen Y and Zhai P 2014 Changing structure of wet periods across Southwest China during 1961–2012 *Clim. Res.* **61** 123–31
- Ciavarella A *et al* 2017 An even newer operational attribution system (unpublished)
- Duan W, He B, Takara K, Luo P, Hu M, Alias N E and Nover D 2015 Changes in precipitation amounts and extremes over Japan between 1901 and 2012 and their connection to climate indices *Clim. Dyn.* **45** 1–20
- Graham R A and Grumm R H 2010 Utilizing normalized anomalies to assess synoptic-scale weather events in the Western United States *Weath. Forecast.* **25** 428–45
- Hartmann D L *et al* 2013 Observations: atmosphere and surface *Climate Change 2013 the Physical Science Basis: Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (Cambridge: Cambridge University Press) pp 159–254 (<https://doi.org/10.1017/CBO9781107415324.008>)
- Hart R and Grumm R H 2001 Using normalized climatological anomalies to rank synoptic-scale events objectively *Mon. Weath. Rev.* **129** 2426–42
- Hirsch R M and Archfield S A 2015 Flood trends: Not higher but more often *Nat. Clim. Change* **5** 198–9
- IPCC 2013 Climate change 2013: the physical science basis *Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* ed T F Stocker *et al* (Cambridge: Cambridge University Press)
- Jiang N, Qian W, Du J, Grumm R H and Fu J 2016 A comprehensive approach from the raw and normalized anomalies to the analysis and prediction of the Beijing extreme rainfall on 21 July 2012 *Nat. Hazards* **84** 1551–67
- Junker N W, Brennan M J, Pereira F, Bodner M J and Grumm R H 2009 Assessing the potential for rare precipitation events with standardized anomalies and ensemble guidance at the hydrometeorological prediction center *Bull. Am. Meteorol. Soc.* **90** 445–53
- Min S K, Zhang X, Zwiers F W and Hegerl G C 2011 Human contribution to more-intense precipitation extremes *Nature* **470** 378–81
- National Academies of Sciences, Engineering, and Medicine 2016 *Attribution of Extreme Weather Events in the Context of Climate Change* (Washington, DC: The National Academies Press) (<https://doi.org/10.17226/21852>)
- Qian W, Fu J, Zhang W and Lin X 2007 Changes in mean climate and extreme climate in China during the last 40 years *Adv. Earth Sci.* **22** 673–84
- Rayner N A, Parker D E, Horton E B, Folland C K, Alexander L V, Row-ell D P, Kent E C and Kaplan A 2003 Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century *J. Geophys. Res.* **108** 4407
- Schneider A, Friedl M A and Potere D 2009 A new map of global urban extent from MODIS satellite data *Environ. Res. Lett.* **4** 044003
- Stott P A *et al* 2016 Attribution of extreme weather and climate-related events *Wiley Interdiscip. Rev. Clim. Change* **7** 23–41
- Taylor K E, Stouffer R J and Meehl G A 2012 An overview of CMIP5 and the experiment design *Bull. Am. Meteorol. Soc.* **93** 485–98
- Walters D *et al* 2017 The met office unified model global atmosphere 6.0/6.1 and JULES global land 6.0/6.1 configurations *Geosci. Model Dev.* **10** 1487–520
- Wang L, Huang Q, Dai A, Guan Z, He J and Wu Z 2011 Inhomogeneous distributions of Meiyu rainfall in the Jiang-Huai basin, and associated circulation patterns *Clim. Res.* **50** 203–14
- Wilks D S 2006 Statistical methods in the atmospheric sciences *International Geophysics Series* vol 91 (Amsterdam: Elsevier Academic) p 627
- Yan Z and Yang C 2000 Geographic patterns of extreme climate changes in China during 1951–1997 *Clim. Environ. Res.* **5** 267–72
- Ying K, Zhao T, Quan X, Zheng X and Frederiksen C S 2015 Interannual variability of autumn to spring seasonal precipitation in eastern China *Clim. Dyn.* **45** 253–71
- Zhai P and Sun A J 1999 Changes of climatic extremes in China *Clim. Change* **42** 203–18
- Zhai P, Zhang X, Wan H and Pan X 2005 Trends in total precipitation and frequency of daily precipitation extremes over China *J. Clim.* **18** 1096–108
- Zhai P, Yu R, Guo Y, Li Q, Ren X, Wang Y, Xu W, Liu Y and Ding Y 2016 The strong El Niño of 2015/16 and its dominant impacts on global and China's climate *J. Meteorol. Res.* **30** 283–97
- Zhang N, Gao Z, Wang X and Chen Y 2010 Modeling the impact of urbanization on the local and regional climate in Yangtze river delta, China *Theor. App. Climatol.* **102** 331–42
- Zhang R and Sumi A 2002 Moisture circulation over East Asia during El Niño episode in Northern winter, spring and autumn *J. Meteorol. Soc. Jpn.* **80** 213–22
- Zhang X, Wan H, Zwiers F W, Hegerl G C and Min S-K 2013 Attributing intensification of precipitation extremes to human influence *Geophys. Res. Lett.* **40** 5252–7
- Zolina O, Simmer C, Gulev S K and Kollet S 2010 Changing structure of European precipitation: longer wet periods leading to more abundant rainfalls *Geophys. Res. Lett.* **37** L06704