

## LJMU Research Online

**Carden, LJ, Saini, P, Seddon, C, Evans, E and Taylor, PJ**

**Shame, social deprivation, and the quality of the voice-hearing relationship.**

<http://researchonline.ljmu.ac.uk/id/eprint/10315/>

### Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Carden, LJ, Saini, P, Seddon, C, Evans, E and Taylor, PJ (2019) Shame, social deprivation, and the quality of the voice-hearing relationship. Psychology and Psychotherapy: Theory, Research and Practice. ISSN 1476-0835**

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)

**Shame, Social Deprivation and the Quality of the Voice-Hearing Relationship**

Carden, L. C.<sup>1</sup>, Saini, P.<sup>2,3</sup>, Seddon, C.<sup>4</sup>, Evans, E.<sup>5</sup>, and Taylor, P. J.<sup>6</sup>

<sup>1</sup> Single Point of Access, Mersey Care NHS Trust, Liverpool, L9 1EP

<sup>2</sup> NIHR CLAHRC NWC, Institute of Psychology, Health & Society, University of Liverpool, Liverpool, L69 3GB

<sup>3</sup> Natural Sciences and Psychology, Liverpool John Moores University, Liverpool, L3 3AF

<sup>4</sup> Liverpool Early Intervention Service, Mersey Care NHS Trust, Liverpool, L7 9NJ

<sup>5</sup> Bootle Community Mental Health Team, Mersey Care NHS Trust, Liverpool, L22 3XR

<sup>6</sup> Division of Psychology & Mental Health, University of Manchester, M13 9PL

Word count (exc. abstract/figures/tables/references Inc. appendix): 4, 959

\*Requests for reprints should be addressed to Dr Louise Carden, Single Point of Access, Clock View Hospital, 2a Oakhouse Park, Liverpool L9 1EP (e-mail: [louise.carden@merseycare.nhs](mailto:louise.carden@merseycare.nhs))

### Abstract

**Objectives:** Many individuals hold different beliefs about the voices that they hear and have distinct relationships with them, the nature of which may determine the distress experienced. Understanding what factors contribute to these beliefs and relationships and consequently the resulting distress is important. The current research examined whether shame and social deprivation, in a sample of adult voice-hearers, were related to the relationships that individuals had with their voices or the beliefs that they held about them. **Design:** The study utilised a cross-sectional, internet-based design. **Methods:** Eighty-seven adult voice-hearers from England were recruited to the online survey. Participants completed measures regarding shame, beliefs about voices and relationships with voices and provided demographic information and postcodes that were used to refer to Index of Multiple Deprivation data (IMD). **Results:** Social deprivation and shame were not associated. Shame was positively associated with variables describing negative voice-hearing beliefs/relationships but not positive voice-hearing beliefs/relationships. Principal component analysis (PCA) on the eight voice-hearing variables yielded two components related to positive and negative voice-hearing qualities. A multiple regression conducted on the two components identified that shame was only associated with negative voice-hearing qualities. **Conclusions:** The results suggest that therapies that target shame may be helpful when working with negative voice-hearing beliefs and relationships. Future research should utilise experimental or longitudinal designs to examine the direction of the relationship.

**Keywords:** Shame; Hearing Voices; Social Deprivation; Relationship

**Practitioner Points:**

- The results contribute to the limited research evidence available regarding the relationship between shame and voice-hearing.
- The results suggest the utility of psychological therapies which focus on shame such as compassion focused therapy and that conceptualise voices interpersonally such as cognitive analytic therapy.
- No conclusions can be made regarding causation. The sample size was relatively small and results cannot be generalised to other areas of the UK.
- Future research should utilise experimental and longitudinal designs to examine the impact of shame on voice-hearing experiences and to examine other factors that may predict shame.

## Introduction

Hearing voices that others cannot hear is a common experience (Beavan, Read, & Cartwright, 2011) that is often associated with distress (Chadwick, Lees, & Birchwood, 2000). Understanding what contributes to positive and negative aspects of the voice-hearing experience is important, as such variables may determine the distress experienced. Research has identified parallels between voice-hearers' interpretations of their voices and the way they perceive themselves and others (Birchwood, Meaden, Trower, Gilbert, & Plaistow, 2000). Individuals who feel more powerless, inferior or of low social rank tend to attribute similar characteristics to their relationship with their voices (*ibid.*). Shame and social deprivation are two factors that are related to social rank (Gilbert & McGuire, 1998; Wilkinson & Pickett, 2009), one internal and subjective, and the other external and objective. This study examines the relationship between these factors and positive and negative aspects of the voice-hearing experience.

The term "hearing voices" has been adopted by user-led groups such as the Hearing Voices Network (Corstens, Longden, McCarthy-Jones, Waddingham, & Thomas, 2014) to describe auditory verbal hallucinations, "any percept like experience which a) occurs in the absence of an appropriate stimulus, (b) has the full force or impact of the corresponding actual (real) perception, and (c) is not amenable to direct and voluntary control by the experiencer" (Slade & Bentall, 1988, p. 23). Hearing voices is a common experience, with a systematic review of the literature identifying that an average of 13.2% of the general population hear voices at some point during their life course (Beavan et al., 2011). Despite some voice-hearers' reporting intimacy and companionship within their voice-hearing relationships (e.g., Nayani & David, 1996; Romme & Escher, 2000) many individuals state that their voices cause them distress (Birchwood et al., 2000; Birchwood et al., 2004; Chadwick & Birchwood, 1994). Understanding what contributes to this distress is important.

Cognitive approaches suggest that individuals' beliefs about the identity and meaning of their voices (e.g., voice as omnipotent and powerful, voice as malevolent and harmful; Chadwick & Birchwood, 1994) has a resulting impact on levels of distress (Peters, Williams, Cooke, & Kuipers, 2012). Distress arising from voice-hearing may be understood according to the individuals' relationship with their voice, specifically where relating was characterised by subordination to a dominant other (Birchwood & Chadwick, 1997). Birchwood and colleagues (2000) examined the voice/voice-hearer relationship drawing on social rank theory (Gilbert & Allan, 1998) and noted that differences in power and rank identified in the voice/voice-hearer relationship were mirrored in the differences observed between voice-hearers and significant others in their social world. This suggests that if a person feels inferior to others in their external world, this dynamic is likely to emerge in terms of how they relate to their voices (i.e., voices as superior or judging, the individual as inferior). Individuals often attribute their voice to others and personify their voices with certain individualities (Chadwick, Birchwood, & Trower, 1996; Leudar, Thomas, McNally, & Glinski, 1997). This has led to the incorporation of interpersonal schemata in theories of voice-hearing (Birchwood et al., 2000; Birchwood et al., 2004).

Relational conceptualisations of the voice-hearing experience have developed beyond dimensions of power and rank, to consider the notion that individuals can form an interpersonal relationship with their voice (Benjamin, 1989), in the same way they form relationships with people in their external world. Birtchnell's theory of relating (1996; 2002) proposes that relating and interrelating occur on orthogonal intersecting axes of proximity (close and distant) and power (upperness and lowerness) and asserts that individuals can relate positively or negatively with regards to any four positions. This framework has been used within studies to examine the voice-hearing relationship. Findings have suggested that relating to voices is associated with external social relationships (Hayward, 2003) and levels of distress, with

greater levels of distress reported in those who attempt to distance themselves from voices that are perceived to be more dominant and intrusive (Sorrell, Hayward, & Meddings, 2010; Vaughan & Fowler; 2004). The Voice and You scale (VAY; Hayward, Denney, Vaughan, & Fowler, 2008) was developed based on relating theory as a measure to assess interrelating between the voice-hearer and their predominant voice.

Shame is an interpersonal emotional state that is characterised by feelings of inferiority, defectiveness and negative evaluation of the self (Feiring, Taska, & Lewis, 2002; Lewis, 1971; Tangney & Dearing, 2003), which has been defined as an emotional manifestation of low social rank – one's sense of status in relation to others (Birchwood et al., 2004; Gilbert et al., 2010). As such, shame could be expected to impact on how voices are perceived, with a mirroring between the emotion of shame and individuals' relationships with their voices. Hence those who feel more shame may also see their voices as more hostile or dominating. Voices are often perceived to be dominant and shaming and or to have access to shaming information about the individuals (Birchwood et al., 2004; Byrne, Trower, Birchwood, Meaden, & Nelson, 2003; Chadwick & Birchwood, 1994; Nayani & David, 1996). Relationships have been identified between the power of the voice and behavioural tendencies associated with shame, notably the desire to escape and hide (Gilbert et al., 2001). Despite growing interest regarding the psychological, evolutionary, and phenomenological aspects of shame and voice-hearing (McCarthy-Jones, 2017; Woods, 2017), little research has investigated shame in relation to specific psychotic symptoms, and none has explored how shame effects the relationship individuals hold with their voices. If relationships with voices mirror external interpersonal patterns then we would predict that shame, which is characterised by perceived inferiority, would be mirrored in a voice-hearing relationship also characterised by inferiority relative to the voice.

Social deprivation can be described as lacking the material and social resources that are

customary in the societies to which individuals belong (Townsend, 1993). It can be conceptualised as an external marker of social rank, much in the way that shame may be an internal marker of positioning or status. Within the literature relationships between social deprivation and shame have been described (Peacock, Bissell, & Owen, 2014; Wilkinson & Pickett, 2009). Psychosis is associated with greater levels of social deprivation (Kirkbride, Jones, Ullrich, & Coid, 2014), though the direction of this relationship remains unclear, and could be bi-directional. Social deprivation may be a product of downward social drift (Goldberg & Morrison, 1963) whereby psychosis may lead to greater social deprivation, but deprivation also increases risk of psychosis (Harrison, Gunnell, Glazebrook, Page, & Kwiecinski, 2001; Read, Bentall, & Fosse, 2009; Wicks, Hjern, Gunnell, Lewis, & Dalman, 2005). Once again we might predict that being in a low status and deprived position in the outside world would again be mirrored by a more negative relationship with voices. It is possible that external deprivation affects the voice-hearing relationship through its impact on emotional states like shame.

For many individuals, the experience of hearing voices can be a major source of distress, notably in terms of the content, meaning ascribed, and the relationship between the hearer and the voice. The current research aims to explore the possible psychosocial determinants of this relationship, focusing on shame and social deprivation as putative correlates of voice relationship. This study may provide an understanding of why voice-hearers feel a certain way in relation to their voices and inform social policy and intervention.

**Hypotheses:**

1. Shame will be positively associated with negative voice-hearing qualities (malevolence, omnipotence, dominance, intrusiveness, and hearer distance).
2. Shame will be negatively associated with positive voice-hearing qualities (benevolence and positive relating).



3. Social deprivation will be positively associated with shame.
4. Shame will mediate the association between social deprivation and negative voice-hearing qualities.

## **Method**

### **Participants**

One-hundred and seventy-one participants were recruited to the online study. Eighty-eight participants completed part of the questionnaire and 73 the whole questionnaire (see Figure 1). Recruitment was through multiple sources to maximise the identification of individuals who hear voices. Posters were placed in Community Mental Health Teams and Early Intervention services in three NHS Trusts (Mersey Care, Cheshire and Wirral and North West Boroughs) and the researcher attended trust locations to disseminate information to professionals. The study was advertised on relevant websites (e.g., Hearing Voices Network, ISPS), social media (Twitter, Facebook) and the researcher attended NHS and Hearing Voices Network hearing voices groups to advertise the research.

Participants must have heard at least one voice, irrespective of any mental health diagnosis. Individuals hearing a single voice or multiple different voices were both eligible for the study. The voice(s) must have occurred for at least one month and must have been a current experience at the time of participation. The voice(s) could produce a word or words, but also other utterances that could be attributed to a being (e.g., laughing, crying). Other auditory hallucinations that could not be related to an individual (e.g., machine noises) were not classed as a voice. The voice(s) may have been perceived as human or non-human (e.g., god) or viewed as a product of psychosis or illness. Only participants aged 18 and above, who lived in England and who could understand or speak English were eligible for recruitment into the study. Adherence to inclusion criteria was determined by participant self-report.

[INSERT FIGURE 1 HERE]

## Measures

### **Social deprivation data.**

Participants provided their full current post code. This information was entered in to GeoConvert (Office for National Statistics, 2015) to refer to the English Indices of Multiple Deprivation data 2015 (IMD; Department for Communities and Local Government, 2015). GeoConvert cross-references the participant postcode with an existing database of deprivation data and the corresponding IMD score, rank and decile is obtained. A total of 38 indicators over seven domains (income, education, health, employment, living environment, access to services and crime) are used to obtain the IMD score, which was used in this case. The greater the IMD score, the more deprived that area is. Participants also answered three scaling questions designed to measure subjective perceptions of social deprivation (in comparison to others in the UK, others in their community, and how deprived others may perceive them to be). Participants were asked to answer the questions according to a sliding scale (0 = lowest standing, to 100 = highest standing). These questions were included as the researchers were interested not only in the level of deprivation in each area but also the perceptions of the participants taking part. Whilst these items have not previously been validated they have face validity, and are similar to other measures of subjective social status (Cundiff, Smith, Uchino, & Berg, 2013).

### **The Experience of Shame Scale (ESS; Andrews, Qian, & Valentine, 2002).**

The ESS is a 25-item self-report questionnaire that measures experiences of shame

(over the past year) in relation to three aspects of shame: characterological shame, behavioural shame, and bodily shame. In the current study, characterological shame was utilised as the overall measure of shame. This type of shame was considered to be most relevant in relation to the focus of the current research. Participants are required to answer items in relation to how they have felt in the past year. Each response is rated on a 4-point scale (1 = not at all, to 4 = very much). The 3-factor structure is supported in the literature in addition to the construct validity and discriminant validity of the ESS total scale and its component subscales (Andrews et al., 2002). In the current study the ESS characterological subscale demonstrated good internal reliability (Cronbach's  $\alpha = .95$ ; McDonald's  $\omega = .95$ ).

**Beliefs About Voices Questionnaire-Revised (BAVQ-R; Chadwick, Lees, & Birchwood, 2000).**

This measure contains 35-items relating to an individual's beliefs about their voices, and the behavioural and emotional responses that they have to them. There are five subscales; three subscales relating to beliefs: omnipotence, malevolence and benevolence; and two subscales relating to an individual's behavioural and emotional responses: resistance or engaging. Only the omnipotence, malevolence and benevolence subscales were used in the current study. Responses are indicated according to a 4-point Likert scale (0 = disagree to 3 = strongly agree). The authors report construct validity with strong negative correlations identified between most subscales. In the current study Cronbach's  $\alpha$  range = .81-.89 and McDonald's  $\omega$  range = .83-.89.

**Voice and You scale (VAY; Hayward, Denney, Vaughan, & Fowler, 2008).**

The VAY is a 29-item measure of the relationship between a voice-hearer and their predominant voice that was developed from the theoretical underpinnings of Birtchnell's (1996, 2002) relating theory. There are four subscales within the VAY, two which contain items regarding the hearers' relationship with their voice (distance and dependence), and two which contain

items relating to the hearer's perception of the voice's relationship with them (dominance and intrusiveness). Responses are indicated according to a 4-point scale (0 = nearly always true, to 3 = rarely true). The authors report good internal reliability (Cronbach's  $\alpha$  range = .92-.77), test-retest reliability ( $r$  = .91-.72), and concurrent validity with other measures of voice-hearing ( $r$  = .87-.48). In the current study the VAY demonstrated good internal reliability (Cronbach's  $\alpha$  range = .84-.94; McDonald's  $\omega$  range = .83-.94).

### **Positively-framed relational items to accompany the Voice and You scale.**

Eight items to capture positive relating to voices (e.g., voices as comforting/entertaining) were developed to be included in the study. With reference to service user advice and to the relevant literature this was noted to be important, yet underrepresented in the measures used. The items were developed with the assistance of two individuals (females aged 25 and 60) with lived experience of hearing voices and upon consultation of existing literature. Responses were indicated according to a 4-point Likert scale (0 = disagree, to 3 = strongly agree). In the present study, the scale demonstrated good internal reliability (Cronbach's  $\alpha$  = .92; McDonald's  $\omega$  = .93).

### **Procedure**

Ethical approval was obtained from the NHS Research Ethics Committee. An online survey was created utilising the Qualtrics survey platform (Qualtrics, 2017). Research suggests that online recruitment methods are superior to offline methods in terms of efficiency and cost (Christensen et al. 2017) and result in larger voice-hearing sample sizes (Berry, Band, Corcoran, Barrowclough, & Wearden, 2007a; Lawrence, Jones, & Cooper, 2010). Those choosing to take part in the survey were asked to read the study information and provide informed consent prior. Participants were then asked to provide demographic information (including the postal code of their current address) before completing the battery of measures. At the end of the study participants were given the option of entering a prize draw and receiving

a summary of the results.

### **Power Calculation and Data Analysis**

According to Fritz & Mackinnon (2007) the sample size required to detect a medium indirect (mediated) effect with 80% power using the bias-corrected bootstrap method was  $n = 71$ . A power calculation was conducted using G\* Power 3.1 (Faul, Erdfelder, Buchner, & Lang, 2009) to compute the achieved power for a sample size of 71 participants for a linear multiple regression with four predictors being tested with a medium effect size based on associations between shame and psychosis identified in the literature. The analysis suggested that power obtained for this sample size would be .99.

Data analysis was conducted using SPSS v24 (IBM, 2016). Study data was prepared by coding the data for the analysis, generating total subscale scores for each measure and conducting mean imputation on data that had less than 20% of data missing from each scale. Non-parametric Spearman's correlational analyses were performed to explore relationships between the variables, as variables were non-normally distributed. To adjust for multiple testing Bonferroni correction was applied. A principal component analysis (PCA) was performed to test the validity of summing the voice-hearing variables and creating total summary scores to capture the shared contribution of the voice-hearing variables. Oblique rotation (promax) was used as components were expected to be correlated. A multiple linear regression was conducted to test the relationship between shame and voice hearing variables. Confidence intervals were generated via bias-corrected bootstrapping with 5000 resamples as tests of assumptions identified non-normally distributed residuals. Cronbach's alpha was supplemented with McDonald's Omega as a measure of internal reliability, due to the problems with the rarely met assumptions of the former index (e.g., equivalent factor loadings; Cho & Kim, 2015). Omega was calculated in Mplus 7.4 (Muthén & Muthén, 1998-2011).

### **Results**

### **Participant Characteristics**

Eighty-seven adult participants were recruited to the study, four participants were excluded due to questionable responses (three failed catch questions designed to identify false responding, and one was from Canada, not the UK). Nine participants left prior to completing the study, four of whom did not complete any of the questionnaires and five participants that did not continue past the first questionnaire. This left a sample of 74 participants, whose age ranged from 18 to 65 years old ( $N = 73$ ;  $M = 37.8$ ;  $SD = 12.4$ ), there were 50 females, 21 males, and 3 individuals that characterised themselves as 'other'. Table 1 provides demographic information. In the remaining sample there was up to 21.6% missing data per variable, with the largest amount of missing data for the subjective social deprivation items (missing for  $n = 9 - 16$ ), and up to 9.5% missing data for the remaining items. The IMD score in the sample ranged from 1 (in the least deprived 10%) to 77 (in the most deprived 10%). Table 1 provides the distribution of deprivation across the sample grouped into 20% brackets. As can be seen there was a broad range within the sample from those living in the most to the least deprived areas.

**[INSERT TABLE 1 HERE]**

### **Principal Components Analysis**

When examining the relationships between social deprivation, shame and the voice-hearing variables, a high degree of intercorrelation was identified amongst the voice-hearing variables. Therefore, it was useful to ascertain if these numerous lower-order subscales could be combined into a smaller number of higher-order variables, capturing key dimensions in voice-hearing quality. To do this a principle component analysis (PCA) with oblique rotation (promax) was conducted on the eight variables from the VAY, the BAVQ-R, and the positively-framed relational items to accompany the Voice and You scale. The Kaiser–Meyer–

Olkin statistic verified the adequacy of the sample for analysis,  $KMO = .79$  ('good' according to Field, 2009). Bartlett's test of sphericity  $\chi^2(28) = 473.62, p < .01$ , indicated that correlations between variables were large enough for PCA. Two components had eigenvalues over 1 and in combination explained 82.3% of the variance. The scree plot also demonstrated inflexions that would justify retaining two components. Table 2 demonstrates the pattern matrix factor loadings after rotation (converging 3 iterations). All standardised component loadings were high ( $> .4$ ; Costello & Osborne, 2005).

The variables that clustered on to Component 1 were voice-hearing beliefs and relationship variables that could be described as representing negative voice-hearing qualities; voice dominance, voice intrusiveness, hearer distance, malevolence, and omnipotence. The variables that clustered on to Component 2 were voice-hearing beliefs and relationship variables that represented positive voice-hearing qualities; benevolence and positive-relating items. The subscales within each of the components were then summed to obtain two new variables, one providing a total score for positive voice-hearing qualities (Cronbach's  $\alpha = .86$ ; McDonald's  $\omega = .88$ ) and the other providing a total score for negative voice-hearing qualities (Cronbach's  $\alpha = .88$ ; McDonald's  $\omega = .90$ ). Hearer dependence was excluded from the summing of the two new scales due to problematic cross-loading across both Components 1 and 2.

**[INSERT TABLE 2 HERE]**

### **Correlational Analyses**

Spearman's correlations were conducted on the eight voice-hearing variables, positive voice-hearing qualities, negative voice-hearing qualities, social deprivation, and shame (Table 3). Consistent with hypothesis one, significant positive associations were identified between shame and several negative voice-hearing variables (dominance, intrusiveness, hearer distance, omnipotence, and malevolence). Results did not support hypothesis two; there was no

association between shame and positive voice-hearing variables (benevolence and positive items). Shame was not associated with social deprivation (Hypothesis 3) and therefore, the hypothesized indirect effect of social deprivation on the quality of the voice-hearing relationship via shame was not supported (Hypothesis 4). However, significant associations were identified between shame and all subjective deprivation items.

**[INSERT TABLE 3 HERE]**

### **Regression Analysis**

Multiple regression analyses were conducted with either positive or negative voice-hearing qualities as the outcome, and number of voices (ordinal variable rated from 1 = “1 voice” to 4 = “10 or more”), length of time hearing voices (rated from 1 = “1-6 months” to 9 = “41 years and above”), and characterological shame entered as predictors (Table 4). This analysis allowed us to examine the independent association that positive voice-hearing qualities and negative voice-hearing qualities had with shame, accounting for their overlapping variance and adjusting for other potential confounders. Assumptions were tested utilising histograms, P-P plots, and scatterplots. No outliers were identified according to a Cook’s distance value of  $< 1$ , a standardized  $DF\beta < 1$ . Residuals were not normally distributed and therefore, bias-corrected and accelerated confidence intervals were generated via bootstrapping with 5000 resamples, to allow inferential testing. The regression model explained 19% of the variance in negative voice-hearing qualities ( $R^2 = .19$ ,  $F(3, 67) = 5.21$ ,  $p < .01$ ) and 18% of the variance in positive voice-hearing qualities ( $R^2 = .18$ ,  $F(3, 67) = 0.74$ ,  $p = .53$ ). Shame was only associated with negative voice-hearing qualities, predicting 18% of the variance. This analysis was repeated adjusting for age and sex (female = 1; other = 0), see Table 4, but produced equivalent findings.

**[INSERT TABLE 4 HERE]**



### **Discussion**

The primary aim of this study was to examine whether shame and social deprivation were associated with the quality of the voice-hearing relationship and if so, whether a mediational model could explain this association. Correlational hypotheses were partially supported. No relationship was identified between shame and social deprivation, yet associations were identified between shame, subjective ratings of deprivation and several of the voice-hearing variables. Additional analyses identified two higher order variables relating to positive and negative voice-hearing qualities. Shame was only associated with negative voice-hearing qualities.

In the current study, objective social deprivation was not identified as an important factor in relation to the beliefs that individuals had regarding their voices or the relationships that they had with them. This is consistent with previous research using IMD data that has identified significant associations between deprivation and paranoia but not auditory hallucinations (Wickham, Taylor, Shevlin, & Bentall, 2014). Objective social deprivation was not related to shame, yet subjective deprivation was significantly associated with both shame and negative voice hearing qualities. Subjective deprivation also had no relationship with objective social deprivation. The relationship between subjective and objective social status is complex and appears to vary with the population studied (Shaked, Williams, Evans & Zonderman, 2016). In the current study, subjective perceptions of social deprivation were moderately associated with shame. This suggests that personal evaluations of material and social resources are more relevant in relation to shame and voice-hearing than the material and social resources available in the area that individuals live within. It may also be that the same cognitive-evaluative process that underlies feelings of shame influenced judgements of subjective deprivation, since both shame and subjective social status relate to common judgements about ones status or positions in the world. Importantly, the subjective social

deprivation scales have not been previously validated, and further evidence of their psychometric quality is needed.

Consistent with hypotheses, several of the negative voice-hearing variables were positively associated with shame. However, there was a high degree of inter-correlation between the belief and relational variables suggesting that they contained a large proportion of shared variance. Other research has also identified large inter-correlations between belief and relational voice-hearing subscales (Sorrell, Hayward, & Meddings, 2010), suggesting that these variables may load onto a set of common dimensions or constructs. To address this question, future studies could use factor analysis, in larger samples, to examine the structure of these constructs.

The current findings support the notion of a mirroring between voice-hearers' experiences of shame and the quality of the voice-hearing relationship. Specifically, that where voice-hearers experience feelings of inferiority and defectiveness in relation to the self, this is mirrored in the voice being experienced as powerful, dominant, harming, and intrusive. This is consistent with research that describes an emotional mirroring of the relationship between the voice-hearer/voice and significant others in their external world (Birchwood et al., 2004). As expected, relationships between shame and voice-hearing were only identified in the context of negative voice-hearing qualities, suggesting that positive voice-hearing qualities are not impacted by shame and can thrive despite the presence of this negative emotion. This supports the notion that those who experience shame and negative voice-hearing qualities may also experience positive voice-hearing qualities. This fits with qualitative accounts of voice-hearers who describe coexisting positive and negative voice-hearing experiences; for example, voices that are perceived to provide companionship, despite being distressing (Mawson, Berry, Murray & Hayward, 2011; Romme & Escher, 2000).

The direction of the relationship between shame and negative voice-hearing qualities remains unclear. It could be that experiences of shame across the life course inform the quality of the voice-hearing relationship. This is consistent with previous research that identifies associations between traumatic and shaming life adversities and voice-hearing (Bentall, Wickham, Shevlin, & Varese, 2012; Longden, Madill, & Waterman, 2012a). However, it could be that the quality of the voice-hearing relationship contributes to and reinforces voice-hearers' experiences of shame and inferiority, in keeping with qualitative accounts (Mawson et al., 2011). In the current study shame was treated as the predictor variable within regression analyses. However, as the study was cross-sectional the direction of effect cannot be inferred and future longitudinal work is needed.

The present study contributes to existing literature that identifies the importance of shame in relation to voice-hearing. The findings suggest that interventions that target shame such as compassion-focused therapy (Gilbert, 2009) or that address negative voice-hearing beliefs such as cognitive behavioural therapy for psychosis (CBTp) may be helpful. Interventions such as acceptance and commitment therapy that help individuals to non-judgmentally acknowledge distressing experiences whilst pursuing valued goals, and mindfulness, which focuses upon changing the nature of the relationship that individuals have with their voices, may also play a role in relation to reducing the emotional dysfunction associated with psychosis (Aust & Bradshaw, 2017; Gumley et al., 2017; White et al., 2011; White et al., 2015). Results suggest the use of therapies that address the interpersonal relationship between the voice-hearer and the voice such as relating therapy (Hayward, Overton, Dorey, & Denney, 2009) or cognitive analytic therapy (CAT; Ryle, 1995), which has several features that suggest its suitability for working with experiences of psychosis (Taylor, Perry, Hutton, Seddon, & Tan, 2014). The findings imply that HVN support groups that facilitate voice-hearers to develop a sense of meaning regarding their experiences in a safe and

supportive environment, where positive and trusting external social relationships can be developed may also be effective (Dillion & Hornstein, 2013; Oakland & Berry, 2015; Payne, Allen, & Lavender, 2017). Indeed, recent research has begun to examine how CBTp and HVN approaches may complement one another (Kay, Kendall, & Dark, 2017). Furthermore, results suggest the importance of clinicians providing a therapeutic space for individuals to discover what their voice/s represent using frameworks such as the Maastricht Interview (Romme & Escher, 2000) and that they support individuals to identify and change the nature of the relationship that they have with their voice/s.

Some limitations were identified in relation to the current study. As previously mentioned, the study design was cross-sectional thus making inference about direction of effect impossible. Despite achieving the minimum sample size recommended by the power analysis, a relatively small sample size was utilised. This prevented more advanced statistical techniques such as structural equation modelling (SEM) that may have allowed for the creation of latent variables and simultaneous consideration of multiple outcomes. The study was conducted online limiting the sample to those who have internet access. This may have resulted in selection bias; however, it is worthy to note that advertisements were placed in some locations with online facilities for those without personal access.

The positively-framed relational voice-hearing questions developed for the purposes of the study were not previously piloted and the psychometric properties were unknown. There were a larger proportion of females in the sample, which may again affect generalisability. Furthermore, there were differences in the time period that the questionnaires utilised examined. However, this is a cross-sectional study and the ESS, VAY and BAVQ-R all ask participants to consider how they felt at the time of the study, with the ESS asking participants to consider the past year in addition to this. The IMD data was published at the time of the study in 2015 with census data from 2012. To examine whether the IMD data would be accurate

for use participants were asked to indicate how long they had been living at their address. Only a small number of participants had lived at their address for less than the time period that had elapsed.

Future research should further delineate the role of shame in relation to hearing voices using experimental or longitudinal research designs. Specifically, this should involve larger sample sizes, should address specific negative relational and belief qualities, and should utilise more advanced statistical techniques that are able to better model the shared variance between specific voice-hearing variables. Furthermore, research should aim to identify other proximal relational and environmental factors that are related to current experiences of shame, and contribute to ongoing negative voice-hearing experiences.

### References

- Andrews, B., Qian, M., & Valentine, J. D. (2002). Predicting depressive symptoms with a new measure of shame: The Experience of Shame Scale. *British Journal of Clinical Psychology, 41*, 29–42. doi: 10.1348/014466502163778
- Aust, J., & Bradshaw, T. (2017). Mindfulness interventions for psychosis: A systematic review of the literature. *Journal of Psychiatric and Mental Health Nursing, 24*, 69–83. doi: 10.1111/jpm.12357
- Beavan, V., Read, J., & Cartwright, C. (2011). The prevalence of voice-hearers in the general population: A literature review. *Journal of Mental Health, 20*(3), 281-292. doi: 10.3109/09638237.2011.562262
- Benjamin, L. S. (1989). Is chronicity the function of the relationship between the person and the auditory hallucination? *Schizophrenia Bulletin, 15*(2), 291-309. doi: 10.1093/schbul/15.2.291
- Bentall, R.P., Wickham, S., Shevlin, M., & Varese, F. (2012). Do specific early-life adversities lead to specific symptoms of psychosis? A study from the 2007 The Adult Psychiatric Morbidity Survey. *Schizophrenia Bulletin, 38*(4), 734-740. doi:10.1093/schbul/sbs049
- Berry, K., Band, R., Corcoran, R., Barrowclough, C., & Wearden, A. (2007a). Attachment styles, earlier interpersonal relationships and schizotypy in a non-clinical sample. *Psychology and Psychotherapy: Theory, Research and Practice 80*(4), 563-576. doi: 10.1348/147608307X188368
- Birchwood, M., & Chadwick, P. (1997). The omnipotence of voices: Testing the validity of a cognitive model. *Psychological Medicine, 27*, 1345-1353. doi: 10.1017/S0033291797005552
- Birchwood, M., Gilbert, P., Gilbert, J., Trower, P., Meaden, A., Hay, J., Murray, E., & Miles,

- J. N. V. (2004). Interpersonal and role related schema influence the relationship with the dominant 'voice' in schizophrenia: A comparison of three models. *Psychological Medicine*, 34, 1571–1580. doi: 10.1017/S0033291704002636
- Birchwood, M., Meaden, A., Trower, P., Gilbert, P., & Plaistow, J. (2000). The power and omnipotence of voices: Subordination and entrapment by voices and significant others. *Psychological Medicine*, 30, 337-344. doi: 10.1017/S0033291799001828
- Birtchnell, J. (1996). *How humans relate: A new interpersonal theory*. Hove: Psychology Press.
- Birtchnell, J. (2002). *Relating in psychotherapy: The application of a new theory*. Hove: Brunner-Routledge.
- Byrne, S., Trower, P., Birchwood, M., Meaden, A. & Nelson, A. (2003). Command hallucinations: Cognitive theory and therapy. *Journal of Cognitive Psychotherapy, An International Quarterly*, 17, 67–84. doi: 10.1891/jcop.17.1.67.58271
- Chadwick, P., & Birchwood, M. (1994). The omnipotence of voices: A cognitive approach to auditory hallucinations. *British Journal of Psychiatry*, 164, 190-201. doi: 10.1192/bjp.164.2.190
- Chadwick, P., Birchwood, M., & Trower, P. (1996). *Cognitive therapy for delusions, voices and paranoia*. Chichester: John Wiley and Sons.
- Chadwick, P., Lees, S., & Birchwood, M. (2000). The revised beliefs about voices questionnaire (BAVQ-R). *British Journal of Psychiatry*, 177, 229-232. doi: 10.1192/bjp.177.3.229
- Cho, E., & Kim, S. (2015). Cronbach's coefficient alpha: Well know but poorly understood. *Organizational Research Methods*, 18, 207-230. doi: 10.1177/1094428114555994.
- Corstens, D., Longden, E., McCarthy-Jones, S., Waddingham, R., & Thomas, N. (2014). Emerging perspectives from the Hearing Voices Movement: Implications for

- research and practice. *Schizophrenia Bulletin*, 40(4), 285–294. doi: 10.1093/schbul/sbu007
- Costello, A. B., & Osborne, J. W. (2005). Best practices in exploratory factor analysis: Four recommendations for getting the most from your analysis. *Practical Assessment, Research & Evaluation*, 10(7), 1-9.
- Creed, P. A., & Muller, J. (2006) Psychological distress in the labour market: Shame or deprivation? *Australian Journal of Psychology*, 58(1), 31-39. doi: 10.1080/00049530500125116
- Christensen, T., Riis, A. H., Hatch, E. E., Wise, L. A., & Marie, G. (2017). Costs and efficiency of online and offline recruitment methods: A web-based cohort study corresponding author. *Journal of Medical Internet research*, 19(3), 1-12. doi: 10.2196/jmir.6716
- Cundiff, J. M., Smith, T. W., Uchino, B. N., & Berg, C. A. (2013). Subjective social status: construct validity and associations with psychosocial vulnerability and self-rated health. *International Journal of Behavioural Medicine*, 20, 148–158. doi: 10.1007/s12529-011-9206-1
- Dillon, J., & Hornstein, G. A. (2017). Hearing voices peer support groups: A powerful alternative for people in distress. *Psychosis*, 5(3), 286-295. doi: 10.1080/17522439.2013.843020
- Department for Communities and Local Government (2015). *The English indices of multiple deprivation 2015*. Retrieved from: <https://www.gov.uk/government/statistics/english-indices-of-deprivation-2015>
- Faul, F., Erdfelder, E., Buchner, A., & Lang, A. G. (2009). Statistical power analyses using G\*Power 3.1: Tests for correlation and regression analyses. *Behavior Research Methods*, 41, 1149-1160. doi: 10.3738/BRM.41.4.1149



- Feiring, C., Taska, L., & Lewis, M. (2002). Adjustment following sexual abuse discovery: The role of shame and attributional style. *Developmental Psychology*, 38, 79-92. doi: 10.1037//0012-1649.38.1.79
- Field, A. (2009). *Discovering statistics using SPSS* (3rd Ed.) London: SAGE.
- Fritz, M. S., & MacKinnon, D. P. (2007). Required sample size to detect the mediated effect. *Psychological Science*, 18(3), 233-239. doi: 10.1111/j.1467-9280.2007.01882.x
- Gilbert, P. (2009). Introducing compassion focused therapy. *Advances in Psychiatric Treatment*, 15, 199-208. doi: 10.1192/apt.bp.107.005264
- Gilbert, P. & Allan, S. (1998). The role of defect and entrapment (arrested flight) in depression: An exploration of an evolutionary view. *Psychological Medicine* 28, 585-598. doi: 10.1017/S0033291798006710
- Gilbert, P., Birchwood, M., Gilbert, J., Trower, P., Hay, J., Murray, E., Meaden, A., Olsen, K. & Miles, J. N. V. (2001). An exploration of evolved mental mechanisms for dominant and subordinate behaviour in relation to auditory hallucinations in schizophrenia and critical thoughts in depression. *Psychological Medicine*, 31, 1117–1127. doi: 10.1017/S0033291701004093
- Gilbert, P., McEwan, K., Irons, C., Bhundia, R., Christie, R., Broomhead, C., & Rockliff, H. (2010). Self-harm in a mixed clinical population: The roles of self-criticism, shame, and social rank. *British Journal of Clinical Psychology*, 49, 563–576. doi: 10.1348/014466509X479771
- Gilbert, P., & McGuire, M. T. (1998). Shame, status, and social roles: Psychobiology and evolution. In P. Gilbert & B. Andrews (Eds.), *Shame: Interpersonal behaviour, psychopathology, and culture* (pp. 99– 125). New York, NY: Oxford University Press.
- Goldberg, E. M., & Morrison, S. L. (1963). Schizophrenia and social class. *British Journal of*

- Psychiatry*, 109, 785–802. doi: 10.1192/bjp.109.463.785
- Gumley, A. I., White, R. G., Briggs, A., Ford, I., Barry, S., Stewart, C., ... McLeod, H. (2017). A parallel group randomised open blinded evaluation of acceptance and commitment therapy for depression after psychosis: Pilot trial outcomes (ADAPT). *Schizophrenia Research*, 183, 143–150. doi: 10.1016/j.schres.2016.11.026
- Harrison, G., Gunnell, D., Glazebrook, C., Page, K., & Kwiecinski, R. (2001). Association between schizophrenia and social inequality at birth: Case-control study. *British Journal of Psychiatry*, 179, 346–350. doi: 10.1192/bjp.179.4.346
- Hayward, M. (2003). Interpersonal relating and voice hearing: To what extent does relating to the voice reflect social relating? *Psychology and Psychotherapy: Theory Research and Practice*, 76, 369–383. doi: 10.1348/147608303770584737
- Hayward, M., Denney, J., Vaughan, S., & Fowler, D. (2008). The Voice and You: Development and psychometric evaluation of a measure of relationships with voices. *Clinical Psychology and Psychotherapy*, 15, 45–52. doi: 10.1002/cpp.561
- Hayward, M., Overton, J., Dorey, T., & Denney, J. (2009). Relating therapy for people who hear voices: A case series. *Clinical Psychology and Psychotherapy*, 16, 216–227. doi: 10.1002/cpp.615
- IBM (2016). *IBM SPSS Statistics for Windows, Version 24.0*. Armonk, NY: IBM Corp.
- Kay, G., Kendall, E., & Dark, F. (2017). Are Hearing Voices Networks compatible with cognitive behavioural therapy for psychosis? *Australian Social Work*, 0(0), 1–12. doi: 10.1080/0312407X.2016.1262883
- Kirkbride, J. B., Jones, P. B., Ullrich, S., & Coid, J. (2014). Social deprivation, inequality, and the neighborhood-level incidence of psychotic syndromes in East London. *Schizophrenia Bulletin* 40, 169–180. doi: 10.1093/schbul/sbs151
- Lawrence, C., Jones, J., & Cooper, M. (2010). Hearing voices in a non-psychiatric

- population. *Behavioural and Cognitive Psychotherapy*, 38(3), 363-373. doi: 10.1017/S1352465810000172
- Leudar, I., Thomas, P., McNally, D., & Glinski, A. (1997). What voices do with words: Pragmatics of verbal hallucinations. *Psychological Medicine*, 27, 885–898. doi: 10.1017/S0033291797005138
- Lewis, H. B. (1971). *Shame and guilt in neurosis*. New York: International Universities Press.
- Longden, E., Madill, A., & Waterman, M.G. (2012a). Dissociation, trauma, and the role of lived experience: Toward a new conceptualization of voice hearing. *Psychological Bulletin*, 138, 28–76. doi: 10.1037/a0025995
- Mawson, A., Berry, K., Murray, C., & Hayward, M. (2011). Voice hearing within the context of the voice hearers' social worlds: An interpretative phenomenological analysis. *Psychology and Psychotherapy: Theory, Research and Practice*, 84, 256–272. doi:10.1348/147608310X524883
- McCarthy-Jones, S. (2017). Is shame hallucinogenic? *Frontiers in Psychology*, 8, 1310. doi: 10.3389/fpsyg.2017.01310
- Muthén, L. K., & Muthén, B. O. (1998-2011). *Mplus User's Guide*. Sixth Edition. Los Angeles, CA: Muthén & Muthén.
- Nayani, T. H., & David, A. S. (1996). The auditory hallucination: A phenomenological survey. *Psychological Medicine*, 26, 177–189. doi: 10.1017/S003329170003381X
- Oakland, L., & Berry, K. (2017). “Lifting the veil”: A qualitative analysis of experiences in Hearing Voices Network groups. *Psychosis*, 7(2), 119-129. doi: 10.1080/17522439.2014.937451
- Office for National Statistics, Postcode Directories (2015): *GeoConvert*. UK Data Service. doi: 10.5257/census/geoconvert-1

- Payne, T., Allen, J., & Lavender, T. (2017) Hearing Voices Network groups: Experience of eight voice hearers and the connection to group processes and recovery. *Psychosis*, 1–11. doi: 10.1080/17522439.2017.1300183
- Peacock, M., Bissell, P., & Owen, J. (2014). Shaming encounters: Reflections on contemporary understandings of social inequality and health. *Sociology*, 48(2), 387–402. doi: 10.1177/0038038513490353
- Peters, E. R., Williams, S. L., Cooke, M. A. & Kuipers, E. (2012). It's not what you hear, it's the way you think about it: Appraisals as determinants of affect and behaviour in voice hearers. *Psychological Medicine*. 42, 1507–1514. doi: 10.1017/S0033291711002650
- Qualtrics [Computer software]. (2017). Retrieved from <https://www.qualtrics.com/>
- Read, J., Bentall, R. P., & Fosse, R. (2009). Time to abandon the bio-bio-bio model of psychosis: Exploring the epigenetic and psychological mechanisms by which adverse life events lead to psychotic symptoms. *Epidemiologia e Psichiatria Sociale*, 18(4), 299-310. doi: 10.1017/S1121189X00000257
- Romme, M. A., & Escher, S. (2000). *Making sense of voices*. London: Mind Publications.
- Ryle, A. (Ed.). (1995). *Cognitive analytic therapy: Developments in theory and practice*. Chichester: Wiley.
- Shaked, D., Williams, M., Evans, M. K., & Zonderman, A. B. (2016). Indicators of subjective social status: Differential associations across race and sex. *SSM - Population Health*, 29(2), 700-707. doi: 10.1016/j.ssmph.2016.09.009
- Slade, P. D., & Bentall, R. P. (1988). *Sensory deception: A scientific analysis of hallucination*. London: Johns Hopkins University Press.
- Sorrell, E., Hayward, M., & Meddings, S. (2010). Interpersonal processes and hearing voices: A study of the association between relating to voices and distress in clinical and non-

- clinical hearers. *Behavioural and Cognitive Psychotherapy*, 38, 127–140. doi: 10.1017/S1352465809990506
- Studenmund, A. H., & Cassidy, H. J. (1987). *Using econometrics: A practical guide*. Boston: Little Brown.
- Tangney, J. P., & Dearing, R. L. (2003). *Shame and Guilt*. Guildford Press: New York.
- Taylor, P., Perry, A., Hutton, P., Seddon, C., & Tan, R. (2014). Curiosity and the CAT: Considering cognitive analytic therapy as an intervention for psychosis. *Psychosis*, 7, 276-278. doi: 10.1080/17522439.2014.956785
- Townsend, P. (1993). *The International Analysis of Poverty*. London: Harvester Wheatsheaf.
- Vaughan, S., & Fowler, D. (2004). The distress experienced by voice hearers is associated with the perceived relationship between the voice hearer and the voice. *British Journal of Clinical Psychology*, 43, 143-153. doi: 10.1348/014466504323088024
- White, R. G., Gumley, A. I., McTaggart, J., Rattrie, L., McConville, D., Cleare, S., McLeod, H. J., & Mitchell, G. (2015). Acceptance and commitment therapy for depression following psychosis: An examination of clinically significant change. *Journal of Contextual Behavioral Science*, 4, 203-209. doi: 10.1016/j.jcbs.2015.06.004
- White, R. G., Gumley, A. I., McTaggart, J., Rattrie, L., McConville, D., Cleare, S., & Mitchell, G. (2011). A feasibility study of acceptance and commitment therapy for emotional dysfunction following psychosis. *Behaviour Research and Therapy*, 49, 901-907. doi: 10.1016/j.brat.2011.09.003
- Wickham, S., Taylor, P., Shevlin, M., & Bentall, R. P. (2014). The impact of social deprivation on paranoia, hallucinations, mania and depression: The role of discrimination social support, stress and trust. *PLoS ONE* 9(8), e105140. doi:10.1371/journal.pone.0105140
- Wicks, S., Hjern, A., Gunnell, D., Lewis, G., Dalman, C. (2005). Social adversity in

childhood and the risk of developing psychosis: A national cohort study. *American Journal of Psychiatry*, 162, 1652–1657. doi: 10.1176/appi.ajp.162.9.1652

Wilkinson, R., & Pickett, K. E. (2009). *The spirit level: Why more equal societies almost always do better*. London: Penguin.

Woods, A. (2017). On shame and voice-hearing. *Medical Humanities*, 0, 1–6.  
doi:10.1136/medhum-2016-011167




Figure 1. Flow chart of participation.

Table 1

*Participant Characteristics*

Variable	<i>n</i>	%
Gender (female)	50	67.6
Employed	33	44.6
Student	11	14.9
Given mental health diagnosis	55	74.3
Taking mental health medication	50	67.6
Deprivation (IMD decile) <sup>a b</sup>		
Bottom 20%	18	24.3
20-40%	17	23.0
60-80%	12	16.0
Top 20%	10	13.5

*Note.* <sup>a</sup> Deciles of relative social deprivation derived from the Indices of Multiple Deprivation, where bottom 20% represents the most deprived. <sup>b</sup> Participant numbers for deprivation data do not equal 74 as a result of missing data.



Table 2

*Factor Loadings for Principal Component Analysis with Promax Rotation of the Eight Voice-Hearing Variables (Pattern Matrix)*

Variable	Component	
	1	2
Omnipotence	<b>.94</b>	.19
Voice intrusiveness	<b>.85</b>	.14
Malevolence	<b>.84</b>	-.26
Voice dominance	<b>.80</b>	-.26
Hearer distance	<b>.44</b>	-.70
Positive items	-.05	<b>.93</b>
Benevolence	-.10	<b>.88</b>

*Note.* Factor loadings > .40 are in boldface and indicate loadings on to each component.

Table 3

*Spearman's Non-Parametric Correlations for the Eight Voice-Hearing Variables, Social deprivation and Shame*

Variable	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1. Voice dominance	1														
2. Voice intrusiveness	.56*** <sup>a</sup>	1													
3. Hearer dependence	.07	.38*** <sup>a</sup>	1												
4. Hearer distance	.65*** <sup>a</sup>	.45*** <sup>a</sup>	-.33**	1											
5. Malevolence	.80*** <sup>a</sup>	.60*** <sup>a</sup>	.07	.59*** <sup>a</sup>	1										
6. Benevolence	-.47*** <sup>a</sup>	-.19	.49*** <sup>a</sup>	-.67*** <sup>a</sup>	-.49*** <sup>a</sup>	1									
7. Omnipotence	.65*** <sup>a</sup>	.66*** <sup>a</sup>	.47*** <sup>a</sup>	.39*** <sup>a</sup>	.75*** <sup>a</sup>	-.20*	1								
8. Positive items	-.43*** <sup>a</sup>	-.05	.63*** <sup>a</sup>	-.63*** <sup>a</sup>	-.43*** <sup>a</sup>	.76*** <sup>a</sup>	-.04	1							
9. Characterological shame	.51*** <sup>a</sup>	.31**	.33**	.24*	.43*** <sup>a</sup>	-.08	.44*** <sup>a</sup>	-.02	1						
10. IMD score	-.07	-.08	-.10	.10	.01	-.12	.07	-.07	-.04	1					
11. Sub dep (UK)	-.18	-.39*** <sup>a</sup>	-.29*	-.10	-.19	.07	-.28*	.03	-.40*** <sup>a</sup>	-.20	1				
12. Sub dep (Community)	-.19	-.50*** <sup>a</sup>	-.44*** <sup>a</sup>	-.10	-.16	.02	-.43*** <sup>a</sup>	-.04	-.35*	-.02	.70*** <sup>a</sup>	1			
13. Sub dep (Others)	-.25	-.43*** <sup>a</sup>	-.36**	-.06	-.26*	.04	-.44*** <sup>a</sup>	.03	-.42*** <sup>a</sup>	-.09	.70*** <sup>a</sup>	.84*** <sup>a</sup>	1		
14. Positive voice qualities	-.47*** <sup>a</sup>	-.11	.62*** <sup>a</sup>	-.70*** <sup>a</sup>	-.48*** <sup>a</sup>	.89*** <sup>a</sup>	-.09	.96*** <sup>a</sup>	-.01	-.07	.04	-.03	.03	1	
15. Negative voice qualities	.91*** <sup>a</sup>	.75*** <sup>a</sup>	.12	.74*** <sup>a</sup>	.92*** <sup>a</sup>	-.51*** <sup>a</sup>	.81*** <sup>a</sup>	-.42*** <sup>a</sup>	.47*** <sup>a</sup>	.03	-.26*	-.31*	-.34**	-.48*** <sup>a</sup>	1

*Note.* \*  $p < .05$ , two-tailed. \*\*  $p < .01$ , two-tailed; <sup>a</sup>  $p < .003$  (alpha adjusted by Bonferroni correction); sub dep = subjective deprivation

Table 4

*Multiple Linear Regressions with Characterological Shame as a Predictor of Voice-Hearing Variables*

Predictor variable	<u>Positive voice-hearing qualities</u>			<u>Negative voice-hearing qualities</u>		
	B	CI (95%)	$\beta$	B	CI (95%)	$\beta$
<u>Model 1</u>						
Shame**	-0.04	-0.33, 0.25	-.03	0.98	0.44, 1.60	.42
Number of voices	2.61	-0.99, 6.13	.18	-0.98	-8.35, 7.55	-.03
Length of time hearing voices	-0.11	-1.29, 1.01	-.02	1.10	-1.00, 3.07	.10
<u>Model 2</u>						
Age	-0.25	-0.52, 0.01	-.27	0.04	-0.63, 0.63	.02
Sex (female)	-1.47	-7.80, 4.44	-.06	-2.34	-12.70, 8.24	-.04

Shame**	-0.11	-0.41, .20	-.10	1.10	.35, 1.72	.43
Number of voices	1.93	-1.63, 5.38	.14	-0.75	-8.74, 8.49	-.03
Length of time hearing voices	0.70	-0.87, 2.31	.13	1.04	-1.67, 3.68	.09

*Note.* \*\*  $p < .01$ , two-tailed; CI = Bias-Corrected and accelerated Confidence Intervals estimated via bootstrapping with 5000 resamples