



LJMU Research Online

Al Groshi, A, Jasim, HA, Evans, AR, Ismail, FMD, Dempster, NM, Nahar, L and Sarker, SD

Growth inhibitory activity of biflavonoids and diterpenoids from the leaves of the Libyan Juniperus phoenicea against human cancer cells

<http://researchonline.ljmu.ac.uk/id/eprint/10627/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Al Groshi, A, Jasim, HA, Evans, AR, Ismail, FMD, Dempster, NM, Nahar, L and Sarker, SD (2019) Growth inhibitory activity of biflavonoids and diterpenoids from the leaves of the Libyan Juniperus phoenicea against human cancer cells. Phytotherapy Research. 33 (8). pp. 2075-2082. ISSN

LJMU has developed [LJMU Research Online](#) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

**Growth inhibitory activity of biflavonoids and
diterpenoids from the leaves of the Libyan *Juniperus
phoenicea* against human cancer cells**

Afaf Al Groshi, Hiba A. Jasim, Andrew R. Evans, Fyaz M. D. Ismail,
Nicola M. Dempster, Lutfun Nahar and Satyajit D. Sarker*

Centre for Natural Products Discovery, School of Pharmacy and Biomolecular Sciences,
Liverpool John Moores University, James Parsons Building, Byrom Street, Liverpool L3
3AF, United Kingdom

Supplementary data

Supplementary Table 1: ¹H NMR data for the biflavonoids, cupressoflavone (**1**), amentoflavone (**2**) and sumaflavone (**3**)

¹ H	Position Chemical Shift δ (ppm), multiplicity and J in Hz		
	1* (300 MHz)	2** (600 MHz)	3** (300 MHz)
3	6.78 (s)	6.66 (s)	6.78 (s)
5	13.22 (s) (OH)	-	13.01 (s) (OH)
6	6.45 (s)	6.22 (br s)	6.16 (d, 2.0)
7	10.33 (s) (OH)	-	13.06 (s) (OH)
8		6.42 (br s)	
2`	7.49 (d, 8.9)	8.00 (d, 2.3)	8.17 (s)
3`	6.74 (d, 8.9)	-	
4`	10.33 (s) (OH)	-	
5`	6.74 (d, 8.9)	7.96 (dd, 8.5, 2.2)	7.0 (d, 8.6)
6`	7.49 (d, 8.9)	7.16 (d, 8.6)	7.92 (d, 8.8)
3``	6.78 (s)	6.64 (s)	6.71 (s)
6``	6.45 (s)	6.40 (s)	
2'''	7.49 (d, 8.9)	7.58 (d, 8.8)	7.63 (d, 8.7)
3'''	6.74 (d, 8.9)	6.74 (d, 8.8)	6.61 (d, 8.6)
5'''	6.74 (d, 8.9)	6.74 (d, 8.8)	6.61 (d, 8.6)
6'''	7.49 (d, 8.9)	7.58 (d, 8.8)	7.63 (d, 8.7)

*= (DMSO-d₆), **= (CD₃OD)

Supplementary Table 2: ^{13}C NMR data of biflavonoids **1** and **2**

Position ^{13}C	Chemical Shift δ (ppm),		Position ^{13}C	Chemical Shift δ (ppm)	
	1* (75 MHz)	2** (150 MHz)		1* (75 MHz)	2** (150 MHz)
2	163.9	166.0	2''	163.9	166.2
3	103.0	103.5	3''	103.0	100.3
4	182.5	184.0	4''	182.5	184.4
5	161.5	163.4	5''	161.5	163.4
6	99.3	100.3	6''	99.3	95.3
7	164.1	166.2	7''	164.1	162.7
8	98.9	95.3	8''	98.9	105.5
9	155.2	159.6	9''	155.2	159.6
10	104.0	105.5	10''	104.0	105.5
1`	121.7	123.2	1'''	121.7	123.4
2`	128.4	133	2'''	128.4	129.0
3`	116.3	122.0	3'''	116.3	117.0
4`	161.3	161.5	4'''	161.3	163.4
5`	116.3	129	5'''	116.3	117.0
6`	128.4	117.0	6'''	128.4	129.0

* = (DMSO- d_6), ** = (CD_3OD)

Supplementary Table 3: ¹H NMR (600 MHz, CDCl₃) for diterpenes, 13-*epi*-cupressic acid (**4**), imbricatolic acid (**5**) and 3-hydroxy sandaracopimaric acid (**6**)

Position ¹ H	Chemical Shift δ (ppm), multiplicity and <i>J</i> in Hz		
	4	5	6
1	1.05 (br s), 1.71(m)	0.95 (m); 1.86 (m)	1.07 (m), 1.21 (m)
2	1.50 (m), 1.71 (m)	1.45 (m); 1.86 (m)	1.38 (m), 1.45 (m)
3	1.04 (dd; 4.8; 13.9) 1.97, (br s)	1.06 (dd; 4.2; 13.3) 2.14 (br s)	4.02 (dd; 4.6, 11.8)
5	1.32 (m)	1.32 (m)	1.53 (m)
6	1.83 (m), 1.89 (m)	1.85 (m), 1.87 (m)	0.92 (m), 1.25 (m)
7	1.86, (m), 2.40 (br s)	1.88 (m), 1.99 (m)	1.86 (m), 2.23 (m)
9	1.50, m	1.91 (m) 2.40 (dd, 2.8, 8.8)	
11	1.32 (m), 1.45, (m)	1.36 (m)	1.38 (m), 1.50 (m)
12	1.32 (m), 1.57, (m)	1.29 (m)	1.2 (m), 1.24 (m)
13		1.34 (m)	-
14	5.91, dd, 10.2, 17.6	1.34 (m)	5.24 (s)
15	5.21, dd, 1.3. 17.2 5.06, dd, 10.7, 0.8	3.67 (m)	5.76 (dd; 10.4, 17.7)
16	1.27, s	0.90 (d, 6.5)	4.89 (d, 10.6), 4.90 (d, 17.5)
17	4.49 (s), 4.83 (s)	4.83 (s), 4.49(s)	1.04
18	1.24 (s)	1.24 (s)	-
19	-	-	1.18 (s)
20	0.59 (s)	0.60 (s)	0.83 (s)

Supplementary Table 4: ^{13}C NMR (150 MHz, CDCl_3) for diterpenes: 13-*epi*-cupressic acid (**4**), imbricatolic acid (**5**) and 3-hydroxy sandaracopimaric acid (**6**)

Position		Chemical Shift δ (ppm)			Position		Chemical Shift δ (ppm)		
^{13}C	4	5	6	^{13}C	4	5	6		
1	39.4	39.1	38.5	11	18.1	21.5	28.1		
2	20.2	20.2	20	12	41.6	40.0	35.9		
3	38.2	36.8	76.6	13	74.1	29.4	38.7		
4	44.5	44.5	54.9	14	145.8	41.0	130.6		
5	56.7	56.7	52.0	15	112.0	61.6	150.0		
6	26.4	26.4	25.7	16	28.3	21.5	111.0		
7	39.0	38.4	36.7	17	107.1	106.6	26.7		
8	148.2	148.6	137.8	18	29.2	30.6	181.5		
9	56.7	57.0	51.6	19	183.2	183.4	11.9		
10	40.9	39.5	38.7	20	12.9	13.1	15.8		