

Title: Average acceleration and intensity gradient of primary school children and associations with indicators of health and wellbeing

Running title: Standardised physical activity metrics & associations with child health

Stuart J. Fairclough¹, Sarah Taylor¹, Alex V. Rowlands^{2,3,4}, Lynne, M. Boddy⁵ and Robert. J. Noonan¹

¹Movement Behaviours, Health, and Wellbeing Research Group, Department of Sport and Physical Activity, Edge Hill University, Ormskirk, UK

²Diabetes Research Centre, University of Leicester, Leicester General Hospital, Leicester, UK

³NIHR Leicester Biomedical Research Centre, UK

⁴Alliance for Research in Exercise, Nutrition and Activity (ARENA), Sansom Institute for Health Research, Division of Health Sciences, University of South Australia, Adelaide, Australia

⁵Physical Activity Exchange, Research Institute for Sport and Exercise Sciences, Liverpool John Moores University, Liverpool, UK

Corresponding author address

Professor Stuart Fairclough, Movement Behaviours, Health, and Wellbeing Research Group, Department of Sport and Physical Activity, Edge Hill University, St Helens Road, Ormskirk, UK; Email: stuart.fairclough@edgehill.ac.uk

Keywords: physical activity, standardised metrics, raw accelerations, GGIR, wrist, youth

Word count: 4814

Removed section from anonymous manuscript version because it potentially identifies the authors (as per email request from Camille on 7th May 2019)

Methods

This is a secondary analysis of data collected in the Active Schools: Skelmersdale PA pilot intervention study (ClinicalTrials.gov registration: NCT03283904). The methods have been described previously (8) but are outlined briefly here. Two hundred and thirty two 9-10 year old participants were recruited from 7 primary schools. The schools were situated in a low socioeconomic status (SES) town in West Lancashire, north-west England, where the prevalence of overweight/obesity is above the national average (13). Ethical approval was granted by Edge Hill University's Research Ethics Committee (reference # SPA-REC-2015-330) and informed consent and assent were provided by the participants' parents/carers, and the participants themselves, respectively. Data collection took place between September and December 2017.

Following collection of baseline measurements, schools were randomly assigned to either intervention (4 schools) or control groups (3 schools). The AS:Sk pilot intervention included eight components which were implemented over 8-weeks.

Abstract

Average acceleration (AvAcc) and intensity gradient (IG) have been proposed as standardised metrics describing physical activity (PA) volume and intensity, respectively. We examined hypothesised between-group PA differences in AvAcc and IG, and their associations with health and wellbeing indicators in children. ActiGraph GT9X wrist accelerometers were worn for 24-h·d⁻¹ over seven days by 145 children aged 9-10. Raw accelerations were averaged per 5-s epoch to represent AvAcc over 24-h. IG represented the relationship between log values for intensity and time. Moderate-to-vigorous PA (MVPA) was estimated using youth cutpoints. BMI z-scores, waist-to-height ratio (WHtR), peak oxygen uptake (VO_{2peak}), Metabolic Syndrome risk (MetS score), and wellbeing were assessed cross-sectionally, and 8-weeks later. Hypothesised between-group differences were consistently observed for IG only ($p<.001$). AvAcc was strongly correlated with MVPA ($r=0.96$), while moderate correlations were observed between IG and MVPA ($r=0.50$) and AvAcc ($r=0.54$). IG was significantly associated with health indicators, independent of AvAcc ($p<.001$). AvAcc was associated with wellbeing, independent of IG ($p<.05$). IG was significantly associated with WHtR ($p<.01$) and MetS score ($p<.05$) at 8-weeks follow-up. IG is sensitive as a gauge of PA intensity that is independent of total PA volume, and which relates to important health indicators in children.

Introduction

Until recently, comparability of data collected using different accelerometer brands was not possible because of the reliance on device-specific ‘counts’, which were based on proprietary algorithms (1). In the last decade, the move towards increased accessibility of raw acceleration signals has greatly increased the potential for cross-device comparability. However, studies using raw accelerations still tend to apply population-specific and protocol-specific thresholds or cutpoints to estimate time spent in different movement intensities (2-4). There is though, no consensus as to the most appropriate raw acceleration cutpoints to estimate time spent in different PA intensities. Application of different cutpoints can result in vastly different estimates of PA (5), which is confusing for interpretation and translation of data for surveillance and intervention evaluation. Most of the studies that have used raw accelerations to describe PA outcomes (6-9) have employed the Euclidean norm minus one g (ENMO) metric to summarise the raw acceleration signal vector magnitude (10). They have also applied acceleration cutpoints from the laboratory calibration study of Hildebrand et al. (2014) (2), which involved a convenience sample of 30, 7-11 year old children. These cutpoints have been used frequently, but are based on a limited number of activities and have not been cross-validated in free-living settings. As such, they may not be appropriate for all youth populations.

Generating further population-specific accelerometer cut-points limits comparability between studies, but this could be overcome by using standardised PA metrics that could maximise data comparability and the potential for data harmonisation (1). Recently, Rowlands (2018) argued that accelerometer metrics should be standardised so that they are meaningful, interpretable, and comparable (1). In particular, it was suggested that raw acceleration data should be used and reported as (i) the average acceleration (i.e., acceleration due to movement, corrected for gravity) as a measure of activity volume, and (ii) the profile of PA intensity, termed the

Intensity Gradient (IG) (1). The IG describes the straight line negative slope of the natural logs of time and acceleration intensity (11). A better IG is reflected by a shallower (i.e., less negative) slope, whereas a steeper (i.e., more negative) slope would reflect an inferior IG (11). The IG reflects the entire intensity profile, rather than small proportions of cutpoint-derived PA (e.g., moderate-to-vigorous intensity PA; MVPA). Further, it does not depend on the bias introduced by cutpoint calibration protocols, as it uses the full range of recorded data (11). Moreover, the IG is more independent of overall PA level, and therefore can be used in combination with average acceleration to describe intensity and volume of the PA profile. This allows the relative importance of PA intensity and volume to be examined in relation to specific health outcomes (1, 11). This information could subsequently be used to inform content and design of health-related interventions. Using the GENEActiv wrist accelerometer, Rowlands et al. (2018) demonstrated that average acceleration and IG each explained unique variance in PA profiles and were independently associated with body fatness and physical function (11). Average acceleration and IG therefore have potential as standardised measures describing PA volume and intensity, respectively, to explore their relative contributions to health, and to allow comparisons between studies where raw acceleration signals have been used. Importantly, this approach removes the complications of the ‘cutpoint conundrum’ which often render meaningful between-study comparisons impossible, provide inconsistent estimates of activity levels, and serve to confuse the evidence base and its interpretation (12). Generating and reporting data on PA volume and intensity as standard metrics would increase comparability between studies, with population-specific interpretation of the data (e.g., estimating time in specific intensities) applied post-analysis by the researchers themselves as well as by others (1, 11).

This study further examines the utility of the average acceleration and IG metrics using the ActiGraph accelerometer in a sample of primary school children. The study sought to address the following objectives:

1. Investigate whether hypothesised differences in PA between sex, weight status, obesity risk, metabolic risk, and cardiorespiratory fitness (CRF) status groups are apparent for average acceleration, IG, and MVPA.
2. Explore the magnitude of associations between average acceleration, IG, and cutpoint-based estimates of sedentary time (ST), light PA (LPA), and MVPA to determine whether the IG is more independent of average acceleration than the cut-point based metrics.
3. Investigate cross-sectionally, whether average acceleration and IG are independently associated with obesity indicators, metabolic risk, CRF, and health-related quality of life (HRQoL), when adjusting for covariates.
4. Examine the associations between baseline PA metrics and the health indicators described in #3 above, measured 8-weeks later.

Methods

This is a secondary analysis of data collected in the Active Schools: Skelmersdale PA pilot intervention study (ClinicalTrials.gov registration: NCT03283904). The methods have been described previously (8) but are outlined briefly here. Two hundred and thirty two 9-10 year old participants were recruited from 7 primary schools. The schools were situated in a low socioeconomic status (SES) town in West Lancashire, north-west England, where the prevalence of overweight/obesity is above the national average (13). Ethical approval was granted by Edge Hill University's Research Ethics Committee (reference # SPA-REC-2015-330) and informed consent and assent were provided by the participants' parents/carers, and

the participants themselves, respectively. Data collection took place between September and December 2017.

Following collection of baseline measurements, schools were randomly assigned to either intervention (4 schools) or control groups (3 schools). The AS:Sk pilot intervention included eight components which were implemented over 8-weeks. The components were active classroom breaks, high-intensity jumping activities, structured exercise videos, running/walking activities, playground activity challenge cards, physical education teacher training, parental newsletters, and PA homework. Control schools continued with their usual timetabled amount of playground breaks and physical education lessons without any additional time allocated for PA participation. No intervention effects were observed for MVPA but sedentary time (ST) decreased in the intervention schools (8).

Measures

Physical activity

Participants wore an ActiGraph GT9X triaxial accelerometer (ActiGraph, Pensacola, FL, USA) on the non-dominant wrist for 24 h·d⁻¹ over seven days. Wrist-worn accelerometers have demonstrated excellent validity against energy expenditure as the criterion measure (14), and in comparison to hip-worn accelerometers (14, 15). ActiGraphs recorded accelerations at 100 Hz, data were downloaded using ActiLife version 6.11.9 (ActiGraph, Pensacola, FL, USA), and saved in raw format as GT3X files, before being converted to raw csv file format for signal processing. These csv files were processed in R (<http://cran.r-project.org>) using GGIR beta v1.6-1 which carried out autocalibration procedures (16), identified non-wear (10), and converted the raw triaxial accelerometer signals into one omnidirectional measure of acceleration (ENMO) (10). Computed average day (labelled ‘AD’ in GGIR) ENMO values

were averaged per 5 s epoch over each of the seven monitored days to represent average acceleration, and were expressed in milligravitational units (mg). Accelerometer non-wear was determined based on the SD and value range of the accelerations at each axis, calculated for 60-min windows with a 15-min sliding window (10). If for at least 2 out of the 3 axes the SD was less than 13 mg or the value range was less than 50 mg, the time window was classified as non-wear (10). By default, GGIR imputed non-wear data by the average at similar time points on other days of the week. Therefore, participants' outcome variables were based on the complete 24-h cycle (i.e., 1440 min). Participants were excluded if the ActiGraph files demonstrated (i) post-calibration error greater than 0.01 g (16), (ii) less than 3 valid days of wear (17), which was defined as at least 16 h d⁻¹ (11), or (iii) missing wear data for any 15-min window over the 24-h cycle, indicated in GGIR by the '24-h cycle <1' variable.

Total PA was expressed as the average acceleration over 24-h. The IG metric was calculated in GGIR following the method described by Rowlands et al. (2018) (11) and was represented by the 'AD_IG' variable in GGIR. The IG is based on the relationship between log values for intensity (i.e., incremental intensity bins, 0-25 mg, 25-50 mg, etc) and time (i.e., accumulated time in each intensity bin), and is always negative, reflecting the drop in time accumulated in increasing intensity bins (11). For each participant, their IG over 24-h, the constant of the linear regression equation, and R^2 value (indication of the goodness of fit of the linear model) were produced, as were time spent in LPA, MVPA, and time spent inactive. We used the only available published ENMO prediction equations to identify cut-points for classifying activity as MVPA (3 metabolic equivalents (METs; child-specific); 200 mg) (2). Inactive time was defined as time accumulated below 50 mg, which is consistent with the previous average acceleration and IG study (11) and recently published sedentary time thresholds (18). LPA was defined as > 50 and < 200 mg.

Obesity-related outcomes

Height was measured using a portable stadiometer (Leicester Height Measure, Seca, Birmingham, UK), and body mass was measured using calibrated scales (813 model, Seca). Body mass index (BMI) was calculated for each participant, BMI z-scores were assigned (19) and IOTF BMI cut-points applied to classify the participants as normal weight or overweight/obese (underweight participants were grouped into the normal weight category) (20). Waist circumference was measured using an anthropometric tape measure, and waist-to-height ratio (WHtR) was calculated as a measure of central obesity (21). A WHtR of ≥ 0.5 was used to categorise participants as at risk or not at risk of central obesity (22). Sex-specific equations were used to predict age from peak height velocity (APHV), as a proxy measure of biological maturation (23). For all measurements the participants wore shorts and t-shirt with shoes removed.

Cardiorespiratory fitness

The 20-m multistage shuttle run test was conducted to provide an estimate of cardiorespiratory fitness (CRF). This test has been used extensively with participants of a similar age to those in the current study (24). The running speed at the last completed lap was used to estimate peak oxygen uptake (VO_2 peak; $\text{ml kg}^{-1} \text{min}^{-1}$) using the Leger et al. prediction equation (25). Participants were classified as having higher or lower CRF levels using the 40th centile for VO_2 peak in European children, which is the normative quintile-based framework cutoff for low to very low fitness (boys: $47.0 \text{ ml kg}^{-1} \text{min}^{-1}$; girls: $44.4 \text{ ml kg}^{-1} \text{min}^{-1}$) (24).

Metabolic health

A metabolic syndrome (MetS) score was calculated to describe metabolic risk using non-invasive variables (26). Z-scores were calculated for WHtR and the inverse of CRF ($1/\text{VO}_2$ peak), summed, then averaged to provide a MetS risk score. This approach has demonstrated sensitivity and specificity of 0.85 in ROC analyses (26) against the International Diabetes Federation definition of MetS encompassing obesity prevalence and elevated levels of triglycerides, HDL-C, blood pressure, and glucose (27). A MetS risk z-score > 0.51 was used to classify participants as low or high risk of MetS (26).

Health-related quality of life

The KIDSCREEN-10 Index questionnaire was used as a measure of global health-related quality of life (HRQoL) (28). KIDSCREEN-10 Index is a 10-item questionnaire which asks participants how they felt in the last week. Items reflect the factors of physical well-being, psychological wellbeing, autonomy and parent relations, peers and social support, and school environment, which are derived from the 27-item version of KIDSCREEN and are presented using a 1-5 Likert scale (29). Raw scores were converted to T-scores using the methodology described in the KIDSCREEN administration manual (28).

Socioeconomic status

Neighbourhood-level SES was calculated for each child using the 2015 Indices of Multiple Deprivation (IMD) (30). The IMD is a UK government-produced deprivation measure for England comprising income, employment, health, education, housing, environment, and crime. IMD rank scores were generated from parent-reported home post codes using the National Statistics Postcode Directory database. Every neighbourhood in England is ranked from one (most deprived area) to 32,844 (least deprived area).

Analyses

Descriptive statistics were calculated for all measures using means (SD) or percentages for continuous and categorical variables, respectively. The main analyses were designed to address each research objective in turn. For objective 1, the dependent variables were average acceleration, IG, and MVPA. Mixed linear models with random intercepts were used to adjust for school-level clustering to compare each dependent variable by sex, weight status, central obesity risk status, MetS risk status, and CRF status. We hypothesised that PA would be greater among boys, participants with normal weight, low central obesity risk, low MetS risk, and higher CRF. Bivariate Pearson correlation coefficients were calculated to address objective 2. For objective 3, separate cross-sectional mixed linear models with random intercepts were constructed accounting for school-level clustering. Model 1 included only the PA metric (i.e., average acceleration or IG). Model 2 was additionally adjusted for sex, maturation, and SES, while Model 3 was further adjusted for the alternate metric (i.e., average acceleration or IG, depending on which was the predictor) to test whether associations were independent of either metric. Multicollinearity was checked using the variance inflation factor (VIF) with a VIF of >5 indicating excessive multicollinearity (31). To allow comparison of the IG results with MVPA, all models were repeated using average acceleration and MVPA as the alternate metrics. For objective 4, the dependent variables for these analyses were average acceleration, IG, and MVPA. Mixed linear models with random intercepts and adjusted for school-level clustering examined the association between the baseline PA metrics and health indicators measured 8-weeks later. Analyses were adjusted for the alternate PA metric, baseline health indicators, group designation (i.e., Control or Intervention group), sex, BMIz, maturation, and SES.

Regression coefficients in the main and interaction models were assessed for significance using the Wald statistic (32). Statistical significance was set at $p < .05$ in all analyses. Analyses for objectives 1, 3, and 4 were performed using MLwiN 2.26 software (Centre for Multilevel Modelling, University of Bristol, UK). IBM SPSS Statistics version 23 (IBM, Armonk, NY) was used to undertake analyses for research objective 2.

Results

Descriptive statistics are presented in Table 1. Baseline accelerometer data were available for 226 of the 232 participants (6 participants were absent on the day the accelerometers were distributed). Forty-one participants wore the accelerometers for $< 16 \text{ h} \cdot \text{d}^{-1}$ for at least 3 days, 39 participants had incomplete accelerometer data (i.e., missing wear data for any 15-min window over the 24-h cycle), and one participant had spurious accelerometer data. These participants were subsequently removed, which resulted in a final analytical baseline sample of 145 participants (62 boys). Almost 70% of the sample were classified as normal weight, 62.8% of them had higher CRF levels, and just over half engaged in at least 60 min MVPA per day. There were no significant differences between included and excluded participants in obesity-related variables, CRF, and HRQoL. Excluded participants were more likely to be girls ($p < .05$) with more advanced somatic maturity ($p < .05$).

TABLE 1

Objective 1. The hypothesised differences in PA metrics between boys and girls were observed for MVPA ($p < .05$ – $p < .001$), average acceleration ($p < .05$ – $p < .001$), and IG ($p < .001$) (Table 2). Hypothesised differences were observed for IG between normal weight and overweight/obese participants ($p < .001$), between those with low and high risk of central

obesity ($p<.001$), those with low and high risk of MetS ($p<.001$), and those with higher and lower CRF ($p<.05$).

TABLE 2

Objective 2. Average acceleration was strongly correlated with MVPA ($r=0.96$) and inactive time ($r=-0.82$) (both $p<.001$), but only moderately with LPA ($r = 0.59$, $p<0.001$) and IG ($r=0.54$, $p<.001$). IG and MVPA were moderately correlated ($r=0.50$, $p<.001$). IG was weakly and non-significantly correlated with inactive time ($r=-0.13$), and LPA ($r=-0.09$).

Objective 3. Results of the models investigating the cross-sectional associations between average acceleration and IG, with health indicators are presented in Table 3. Significant effects for average acceleration or IG, independent of the alternate metric (Model 3), indicated whether volume or intensity were most important for a given health indicator. Average acceleration was significantly associated with BMIz, CRF, and HRQoL in the first (unadjusted) and second (adjusted) models. Average acceleration was not significantly associated with BMIz, WHtR, CRF, and MetS score in the third models when IG was included, indicating that the associations between average acceleration and these health indicators were not independent of IG. Conversely, the association between average acceleration and HRQoL was independent of IG in model 3, indicating that PA volume, rather than intensity was more important for HRQoL. In unadjusted and adjusted models, IG was negatively significantly associated with BMIz, WHtR, and MetS risk score, and positively significantly associated with CRF, HRQoL. These associations were significant independent of average acceleration in the third models, with the exception of HRQoL, indicating that PA intensity rather than volume was most important for the physical health indicators. When average acceleration and MVPA were included as the

alternate PA metrics, MVPA was positively associated with CRF and HRQoL in the unadjusted models and the adjusted models 2 (Table S1). This significant association was not observed in model 3 for HRQoL, which demonstrated that the association with MVPA was not independent of average acceleration, while the significant association between CRF and MVPA was maintained in model 3. MVPA was also significantly associated with MetS score in model 3, indicating that this association was independent of average acceleration. Average acceleration was significantly associated with BMIz, WHtR, CRF, and MetS score when adjusted for MVPA in the third models. In all analyses the VIF values ranged from 1.02 (IMD rank) to 4.26 (sex).

TABLE 3

Objective 4. When baseline and follow-up health indicator data were merged, nine participants were lost due to absence on the day of data collection. This resulted in an analytical sample of 136 participants. WHtR and MetS score at follow-up were significantly associated with baseline IG, independent of average acceleration (Table 4). Specifically, significant inverse associations were observed between baseline IG and follow-up WHtR ($p<.01$) and MetS score ($p<.05$), indicating that PA intensity at baseline was more important than volume for follow-up WHtR and MetS score. When the analyses were repeated with baseline average acceleration and MVPA as the alternate metrics, no significant associations were observed with any health indicators at follow-up (Table S2). The Beta values of the health indicators were greatest when baseline IG was the predictor variable compared to average acceleration and MVPA.

TABLE 4

An example of translation and interpretation of the descriptive results is presented in Figure 1, which shows the activity profile for the sample categorised by IG tertile. Acceleration is described according to thresholds of 0-49 mg (inactive time), 50-199 mg (pottering/slow walking; LPA), 200-699 mg (brisk walking/jogging; MPA), and 700 mg+ (slow to fast running; VPA). The time spent inactive is at the base of each column. The time spent inactive and in pottering/slow walking was similar across tertiles. For participants in the High IG tertile, ~12 min and ~20 min more time was spent in activities equivalent to brisk walking/jogging, compared to participants in the Medium and Low IG tertiles, respectively. Moreover, High IG participants accumulated almost twice as much time in the highest intensity acceleration activities (i.e., slow-to-fast running), than those in the medium tertile, and three times as much time as peers in the low IG group.

FIGURE 1

Discussion

This is the first study to use the ActiGraph GT9X wrist accelerometer with primary-school aged children, to investigate the utility of average acceleration and IG relative to hypothesised between-group differences in PA, and in relation to associations with cutpoint-based PA metrics, and health indicators. The higher average acceleration (45.4 mg) and lower IG (-1.96) values observed in our primary school sample, compared to previously reported values for adolescent girls (36.3 mg and -2.47) and adults (22.1 mg and -3.11) (11) are consistent with expected age-related differences in PA (33, 34). When we investigated hypothesised PA differences between dichotomised groups defined by weight status, obesity risk, MetS risk, and CRF status, significant differences were observed for IG in all analyses, whereby the ‘healthier’ groups had more favourable (i.e., shallower) intensity profiles, which are indicative of

engagement in relatively higher PA intensities. When the analyses were repeated with average acceleration and MVPA, no between-group differences were evident. The significant differences in IG in all analyses offers support for the potentially greater sensitivity of this metric to detect between group differences in PA. This is possibly because the IG reflects the full intensity spectrum and uses all of the acceleration information available (35), compared to cutpoint-based methods which are subject to greater sources of error (36) and which only represent a small proportion of the day (e.g., 63.9 min of MVPA or 4% of the day in this sample).

The magnitudes of the correlations between IG and inactivity, LPA, and MVPA ($r = -0.09 - 0.50$) were smaller than those observed between average acceleration and the cutpoint-based outcomes ($r = 0.59 - 0.96$). These analyses confirm previous work (11), showing that IG is more independent of average acceleration (i.e., volume of PA) than the cutpoint-based outcomes. Furthermore, the stronger correlations between IG and MVPA compared to those with LPA and inactivity, demonstrate the utility of IG as a PA metric that captures higher intensity PA, which has greatest health benefits (6, 35). The cross-sectional relationships between the PA metrics and indicators of adiposity mirrored those reported in adolescent girls (11) whereby in adjusted analyses, IG was significantly associated with BMI_z and WHtR, independent of average acceleration. The same outcome was observed with CRF and MetS score as the dependent variables, indicating that PA intensity is more important than PA volume for these health indicators. In adjusted models, average acceleration was significantly associated with BMI_z and CRF but these relationships were no longer significant when IG was added. In contrast, when average acceleration and MVPA (rather than IG) were included in the models, a significant independent association was only observed between MVPA and MetS score. This is consistent with studies using ActiGraph counts which demonstrated time spent

in higher intensities of PA were most strongly associated with cardiometabolic risk (37, 38). These findings provide further evidence of the sensitivity of IG as an indicator of PA intensity that is relatively independent of total volume of PA (i.e., average acceleration), and which relates to important indicators of physical health in children.

Further interpretation of these findings is possible by considering the unit change in health indicators relative to the change in IG. The SD of the IG in our sample was 0.14 which we employed with the final regression models to demonstrate predicted change in the health indicators (represented by the Beta values in Table 3). For example, a 0.14 increase in IG would be reflected by the following changes: BMIz Δ -0.57, WHtR Δ -0.03, VO₂ peak Δ +1.93 ml·kg⁻¹·min⁻¹, and MetS score Δ -0.23. Such a change in BMIz is greater than reductions reported in intervention studies among obese (39) and non-obese children (40). Moreover, the equivalent change in CRF for a 0.14 increase in IG would be sufficient to shift a 10-year old child up approximately two deciles of recently published normative VO₂ peak values (24). Thus, improved engagement in higher intensity activities (for example through intervention programming, active play, sports participation, etc) would be reflected by shallower intensity profiles represented by the IG, which are associated with meaningful and favourable changes in physical health indicators. In keeping with recent recommendations, there is a need to provide further translation examples so these new PA metrics are interpretable and user-friendly (1). For example, applying the procedure described by Rowlands et al. (11) children in the current sample would need to replace time spent at the average acceleration with brisk walking for 2-h, slow running for 24-min, or medium-paced running for 19-min, accumulated across the day, in order to increase their average acceleration by 1 SD (13.1 mg). Such changes could be achieved through increased participation in daily PA opportunities such as active school commuting (41) (i.e., brisk walking), and school-based activities such as active recess

play (42) and co-curricular activities like running programmes that are becoming increasingly popular in primary school settings (43) (i.e., slow and medium-paced running). For each child, such changes would have an impact on their IG values, with the greatest impact coming from the more intense activities (i.e. running) (11), as described in Figure 1. Therefore, running (or activities of an equivalent intensity) could be recommended for BMIz, WHtR, MetS score, and CRF, which demonstrated an independent effect of IG. Further translation of incremental intensity distributions using health-related acceleration thresholds or indicative activity modes, could also be used to aid public health messaging and intervention programme design.

The analyses of baseline PA metrics relative to the health indicators measured 8-weeks later resulted in significant associations between IG and WHtR and MetS score. When the Beta values were summed by IG SD of 0.14, this resulted in predicted changes of -0.01 WHtR units and -0.07 MetS score. Although such changes are favourable, the short follow-up period limits how meaningful the magnitude of these associations are. Longitudinal studies of at least 2-years have demonstrated prospective associations between MVPA and decreased fat mass (44), and metabolic risk (45), between absence of organised sport participation and increased BMIz (46), and between VPA and decreases in BMIz and waist circumference, and increases in CRF (47). These prospective associations reinforce the importance of higher intensity PA for health in children, and our findings show that IG is sensitive to capture such higher intensity PA, independent of PA volume, albeit over a relatively short follow-up period.

The association between HRQoL and IG was not independent of average acceleration, while the opposite was true when average acceleration was the metric of interest. This suggests that simply moving more (i.e., increasing PA volume), irrespective of intensity was positively associated with increased HRQoL. PA metrics have seldom been studied in relation to

wellbeing, and this is the first study to examine the utility of the IG in relation to HRQoL as a wellbeing indicator. A recent systematic review reported that there were inconclusive relationships between PA and wellbeing, mainly due to lack of consistent PA and wellbeing outcome measures between studies, with greatest variability in the latter (48). For example, it was found that children who self-reported meeting the 60 min·d⁻¹ MVPA guideline scored higher on the KIDSCREEN-52 dimensions of self-perceptions, social acceptance, and social support (49). A similar finding was observed among youth who self-reported time spent in various sports (deemed equivalent to MVPA) and completed the PedsQL HRQoL questionnaire (50). In contrast, an intervention study where children wore a hip-mounted ActiGraph and completed the KIDSCREEN-27 reported increased MVPA in the control group, yet no corresponding changes in HRQoL (51). Further work is needed to better understand the inter-relationships between objective PA metrics and HRQoL. This would be helped by more consistent choice of measures, which reinforces support to the call for the use of average acceleration and IG as standard PA metrics (1).

This study is limited by the modestly-sized sample that was located in a low SES English town, which may inhibit generalisability of the findings to other populations. Moreover, most of the analyses were cross-sectional, which prohibits conclusions about causality between the PA metrics and health indicators. The cutpoints used to determine MVPA may have been subject to population-specific and protocol-specific biases, which could have influenced the accuracy of the reported MVPA estimates. Further, although accelerometer wear averaged 19.2 hours·valid d⁻¹ for 5.2 valid days we cannot be certain that the wear time criteria applied were suitable for use with the average acceleration and IG metrics. Future work may be needed to establish the wear time criteria required to reliably estimate typical 7-day values for these new metrics. The 8-weeks follow-up measures were a strength of the study, but this period may not

have been long enough to make meaningful inferences about the associations with the baseline PA metrics. Other strengths included the use of mixed linear models to account for school-level variance and the inclusion of known correlates of PA in the models. Furthermore, the study reported associations between the PA metrics and a range of important physical and mental health indicators.

Conclusions

This is the first study to examine the utility of the recently introduced average acceleration and IG metrics using the wrist-worn ActiGraph monitor with primary-school aged children. Significant differences in IG were observed between sex, weight status, central obesity risk, and CRF status groups. Moreover, IG was significantly associated with BMI_z, WHtR, CRF, and MetS score independent of average acceleration. The magnitude of these associations reflected meaningfully beneficial changes in health indicators. Significant associations of a smaller magnitude were apparent between baseline IG and WHtR and MetS score at 8-weeks follow-up. The results provide further evidence of the utility of average acceleration and IG to describe children's PA, and go beyond those reported previously reported, by including health indicators reflecting CRF, metabolic risk, and HRQoL, rather than just obesity-related measures. The IG can be considered a meaningful PA metric that is sensitive as a gauge of PA intensity, that is independent of total volume of PA, and which relates to important indicators of physical health in children.

Discosure of interest

The authors declare no conflicts of interest.

Data access statement

The data that support the findings of this study are available at <https://osf.io/tfpk9/>.

Acknowledgements (including author contributions)

We are grateful to the children and teachers for their participation in the project, and West Lancashire Sport Partnership for their assistance with the data collection. The study was funded by West Lancashire Sport Partnership, West Lancashire Leisure Trust, and Edge Hill University. AVR is with the NIHR Leicester Biomedical Research Centre, and the Collaboration for leadership in Applied Health Research and Care (CLAHRC) East Midlands. The views expressed are those of the authors and not necessarily those of the NHS, NIHR, or Department of Health. SJF conceived and designed the study, performed the data analyses, and drafted the manuscript. ST contributed to the study design, collected the data, and edited and commented on the manuscript. AVR advised on the data processing and analyses, and edited and commented on the manuscript. LMB advised on the data analyses and edited and commented on the manuscript. RJN contributed to the study design, edited and commented on the manuscript. All authors had final approval of the submitted manuscript.

References

1. Rowlands AV. Moving forward with accelerometer-assessed physical activity: two strategies to ensure meaningful, interpretable, and comparable measures. *Pediatr Exerc Sci.* 2018;30:450-6.
2. Hildebrand M, Van Hees VT, Hansen BH, Ekelund U. Age-group comparability of raw accelerometer output from wrist- and hip-worn monitors. *Med Sci Sports Exerc.* 2014;46(9):1816-24.
3. Phillips LR, Parfitt G, Rowlands AV. Calibration of the GENEActiv accelerometer for assessment of physical activity intensity in children. *J Sci Med Sport.* 2012;16(2):124-8.
4. Schaefer CA, Nigg CR, Hill JO, Brink LA, Browning RC. Establishing and evaluating wrist cutpoints for the GENEActiv accelerometer in youth. *Med Sci Sports Exerc.* 2014;46:826-33.
5. Bornstein DB, Beets MW, Byun W, Welk G, Bottai M, Dowda M, et al. Equating accelerometer estimates of moderate-to-vigorous physical activity: In search of the Rosetta Stone. *J Sci Med Sport.* 2011;14(5):404-10.
6. Fairclough SJ, Dumuid D, Taylor S, Curry W, McGrane B, Stratton G, et al. Fitness, fatness and the reallocation of time between children's daily movement behaviours: an analysis of compositional data. *Int J Behav Nutr Phys Activity.* 2017;14(1):64.
7. Fairclough SJ, Noonan R, Rowlands AV, Van Hees V, Knowles Z, Boddy LM. Wear compliance and activity in children wearing wrist- and hip-mounted accelerometers. *Med Sci Sports Exerc.* 2016;48(2):245-53.
8. Taylor S, Noonan R, Knowles Z, Owen M, McGrane B, Curry W, et al. Evaluation of a pilot school-based physical activity clustered randomised controlled trial—Active Schools: Skelmersdale. *Int J Environ Res Public Health.* 2018;15(5):doi:10.3390/ijerph15051011.

- 531 9. Rowlands AV, Harrington DM, Bodicoat DH, Davies MJ, Sherar LB, Gorely T, et al.
532 Compliance of adolescent girls to repeated deployments of wrist-worn accelerometers. *Med*
533 *Sci Sports Exerc.* 2018;50(7):1508-17.
- 534 10. van Hees VT, Gorzelniak L, León EC, Eder M, Pias Mo, Taherian S, et al. Separating
535 movement and gravity components in an acceleration signal and implications for the
536 assessment of human daily physical activity. *PLoS ONE.* 2013;8(4):e61691. doi:
537 10.1371/journal.pone.0061691.
- 538 11. Rowlands AV, Edwardson CL, Davies MJ, Khunti K, Harrington DM, Yates T.
539 Beyond cut-points: accelerometer metrics that capture the physical activity profile. *Med Sci*
540 *Sports Exerc.* 2018;50(6):1323-32.
- 541 12. Migueles JH, Cadenas-Sanchez C, Tudor-Locke C, Löf M, Esteban-Cornejo I,
542 Molina-Garcia P, et al. Comparability of published cut-points for the assessment of physical
543 activity: Implications for data harmonization. *Scand J Med Sci Sports.* 2019;29:566-74.
- 544 13. Collins S. The Seven Wards: A Focus on Skelmersdale. Preston, England: Lancashire
545 County Council; 2015.
- 546 14. Esliger DW, Rowlands AV, Hurst TL, Catt M, Murray P, Eston RG. Validation of the
547 GENEa accelerometer. *Med Sci Sports Exerc.* 2011;43(6):1085-93.
- 548 15. Rowlands AV, Rennie K, Kozarski R, Stanley RM, Eston RG, Parfitt GC, et al.
549 Children's physical activity assessed with wrist- and hip-worn accelerometers. *Med Sci*
550 *Sports Exerc.* 2014;46(12):2308-16.
- 551 16. van Hees VT, Fang Z, Langford J, Assah F, Mohammad A, da Silva IC, et al.
552 Autocalibration of accelerometer data for free-living physical activity assessment using local
553 gravity and temperature: an evaluation on four continents. *J Appl Physiol.* 2014;117(7):738-
554 44.

555 17. Rich C, Geraci M, Griffiths L, Sera F, Dezateux C, Cortina-Borja M. Quality control
556 methods in accelerometer data processing: defining minimum wear time. PLoS ONE.
557 2013;8(6):e67206. doi: 10.1371/journal.pone.0067206.

558 18. Hurter L, Fairclough SJ, Knowles Z, Porcellato L, Cooper-Ryan A, Boddy L.
559 Establishing raw acceleration thresholds to classify sedentary and stationary behaviour in
560 children. Children. 2018;5(12):172. doi: 10.3390/children5120172.

561 19. Cole T, Freeman J, Preece M. Body mass index reference curves for the UK, 1990.
562 Arch Dis Child. 1995;73(1):25 - 9.

563 20. Cole T, Bellizzi M, Flegal K, Dietz W. Establishing a standard definition for child
564 overweight and obesity worldwide: international survey. Br Med J. 2000;320:1240 - 3.

565 21. Mokha JS, Srinivasan SR, DasMahapatra P, Fernandez C, Chen W, Xu J, et al. Utility
566 of waist-to-height ratio in assessing the status of central obesity and related cardiometabolic
567 risk profile among normal weight and overweight/obese children: The Bogalusa Heart Study.
568 BMC Pediatr. 2010;10(1):73. doi: 10.1186/1471-2431-10-73.

569 22. Mehta SK. Waist circumference to height ratio in children and adolescents. Clinical
570 Pediatr. 2015;54(7):652-8.

571 23. Moore SA, McKay HA, Macdonald H, Nettleford L, Baxter-Jones ADG, Cameron
572 N, et al. Enhancing a somatic maturity prediction model. Med Sci Sports Exerc.
573 2015;47(8):1755-64.

574 24. Tomkinson GR, Carver KD, Atkinson F, Daniell ND, Lewis LK, Fitzgerald JS, et al.
575 European normative values for physical fitness in children and adolescents aged 9–17 years:
576 results from 2 779 165 Eurofit performances representing 30 countries. Br J Sports Med.
577 2018;52(22):1445-14563.

578 25. Leger LA, Mercier D, Gadoury C, Lambert J. The multistage 20 metre shuttle run test
579 for aerobic fitness. J Sports Sci. 1988;6:93-101.

580 26. Andersen LB, Lauersen JB, Brond JC, Anderssen SA, Sardinha LB, Steene-
581 Johannessen J, et al. A new approach to define and diagnose cardiometabolic disorder in
582 children. *J Diabetes Res.* 2015;2015:10. doi: 10.1155/2015/539835.

583 27. Zimmet P, Alberti KGM, Kaufman F, Tajima N, Silink M, Arslanian S, et al. The
584 metabolic syndrome in children and adolescents – an IDF consensus report. *Pediatr Diabetes.*
585 2007;8(5):299-306.

586 28. KIDSCREEN Group Europe. The KIDSCREEN Questionnaires - Quality of life
587 questionnaires for children and adolescents. Handbook. Lengerich: Pabst Science Publishers;
588 2006.

589 29. Ravens-Sieberer U, Gosch A, Rajmil L, Erhart M, Bruil J, Duer W, et al.
590 KIDSCREEN-52 quality-of-life measure for children and adolescents. *Expert Rev*
591 *Pharmacoeconomics & Outcomes Res.* 2005;5(3):353-64.

592 30. Department for Communities and Local Government. The English Indices of
593 Deprivation. Statistical Release. London: DCLG; 2015.

594 31. Montgomery D, Peck E, Vining G. *Introduction to Linear Regression Analysis.* New
595 York: John Wiley and Sons, Inc.; 2001.

596 32. Twisk JWR. *Applied Multilevel Analysis.* Cambridge: Cambridge University Press;
597 2006.

598 33. Farooq MA, Parkinson KN, Adamson AJ, Pearce MS, Reilly JK, Hughes AR, et al.
599 Timing of the decline in physical activity in childhood and adolescence: Gateshead
600 Millennium Cohort Study. *Br J Sports Med.* 2017;52:1002-6.

601 34. Doherty AR, Jackson D, Hammerla N, Plötz T, Olivier P, Granat MH, et al. Large
602 scale population assessment of physical activity using wrist worn accelerometers: The UK
603 Biobank Study. *PLoS ONE.* 2017;12(2):e0169649. doi: 10.1371/journal.pone.0169649.

604 35. Aadland E, Andersen LB, Anderssen SA, Resaland GK, Kvalheim OM. Associations
605 of volumes and patterns of physical activity with metabolic health in children: A multivariate
606 pattern analysis approach. *Prev Med.* 2018;115:12-8.

607 36. Troiano RP, McClain JJ, Brychta RJ, Chen KY. Evolution of accelerometer methods
608 for physical activity research. *Br J Sports Med.* 2014;48(13):1019-23.

609 37. Anderson CB, Hagstromer M, Yngve A. Validation of the PDPAR as an adolescent
610 diary: Effect of accelerometer cut points. *Med Sci Sports Exerc.* 2005;37:1224-30.

611 38. Tarp J, Child A, White T, Westgate K, Bugge A, Grøntved A, et al. Physical activity
612 intensity, bout-duration, and cardiometabolic risk markers in children and adolescents. *Int J*
613 *Obes.* 2018;42:1639-50.

614 39. Ho M, Garnett SP, Baur L, Burrows T, Stewart L, Neve M, et al. Effectiveness of
615 lifestyle interventions in child obesity: systematic review with meta-analysis. *Pediatr.*
616 2012;130(6):e1647-71.

617 40. Fairclough SJ, Hackett A, Davies I, Gobbi R, Mackintosh K, Warburton G, et al.
618 Promoting healthy weight in primary school children through physical activity and nutrition
619 education: a pragmatic evaluation of the CHANGE! randomised intervention study. *BMC*
620 *Public Health.* 2013;13(1):626. doi: 10.1186/1471-2458-13-626.

621 41. Cooper AR, Page AS, Wheeler BW, Griew P, Davis L, Hillsdon M, et al. Mapping
622 the walk to school using accelerometry combined with a global positioning system. *Am J*
623 *Prev Med.* 2010;38(2):178-83.

624 42. Ridgers ND, Salmon J, Parrish A-M, Stanley RM, Okely AD. Physical activity during
625 school recess: a systematic review. *Am J Prev Med.* 2012;43(3):320-8.

626 43. Chesham RA, Booth JN, Sweeney EL, Ryde GC, Gorely T, Brooks NE, et al. The
627 Daily Mile makes primary school children more active, less sedentary and improves their

628 fitness and body composition: a quasi-experimental pilot study. *BMC Med.* 2018;16(1):64.
629 doi: 10.1186/s12916-018-1049-z.

630 44. Riddoch CJ, Leary SD, Ness AR, Blair SN, Deere K, Mattocks C, et al. Prospective
631 associations between objective measures of physical activity and fat mass in 12-14 year old
632 children: the Avon Longitudinal Study of Parents and Children (ALSPAC). *Br Med J.*
633 2009;339: doi: 10.1136/bmj.b4544.

634 45. Metcalf BS, Voss LD, Hosking J, Jeffery AN, Wilkin TJ. Physical activity at the
635 government-recommended level and obesity-related health outcomes: a longitudinal study
636 (Early Bird 37). *Arch Dis Child.* 2008;93(9):772-7.

637 46. Koning M, Hoekstra T, de Jong E, Visscher TLS, Seidell JC, Renders CM.
638 Identifying developmental trajectories of body mass index in childhood using latent class
639 growth (mixture) modelling: associations with dietary, sedentary and physical activity
640 behaviors: a longitudinal study. *BMC Public Health.* 2016;16(1):1128. doi: 10.1186/s12889-
641 016-3757-7.

642 47. Carson V, Rinaldi RL, Torrance B, Maximova K, Ball GDC, Majumdar SR, et al.
643 Vigorous physical activity and longitudinal associations with cardiometabolic risk factors in
644 youth. *Int J Obes.* 2013;38:16-21.

645 48. Rafferty R, Breslin G, Brennan D, Hassan D. A systematic review of school-based
646 physical activity interventions on children's wellbeing. *Int Rev Sport Exerc Psychol.*
647 2016;9:215-30.

648 49. Breslin G, Gossrau-Breen D, McCay N, Gilmore G, Macdonald L, Hanna D. Physical
649 activity, gender, weight status, and wellbeing in 9- to 11-year-old children: a cross-sectional
650 survey. *J Phys Act Health.* 2012;9(3):394-401.

- 651 50. Gopinath B, Hardy LL, Baur LA, Burlutsky G, Mitchell P. Physical activity and
652 sedentary behaviors and health-related quality of life in adolescents. *Pediatr.*
653 2012;130(1):e167-e74.
- 654 51. Ha AS, Burnett A, Sum R, Medic N, Ng JYY. Outcomes of the rope skipping ‘STAR’
655 programme for schoolchildren. 2015;45(1):233-240.
- 656

657 Table 1. Descriptive characteristics of the study sample

658 Table 2. Between-group differences in MVPA, average acceleration, and intensity gradient

659 Table 3. Cross-sectional associations between the physical activity metrics and health

660 indicators

661 Table 4. Associations between baseline physical activity metrics and health indicators

662 measured 8-weeks later

663 Figure 1. Accumulated time spent in acceleration ranges of participants categorised by

664 intensity gradient tertile

665