



LJMU Research Online

Maiorana, AJ, Naylor, LH, Dongelmans, S, Jacques, A, Thijssen, DHJ, Dembo, L, O'Driscoll, G and Green, DJ

Ventilatory efficiency is a stronger prognostic indicator than peak oxygen uptake or body mass index in heart failure with reduced ejection fraction.

<http://researchonline.ljmu.ac.uk/id/eprint/11111/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Maiorana, AJ, Naylor, LH, Dongelmans, S, Jacques, A, Thijssen, DHJ, Dembo, L, O'Driscoll, G and Green, DJ (2019) Ventilatory efficiency is a stronger prognostic indicator than peak oxygen uptake or body mass index in heart failure with reduced ejection fraction. European Journal of

LJMU has developed [LJMU Research Online](http://researchonline.ljmu.ac.uk/) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

Ventilatory efficiency is a stronger prognostic indicator than peak oxygen uptake or BMI in heart failure with reduced ejection fraction

Andrew J Maiorana^{a,b}
Louise H Naylor^{a,c}
Stijn Dongelmans^{c,d}
Angela Jacques^b
Dick H.J. Thijssen^{d,e}
Lawrence Dembo^a
Gerry O'Driscoll^f
Daniel J. Green^c

^a*Advanced Heart Failure & Cardiac Transplant Service; Department of Allied Health, Fiona Stanley Hospital, Australia*

^b*School of Physiotherapy & Exercise Science, Curtin University, Australia.*

^c*School of Human Sciences (Exercise and Sport Science), The University of Western Australia, Australia*

^d*Radboud Institute for Health Sciences, Radboud University Medical Center, the Netherlands*

^e*Research Institute for Sport & Exercise Sciences, Liverpool John Moores University, UK*

^f*Perth Cardiovascular Institute, Nedlands, Australia*

Sources of support

DJG was supported by a National Health and Medical Research Council Principal Research Fellowship (PRF) 1080914.

DHJT was supported by the Netherlands Heart Foundation (2009T064).

Disclosures

None.

Author for correspondence and request for reprints:

Associate Professor Andrew Maiorana, School of Physiotherapy and Exercise Science
Curtin University, GPO Box U1987, Perth, Western Australia 6845
Telephone: +618 9266 9225
Fax: +618 9266 3699
Email: A.Maiorana@curtin.edu.au

Word count

1234

Chronic heart failure (HF) is a complex condition associated with poor prognosis.¹ In HF with reduced ejection fraction (HFrEF), $\dot{V} O_{2peak}$ is a strong prognostic indicator for mortality, while at extremes of age, expressing $\dot{V} O_{2peak}$ relative to age-predicted values is recommended.² However, it can be difficult to achieve a peak exercise test and in such cases the relationship between ventilation (\dot{V}_E) and carbon dioxide production ($\dot{V} CO_2$) may be able to guide prognosis. An elevated $\dot{V}_E/\dot{V} CO_2$ slope is associated with poor prognosis,² while the exact cut point might be modified by age.³

There is evidence in HF that having a high body mass index (BMI) may paradoxically confer ‘protection’ against mortality.^{4,5} Studies examining these prognostic indicators in HF suggest that this “obesity paradox” may be evident in those with low cardiopulmonary fitness ($\dot{V} O_{2peak}$), but less apparent when fitness is preserved.⁶⁻⁹ We hypothesised that, in patients with HFrEF, elevated $\dot{V}_E/\dot{V} CO_2$ slope would outperform $\dot{V} O_{2peak}$ and BMI as a predictor of all-cause mortality, and that those with a low $\dot{V}_E/\dot{V} CO_2$ slope would have worse outcomes if they were also in the lower range for BMI.

A retrospective analysis was conducted on cardiopulmonary exercise testing (CPET) data from 312 HFrEF patients between 1997 and 2014. The study complied with the Declaration of Helsinki and was approved by Royal Perth Hospital Ethics Committee (REG 14-068). $\dot{V} O_{2peak}$ was assessed using an incremental CPET performed on a treadmill. Regression line of \dot{V}_E and $\dot{V} CO_2$ production was used to calculate $\dot{V}_E/\dot{V} CO_2$ slope. Patients were characterized as having “higher fitness” or “lower fitness” with cut points for $\dot{V} O_{2peak}$ at $14\text{ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$, for age-predicted $\dot{V} O_{2peak}$ at 50% and for $\dot{V}_E/\dot{V} CO_2$ slope at 35. They were further characterised as having ‘normal BMI’ ($18.5\text{-}25.0\text{kg}\cdot\text{m}^{-2}$) or ‘high BMI’ ($>25\text{kg}\cdot\text{m}^{-2}$). This gave

four classifications: 'normal BMI and lower fitness (or slope)', 'high BMI and lower fitness', 'normal BMI and higher fitness', or 'high BMI and higher BMI'.

All-cause mortality was documented from hospital records and a mortality database to May 2016. Patients that received a cardiac transplant or ventricular assist device were censored. Baseline characteristics were calculated for the four fitness-overweight groups with one-way analysis and chi-square tests were conducted. Kaplan-Meier analysis was used to produce survival functions, which were compared between the groups using Log rank tests. Cox regression models were used for calculating hazard ratios (HR), using the fitness indicators as variables, expressed with 95% confidence intervals.

Clinical characteristics of groups based on $\dot{V} O_{2peak}$ ($ml \cdot kg \cdot min^{-1}$) and BMI are reported in Table 1. All 3 fitness indicators were independent predictors of mortality at 2 ($p < 0.01$) and 5 years ($P < 0.001$). Having higher or lower BMI was not an independent predictor of mortality. Cox regression showed having a high $\dot{V}_E / \dot{V} CO_2$ slope as the strongest predictor of mortality with HR of 6.83 (2.75-16.97) and 4.13 (2.34-7.25) for 2 and 5 years. Figure 1 shows survival for the different groups. Patients who had higher fitness based on $\dot{V} O_{2peak}$ had better survival outcomes than their unfit counterparts, regardless of BMI. Whereas in the lower fitness group, those who had higher BMI had significantly higher survival those with normal BMI. When $\dot{V}_E / \dot{V} CO_2$ (\geq or < 35) was used, lower slope patients had better survival than their higher slope counterparts, regardless of their BMI. $\dot{V}_E / \dot{V} CO_2$ slope did not discriminate between the BMI categories in terms of survival.

Our findings confirm that fitness, and particularly ventilatory inefficiency (high \dot{V}_E/\dot{V}_{CO_2} slope), predicted mortality, whereas BMI did not. Those with elevated \dot{V}_E/\dot{V}_{CO_2} (≥ 35) had lower survival, regardless of their BMI. These data confirm the prognostic significance of cardiopulmonary fitness and submaximal exercise derived ventilatory inefficiency.

High BMI has been reported to convey a survival advantage in those with established HF and lower fitness.^{4,5,9} While we found this to be the case when fitness was stratified using $\dot{V}O_{2peak}$, it was less apparent when using ventilatory efficiency. Our results add to the growing evidence that suggest the obesity paradox is mostly due to confounding factors.^{4,7,8}

The \dot{V}_E/\dot{V}_{CO_2} slope has demonstrated to confer predictive capacity which may exceed that associated with $\dot{V}O_{2peak}$.² our results indicate stronger predictive capacity for \dot{V}_E/\dot{V}_{CO_2} slope over $\dot{V}O_{2peak}$. These findings contribute to evidence that ventilatory inefficiency, derived from submaximal stages of a graded exercise test, are a valuable component of prognostic assessment for patients with HF.

Declaration of conflicting interests

The Authors declare that there is no conflict of interest

References

1. Benjamin EJ, Blaha MJ, Chiuve SE, et al. Heart Disease and Stroke Statistics—2017 Update: A Report From the American Heart Association. *Circulation* 2017; 135: e146–e603. DOI: 10.1161/cir.0000000000000485.
2. Mehra MR, Canter CE, Hannan MM, et al. The 2016 International Society for Heart Lung Transplantation listing criteria for heart transplantation: A 10-year update. *J Heart Lung Transpl* 2016; 35: 1-23. DOI: 10.1016/j.healun.2015.10.023.
3. Kato Y, Suzuki S, Uejima T, et al. Relationship between the prognostic value of ventilatory efficiency and age in patients with heart failure. *Eur J Prev Cardiol* 2018; 25: 731-739. 2018/02/13. DOI: 10.1177/2047487318758775 [doi].
4. Pandey A, Patel KV and Lavie CJ. Obesity, Central Adiposity, and Fitness: Understanding the Obesity Paradox in the Context of Other Cardiometabolic Parameters. *Mayo Clin Proc* 2018; 93: 676-678. 2018/06/06. DOI: S0025-6196(18)30314-8 [pii] 10.1016/j.mayocp.2018.04.015 [doi].
5. Horwich TB, Fonarow GC and Clark AL. Obesity and the obesity paradox in heart failure. *Progr Cardiovasc Dis* 2018; 61: 151-156. DOI: <https://doi.org/10.1016/j.pcad.2018.05.005>.
6. Clark AL, Fonarow GC and Horwich TB. Impact of Cardiorespiratory Fitness on the Obesity Paradox in Patients With Systolic Heart Failure. *Am J Cardiol* 2015; 115: 209-213. DOI: 10.1016/j.amjcard.2014.10.023.
7. Lavie CJ, Cahalin LP, Chase P, et al. Impact of Cardiorespiratory Fitness on the Obesity Paradox in Patients With Heart Failure. *Mayo Clin Proc* 2013; 88: 251-258. DOI: 10.1016/j.mayocp.2012.11.020.

8. Piepoli MF, Corra U, Veglia F, et al. Exercise tolerance can explain the obesity paradox in patients with systolic heart failure: data from the MECKI Score Research Group. *Eur J Heart Fail* 2016; 18: 545-553. 2016/05/03. DOI: 10.1002/ejhf.534 [doi].
9. McAuley PA, Keteyian SJ, Brawner CA, et al. Exercise Capacity and the Obesity Paradox in Heart Failure: The FIT (Henry Ford Exercise Testing) Project. *Mayo Clin Proc* 2018; 93: 701-708. 2018/05/08. DOI: S0025-6196(18)30115-0 [pii] 10.1016/j.mayocp.2018.01.026 [doi].
10. Lund LH, Khush KK, Cherikh WS, et al. The Registry of the International Society for Heart and Lung Transplantation: Thirty-fourth Adult Heart Transplantation Report—2017; Focus Theme: Allograft ischemic time. *J Heart Lung Transpl* 2017; 36: 1037-1046. DOI: 10.1016/j.healun.2017.07.019.

Table 1. Characteristics of individuals with heart failure (HF) at the time of testing stratified based on $\dot{V} O_{2peak}$ ($ml \cdot kg^{-1} \cdot min^{-1}$) and BMI.

	Normal BMI, higher fitness (n=67)	Normal BMI, lower fitness (n=30)	BMI, High higher fitness (n=146)	BMI, High lower fitness (n=69)	BMI, $P < 0.05^*$ $P < 0.01^{**}$
Male, n (%)	50 (68)	19 (63)	129 (89)	58(84)	NF vs HF*;NL vs HF*; NL vs HL*
Age, yrs (SD)	47 (15)	56 (15)	56 (10)	51(13)	NS
BMI, $kg \cdot m^{-2}$ (SD)	22.4 (1.7)	22.0 (1.6)	30.3 (3.6)	29.6 (3.3)	NF vs HF**; HL vs NL**
$\dot{V} O_{2peak}$, $ml \cdot kg^{-1} \cdot min^{-1}$ (SD)	21.1 (5.9)	11.1 (2.0)	20.4 (5.1)	11.3 (2.0)	NF vs NL**; HF vs HU**
$\dot{V} O_{2peak}$ % age-pred. (SD)	60 (15.0)	37 (8.2)	60 (12.9)	36 (6.8)	NF vs NL** HF vs HL**
$\dot{V} E / \dot{V} C O_2$ slope	34.4 (7.9)	48.9 (14.2)	30.9 (5.8)	39.9 (9.11)	NF vs NL**; HF vs HL**
Peak HR bpm (SD)	132 (36)	108 (26)	138 (33)	106 (33)	NF vs NL**; HF vs HL**
Peak SBP mmHg (SD)	121 (32)	111 (14)	142 (20)	114 (26)	NF vs NL**; HF vs HL**
Peak RER (SD)	1.11(0.07)	1.12 (0.07)	1.12 (0.07)	1.11(0.09)	NS
LVEF % (SD)	27 (12)	27 (16)	25 (10)	29 (12)	NS

NYHA class, (<i>SD</i>)	2.7 (0.84)	3.2 (0.58)	2.5 (0.84)	3.1 (0.79)	NL vs NF*; HL vs HF**; NF vs HF*; NL vs HF*;
Ischaemic aetiology, %	25	28	30	54	HL vs HF; NF, NL**
Hx hypertension, %	10	17	19	46	HL vs HF, NF, NL**; HF vs NF*; HL vs NL*
Hx type 2 diabetes, %	3	14	16	33	HL vs HF, NF, NL**
Hx hypercholesterolemia, %	9	7	17	30	HL vs HF, NF, NL**

NF, normal BMI, higher fitness; NL, normal BMI lower fitness; HF, high BMI, higher fitness; HL, high BMI, lower Fitness; SD, standard deviation; NS, not significant; BMI, body mass index; $\dot{V} O_2$ peak, peak oxygen consumption; $\dot{V} E/\dot{V} C O_2$ slope, ratio ventilation:carbon dioxide output; HR, heart rate; bpm, beats per minute; SBP, systolic blood pressure, RER, respiratory exchange ratio; DBP, diastolic blood pressure; LVEF, left ventricular ejection fraction; NYHA, New York Heart Association; Hx, history

Figure 1. (A) The effect of fitness, expressed as $\dot{V} O_{2peak} \geq$ or $< 14 \text{ ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ in patients who had high or low BMI, on mortality over 5 years. Higher fitness groups had lower mortality vs lower fitness groups, regardless of BMI, at both 2 and 5 years follow-up ($P<0.01$). The lower fitness group with high BMI had lower mortality at 2 ($P=0.008$) and 5 years ($P=0.037$) than the lower fitness with low BMI group.

(B) The effect fitness, expressed as $\dot{V} O_{2peak} \geq$ or $< 50\%$ age-predicted in patients who had high or low BMI, on mortality over 5 years. Higher fitness groups had lower mortality than lower fitness groups, regardless of BMI at 2 and 5 years ($P<0.01$). The lower fitness group with high BMI had lower mortality at 2 ($P=0.008$) but not 5 years ($P=0.177$) than the lower fitness with low BMI group.


(C) The effect of ventilatory efficiency, expressed relative to $\dot{V}_E/\dot{V} CO_2 \geq$ or < 35 , in patients who had high or low BMI on all events over 5 years. Low slope subjects had lower mortality than a high slope subjects, regardless of whether they had normal BMI ($P<0.001$) or high BMI ($P<0.05$), over both 2 and 5 years.

ISHLT and RPH AHFCTS post-transplant survival statistics are superimposed.¹⁰


BMI, Body mass index; ISHLT, International Society for Heart & Lung Transplantation; RPH AHFCTS, Royal Perth Hospital Advanced Heart Failure and Cardiac Transplant Service; $\dot{V}_E/\dot{V} C O_2$ slope, ratio ventilation:carbon dioxide output.

Figure 1

A.



B.



C.

