

# Clinical Physiology and Functional Imaging

Published for the Scandinavian Society of Clinical Physiology and Nuclear Medicine

## Effects of post-exercise cooling on heart rate recovery in normotensive and hypertensive men

Journal:	<i>Clinical Physiology and Functional Imaging</i>
Manuscript ID	CPF-2019-0073.R1
Manuscript Type:	Original Article
Date Submitted by the Author:	01-Oct-2019
Complete List of Authors:	<p>Peçanha, Tiago; University of Sao Paulo, Exercise Hemodynamic Laboratory, School of Physical Education and Sport</p> <p>Low, David; Liverpool John Moores University Faculty of Arts Professional and Social Studies, Sport and Exercise Sciences</p> <p>Brito, Leandro; University of Sao Paulo, Exercise Hemodynamic Laboratory, School of Physical Education and Sport</p> <p>Fecchio, Rafael; University of São Paulo, School of Physical Education and Sport</p> <p>Sousa, Patrícia; University of Sao Paulo, Exercise Hemodynamic Laboratory, School of Physical Education and Sport</p> <p>Silva-Junior, Natan; University of Sao Paulo, Exercise Hemodynamic Laboratory, School of Physical Education and Sport</p> <p>Abreu, Andrea; University of Sao Paulo, General Hospital, Faculty of Medicine</p> <p>Silva, Giovanio; University of Sao Paulo, General Hospital, Faculty of Medicine</p> <p>Mion-Junior, Décio; University of Sao Paulo, General Hospital, Faculty of Medicine</p> <p>Forjaz, Cláudia; University of Sao Paulo, Exercise Hemodynamic Laboratory, School of Physical Education and Sport</p>
Key Words:	heart rate variability, thermoregulation, autonomic nervous system, parasympathetic, hypertension, blood pressure

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**1 Effects of post-exercise cooling on heart rate recovery in normotensive and**  
**2 hypertensive men**

3 Tiago Peçanha<sup>1</sup>, David Low<sup>2</sup>, Leandro Campos de Brito<sup>1</sup>, Rafael Yokoyama Fecchio<sup>1</sup>,  
4 Patrícia Nascimento de Sousa<sup>1</sup>, Natan Daniel da Silva-Júnior<sup>1</sup>, Andrea Pio de Abreu<sup>3</sup>,  
5 Giovanio Vieira da Silva<sup>3</sup>, Décio Mion-Junior<sup>3</sup>, Cláudia Lúcia de Moraes Forjaz<sup>1</sup>.

6  
7 **Authors' affiliations:**

- 8 1- Exercise Hemodynamic Laboratory, School of Physical Education and Sport,  
9 University of São Paulo, São Paulo, Brazil.  
10 2 - Research Institute for Sport and Exercise Sciences, Liverpool John Moores  
11 University, Liverpool, UK.  
12 3 - Hypertension Unit, General Hospital, University of São Paulo, São Paulo, Brazil.

13  
14 **Establishment where the work was performed:** Exercise Hemodynamic Laboratory,  
15 School of Physical Education and Sport, University of São Paulo, São Paulo, Brazil.  
16 Av. Prof. Mello Moraes, 65 , São Paulo, Brazil. ZIP Code: 05508-030. Phone: +55+11  
17 30918792. E-mail: [cforjaz@usp.br](mailto:cforjaz@usp.br)

18  
19 **Short title: Post-exercise cooling in hypertensives**

20 **Word count: 3533**

21 **Number of display items: 5**

22  
23 **Corresponding author:** Tiago Peçanha. Exercise Hemodynamic Laboratory, School  
24 of Physical Education and Sport, University of São Paulo. Av. Prof. Mello Moraes, 65 ,  
25 São Paulo, Brazil. ZIP Code: 05508-030. Phone: +55+11 30918792. E-mail:  
26 [pecanhatiago@gmail.com](mailto:pecanhatiago@gmail.com).

## SUMMARY

**Background:** Post-exercise heart rate recovery (HRR) is determined by cardiac autonomic restoration after exercise and is reduced in hypertension. Post-exercise cooling accelerates HRR in healthy subjects, but its effects in a population with cardiac autonomic dysfunction, such as hypertensives (HT), may be blunted. This study assessed and compared the effects of post-exercise cooling on HRR and cardiac autonomic regulation in HT and normotensive (NT) subjects. **Methods:** Twenty-three never-treated HT ( $43 \pm 8$  ys) and 25 NT ( $45 \pm 8$  ys) men randomly underwent two exercise sessions (30 min of cycling at  $70\% \text{VO}_{2\text{peak}}$ ) followed by 15 min of recovery. In one randomly allocated session, a fan was turned on in front of the subject during the recovery (cooling), while in the other session, no cooling was performed (control). HRR was assessed by heart rate reductions after 60 (HRR60s) and 300s (HRR300s) of recovery, short-term time constant of HRR (T30), and the time constant of the HRR after exponential fitting (HRR $\tau$ ). HRV was assessed using time- and frequency-domain indices. **Results:** HRR and HRV responses in the cooling and control sessions were similar between the HT and NT. Thus, in both groups, post-exercise cooling equally accelerated HRR (HRR300s =  $39 \pm 12$  vs.  $36 \pm 10$  bpm,  $p \leq 0.05$ ) and increased post-exercise HRV (lnRMSSD =  $1.8 \pm 0.7$  vs.  $1.6 \pm 0.7$  ms,  $p \leq 0.05$ ). **Conclusion:** Differently from the hypothesis, post-exercise cooling produced similar improvements in HRR in HT and NT men, likely by an acceleration of cardiac parasympathetic reactivation and sympathetic withdrawal. These results suggest that post-exercise cooling equally accelerates HRR in hypertensive and normotensive subjects.

49

Keywords: heart rate variability, thermoregulation, autonomic nervous system, parasympathetic, hypertension, blood pressure

INTRODUCTION

Post-exercise heart rate recovery (HRR) is a non-invasive tool to assess cardiac autonomic function (Cole, *et al.* 1999; Imai, *et al.* 1994). HRR presents a biphasic behavior, with an initial fast decay determined by parasympathetic reactivation (Imai, *et al.* 1994) and a second slow decay that is determined by sympathetic withdrawal and further parasympathetic reactivation (Perini, *et al.* 1989).

A reduced HRR reflects autonomic dysfunction, and has been observed in several chronic diseases (Peçanha, *et al.* 2014) and is associated with poor prognosis (Cole, *et al.* 1999). Several strategies have been used to improve cardiac autonomic restoration after exercise and, thus, to accelerate HRR (Al Haddad, *et al.* 2010; de Oliveira Ottone, *et al.* 2014; Leicht, *et al.* 2009). Of note, Leicht *et al.* (2009) observed a faster HRR when subjects were exposed to a fan with water spray or ice pack application during the post-exercise period. Likewise, Al Haddad *et al.* (2010) observed similar effects using cold water immersion. These strategies are based on the recognized relationship between the autonomic control of HRR and thermoregulation (e.g., heat stress-induced impairments in HRR) (Peçanha, *et al.* 2017b). It should be noted, however, that these cooling studies were conducted with healthy subjects, and little is known regarding the effects of such strategies on HRR and its regulation in subjects with cardiovascular disorders usually associated with autonomic dysfunction and reduced HRR.

Hypertension is a highly prevalent cardiovascular disease characterized by the presence of autonomic dysfunction, i.e., a decrease in parasympathetic and an increase in sympathetic nerve activities (Mancia & Grassi 2014). This autonomic imbalance culminates with reduced HRR and increased cardiovascular risk in hypertensive (HT) when compared to normotensive (NT) subjects (Erdogan, *et al.* 2011). In light of the benefits of post-exercise cooling on HRR in healthy subjects, one could hypothesize that this strategy could be as effective in accelerating HRR in HT.

However, previous evidence suggests negative cardiovascular responses to cooling in hypertension. Greaney et al. (Greaney, *et al.* 2017) reported exaggerated increases in sympathetic nerve activity during whole body cooling in HT, and this response may mitigate the potential beneficial effect of post-exercise cooling on HRR in HT, which to the best of our knowledge has not been investigated yet.

Thus, this study was designed to assess the effects of post-exercise cooling with a fan on HRR in never-treated HT and normotensive (NT) men. The hypothesis was that the acceleration of HRR promoted by cooling would be blunted in HT compared with NT. To better clarify the mechanisms underlying the responses of HRR to post-exercise cooling in NT and HT, the effects of such a strategy on post-exercise heart rate variability (HRV) and blood pressure (BP) were also evaluated.

## MATERIAL AND METHODS

### Subjects

The participants were middle-aged (30–60 years) and physically inactive HT (systolic/diastolic BP between 140/90 and 159/99 mmHg) and NT (i.e. systolic/diastolic BP <120/80 mmHg) men. The exclusion criteria included smoking; established cardiovascular diseases; body mass index  $\geq 35$  kg/m<sup>2</sup>; use of medications that could directly affect cardiovascular responses to exercise; and abnormal resting or exercise electrocardiogram. Additionally, HT subjects had never been treated with antihypertensive drugs and had no target organ damage or secondary hypertension. After a detailed explanation of the experimental procedures, subjects provided their written informed consent. This study was conducted in accordance with the Declaration of Helsinki and was approved by local Institutional research ethics committee (281.905/2013).

### Preliminary assessment

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

105 Health status was investigated through a detailed anamnesis. Resting BP was  
106 measured by a mercury sphygmomanometer (Uniteq, São Paulo, Brazil), three times  
107 after 5-min seated rest in two distinct visits to the laboratory (Chobanian, *et al.* 2003).  
108 All HT subjects also underwent the routine screening of the Hypertension Unit of the  
109 General Hospital of the University of São Paulo to detect target organ damage,  
110 secondary hypertension, and/or other clinical conditions that preclude exercise  
111 participation.

112 On a separate day, subjects performed a maximal cardiopulmonary exercise  
113 test with assessments of resting and exercise electrocardiograms. This test was  
114 performed on a cycle ergometer (Computrainer, RacerMate, Seattle, USA) with an  
115 initial workload of 50 Watts and increments of 20 Watts every 3 min until volitional  
116 exhaustion. Immediately after exercise, workload was reduced to 50 Watts, and the  
117 subjects completed a 5 min recovery. During the test, ventilatory variables were  
118 continuously measured using a metabolic cart (CPX-Ultima, Medical Graphics  
119 Corporation, Minnesota, USA). Peak oxygen consumption ( $VO_{2peak}$ ) and HR ( $HR_{peak}$ )  
120 were determined by the maximal values attained at the end of exercise (average of  
121 30s). HRR was assessed through the calculation of the HRR60s index (i.e.  $HR_{peak} -$   
122 HR at 60s of recovery) (Peçanha, et al. 2014).

123 Experimental protocol

124 All subjects randomly underwent two experimental sessions conducted in the  
125 morning of two separate days, with a minimum interval of 48h between sessions.  
126 Temperature and humidity of the room were kept constant across the sessions (20–  
127 22°C and 75–80%). Subjects were instructed to arrive at the laboratory in a fasted  
128 state, and to avoid alcohol and exercise for 24h and caffeine ingestion for 12h prior to  
129 the sessions. In each session, they received a standardized meal (two 25g cereal bars  
130 and 50ml of juice) and experiment began 30min after the ingestion of the meal. The  
131 experimental sessions started with resting measurements in the seated position for

10min and, then the subjects were submitted to 30min of cycle ergometer exercise (CompuTrainer, RacerMate, Seattle, USA) at 70% $\text{VO}_{2\text{peak}}$ . After exercise, subjects immediately stopped pedaling and remained seated on the cycle ergometer for a 15min recovery. In one of the sessions, an industrial fan was turned on in front (~1 meter) of the subjects during all of the recovery period (cooling session), while in the other session, the recovery was performed without fanning (control session).

#### Experimental Measurements

$T_c$  was measured in 10-s intervals via a telemetric temperature pill system (CorTemp®, HQInc., Palmetto, USA) ingested, at least, 2h before the experiments (Byrne & Lim 2007). HR was continuously measured using a three-lead electrocardiogram (EMG System, São Paulo, Brazil) and beat-by-beat BP was obtained using finger photoplethysmography (Finometer, FMS, Arnhem, The Netherlands). These signals were continuously acquired online (Windaq, Dataq Instruments, Ohio, USA; 500 Hz/channel). Mean  $T_c$ , HR and SBP were calculated for rest (5–10 min), exercise (25–30min), and immediate (0–5min) and late (10–15min) recovery periods. In addition,  $\text{VO}_2$  was continuously measured during exercise with a metabolic cart (CPX-Ultima, Medical Graphics Corporation, Minnesota, USA).

#### Cardiovascular Autonomic Assessment

Preprocessing procedures. HR signals were exported to HeartScope software (A.M.P.S. LLC, New York, USA) for the generation of RR intervals (RRi) time series. These series were visually inspected and occasional misdetections were corrected. Likewise, ectopic beats were identified and replaced with interpolated RRi values (<2% of the total signal).

Heart Rate Recovery Analysis. HRR was assessed as previously reported (Peçanha, *et al.* 2016). The following indices were calculated: a) HRR60s and HRR300s, i.e. the absolute differences between peak HR (mean of the last 60s of exercise) and the HRs obtained, respectively, at 60 and 300s of recovery; b) T30, i.e. the short-term time

constant of HRR obtained from the negative reciprocal of the linear regression line between the log-transformed HR and the first 30 s of recovery (Imai et al., 1994) and; c) HRRt, i.e. the time-constant of HRR after exponential fitting of the HR during the entire 300s of recovery.

Heart Rate Variability. HRV was assessed at rest, during immediate recovery (0–5min) and during late recovery (10–15min) after exercise. For the immediate recovery period, given the non-stationary behavior of RRI, the assessment of HRV was performed using a time-varying approach (Goldberger, *et al.* 2006). Firstly, the RRI time series of recovery was filtered using a median filter operation. Then, HRV was assessed through the calculation of RMSSD (root mean square of successive differences in RRI) and RMS (root mean square residual of RRI) indices, on successive non-overlapped 30s segments, during the entire 5min immediate recovery.

At rest and late recovery, given the relative stationarity of the cardiovascular signals, HRV was assessed via spectral analysis using the Heart Scope software (A.M.P.S. LLC, New York, USA) and following international Task Force recommendations(1996). The power spectral density analyses of RRI (250-300 beats) were performed using the autoregressive method and the spectral components were calculated via the Levinson-Durbin recursion employing Akaike's criteria for choosing the order of the model (Malliani, *et al.* 1991). Low- (LF: 0.04–0.15Hz) and high-frequency (HF: 0.15–0.4Hz) components of RRI variability were expressed in normalized units (Task-Force 1996).

#### Statistical Analysis

Based on an expected Cohen's d effect size of 0.87 of post-exercise cooling on HRR60s (Al Haddad, et al. 2010), the sample size required to achieve a power of 90% and an  $\alpha$  level of 5% was 32 subjects (i.e. 16 subjects per group) (G\*Power v. 3.1.9.2, Universität Kiel, Germany).

## RESULTS

Responses to exercise in the experimental sessions were also similar between groups. There were no differences in HR or  $\text{VO}_2$  between groups and sessions. In both groups and sessions, exercise intensity corresponded to approximately 70% of  $\text{VO}_{2\text{peak}}$  (Table 2).

## Core Temperature and Hemodynamics

Figure 1 displays  $T_c$ , HR and SBP responses throughout the experimental protocol. There were no differences between the groups in  $T_c$  and HR at any time point,



>>>>>>>>>>>>>>>>>>FIGURE 3>>>>>>>>>>>>>>>>>>

[illegible]

To the best of our knowledge, this is the first study to assess the effects of post-exercise cooling on HRR and autonomic regulation in a population with autonomic dysfunction and increased cardiovascular risk, such as HT men (Mancia & Grassi

2014). The presence of autonomic dysfunction in the HT group was shown by their increased resting LF- and decreased HF-HRV as well as by the reduced HRR60s after the maximal exercise test in comparison with NT consistent with other reports (Aneni, *et al.* 2014; Best, *et al.* 2014; Erdogan, *et al.* 2011; Mancina & Grassi 2014; Pagani & Lucini 2001).

Based on recent evidence showing exaggerated sympathetic responses to cooling in hypertension (Greaney, *et al.* 2017), the hypothesis of the present study was that the previously reported benefits of cooling accelerating HRR would be blunted in HT compared with NT. On the contrary, the present results showed that cooling induced similar changes in HRR in both groups, suggesting that the presumed cooling-induced sympathetic overactivity in HT did not occur. This discrepancy may be due to the fact that in the present study cooling was used after exercise, i.e., a condition in which body temperature was elevated, while in the previous study cooling was employed at rest, when body temperature is normal. Thus, the present results suggest that HT do not have an exaggerated sympathetic response to cooling when this strategy is employed after exercise.

Based on HRR and HRV indices used in the present study, it is possible to suggest the autonomic mechanisms behind the improved HRR observed in both groups in the cooling session. T30, RMSSD and HF are indices accepted as mainly dependent on parasympathetic reactivation (Goldberger, *et al.* 2006; Hayano, *et al.* 1991; Imai, *et al.* 1994), while, despite some criticism (Goldstein, *et al.* 2011), HRR300s, RMS and LF indices are considered, at least in part, markers of sympathetic modulation (Malliani, *et al.* 1991; Ng, *et al.* 2009; Peçanha, *et al.* 2017a). Thus, the lower T30 and LF observed in the cooling session together with the higher RMSSD, HF, HRR300s and RMS suggest that the cooling strategy likely accelerated both cardiac parasympathetic reactivation and sympathetic withdrawal after exercise,

292 supporting both mechanisms as responsible for the cooling-induced faster HRR in both  
293 NT and HT subjects.

294 Besides accelerating HRR, the cooling protocol also increased SBP during  
295 recovery. In fact, cooling prevented the decrease in SBP below pre-exercise levels  
296 (i.e., post-exercise hypotension) that was observed at 15 min of recovery in the control  
297 session. The mechanisms behind this divergent BP response to the sessions are  
298 beyond the scope of this study, however, the existing literature might help to elucidate  
299 them. In the control session recovery, the thermoregulatory-induced skin vasodilatation  
300 alongside the interruption of the muscle pump might have reduced vascular resistance  
301 and stroke volume, promoting a decrease in BP (Carter, *et al.* 1999; Peçanha, *et al.*  
302 2017b). So, the increased HR (i.e. reduced HRR) observed in the control session might  
303 partly reflect the baroreflex's attempt to counteract a BP decrease (Peçanha, *et al.*  
304 2017b). On the other hand, in the cooling session, despite the interruption of the  
305 muscle pump, skin vasodilatation was reduced by the effect of the fan facilitating heat  
306 loss via convection (Barwood, *et al.* 2009), which might have limited the decrease in  
307 vascular resistance, keeping central blood volume and BP higher than in the control  
308 session. As a response, the baroreflex might have produced a greater reactivation of  
309 cardiac parasympathetic and withdrawal of cardiac sympathetic modulations,  
310 accelerating HRR. Therefore, it seems reasonable to suggest that the effect of post-  
311 exercise cooling on HRR and autonomic regulation probably involves its effects on BP.

312 Results of the present study may have relevant clinical impact. As HT subjects  
313 present impaired autonomic responses after exercise (Greaney, *et al.* 2014; Peçanha,  
314 *et al.* 2016), any strategy that could ameliorate this response has clinical importance,  
315 promoting a likely preventive effect on acute cardiac events (Cole, *et al.* 1999; Nishime,  
316 *et al.* 2000). In this sense, the present results showed the clinical applicability of post-  
317 exercise cooling with a fan in a population with recognized autonomic dysfunction and  
318 increased cardiovascular risks (Mancia & Grassi 2014). Other studies with healthy

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

319 subjects have also reported positive effects of different strategies for post-exercise  
320 cooling, such as ice packs (Leicht, et al. 2009) or cold water immersion (Al Haddad, et  
321 al. 2010). However, the use of a fan, may be more practical and safe, especially in  
322 subjects with cardiovascular disease and/or risk factors, since cold water immersion  
323 and ice use can trigger cardiac arrhythmia (Shattock & Tipton 2012) and considerably  
324 increase vasoconstriction and BP (Mawhinney, et al. 2017; Sramek, et al. 2000).

325         This study has some limitations. Firstly, it only involved men, and as  
326 thermoregulatory responses might differ between genders (Kaciuba-Uscilko & Grucza  
327 2001), future investigations should be conducted with women. Other limitations include  
328 the absence of hydration control prior to the sessions. However, subjects in the present  
329 study presented similar pre-exercise body masses in both sessions (NT=87±11 vs.  
330 86±11 kg and HT=91±15 vs. 92±16 kg for the cooling and the control sessions,  
331 respectively), which suggests no difference in hydration status. Finally, the results of  
332 the present study are limited to the first 15 min of the post-exercise period. Assessment  
333 of cardiac autonomic variables for a longer period of recovery might help to improve the  
334 comprehension of the benefits of cooling on cardiac autonomic restoration.

335  
336 **CONCLUSION**

337         In conclusion, post-exercise cooling with a fan equally reduced core  
338 temperature and improved HRR in never-treated HT and NT men. The effects of post-  
339 exercise cooling accelerating HRR appeared to be promoted by accelerated post-  
340 exercise cardiac parasympathetic reactivation and sympathetic withdrawal, which might  
341 be partially related to the increased SBP with cooling. The positive effects of post-  
342 exercise cooling on HRR and post-exercise HRV in HT subjects highlight the clinical  
343 applicability of such a strategy for populations with increased cardiovascular risk and  
344 cardiac autonomic dysfunction.

345

**Acknowledgements**

The authors thank Rhenan Bartels for his assistance with the mathematical procedures, and all of the volunteers for their willingness to participate in this study.

This study was supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo under Grants 2013/04997-0, 2013/05519-4 and 2015/15466-0; Conselho Nacional de Desenvolvimento Científico e Tecnológico under Grant 304003-2014-0; and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES-PROEX) – Finance Code 001.

**Conflict of interest**

The authors declare no conflicts of interest.

REFERENCES

Al Haddad H, Laursen PB, Chollet D, Lemaitre F, Ahmaidi S and Buchheit M. Effect of cold or thermoneutral water immersion on post-exercise heart rate recovery and heart rate variability indices. *Auton Neurosci* (2010); **156**: 111-116.

Aneni E, Roberson LL, Shaharyar S, Blaha MJ, Agatston AA, Blumenthal RS, Meneghelo RS, Conceicao RD, Nasir K and Santos RD. Delayed heart rate recovery is strongly associated with early and late-stage prehypertension during exercise stress testing. *Am J Hypertens* (2014); **27**: 514-521.

Barwood MJ, Davey S, House JR and Tipton MJ. Post-exercise cooling techniques in hot, humid conditions. *Eur J Appl Physiol* (2009); **107**: 385-396.

Best SA, Bivens TB, Palmer MD, Boyd KN, Galbreath MM, Okada Y, Carrick-Ranson G, Fujimoto N, Shibata S and Hastings JL. Heart rate recovery after maximal exercise is blunted in hypertensive seniors. *Journal of Applied Physiology* (2014); **117**: 1302-1307.

Byrne C and Lim CL. The ingestible telemetric body core temperature sensor: a review of validity and exercise applications. *Br J Sports Med* (2007); **41**: 126-133.

Carter R, 3rd, Watenpaugh DE, Wasmund WL, Wasmund SL and Smith ML. Muscle pump and central command during recovery from exercise in humans. *J Appl Physiol* (1999); **87**: 1463-1469.

Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL, Jr., Jones DW, Materson BJ, Oparil S, Wright JT, Jr. and Roccella EJ. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. *JAMA* (2003); **289**: 2560-2572.

Cole CR, Blackstone EH, Pashkow FJ, Snader CE and Lauer MS. Heart-rate recovery immediately after exercise as a predictor of mortality. *N Engl J Med* (1999); **341**: 1351-1357.

- de Oliveira Ottone V, de Castro Magalhães F, de Paula F, Avelar NCP, Aguiar PF, da  
Matta Sampaio PF, Duarte TC, Costa KB, Araújo TL, Coimbra CC, Nakamura  
FY, Amorim FT and Rocha-Vieira E. The Effect of Different Water Immersion  
Temperatures on Post-Exercise Parasympathetic Reactivation. *PLoS One*  
(2014); **9**: e113730.
- Erdogan D, Gonul E, Icli A, Yucel H, Arslan A, Akcay S and Ozaydin M. Effects of  
normal blood pressure, prehypertension, and hypertension on autonomic  
nervous system function. *Int J Cardiol* (2011); **151**: 50-53.
- Goldberger JJ, Le FK, Lahiri M, Kannankeril PJ, Ng J and Kadish AH. Assessment of  
parasympathetic reactivation after exercise. *Am J Physiol Heart Circ Physiol*  
(2006); **290**: H2446-2452.
- Goldstein DS, Benth O, Park MY and Sharabi Y. Low-frequency power of heart rate  
variability is not a measure of cardiac sympathetic tone but may be a measure  
of modulation of cardiac autonomic outflows by baroreflexes. *Exp Physiol*  
(2011); **96**: 1255-1261.
- Greaney JL, Kenney WL and Alexander LM. Sympathetic function during whole body  
cooling is altered in hypertensive adults. *Journal of Applied Physiology* (2017);  
**123**: 1617-1624.
- Greaney JL, Matthews EL, Boggs ME, Edwards DG, Duncan RL and Farquhar WB.  
Exaggerated exercise pressor reflex in adults with moderately elevated systolic  
blood pressure: role of purinergic receptors. *Am J Physiol Heart Circ Physiol*  
(2014); **306**: H132-141.
- Hayano J, Sakakibara Y, Yamada A, Yamada M, Mukai S, Fujinami T, Yokoyama K,  
Watanabe Y and Takata K. Accuracy of assessment of cardiac vagal tone by  
heart rate variability in normal subjects. *Am J Cardiol* (1991); **67**: 199-204.
- Imai K, Sato H, Hori M, Kusuoka H, Ozaki H, Yokoyama H, Takeda H, Inoue M and  
Kamada T. Vagally mediated heart rate recovery after exercise is accelerated in

- athletes but blunted in patients with chronic heart failure. *J Am Coll Cardiol* (1994); **24**: 1529-1535.
- Kaciuba-Uscilko H and Grucza R. Gender differences in thermoregulation. *Current Opinion in Clinical Nutrition & Metabolic Care* (2001); **4**: 533-536.
- Leicht AS, Sinclair WH, Patterson MJ, Rudzki S, Tulppo MP, Fogarty AL and Winter S. Influence of postexercise cooling techniques on heart rate variability in men. *Exp Physiol* (2009); **94**: 695-703.
- Malliani A, Pagani M, Lombardi F and Cerutti S. Cardiovascular neural regulation explored in the frequency domain. *Circulation* (1991); **84**: 482-492.
- Mancia G and Grassi G. The autonomic nervous system and hypertension. *Circ Res* (2014); **114**: 1804-1814.
- Mawhinney C, Jones H, Low DA, Green DJ, Howatson G and Gregson W. Influence of cold-water immersion on limb blood flow after resistance exercise. *Eur J Sport Sci* (2017); **17**: 519-529.
- Ng J, Sundaram S, Kadish AH and Goldberger JJ. Autonomic effects on the spectral analysis of heart rate variability after exercise. *Am J Physiol Heart Circ Physiol* (2009); **297**: H1421-1428.
- Nishime EO, Cole CR, Blackstone EH, Pashkow FJ and Lauer MS. Heart rate recovery and treadmill exercise score as predictors of mortality in patients referred for exercise ECG. *JAMA* (2000); **284**: 1392-1398.
- Pagani M and Lucini D. Autonomic dysregulation in essential hypertension: insight from heart rate and arterial pressure variability. *Auton Neurosci* (2001); **90**: 76-82.
- Peçanha T, Bartels R, Brito LC, Paula-Ribeiro M, Oliveira RS and Goldberger JJ. Methods of assessment of the post-exercise cardiac autonomic recovery: A methodological review. *International Journal of Cardiology* (2017a); **227**: 795-802.

- 1  
2  
3  
4 437 Peçanha T, de Brito LC, Fecchio RY, de Sousa PN, da Silva Junior ND, de Abreu AP,  
5  
6 438 da Silva GV, Mion-Junior D and Forjaz CLdM. Metaboreflex activation delays  
7  
8 439 heart rate recovery after aerobic exercise in never-treated hypertensive men. *J*  
9  
10 440 *Physiol* (2016); **594**: 6211-6223.  
11  
12 441 Peçanha T, Forjaz CLdM and Low DA. Passive Heating Attenuates Post-exercise  
13  
14 442 Cardiac Autonomic Recovery in Healthy Young Males. *Frontiers in*  
15  
16 443 *Neuroscience* (2017b); **11**.  
17  
18 444 Peçanha T, Silva-Junior ND and Forjaz CL. Heart rate recovery: autonomic  
19  
20 445 determinants, methods of assessment and association with mortality and  
21  
22 446 cardiovascular diseases. *Clin Physiol Funct Imaging* (2014); **34**: 327-339.  
23  
24 447 Perini R, Orizio C, Comande A, Castellano M, Beschi M and Veicsteinas A. Plasma  
25  
26 448 norepinephrine and heart rate dynamics during recovery from submaximal  
27  
28 449 exercise in man. *Eur J Appl Physiol Occup Physiol* (1989); **58**: 879-883.  
29  
30 450 Shattock MJ and Tipton MJ. 'Autonomic conflict': a different way to die during cold  
31  
32 451 water immersion? *J Physiol* (2012); **590**: 3219-3230.  
33  
34 452 Sramek P, Simeckova M, Jansky L, Savlikova J and Vybiral S. Human physiological  
35  
36 453 responses to immersion into water of different temperatures. *Eur J Appl Physiol*  
37  
38 454 (2000); **81**: 436-442.  
39  
40 455 Task-Force. Heart rate variability. Standards of measurement, physiological  
41  
42 456 interpretation, and clinical use. Task Force of the European Society of  
43  
44 457 Cardiology and the North American Society of Pacing and Electrophysiology.  
45  
46 458 *Eur Heart J* (1996); **17**: 354-381.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

TABLES

Table 1. Baseline data

	NT	HT	P
	(n = 25)	(n = 23)	
Age (years)	45 ± 8	43 ± 8	0.29
Body mass index (kg.m <sup>-2</sup> )	28.6±2.7	29.4±3.6	0.37
SBP (mmHg)	115±4	142±9	< 0.01
DBP (mmHg)	77±2	96±3	< 0.01
VO <sub>2peak</sub> (ml.kg <sup>-1</sup> .min <sup>-1</sup> )	26.6±4.2	25.1±4.4	0.24
HR <sub>peak</sub> (bpm)	169±11	169±16	0.93

Values are presented as mean ± SD. SBP = systolic blood pressure. DBP = diastolic blood pressure. VO<sub>2peak</sub> = peak oxygen uptake achieved during the maximal exercise test. HR<sub>peak</sub> = peak heart rate achieved during the maximal exercise test.

Table 2. Physiological variables measured during experimental sessions in normotensive (NT) and hypertensive (HT) groups.

	NT		HT		P	P	P
	(n = 25)		(n = 23)		(session)	(group)	(session*group)
	Control	Cooling	Control	Cooling			
HR (bpm)	128±13	129±12	130±13	129±13	0.97	0.73	0.73
HR (%peak)	76±6	76±5	77±7	77±7	0.99	0.65	0.70
VO <sub>2</sub> (ml.kg <sup>-1</sup> .min <sup>-1</sup> )	17.3±2.4	17.7±2.6	16.4±3.1	16.0±3.1	0.63	0.12	0.07
VO <sub>2</sub> (% peak)	71±4	72±6	69±5	68±5	0.95	0.07	0.30

Values are presented as mean ± SD. VO2 = oxygen uptake. HR = heart rate.

## FIGURE LEGENDS

**Fig. 1** Core temperature ( $T_c$ ; 1a), heart rate (HR; 1b) and systolic blood pressure (SBP; 1c) measured at rest, during exercise and at 5 and 15 min of recovery in the cooling and control sessions in the normotensive (N;  $n = 25$ ) and the hypertensive (HT;  $n = 23$ ) groups. \* =  $p \leq 0.05$  vs. rest. \$ =  $p \leq 0.05$  vs. exercise. & =  $p \leq 0.05$  vs. 5 min of recovery. ‡ =  $p \leq 0.05$  vs. control session. On panel c, groups were different in all time points (i.e., main effect of group). The connecting lines were removed to improve visualization.

**Fig. 2** Heart rate recovery indices assessed after the control and the cooling sessions in the normotensive (NT;  $n = 25$ ) and hypertensive (HT;  $n = 23$ ) groups. HRR60s – heart rate decrease at 60s of recovery (3a); T30 = short time-constant of heart rate recovery (3b); HRR300s – heart rate decrease at 300s of recovery (3c); HRRt = long time-constant of heart rate recovery (3d). ‡ =  $p \leq 0.05$  vs. control session

**Fig. 3** RMSSD (4a) and RMS (4b) indices of heart rate variability assessed in segments of 30 s in the first 5 min of recovery after the control and the cooling sessions in the normotensive (NT;  $n = 25$ ) and the hypertensive (HT;  $n = 23$ ) groups. RMSSD = square root of the mean of the sum of the squares of differences between adjacent normal RRI, RMS = root mean square of residual of RRI. ‡ =  $p \leq 0.05$  vs. control session. # =  $p \leq 0.05$  vs. 30s

**Fig. 4** Spectral indices of heart rate variability (HRV) assessed at rest and at the late recovery (Late Rec) of the control and the cooling sessions in the normotensive (NT;  $n = 25$ ) and the hypertensive (HT;  $n = 23$ ) groups. LF (nu) = low frequency component of HRV assessed in normalized units (5a); HF (nu) = high frequency component of HRV assessed in normalized units (5b). \* =  $p \leq 0.05$  vs. rest. † =  $p \leq 0.05$  vs NT. ‡ =  $p \leq 0.05$  vs. control session. The connecting lines were removed to improve visualization




Figure 1

130x184mm (300 x 300 DPI)




Figure 2

133x132mm (300 x 300 DPI)




Figure 3

172x167mm (300 x 300 DPI)




Figure 4

1761x431mm (96 x 96 DPI)