

Social and Mental Health Predictors of Emergency Department and General Practitioner Usage

Dr Pooja Saini, School of Natural Sciences and Psychology, Liverpool John Moores University

Dr Jason McIntyre, School of Natural Sciences and Psychology, Liverpool John Moores University

Prof Rhiannon Corcoran, Institute of Psychology, Health and Society, University of Liverpool

Konstantinos Daras, Public Health, University of Liverpool

Clarissa Giebel, NIHR CLAHRC NWC, University of Liverpool

Elizabeth Fuller, Public Advisor, NIHR CLAHRC NWC

Jane Shelton, Public Advisor, NIHR CLAHRC NWC

Timothy Wilson, Public Advisor, NIHR CLAHRC NWC

Terence Comerford, Public Advisor, NIHR CLAHRC NWC

Prof Rajan Nathan, Cheshire and Wirral NHS Foundation Trust

Prof Mark Gabbay, NIHR CLAHRC NWC, University of Liverpool

Correspondence: Dr Pooja Saini, School of Natural Sciences and Psychology, 2.28b James Parsons Tower, Liverpool John Moores University, Byrom Street, Liverpool, L3 3AF.
Email: P.Saini@ljmu.ac.uk

Abstract

Background: High demand for health services is an issue of current importance in England, in part, because of the rapidly increasing use of the Accident and Emergency (A&E) and GP practices for mental health issues and the high cost of these services.

Aim: To examine the social determinants of health service utilisation in people with mental health issues.

Design and Setting: Twenty-eight neighbourhoods (5000 – 10,000 people) on the North West Coast of England, which differed in relative levels of deprivation.

Method: A comprehensive public health survey was conducted, comprising questions on housing, physical health, mental health, lifestyle, social issues, environment, work and finances. Poisson regression models assessed the effect of mental health comorbidity, mental and physical health comorbidity, and individual symptoms on A&E and General Practice (GP) attendances, adjusting for relevant socioeconomic and lifestyle factors.

Results: People who had both a physical *and* mental health condition reported attending A&E (RR = 4.63, 95%CI 2.86 to 7.51) and GPs (RR = 3.82, 95%CI 3.15, 4.62) more frequently than all other groups. Having a higher number of mental health symptoms was associated with higher GP and A&E service usage. Depression was the strongest symptom predictor of A&E attendance (RR = 1.41, 95%CI 1.05 to 1.90), while anxiety was the strongest symptom predictor of GP attendance, RR = 1.19, 95%CI 1.03 to 1.38.

Conclusion: Mental health comorbidities increase risk of attendances to both A&E and GPs. Further research into the social attributes that contribute to reduced A&E and GP attendance rates is required.

Keywords: Mental health, Social care, Emergency Department, Primary Care, Service usage

1

2

3 **How this fits in**

4 • The current A&E ‘crisis’ in England is linked to health inequality as people living in more
5 deprived areas use NHS services much more.

6 • Mental health patients are at risk of repeated hospital admissions to hospital and increased
7 number of GP attendances.

8 • The National Institute of Health Research Collaboration for Leadership in Applied Health
9 Research and Care North West Coast (NIHR CLAHRC NWC) Household Health Survey
10 (HHS) provides information on the rate of usage of services within regions of the North
11 West Coast.

12 • Mental health comorbidities increase risk of attendances to both A&E and GPs. Depression
13 predicted higher A&E attendance and anxiety predicted higher GP attendance when
14 adjusting for physical health and socio-economic status.

15

16

17

18

Introduction

The UK has the second-highest levels of economic inequality in the European Union.¹ High demand for emergency admissions is an issue of pressing importance in England because of the rapidly increasing use of the Accident and Emergency (A&E) for mental health issues and the high cost of these services.² The current A&E ‘crisis’ in England is linked to health inequality as people living in more deprived areas use NHS services substantially more than people in less deprived areas.³⁻⁶ Patients from deprived areas also appear to attend A&E for less serious conditions^{4,6-7} and present at A&E nearly two and a half times more than those in the least deprived areas for preventable emergency hospitalisations.⁴ Furthermore, people with mental health issues, substance misuse and/or long-term health conditions are most likely to attend A&E⁵⁻⁶ and are at risk of repeated hospital admissions.⁷ Indeed, It has been reported that in 2017, the number of people going to A&E for mental ill-health had risen by nearly half since 2011-12.^{8,9}

In primary care, patients consult General Practitioners (GP) on average six times per year.¹⁰ More frequent attendance in the previous year has been noted for patients who died by suicide compared to patients who remained alive.¹¹ However, frequent attendance in general practice is associated with excessive service use.¹² Frequent attenders have been shown to have higher rates of common mental health difficulties¹³ including depression,¹⁴⁻¹⁵ anxiety,¹⁶ and somatic disorders.¹⁷⁻¹⁹ Additionally, higher consultation rates within primary care have been reported for those living in the most deprived quintile (21.5%) compared with the least deprived quintile (16.1%);²⁰ suggesting a link between deprivation and common mental health issues. Asaria and colleagues³ found that while increasing accessibility to GPs in deprived areas reduced socioeconomic inequalities in primary care access, it only resulted in modest reductions in A&E usage.

More knowledge is needed about the determinants of A&E and primary care service utilisation for mental health problems and how this compares across areas of differing deprivation. Moreover, it is important to consider lifestyle, socioeconomic, and accessibility factors when identifying determinants as these may underpin relationships between mental health and service usage. Due to the disparities that exist in the way people use health services, potential gains may be made by addressing health inequalities, particularly for people attending with mental health issues. To co-ordinate care effectively within a given area, up-to-date information about local healthcare utilization for mental health issues is required. In this study, we use data collected by The National Institute of Health Research Collaboration for Leadership in Applied Health Research and Care North West Coast (NIHR CLAHRC-NWC) Household Health Survey (HHS) to explore self-reported service usage within regions of the NWC and to examine patient characteristics that predict attendance at A&E departments and GP surgeries.

The specific aims of the study were to:

1. Assess the relationship between mental health, A&E and primary care attendance;
2. Quantify the extent to which comorbidities relate to service usage;
3. Identify individual mental health symptoms that predict service usage.

Method

Participants and design

The NIHR CLAHRC NWC HHS is comprehensive quantitative public health survey co-produced with public and patient advisors, Local Authorities, NHS clinicians, and university partners. Twenty-eight neighbourhoods were surveyed using random probability sampling across

the NWC of England, targeting 20 high deprivation neighbourhoods identified by Local Authority partners and eight less deprived areas to serve as comparison neighbourhoods. Each neighbourhood had a population of 5000-10,000 people and the majority of areas were defined by electoral ward boundaries. The survey assessed demographic, socioeconomic, physical health, mental health, and lifestyle factors. We selected a subset of these variables for the present analysis. In total, 4319 people were recruited via door-knocking. Research teams conducted face-to-face interviews and recorded responses on tablets. Interviews were conducted between August 2015 and January 2016. Interview teams worked at varying times of day to ensure the sample was representative. This resulted in 55% of interviews being conducted on weekends or after 4pm on weekdays, and 45% of interviews conducted before 4pm on weekdays. The adjusted response rate was 61%. The sample comprised 1854 (43%) men and 2465 (57%) women whose ages ranged from 18 to 95 years ($M = 49.12$, $SD = 19.13$). The majority of participants (89%) indicated that they were of White European ethnic background. Compared to census data for the North West of England, the sample was biased towards female (our sample: 57%, census: 51%) and BME (our sample: 11%, census: 8%) participants.²¹ Participants were reimbursed with a £10 voucher in return for their participation. A detailed description of the sampling procedure is available elsewhere.²²

Measures

Our outcome variable was defined as the number of times respondents reported attending an A&E department or their GP over the previous 12 months. Measures of socioeconomic conditions included education, employment, financial hardship, change in financial circumstances and housing quality. Physical health was assessed with the four physical health dimensions of the EuroQuol five-dimension scale (EQ-5D-5L),²³ which included mobility, self-care, engagement in usual

activities and pain. Mental and physical health comorbidity was assessed by asking participants to indicate whether they had any physical or mental health conditions (Yes/No), and then if they responded yes, to indicate which condition or conditions they had from a list of physical and mental health conditions.

Four mental health symptoms (depression, anxiety, paranoia and hallucinations) were assessed using a series of validated instruments. Depression was measured using the nine-item Patient Health Questionnaire.²⁴ Anxiety was measured using the seven-item Generalised Anxiety Disorder scale.²⁵ Paranoia was measured using the persecution subscale of the persecution and deservedness scale (PaDS-5) for symptoms of paranoia.²⁶⁻²⁷ Auditory Verbal Hallucinations (AVH) were assessed using a single item from the Launay-Slade Hallucination Scale: “I have been troubled by hearing voices in my head”.²⁸ When comparing severe to non-severe symptoms, we utilised the published clinical cut-offs for severe symptoms of Depression (> 14 on PHQ-9)²⁴ and Anxiety (> 14 on GAD-7).²⁵ There are no validated clinical cut-offs for the PaDS-5 paranoia scale,²⁷ so we categorised people as severe if they scored above the midpoint. People were categorized as experiencing severe AVH if they agreed or strongly agreed with the AVH item.²⁸ Mental health comorbidity was calculated by summing the number of conditions where someone met the severe criterion.

Alcohol consumption was measured by participants indicating if they ever drank alcohol, and if so, how many alcoholic drinks (which we converted to alcohol units dependent on type of alcohol) they had consumed over the past seven days.²⁹ Practical social support and social contact were assessed based on the level of agreement with the statements ‘If I needed help, there are people who would be there for me’ and ‘If I wanted company or to socialise, there are people I can

call on'. (more detailed information on each of the survey variables is given in the online supplementary Appendix A_Table 1).

An index of Multiple Deprivation (IMD) score was also entered for each participant based on their Lower Level Super Output Area (LSOA). The IMD is calculated based on indices of deprivation across seven domains: income, employment, education, health and disability, crime, housing, and living environment. Access to GP and A&E departments was assessed with a distance measure (km) calculated using the Routino open source tool.⁹ We calculated the shortest road distance between the centre of each postcode in the sampled area and each health facility.³⁰ The average distance to GP clinics and A&E departments for all LSOAs was then calculated and linked to the LSOA of each participant's residence.

Preliminary analyses and data preparation

Descriptive statistics and bivariate correlations between mental health symptoms and healthcare service usage are reported (see Supplementary Appendix B, Table 2). On average, participants attended A&E .69 (SD = 2.74) times and a GP 5.5 (SD = 15.05) times in the previous 12 months. The mode was 0 for both A&E attendance (n = 3212) and GP attendance (n = Figure 1 shows the mean levels of attendances for GP and A&E services among people classified as having severe symptoms alongside the total sample mean. Listwise deletion was employed to account for missing values in all analyses. The level of missing data was very low at less than 5% for all variables.

For the regression analyses predicting A&E and GP attendance, we constructed poisson regression models that controlled for potential demographic, socioeconomic, lifestyle, health, and

healthcare access confounds. Rate ratios (RR) were calculated for each variable (see tables 1-3) and standard errors were adjusted to account for the multi-stage nature of the sampling procedure. The model was also weight-adjusted to account for demographic variation in non-response.

Results

The relationship between physical and mental health co-morbidity and service use

Figures 2 shows mean A&E and GP attendances as a function of comorbidity. Results of the poisson regression analysis showed that comorbidity significantly predicted greater A&E and GP attendance over the previous 12 months. Specifically, relative to having no health conditions, having only a mental health condition was associated with a 2.1 times higher rate of A&E attendance (RR = 2.10, 95%CI = 1.33 to 3.31) and 2.5 times higher rate of GP visits (RR = 2.49, 95%CI = 2.03 to 3.04). Having only a physical health condition(s) was associated with a 2.7 times higher rate of A&E attendance (RR = 2.65, 95%CI = 1.78 to 3.95) and a 2.4 times higher rate of attending a GP (RR = 2.43, 95%CI = 2.10 to 2.81). Reporting at least one physical health condition and at least one mental health condition was associated with a 4.6 times higher rate of A&E attendance (RR = 4.64, 95%CI 2.86 to 7.51) and 3.8 times higher rate of attending a GP (RR = 3.82, 95%CI = 3.16 to 4.62). Details of the coefficients for the full model are reported in Table 1.

The relationship between mental health comorbidity and service usage

Comorbidity was operationalised as the numbers of mental health symptoms for which the participant met the “severe” criteria as defined earlier. We constructed separate models for each health service. Mean attendances for each level of comorbidity are shown in Figure 4. Models including coefficients for all variables are provided in Table 2.

Mental health comorbidity was associated with higher A&E attendance for people with four or more severe symptoms. Specifically, people with four severe symptoms were 2.5 times more likely to attend A&E (RR = 2.54, 95%CI 1.43 to 2.4.52) compared to people with no severe symptoms. For GP usage, having one (RR = 1.16, 95%CI 1.01 to 1.32), two (RR = 1.92, 95%CI 1.31, 2.82), three (RR = 1.51, 85%CI 1.23, 1.87), or four (RR = 2.19, 95%CI 1.50 to 3.20) severe symptoms was associated with elevated rates of GP attendance relevant to having no severe symptoms.

Mental health symptoms as predictors of A&E and GP attendance

Details of the coefficients for all variables are reported in Table 3. Of relevance to our key research question, depression was the only mental health symptom that was significantly associated with A&E attendance (RR = 1.41, 95% 1.05 to 1.90) and anxiety was the only symptom significantly associated with GP attendance (RR = 1.19, 95% 1.03 to 1.38), while controlling for health, socioeconomic, alcohol consumption and healthcare access variables.

Discussion

When controlling for a range of potential confounds, physical and mental health comorbidity was an important risk-factor for A&E and primary care attendance, as was mental health comorbidity alone. Specifically, having both a physical and mental health condition was associated with a three-fold increase in GP attendance and a four-fold increase in A&E attendance. Rates of both A&E attendance more than doubled when people had four severe mental health symptoms. However, GP attendance rates were significantly elevated with any number of severe mental health symptoms compared to having no severe symptoms. When examining individual symptoms, we found

depression was associated with more A&E attendances, while anxiety was associated with greater GP attendance. Along with health status, being aged 18-24, from a white ethnic background, and living in disadvantaged circumstances were consistently associated with increased service usage, and living closer to an A&E department predicted greater A&E attendance in all analyses.

6Comparison with existing literature

Being younger and not in employment were the main significant socioeconomic or demographic risk-factors of A&E attendance in this sample, as has been found in previous studies.^{5,30} Having problems with self-care, usual daily activities and pain were all significant physical health predictors. As previously reported,³¹ living closer to an A&E department was associated with more A&E attendances but distance to GP was not associated with A&E attendance in non-depressed people.

In line with previous research,³² the present study confirms that people with anxiety visit GPs more frequently. The association between anxiety and use of GP services may be due to acute exacerbation of anxiety becoming intolerable, and thus patients turn to GPs as their first point of support. . This is in contrast, to severe depression, which is associated with becoming withdrawn from society, particularly for men.³³ Alternatively, severe depression may be viewed as a more urgent concern by carers and relatives due to concerns about imminent risk of self-harm, which in turn may lead to A&E visits.³⁴ Another reason, could be the feeling of acute anxiety being mistaken for other physical health symptoms³⁵ and this misattribution prompting increased GP visiting where patients are likely to be seen quicker and by doctors they know, therefore reducing further anxiety brought on by the uncertainty of the length of wait and the social interaction.

1
2 Additionally, age, ethnicity, unemployment, and vocational or professional qualifications were
3 associated with A&E attendance. Previously, ethnicity was shown to be a significant factor for
4 higher A&E attendance in Black and Asian minority groups compared to white majority groups;⁵
5 but in this sample, there were significantly more people from white backgrounds attending A&E.
6 Earlier studies have shown that people from white backgrounds used A&E more than those from
7 non-white backgrounds³⁶ although the studies were based on relatively small geographical areas
8 with high proportions of particular non-white ethnic groups and therefore with limited
9 generalisability to other areas of the UK.

10
11 People with only a physical health condition attended A&E more frequently in the previous 12
12 months than people with only a mental health condition. People who had both a physical *and*
13 mental health condition, attended A&E and primary care more frequently than all other groups. Co-
14 morbid mental health symptoms were also associated with more use of both healthcare services.
15 However, it was only the people with four or more severe mental health symptoms, and therefore
16 the most complex needs, who attended A&E more frequently compared to people with no severe
17 symptoms. Conversely, having any number of mental health symptoms increased people's rate of
18 GP attendance relative to having no severe symptoms. Both results are consistent with previous
19 studies that have reported high levels of A&E and GP attendance for mental health patients with
20 depression and anxiety in the year prior to suicide^{32,34} However, further research is needed on data
21 that examines reasons for service usage to understand why comorbidity might differentially
22 influence A&E and GP attendance rates.

1Strengths and limitations

2

3 This study contributes to the knowledge that patients with comorbid health conditions visit both
4 A&E and GPs more than patients with a single condition. Our finding that distance to services
5 predicted service usage even when controlling for other potential confounds also supports this
6 assertion. Previous research suggests that there a large proportion of attendances at A&E are
7 avoidable,¹² and mental health patients were identified as one of the main groups contributing to
8 increasing demand within A&E.³⁷

9

10 A strength of using self-reported survey data is the reliability and accuracy of the information
11 provided.³⁸ A review conducted by Legget and colleagues³⁹ about self-reported questionnaires on
12 resource utilisation showed good agreement with administrative data such as electronic records,
13 although, visits to general practitioners, outpatient days, and nurse visits had poorer agreement.
14 Overall, self-reported questionnaires were concluded to be a valid method of collecting data on
15 healthcare resource utilisation, however there is an issue of recall bias, particularly after a length of
16 time.³⁸ Whilst the survey collected data from a wide geographical area from both relatively
17 disadvantaged and advantaged areas, thereby increasing its representativeness, some limitations need
18 to be considered. Considering the focus on people living in disadvantaged neighbourhoods, data are
19 limited to those with a fixed address. Thus, the survey was not able to capture the most disadvantaged
20 groups in the population such as homeless people and unregistered migrants.

21

22 This study used data from the North West Coast of England, which limits generalisability to other
23 regions and populations. While the causal directions proposed here seem the most plausible, due to

the cross-sectional design we cannot rule out the possibility that contact with services may in some instances worsen people's mental or physical health.

Implications for research and/or practice

More work is needed on whether other available services for severe depression will reduce the number of patients attending A&E through a proactive approach to managing patients, particularly in disadvantaged groups and areas with high prevalence of comorbid health conditions. Raising public awareness of the alternatives to A&E for patients with severe depression, particularly in more deprived areas, may reduce the number of attendances; however, alternatives need to be accessible. The findings from this research suggest that more expertise in A&E and in GP practices for severe depression is needed. A recent report⁴⁰ suggests that in-house solutions could include co-locating GP services, use of senior doctors at the front door of the A&E, bringing specialist staff closer to the front door and improved mental health liaison teams. However, to date, there is no evidence to suggest that any of these interventions would improve flow and therefore reduce the number of A&E attendances. Larger and better-designed studies are necessary to address some of these questions. Future interventions to improve access to primary care may be most effectively targeted towards younger adults given the increase in attendances among 18 to 24 year olds. In 2012–2013, 10% of all A&E attendances (1.8 million attendances) were for adults aged 18 to 25 years.

These findings validate the recommended 2016 NICE guidance, which includes a focus on comorbidity and aims to provide a range of coordinated services that address people's wider health and social care needs, such as poor housing.⁴⁰ While not the focus of the present study, the findings

highlight that further research is needed into the social determinants of service usage. The association between higher attendance rates and poorer housing, for example, could be attributed to the condition of social housing and how this impacts on mental health or exacerbates long term physical health conditions.³⁰

References

1. OECD. *Income inequality (indicator)*, 2019. doi:10.1787/459aa7f1-en (accessed on 17 January 2019).
2. Health & Social Care Information Centre. *Hospital episode statistics: accident and emergency attendances in England (experimental statistics) 2011–12*, 2013. (accessed on 17 January 2019). <https://digital.nhs.uk/data-and-information/publications/statistical/hospital-accident-and-emergency-activity/2011-12>
3. Asaria, M., Ali, S., Doran, T., et al. How a universal health system reduces inequalities – Lessons from England. *J Epidemiol Community Health*, 2016; 70: 637-643.
4. Cookson, R., Asaria, M., Ali, S., et al. Health Equity Indicators for the English NHS: a longitudinal whole-population study at the small-area level. *HS&DR*, 2016; 4: 26.
5. Scantlebury, R., Rowlands, G., Durbaba, S., et al. Socioeconomic deprivation and accident and emergency attendances: cross-sectional analysis of general practices in England. *Br J Gen Pract*, 2015; 65:e649–e654.
6. McCormick, B., Hill P.S., Poteliakoff, E. *Are Hospital Services Used Differently in Deprived Areas? Evidence to Identify Commissioning Challenges*. Centre for Health Service Economics and Organisation Working Paper No 2. Oxford: Centre for Health Service Economics and Organisation; 2012.

7. Bryce, A. NHS GGC A&E Attendances Steering Group: *Inequalities, Health and A&E Response*. NHS GGC Corporate Inequalities Team, 2012. (accessed on 17 January 2019).
http://www.equalitiesinhealth.org/public_html/documents/AEFinalReportwrefs23march2011.pdf
8. Purdy S. Avoiding hospital admissions. *What does the research evidence say? The Kings Fund* December 2010. (Accessed 17 January 2019).
http://www.kingsfund.org.uk/publications/avoiding_hospital.html NHS Digital, 2017.
9. Hospital Accident and Emergency Activity 2015-2016. Health and Social Care Information Centre, London: NHS Digital, 2017.
10. Department of Health. *Trends in Consultation Rates in General Practice - 1995-2009, 2009..*
<https://digital.nhs.uk/data-and-information/publications/statistical/trends-in-consultation-rates-in-general-practice/trends-in-consultation-rates-in-general-practice-1995-2009>
11. National Confidential Inquiry into Suicide and Homicide by People with Mental Illness (NCISHa). *Suicide in Primary Care in England: 2002-2011*. Uni of Man, Manchester, 2014.
12. Steventon, A., Deeny, S., Friebe, R., Gardner, T., Thorlby, R. *Briefing: Emergency hospital admissions in England: which may be avoidable and how?* The Health Foundation, 2018.
13. Norton, J., Engberink, A. O., Gandubert, C., et al. Health Service Utilisation, Detection Rates by Family Practitioners, and Management of Patients with Common Mental Disorders in French Family Practice. *Can J Psychiatry*, 2017; 62(8), 521–530.
14. Dowrick, C.F., Bellon, J.A., Gomez, M.J. GP frequent attendance in Liverpool and Granada: the impact of depressive symptoms. *Br J Gen Pract*, 2000; 50, 361-365.
15. Gili, M., Luciano, J.V., Serrano, M.J. et al. Mental disorders among frequent attenders in primary care: a comparison with routine attenders. *J Nerv Ment Dis*, 2011; 199:744–9.

- 1 16. Smits, F.T., Brouwer, H.J., Zwinderman, A.H. et al. Why do they keep coming back?
2 Psychosocial etiology of persistence of frequent attendance in primary care: a prospective
3 cohort study. *J Psychosom Res*; 2014; 77: 492–503.
- 4 17. Norton, J., David, M., de Roquefeuil, G., et al. Frequent attendance in family practice and
5 common mental disorders in an open access health care system. *J Psychosom Res*. 2012; 72(6):
6 413–418.
- 7 18. Ferrari, S., Galeazzi, G.M., MacKinnon, A., Rigatelli, M. Frequent attenders in primary care:
8 Impact of medical, psychiatric and psychosomatic diagnoses. *Psychother Psychosom*, 2008;
9 77:306–314.
- 10 19. Reid, S., Wessely, S., Crayford, T. & Hotopf, M. Frequent attenders with medically unexplained
11 symptoms: service use and costs in secondary care. *Brit J Psychiatry*, 2002; 180(3): 248-253.
- 12 20. Mukhtar, T.K., Bankhead, C., Stevens, S., et al. Factors associated with consultation rates in general
13 practice in England, 2013–2014: a cross-sectional study. *Br J Gen Pract*, 2018; 68 (670): e370-
14 e377.
- 15 21. Giebel, C., McIntyre, J. C., Daras, K., et al. What are the social predictors of accident and
16 emergency attendance in disadvantaged neighbourhoods? Results from a cross-sectional
17 household health survey in the north west of England. *BMJ open*, 2019; 9(1), e022820.
- 18 22. McIntyre, J. C., Wickham, S., Barr, B. & Bentall, R. P. Social Identity and Psychosis:
19 Associations and Psychological Mechanisms. *Schizophr Bull*, 2017. doi: 10.1093/schbul/sbx110
- 20 23. van,Hout B., Janssen,M.F., Feng,Y.S., et al. Interim scoring for the EQ-5D-5L: mapping the
21 EQ-5D-5L to EQ-5D-3L value sets. *Value Health*, 2012; 15(5): 708-15.
- 22 24. Kroenke, K. & Spitzer, R.L. The PHQ-9: a new depression diagnostic and severity measure.
23 *Psychiatr ann*, 2002; 32(9), 509-515.

25. Spitzer, R. L., Kroenke, K., Williams, J. B. & Löwe, B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch intern med*, 2006; 166(10): 1092-1097.
26. Melo, S., Corcoran, R., Shryane, N. & Bentall, R.P. The persecution and deservedness scale. *PAPTRAP*, 2009; 82(3): 247-260.
27. Elahi, A., Algorta, G. P., Varese, F., McIntyre, J. & Bentall, R. Do paranoid delusions exist on a continuum with subclinical paranoia? A multi-method taxometric study. *Schizophr Res*, 2017; 190:77-81.
28. Launay, G. & Slade, P. The measurement of hallucinatory predisposition in male and female prisoners. *Pers Individ Dif*, 1981; 2(3), 221-234.
29. Department of Health. *UK Chief Medical Officers' Low Risk Drinking Guidelines*, 2016.
https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/545937/UK_CMOs_report.pdf (accessed on 17 December 2018)
30. Mental Health Taskforce. The five year forward view for mental health [online]. NHS England website, 2016 Available at: www.england.nhs.uk/mentalhealth/taskforce/ (accessed on 18 November 2018).
31. Tian, Y. Data briefing: emergency hospital admissions for ambulatory care-sensitive conditions. King's Fund report, 2012.
https://www.kingsfund.org.uk/sites/default/files/field/field_publication_file/data-briefing-emergency-hospital-admissions-for-ambulatory-care-sensitive-conditions-apr-2012.pdf (accessed 28 Sep 2018).
32. Pearson, A., Saini, P., Da Cruz, D., et al. Primary care contact prior to suicide in individuals with mental illness. *Br J Gen Pract*, 2009; 59(568): 825-32.

33. Oliver, M.I., Pearson, N., Coe, N. & Gunnell, D. Help-seeking behaviour in men and women with common mental health problems: cross-sectional study. *Br J Psychiatry*, 2005; 186: 297-301.
34. Da Cruz, D., Pearson, A., Saini, P., et al. Emergency department contacts prior to suicide in mental health patients. *Emerg Med J*, 2010; 28:467-471. doi:10.1136/emj.2009.081869
35. Jacob K. S. The diagnosis and management of depression and anxiety in primary care: the need for a different framework. *Postgrad med j*, 2006; 82 (974), 836–839. doi:10.1136/pgmj.2006.051185
36. Baker R., Bankart M.J., Rashid A., et al. Characteristics of general practices associated with emergency-department attendance rates: a cross-sectional study. *BMJ Qual Saf*; 2011; 20(11): 953–958.
37. Mason, S., O’Keeffe, C., Jacques, R., Rimmer, M., Ablard, S. Perspectives on the reasons for Emergency Department attendances across Yorkshire and the Humber Final Report, 2017. https://www.sheffield.ac.uk/polopoly_fs/1.730630!/file/CLAHRC_BMA_Final_Report.pdf (accessed on 17 January 2019)
38. Stone, A.A., Bachrach, C.A., Jobe, J.B., et al. *The Science of Self-Report: Implications for Research and Practice*, Mahwah, NJ: Erlbaum, 2000.
39. Leggett, L.E., Khadaroo, R.G., Holroyd-Leduc, J., et al. Measuring resource utilization: a systematic review of validated self-reported questionnaires. *Medicine*. 2016; 95(10): e2759. pmid:26962773.
40. NICE guideline. Coexisting severe mental illness and substance misuse: community health and social care services. [nice.org.uk/guidance/ng58](https://www.nice.org.uk/guidance/ng58), 2016. (accessed on 17 January 2019) <https://www.nice.org.uk/guidance/GID-QS10078/documents/briefing-paper>

Funding

This work was supported by The National Institute for Health Research, Collaboration for Leadership and Health Research and Care North West Coast (NIHR CLAHRC NWC). The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

Ethical approval

The research was approved by the University of Liverpool Committee on Research Ethics (Ref: RETH00836 & IPHS-1516-SMC-192). Participants provided informed consent before completing the study.

Competing Interests

The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

Conflict of interests

All authors declare no conflict of interest.

Author contributions

The study was conceived by PS and MG. Statistical analysis and interpretation of the data was undertaken by JM and reviewed by all of the authors, who also all contributed to the drafting and revision of the manuscript. All authors have approved the final version of the manuscript.

Acknowledgements

The authors thank the NIHR CLAHRC NWC public advisors: Christopher Whittle, Farheen Yameen and Mersey Care NHS Trust Clinicians: Dr Niall Campbell and Dr Cecil Kullu who contributed at the meetings for this research paper.

1

2