



## LJMU Research Online

**Nurunnabi, TR, Sabrin, F, Sharif, DI, Nahar, L, Sohrab, MH, Sarker, SD, Rahman, SMM and Billah, MM**

**Antimicrobial activity of endophytic fungi isolated from the mangrove plant *Sonneratia apetala* (Buch.-Ham) from the Sundarbans mangrove forest**

<http://researchonline.ljmu.ac.uk/id/eprint/12090/>

### Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Nurunnabi, TR, Sabrin, F, Sharif, DI, Nahar, L, Sohrab, MH, Sarker, SD, Rahman, SMM and Billah, MM (2020) Antimicrobial activity of endophytic fungi isolated from the mangrove plant *Sonneratia apetala* (Buch.-Ham) from the Sundarbans mangrove forest. *Advances in Traditional Medicine*.**

LJMU has developed [LJMU Research Online](#) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)

<http://researchonline.ljmu.ac.uk/>

# **Antimicrobial activity of Endophytic Fungi Isolated from the Mangrove Plant *Sonneratia apetala* (Buch.-Ham) from the Sundarbans Mangrove Forest**

Tauhidur R Nurunnabi<sup>1,2</sup>, Farah Sabrin<sup>3</sup>, Dilara I Sharif<sup>4</sup>, Lutfun Nahar<sup>1</sup>, Md. H Sohrab<sup>5</sup>,  
Satyajit D Sarker<sup>1</sup>, S. M. Mahbubur Rahman<sup>2\*</sup> and Md. Morsaline Billah<sup>2\*</sup>

✉ Dr. Md Morsaline Billah

morsaline@yahoo.com

<sup>1</sup>Centre for Natural Products Discovery, School of Pharmacy and Biomolecular Sciences,  
Liverpool John Moores University, James Parsons Building, Byrom Street, Liverpool L3 3AF,  
UK

<sup>2</sup>Biotechnology and Genetic Engineering Discipline, Khulna University, Khulna, Bangladesh

<sup>3</sup>Department of Biotechnology and Genetic Engineering, Mawlana Bhashani Science and  
Technology University, Santosh, Tangail-1902, Bangladesh

<sup>4</sup>Department of Genetic Engineering and Biotechnology, Jagannath University, Dhaka,  
Bangladesh

<sup>5</sup>Pharmaceutical Sciences Research Division, BCSIR Laboratories, Dhaka

## **Short Title**

Antimicrobial Activity of endophytic fungi from *Sonneratia apetala*

## **Acknowledgements**

The research was funded by Robert S McNamara Fellowship Programme from the World Bank.

## **Abstract**

Endophytic fungi reside in the intercellular space of plant nourished by the plant. In return, they provide bioactive molecules which can play critical roles on plant defense system. Fifty six endophytes were isolated from the leaves, root, bark and fruits of *Sonneratia apetala*, a pioneer mangrove plant in the Sundarbans, Bangladesh. A total of 56 isolates were obtained and 12 different species within 8 genera were identified using morphological and molecular characteristics. Antimicrobial activity of Ethyl Acetate (EtOAc) and Methanolic (MeOH) extracts of these 12 different species were analyzed by resazurin assay and the Minimum Inhibitory Concentrations (MICs) were determined. The fungal extracts showed antimicrobial activities against more than one tested bacterium or fungus among 5 human pathogenic microbes, i.e. *Escherichia coli* NCTC 12241, *Staphylococcus aureus* NCTC 12981, *Micrococcus lutus* NCTC 7508, *Pseudomonas aeruginosa* NCTC 7508 and *Candida albicans* ATCC 90028. Overall, Methanolic extracts showed greater activity than that of Ethyl Acetate extracts. Of the isolates identified, *Colletotrichum gloeosporioides*, *Aspergillus niger* and *Fusarium equiseti* were the most active isolates and showed activity against microorganisms under investigation. Methanolic extracts of *C. gloeosporioides* and *A. niger* showed the lowest MIC (0.0024 mg/mL) against *Pseudomonas aeruginosa*. The study indicates that endophytic fungi isolated from *S. apetala* species possess potential antimicrobial properties, which could be further investigated.

**Key words:** Antimicrobial, Endophytic fungi, Mangrove plants, *Sonneratia apetala*, The Sundarbans.

## **Introduction**

The appearance of new diseases such as Influenza, SARS, and H1N1 has become a foremost challenge to the human health. Majority of these newly emerged diseases are caused by microorganisms and causative microbes are increasingly becoming drug resistant over time (Bhatia and Narain 2010). To combat such infectious diseases, novel bioactive compounds from plants and microorganisms could provide the best alternative as source of potential and promising drugs (Morens et al. 2004). Although, plant is the major source of bioactive compounds, nevertheless, endophytes could play a vital role in search of new bioactive compounds (Jalgaonwala et al. 2011). Endophytic fungi colonize healthy plant tissues without any disease symptoms and protect plants from herbivores and pathogens as because they produce bioactive secondary metabolites in the host tissues. It is reported that, endophytic fungi could produce similar or the same bioactive metabolites as its host plant. The environmental conditions of plants habitat also have some effects on the endophytes present inside the plants (Sette et al. 2006).

Mangroves provide a unique and dominant ecosystem comprised of intertidal marine plants, mostly trees, predominantly bordering margins of tropical coastlines around the world. These halophytic (salt tolerant) plants thrive in intertidal areas that receive daily inundation between mean sea levels and the highest astronomical tides. It has been well established that environment where the plant leaves has enormous effect on the diversity of the metabolites produce by the entophytes (Tomlinson 1986; Duke et al. 1998). With this end in view, *S. apetala*, has been selected for the present study because this plant has hardly been considered for its bioactive endophytic fungi and is one of the most dominant species in the Sundarbans mangrove forest. Thus, the present study was conducted to investigate the abundance of endophytic fungi associated with *S. apetala* found in the Sundarbans, Bangladesh and evaluate antimicrobial activity of the fungal endophytes against human pathogens. Screening of endophytic fungi with inhibitory activities against pathogenic bacteria or fungi might lead to potential novel natural products with higher antimicrobial activity.

## **Materials and Methods**

### **Collection of plant samples**

Healthy roots, bark, fruits and leaves of *S. apetala* without any sign of infections were collected from the mangrove forest, the Sundarbans, Khulna, Bangladesh. During sample collection, the plants were identified based on their morphological characteristics outlined by Zabala (1990).

## **Isolation of Endophytic Fungi**

Isolation of endophytic fungi was carried out according to the procedure described by Nurunnabi et al. (2018). Briefly, plant materials were washed thoroughly in sterile water to remove extraneous substances, followed by surface sterilization (sequentially immersing in 70% ethanol for 30 sec and 5% sodium hypochlorite solution for 1 min) and finally rinsed with sterile distilled water. The samples were surface-dried with sterile filter paper (Hormazabal and Piontelli 2009). Roots, bark, fruits and leaves were cut into 0.5 cm × 0.5 cm pieces and placed in petri dishes containing aqueous agar (1.5% agar-agar in distilled water) supplemented with antibiotic streptomycin (0.03mg/mL) and incubated at  $28 \pm 2^\circ\text{C}$  until initiation of fungal growth. The tips of the fungal hyphae were collected from the aqueous agar and placed on Potato Dextrose Agar (PDA) medium. After several days of incubation, the purity of each fungal culture was determined by assessing the colony morphology. The isolates were cultured on PDA media to obtain pure cultures and then transferred into PDA slants for preservation at  $4^\circ\text{C}$ .

## **Microscopic and Molecular Identification of the Fungal Isolates**

Both morphological and molecular techniques were employed to identify the isolated endophytic fungi. Morphological identification was performed according to the procedure described by Qadri et al. (2013). Fungal genomic DNA was isolated according to the protocol supplied along with the DNA isolation kit (Cat-26200; NORGEN BIOTEK Corp., 3430 Schmon Parkway, Thorold, ON, Canada). Forward primer ITS4 (5'TCCTCCGCTTATTGATATGC3') and reverse primer ITS5 (5'GGAAGTAAAAGTCGTAACAAGG 3') were used to amplify the internal transcribed spacer (ITS) region of the fungi using Polymerase Chain Reaction (Sette *et al.* 2006). The PCR reaction was performed in 50 $\mu\text{L}$  reaction volume comprising 10 $\mu\text{L}$  of 5X reaction buffer, 3 $\mu\text{L}$  of 1.5M  $\text{MgCl}_2$ , 5 $\mu\text{L}$  of 10mM dNTP, 3 $\mu\text{L}$  of 100pmol primers, 26 $\mu\text{L}$  of distilled water and 3 $\mu\text{L}$  (2ng/ $\mu\text{L}$ ) of template DNA. The PCR amplification was performed using 2720 PCR Thermo Cycler and the temperature profile was maintained as initial denaturation at  $94^\circ\text{C}$  for 5 min followed by 40 cycles (denaturation  $94^\circ\text{C}$  for 30 sec, annealing  $51^\circ\text{C}$  for 30 sec and extension  $72^\circ\text{C}$  for 2 min). The final extension was carried out at  $72^\circ\text{C}$  for 5 min and held at  $4^\circ\text{C}$ . The PCR amplification of the ITS region was verified by electrophoresis (10 $\mu\text{l}$ ) and the rest of the PCR products (40 $\mu\text{l}$ ) were purified according to requirement of Cambridge Genomic Services (The University of Cambridge, UK) for DNA sequencing. Multiple BLASTN searches against the sequence were made at the National Center for Biotechnology Information (NCBI). Consensus sequences were submitted in the GenBank. Primarily,

Clustal Omega was employed for Multiple Sequence Alignment and further followed by trimming using trimAI tool for later alignment. Phylogenetic analysis performed with the neighbor joining method using MEGA7.0 software.

### **Extraction of secondary metabolites**

Extraction of the secondary metabolites was carried out according to the Nurunnabi et al. (2018). The fungal isolate was grown in (5 × 250mL) conical flasks containing potato dextrose broth (PDB) for 28 days. Culture broth was separated from the mycelium by filtration (Whatman® qualitative filter paper, Grade 1; Sigma-Aldrich, USA) and the filtrates were extracted three times with an equal volume of EtOAc in a separating funnel. The mycelium was dissolved in methanol under dark condition for two days and the mycelium was separated by filtration. Similarly, methanolic extracts were also prepared and both types of extracts were evaporated under reduced pressure at 40-45°C using a rotary evaporator to obtain crude extracts.

### **Antimicrobial screening**

The EtOAc and Methanolic extract of all 12 isolated fungi were tested for their potential antimicrobial activity against two Gram-positive, i.e., *Staphylococcus aureus* NCTC 12981 and *Micrococcus luteus* NCTC 7508, and two Gram-negative, i.e., *Escherichia coli* NCTC 12241 and *Pseudomonas aeruginosa* NCTC 12903 bacterial strains as well as against a fungal strain, *Candida albicans* ATCC 90028 using the resazurin 96-well microtitre plate based *in vitro* antimicrobial assay (Sarker et al. 2007). For bacterial strains, Ciprofloxacin was used as a positive control and nystatin for *C. albicans*. Resazurin solution (4mg of resazurin was dissolved in 20ml of sterile distilled water) was used in this assay as an indicator of cell growth. Briefly, sterile 96 well plates were prepared and labeled under aseptic conditions. A volume of 100µl of test material in 10% (v/v) DMSO (10mg/ml for crude extracts) was pipetted into the first row of the plate and 50 µl of normal saline was added to all other wells. Serial dilutions were performed in such that each well had 50µl of the test material in serially descending concentrations. 30µl nutrient broth and 10µl of resazurin indicator solution were added to each well. Finally, 10µl of bacterial suspension ( $5 \times 10^5$ cfu/ml) was added to each well. To prevent the bacterial culture dehydration, each plate was wrapped loosely with cling film. Each plate had a column with a broad-spectrum of antibiotic as positive control (usually ciprofloxacin in serial dilution) a column with all solutions with the exception of the bacterial solution adding 10µl of nutrient broth instead. The plates were prepared in triplicates, and placed in an incubator set at 37°C for 18-24 hours. The color change was then assessed visually. Any color changes from purples to pink or colorless were

recorded as positive. The lowest concentration at which color change occurred was taken as the Minimum Inhibitory Concentration (MIC) value. The average of three values was calculated and that was the MIC for the test materials.

## Calculation and Statistical Analysis

Isolation rate (IR) was calculated using the formula of Gong and Guo (2009). Isolation rate (IR) was used to demonstrate the fungal richness in a given sample of plant tissue which was counted as the number of isolates obtained from plant segments divided by the total number of segments incubated. The statistical analysis was analyzed using Graph Pad Prism, var. 6.0.

$$IR = \frac{\text{the total number of isolates yielded in a given trial}}{\text{The totaao number of samples in that trial}} \text{----- (i)}$$

Colonization rate (CR) expressed in percentage, was calculated as the total number of Colonization rate (CR), expressed in percentage, which was calculated as the total number of plant tissue segments infected by fungi divided by the total number of segments incubated, was used to indicate comparison of degrees of different tissues infected by endophytic fungi. Isolation rate (IR) was used to demonstrate the fungal richness in a given sample of plant tissue which was counted as the number of isolates obtained from plant segments divided by the total number of segments incubated (Gong and Guo 2009). The statistical analysis was analyzed by Graph Pad Prism.

## Results

### Isolation and Identification of Endophytic fungi

A total of 58 endophytic fungi were isolated from 120 tissue segments (30 segments from Root, Bark, Leaves and Fruits each) from *S. apetala*. The isolation rates for Root, Bark, Fruits and Leaves were 46, 43, 33 and 70%, respectively. The isolation rates of endophytic fungi from leaves were higher than that of bark, root and fruits. On the basis of morphological analysis (Fig. 1), 12 different types of endophytes were selected for molecular identification. Genomic DNA of all 12 endophytic fungi was extracted and multiplied by ITS4 and ITS5 universal primers. After gel electrophoresis, the PCR amplified products of about 500 bases were obtained and presented in Fig. 2.

DNA sequences obtained from amplification of ITS region were submitted to Genbank and accession numbers for each fungus were obtained. A range of 520 to 602 nucleotide pair sequences were obtained. *Neopestalotiopsis chrysea* has the shortest length of ITS region of 520 base pair in contrast to *Aspergillus niger*

which was 602 base pair. Table 1 represents the sequence data of isolated endophytes from *S. apetala*, where 99 to 100% BLAST match sequences were obtained.

### **Pair wise genetic distance and phylogenetic tree construction**

Table 2 represents the pair-wise genetic distances estimated from sequenced data. The pair-wise genetic distances ranged from 0.376 to 0.927. The maximum pair-wise genetic distance (0.927) was found between SaL11 (*A. niger*) and SaR7 (*F. equiseti*). The result can be justifying by reason that they both are from different genus. Although three *Alternaria* spp. and two *Cladosporium* species were present but the maximum similarity was found in between SaL5 (*Alternaria arborescens*) and SaF5 (*Epicoccum nigrum*). A UPGMA tree was constructed (Fig. 3) from these distance matrix using the software PUAP 4.0. The UPGMA tree of 12 isolated endophytic fungi shows two main clusters. Cluster one consisted with eight fungi viz. SaL11 (*A. niger*), SaL2 (*Alternaria brassicicola*), SaR7 (*F. equiseti*), SaR9 (*Cladosporium cladosporioides*), SaR1 (*Xenoacremonium recifei*), SaL13 (*Cladosporium perangustum*), SaL9 (*Colletotrichum glosporides*) and SaF1 (*Cladosporium tenuissimum*). The second cluster was consisted with four fungi and they are SaL5 (*A. arborescens*), SaF5 (*E. nigrum*), SaL4 (*A. tenuissima*) and SaB6 (*Neopestalotiopsis chrysea*).

### **Antimicrobial Activities of Ethyl Acetate and Methanolic extracts**

Methanolic Extracts from *C. cladosporioides*, *C. perangustum*, *C. tenuissimum*, *F. equiseti*, *A. brassicicola*, *A. arborescens*, *C. glosporides*, *N. chrysea* and *A. niger* showed effect on all the tested microorganisms and for the Ethyl Acetate extract *F. equiseti*, *C. glosporides* and *A. niger* showed activity against all tested microorganisms (Table 3). On average, the methanolic extracts seem to be the more effective on the tested microorganisms. Methanolic and Ethyl Acetate extract from *F. equiseti*, *C. glosporides* and *A. niger* showed comparatively greater antimicrobial activity against all tested microorganisms. Methanolic extracts from *C. glosporides* and *A. niger* confirm the highest MIC (0.00024 mg/mL) for *S. aureus* (NCTC 12981). All the fungal extract showed activity against *P. aeruginosa* (NCTC 12903), *S. aureus* (NCTC 12981) and *M. luteus* (NCTC 7508) extracts showed lesser activity against *E. coli* (NCTC 12241) and *C. albicans* (ATCC 90028), respectively.

### **Discussion**

The present study has revealed that *S. apetala* harbors a lot of endophytic fungi that produces potent antimicrobial substances. The findings of this study also suggest that endophytes from harsh and competitive environments, such



as mangrove ecosystem, might be an attractive source for new anti-infective compounds. Xing et al. has recently reported *Mycosphaerella* spp., *Fusarium* spp. *Glomerella* spp., *Phomopsis* spp., and *Pestalotiopsis microspore* from *S. apetala* (Xing et al. 2011). The present work is the first report of endophytic fungi *A. tenuissima*, *A. brassicicola*, *A. arborescens*, *C. cladosporioides*, *C. perangustum*, *C. tenuissimum*, *E. nigrum*, *N. chrysea*, *X. recifei* and *A. niger*. In the present study, *Fusarium* sp. inhibited both bacteria and fungi with lowest MIC, which was in agreement with the information reported in a previous study that demonstrated *Fusarium* sp. isolated from *Saussurea involucrata* with broad antimicrobial spectrum (Lv et al. 2010). In another study, endophytic fungi isolates from twelve mangrove species were found to be diverse. Similar to our study, the study reported the extraction of fungal broth by ethyl acetate while the mycelia were extracted using methanol, hexane and ethyl acetate in sequence, respectively (Buatong et al. 2011). Ethyl acetate is widely used in extraction of endophytic fungal cultures (Bhardwaj et al. 2015; Minarni et al. 2017) followed by methanol. As a solvent, ethyl acetate solvent possesses medium polarity so that it has the ability to dissolve both polar and non-polar active compounds and methanol solvent being a polar solvent can dissolve almost all organic compounds, even polar, semi polar and non-polar (Rahmawati et al. 2018). However, apart from separation of compounds, the solvents had no effect on antimicrobial potential of the fungal extracts as the extracts were evaporated under reduced pressure using rotary evaporator.

In the present study, we have identified the endophytic fungus through morphological analysis, however, this type of identification is presumptive and requires fair bit of experience. Therefore, in addition to this, we chose nucleotide sequence analysis of ITS region because it lies between two highly conserved genes coding for 18S and 28S rRNA. The ITS regions circumscribe two non-coding regions ITS1 and ITS2, which are separated by the highly conserved 5.8S rRNA gene (Crouch et al. 2009). The ITS1 and ITS2 regions are more variable than the other adjacent rRNA gene sequences (Ciardo et al. 2006). Therefore, offer a better identification of closely related species. By sequence determination and comparison with the sequences of Genbank data base, all the strains identified to the species level (99 to 100 percent homology to the best-matching reference sequence).

In this study, we successfully isolated and identified 12 different species of endophytic fungi belonging to 8 different genera. All the isolates are moderately active against tested microorganisms. However, further studies could be initiated with *F. equiseti*, *C. glosporides* and *A. niger* for potential bioactive compounds as our results showed promising activities with these endophytic fungi. The findings of this study also suggests that endophytes

from harsh and competitive environments, such as mangrove ecosystem, might be an attractive source for bio-prospecting of new antimicrobial compounds.

## Acknowledgements

The research was funded by Robert S McNamara Fellowship Programme from the World Bank.

## Conflict of Interests

The authors declare no conflict of interests.

## References

- Buatong J, Phongpaichit S, Rukachaisirikul V, Sakayaroj J (2011) Antimicrobial activity of crude extracts from mangrove fungal endophytes. *World J Microbiol Biotechnol* 27(12): 3005-3008.
- Bhardwaj A, Sharma D, Jadon N, Agrawal PK (2015) Antimicrobial and phytochemical screening of endophytic fungi isolated from spikes of *Pinus roxburghii*. *Arch Clin Microbiol* 6(3): 1-9.
- Bhatia R, Narain JP (2010) The growing challenge of antimicrobial resistance in the South-East Asia Region-Are we losing the battle?" *Ind J Med Res* 135(5): 482-486. <https://dx.doi.org/10.4103%2F0971-5916.73313>
- Ciaro DE, Schar G, Bottger EC, Altwegg M, Bosshard PP (2006) Internal Transcribed Spacer Sequencing versus Biochemical Profiling for Identification of Medically Important Yeasts. *J Clin Microbiol* 44(1): 77-84. <https://doi.org/10.1128/JCM.44.1.77-84.2006>
- Crouch JA, Clarke BB, White JW, Hillman BI (2009) Systematic analysis of the falcate-spored gramicolous *Colletotrichum* and a description of six new species from warm-season grasses. *Mycologia* 101: 717-732. <https://doi.org/10.3852/08-230>
- Duke NC, Benzie JAH, Goodall JA, Ballmen TR (1998) Genetic structure and evolution of species in the mangrove genus *Avicennia* (Avicenniaceae) in the indo-west pacific. *Evolution* 52(6): 1612-1626. <https://doi.org/10.1111/j.1558-5646.1998.tb02242.x>
- Gong LJ, Guo SX (2009) Endophytic fungi from *Dracaena cambodiana* and *Aquilaria sinensis* and their antimicrobial activity. *Afr J Biotechnol* 8(5): 731-736.

- Hormazabal E, Piontelli E (2009) Endophytic fungi from Chilean native gymnosperms: antimicrobial activity against human and phytopathogenic fungi. *World J Microbiol Biotechnol* 25: 813–819. <https://doi.org/10.1007/s11274-008-9953-6>
- Jalgaonwala R, Mohite B, Mahajan RA (2011) Review: Natural products from plant associated endophytic fungi. *Journal of microbiology and biotechnology research* 1: 21–32.
- Lv YL, Zhang FS, Chen J, Cui JL, Xing YM, Li XD, Guo SX(2010) Diversity and antimicrobial activity of endophytic fungi associated with the alpine plant *Saussurea involucreata*. *Biol Pharm Bull* 33(8): 1300–1306. <https://doi.org/10.1248/bpb.33.1300>
- Minarni, MadeArtika I, Julistiono H, Bermawie N, IdaRiyanti E, Hasim, ZainalHasan AE (2017) Anticancer activity test of ethyl acetate extract of endophytic fungi isolated from soursop leaf (*Annona muricata* L.). *Asian Pac J Trop Med* 10(6): 566-571.
- Morens DM, Folkers GK, Fauci AS (2004) The challenge of emerging and reemerging infectious diseases. *Nature* 430: 242–249. <https://doi.org/10.1038/nature02759>
- Nurunnabi TR, Nahar L, Al-Majmaie S, et al. (2018). Anti-MRSA activity of oxysporone and xylitol from the endophytic fungus *Pestalotia sp.* growing on the Sundarbans mangrove plant *Heritiera fomes*. *Phyther Res* 32(2): 348-354. <https://doi.org/10.1002/ptr.5983>
- Rahmawati NI, Sunarva S, Rumidatul S (2018) Exploration of potential bioactive compounds of endophytic microbial culture isolated from gall rust sengon (*Falcataria moluccana*). *J Pharm Res Sci* 10(11): 156-169
- Qadri M, Johri S, Shah BA, Khajuria A, Sidiq T, Lattoo SK, Abdin MZ, Riyaz-Ul-Hassan S (2013) Identification and bioactive potential of endophytic fungi isolated from selected plants of the Western Himalayas. *Springer Plus* 2: 1-14. <https://doi.org/10.1186/2193-1801-2-8>
- Sarker SD, Nahar L, Kumarasamy Y (2007) Microtitre plate-based antibacterial assay incorporating resazurin as an indicator of cell growth, and its application in the *in vitro* antibacterial screening of phytochemicals. *Methods* 42: 321-324. <https://doi.org/10.1016/j.ymeth.2007.01.006>
- Sette L, Passarini M, Delarmelina C, Salati F, Duarte M (2006) Molecular characterization and antimicrobial activity of endophytic fungi from coffee plants. *World. J. Microbiol. Biotechnol* 22(11): 1185-1195. <https://doi.org/10.1007/s11274-006-9160-2>
- Tomlinson PB (1986) *The Botany of Mangroves*. Cambridge University Press, Cambridge.

Xing XK, Chen J, Xu MJ, Lin WH, Guo SX (2011) Fungal endophytes associated with *Sonneratia* (Sonneratiaceae) mangrove plants on the south coast of China. *Forest Pathology* 41: 334–340. <https://doi.org/10.1111/j.1439-0329.2010.00683.x>

Zabala NQ (1990) *Silviculture of species. Development of professional education in the Forestry Sector of Bangladesh.* Institute of Forestry, Chittagong University, Bangladesh, pp 96-150.