



LJMU Research Online

Montirosso, R and McGlone, F

The body comes first. Embodied reparation and the co-creation of infant bodily-self.

<http://researchonline.ljmu.ac.uk/id/eprint/13043/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Montirosso, R and McGlone, F (2020) The body comes first. Embodied reparation and the co-creation of infant bodily-self. Neurosci Biobehav Rev, 113.

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

The body comes first Embodied reparation and the co-creation of infant bodily-self.

Rosario Montirosso^{1*}rosario.montirosso@lanostrafamiglia.it Francis McGlone^{2,3}

¹ *0-3 Center for the at-Risk Infant*, Scientific Institute IRCCS Eugenio Medea, Bosisio Parini (Lecco), Italy

² *School of Natural Sciences and Psychology*, Liverpool John Moores University, Liverpool, UK

³ *Institute of Psychology Health & Society*, University of Liverpool, Liverpool, UK

*Corresponding author: Rosario Montirosso, *0-3 Center for the at-Risk Infant*, Scientific Institute, IRCCS E. Medea, Via don Luigi Monza 20, 23842 Bosisio Parini (Lecco), Italy – Phone: +39-031-877494; E-mail:

Highlights

- Infant bodily-self emerges in early embodied interactions with mother
- Embodied interactions typically fluctuate between attuned and misattuned states
- Embodied reparation is achieved through tactile contact and maternal interoceptive sensitivity
- These elements contribute to co-create the infant bodily-self

Introduction

The bodily-self is a compelling certitude which supports the ability to perceive our body as separate from other entities and provides the basis of self-perception. This in turn is crucial for engaging in social interactions, interpreting and predicting the thoughts, actions and behaviors of others (Meltzoff, 2017). Bodily-self is a concept that involves not only neural representation and modulation of physiological homeostasis, but also the perception of body ownership (Seth, 2013). In adults, information regarding the state of the bodily-self depends upon exteroceptive somatosensory, visual, and vestibular signals, as well as those originating interoceptively, which provide inputs and sensations about the state of one's own body, such as pain, temperature, itch, respiration, cardiac signals, hunger etc., which are integrated to generate bodily-self representation (Riva, 2017). While there are no conclusive findings about the exact timing, developmentally, of the emergence of the sense of a bodily-self (Slaughter and Brownell, 2011), recent studies indicate that processing of information from the body begins during the first year of life and more likely a process that starts *in-utero* (Zieber et al., 2015). Studies on the ontogeny of human body perception in infants suggest that a pre-reflective, non-conceptual form of bodily

self-awareness, a primitive sense of self (i.e., 'minimal self'), is already present in the first months of life (Gallagher, 2000; Damasio, 2010).

The formation of bodily-self is hypothesized to be driven and progressively refined by learning about sensorimotor contingencies via interactions between brain-body-environment (Georgie et al., 2019) and self-generated actions (i.e., self-exploration, Di Mercurio et al., 2018). Interestingly, it has been suggested by Marshall and Meltzoff (2015) that sensory/perceptual experience of one's own body involves relational and emotional exchanges. For example, research on the μ rhythm, recorded electroencephalography (EEG), suggests that infants are provided with a mirror mechanism that enables them to generate, at the cortical level, a bidirectional representation of both other-action perception and their own-action execution (Caspers et al., 2010), and that they can activate sensorimotor correspondences between their own body parts and the body parts of others (Meltzoff et al., 2018). Interestingly, alterations in early interpersonal experience, such as preterm birth, might contribute to deficits in the bidirectional sensorimotor representation of both other-action perception and an infant's own-action execution (Montirosso et al., 2019). This evidence suggests that during infancy the bodily-self is not an entity dependent solely on inherited neuro-maturational processes (e.g., brain-body interactions), rather it is dependent upon relational experiences of body-to-body interactions (i.e., embodied interactions) which provide the basis for the constitution of the minimal-self, a necessary and vital precursor of subsequent 'self-other' distinctions (Fotopoulou and Tsakiris, 2017; Ciaunica and Crucianelli, 2019).

From birth, an infant is engaged in body-to-body interactions with their mother, most of which are part of routines such as having a nappy changed, being fed, playful exchanges etc. During these embodied interactions, by countless multimodal reciprocal exchanges, mother and infant need to coordinate their behaviors and bodies, and both contribute to interpersonal

rhythmic cycles of co-regulation (an interpersonal process variously termed ‘synchrony’, ‘mutual regulation’, or ‘attunement’; Tronick, 2004; DiCorcia and Tronick, 2011). Within these cycles of mutual regulation, face-to-face exchanges, physical proximity and tactile interactions have a key role for supporting bio-behavioral synchrony (Atzi and Gendron, 2017), that is the coordination of physiological, neuroendocrine and behavioral exchanges which that are known to be important for the building of the infant's social development (Feldman, 2012; Feldman et al., 2014). Indeed, accumulating evidence has documented that during the first year of life an infant experiences implicit synchrony between its own bodily activities and the mother's, as evidenced by EEG activity, heart rhythm, and neuroendocrine responses (Winberg, 2005; Leong et al., 2017; Van Puyvelde et al., 2019). While a number of studies suggest that the dyad's ability of co-regulation is important for the development of well-functioning stress regulation (DiCorcia and Tronick, 2011; Feldman et al., 2012; Welch, 2016), more recent views hypothesize that this interpersonal process, which unfolds during early interactions, might support the infant's bodily-self perception (Ciaunica and Crucianelli, 2019).

Recognizing the centrality of a bio-behavioral synchrony perspective being embedded in early embodied interactions leads, at least in part, to a couple of critical questions. First, although mothers and infants are able to reciprocally adjust their own physiological activities (Van Puyvelde et al., 2015), above all when they are in close body contact, it should be recognized that a constant bio-behavioral synchrony is a rather improbable phenomenon. In agreement with evidence from research on early social-interaction dynamics (Tronick, 2004), it is reasonable to assume that during embodied interactions co-regulation states (referred to as attunement) and non co-regulation states (misattunement), alternate. This poses the question: how does each partner's bodily activity, dynamically and continuously, make possible the shift from misattunement to attunement? Second, if the cycles of mutual regulation are a process which develops during

embodied interactions, then we need to consider the role played by the mother's body and not only focus on the infant's body. Specifically, it would be of interest to understand if, and to which extent, the mother's body contributes to how parents interpret and respond to their infant's bodily signals (so-called, maternal sensitivity). Indeed, it remains unclear how infants and their mothers reach bio-behavioral synchrony and what is the contribution of the maternal body in this process. Here, we discuss recent evidence which allows the examination of bio-behavioral synchrony in early interactions indicating that mothers' interact with infants using: i) tactile contact behaviors (e.g., skin-to-skin, touch) and ii) interoceptive sensitivity (i.e., ability to perceive internal input about the state of one's own body). We propose that, along with other interactive processes (Markova et al., 2019), bio-behavioral synchrony is achieved between the dyad through tactile contact behaviors as much as it may be induced by maternal interoceptive sensitivity, which supports moment-by-moment adaptation of the mother to interactive demands and the infant's needs. Although these elements are likely interwoven, the focus here is not to analyze their potential links, rather to describe how they might contribute to bio-behavioral synchrony. We conclude by presenting the hypothesis that embodied interactions promote not only interpersonal synchrony, but also co-creates the infant bodily-self.

Tactile interactions between parent and infant are pervasive, playing a central role for an infant's developing social, emotional, and physical needs (Moszkowski et al., 2009; Jean, Stack, and Fogel, 2009; Stack and Jean, 2011). For example, recent work has documented that touch during the neonatal period is associated with improved physiological and behavioral indices of emotional reactivity in infants of depressed mothers (Sharp et al., 2012). Interestingly, immediate post-natal tactile stimulation and physical contact reduce newborns' crying and distress and support newborn adaption to life outside of the womb (Winberg, 2005). A recent study found that mothers' touching behaviors in the first minutes to an hour after birth were predictive of a

mother's touching behaviors 3-months later (Mercuri et al., 2019). In this latter, the frequency of immediate post-natal maternal touch was associated with more soothing, regulating maternal touch observed after an age-appropriate experimental stress induction procedure, the Face-to-Face Still-Face paradigm, (i.e. stress resulting from a temporary parent's unresponsiveness; Tronick et al., 1978). Furthermore, recent evidence suggests epigenetic mechanisms influencing stress response are associated with tactile contact experience in full-term infants (Lester et al., 2018; Murgatroyd et al., 2015), while adverse stress-related experience, which also implies an altered maternal contact-related experience during the early stage of life (e.g., prematurity), can have an impact on the DNA methylation, stress response and endocrine regulation, short- (Montirosso et al., 2016; Provenzi et al., 2016) and long-term (Provenzi et al., 2019).

Nevertheless, the importance of maternal touch to an infant's developing needs goes well beyond these aspects. Touch is a multidimensional and dynamic system and has a unique role on perception of body ownership in adults (Tsakiris, 2017). Specifically, touch could enhance the early subjective feelings of body ownership, supporting multisensory integration (Crucianelli et al., 2018). Indeed, there is now evidence for multimodal integration related to body perception in the early period of life. Recent research finds newborns are able to detect multisensory integration of body-related information both for the temporal (Filippetti et al., 2013) and spatial properties (Filippetti et al., 2015) of visual-tactile stimulation. At 4 months of age infants are sensitive to tactile-auditory spatio-temporal correspondences (Thomas et al., 2017). Even more remarkably, 5 month old infants looked longer at an animated character that was moving in asynchrony with their own cardiac rhythm, suggesting that even at this early age, infants showed an implicit sensitivity to interoceptive signals, and an ability to integrate these interoceptive signals with external visual-auditory stimuli (Maister et al., 2017). However, the infant's brain is not mature enough for a complete multisensory integration, and recent studies suggest development of this

process is not predetermined (Burr and Gori, 2012; Stein, Stanford and Rowland, 2014), rather it improves and becomes increasingly more specialized with experience (Lewkowicz and Bremner, 2020) and social interaction with physical contact would facilitate multisensory integration. In a study, 8-months-old infants heard pseudo-words both while they were tickled (multimodal audio–tactile condition) and not tickled (unimodal audio condition). After this phase, brain activity (i.e., EEG) was measured during the perception of the same words. Compared to the audio condition, the audio–tactile condition was associated with higher brain activity within the left temporal regions (Tanaka, Kanakogi, Kawasaki and Myowa, 2018). Authors suggest that their findings provide neural evidence that the integration of audio-tactile information in infant’s brain is facilitated by social interaction involving physical contact with others. Thus, along with other processes involved in brain maturation, interpersonal tactile stimulation could be implicated in integrating information between modalities, which in turn would support the development of bodily self-perception in the infant (Fotopoulou and Tsakiris, 2017; Crucianelli and Filippetti, 2018).

While accumulating evidence supports the fundamental role of gentle touch during parent-infant interactions, little is known about the role of parental interoception. However, how interoception is defined and quantified remains a matter of debate (Garfinkel et al., 2015). From a general point of view interoception refers to the ability to perceive physiological changes, especially from internal organs such as the viscera, heart, lungs, and skin (Craig 2002). While its importance for survival is evident, an increasing number of studies suggest that, along with many different external and bodily sensory inputs, interoception might be crucial for the development and function of perception, higher-order cognitive processes and affect (Allen and Tsakiris, 2019). For example, interoceptive accuracy (i.e., the extent to which one can correctly detect one’s own interoceptive signals) has been found to be associated with ability in decision-making (Dunn et al.

2010) and self-regulation (Herbert et al. 2007). Even more important for the aims of the current paper, interoceptive accuracy has been found to be associated with the ability to recognize emotion (i.e., empathy; Fukushima et al., 2011) and to contribute to the perception of the self (Fotopoulou and Tsakiris, 2017). These findings are in line with an embodied cognition theoretical view, which suggests that higher-order cognitive processes are situated in bodily systems, highlighting a putative role of interoception in both self- and other-perception (Gallese, 2007; Tsakiris, 2017). In the light of this evidence, it is reasonable to assume that parental interoception might play a crucial implicit role in ‘mother sensitivity’. In fact, to ‘read’ infant’s bodily signals and to link these cues with specific emotional meanings and appropriate responses, mothers need to have awareness (even if in an implicit way) of their own body experiences (see: Tylka, Lumeng, and Eneli, 2015; for a similar proposition in relation to the development of intuitive eating behaviors).

To analyze this view, we will first discuss the emergence of bodily-self in infants and the links with neuroanatomical and neurophysiological pathways of tactile stimulation. We will also present the role of affectionate touch in mother-infant exchanges. Moreover, beside affectionate touch, early exchanges are mediated by embodied interactions, in which both the mother's and infant’s body dynamically changes their own interoceptive states. We will argue that the mother’s state of interoceptive sensitivity serves as a foundation for responsiveness to her infant’s body signals. According to this perspective, an infant’s subjective experience of bodily-self may not only be localized within the infant (Ciaunica and Crucianelli, 2019), but could rather be conceived as a process of shared inter-corporeality in which the mother plays a critical role in co-creating the infant’s bodily-self.

The emergence of a bodily-self in infants

The newborn’s body is a highly multisensorial ‘entity’ continuously sensing a cacophony of tactile, visual, auditory, vestibular, olfactory, gustatory, proprioceptive and motor signals (André et

al., 2018). Bodily sensations, of interoceptive and exteroceptive origin, are associated with the need of regulating homeostasis and the behavioral drives that are necessary to maintain the integrity of the body (Craig, 2002). Previous studies have suggested the mechanisms through which visual, vestibular, proprioceptive and tactile inputs are integrated in different levels of body awareness (Gentsch et al., 2016; Fotopoulou and Tsakiris, 2017). The mechanisms underpinning somatosensory integration with other external environmental signals are thought to support a non-conceptual and basic form of bodily-self (Ehrsson, Holmes and Passingham, 2005; Ionta et al., 2011; Tsakiris, 2010), likely associated with a sense of agency and body ownership (Riva, 2017; Seth and Tsakiris, 2018).

Human newborns have a basic sense of self (Filippetti et al., 2015) and a rudimentary distinction between representations of their bodies and those of other people (Rochat, 2003). Through self-exploration and perception of contingent sensorimotor information infants learn what belongs to their body (Gergely and Watson, 1999; Filippetti et al., 2013). In fact, synchronous visual-proprioceptive and visual-tactile stimulation is uniquely 'self-specifying'. The infant can use such contingencies to identify self-performed actions very early in life. For example, in the first hours of life infants look longer at a video of another newborn's face if it is stroked by a brush when their own face is brushed synchronously, rather than asynchronously (Filippetti et al., 2013). Importantly, this preference disappeared in both synchronous and asynchronous tactile stimulation when the newborns look at an infant's face in an inverted presentation, suggesting that they may have a predisposition for specific naturalistic body patterns (Filippetti et al., 2013). Likewise, 7- and 9-month-old infants prefer to look at a video of a doll's legs being stroked synchronously, rather than asynchronously, with their own legs (Zmyj et al., 2011). The difference is in the temporal synchrony or spatial congruency between proprioceptive input (where infants feel the body to be) and visual stimulation (where the infants see the body to be), suggesting that

they behave differently towards sensations that originate from their own body versus those that are caused by the external stimulation. Other evidence shows that visual-tactile contingencies may have a specific role in an infants' perception of their own body. Using magnetoencephalography (MEG), Meltzoff and colleagues (2018) investigated the neural representation of body parts in 7-month-old infants. Somatosensory cortex was activated when their own body was touched, but also when infants visually perceived touch to someone else's body. The authors suggest that cortical representations of body parts may underlie connections between self and other. Overall, these findings imply that an infants' multisensory integration of body-related information, including the integration of tactile stimuli with vision and proprioception, could provide the basis for establishing an early distinction between one's body and that of others (Cascio et al., 2012).

Bodily self-perception and interoception: The role of the insula

Although the integration of exteroceptive signals (i.e., proprioceptive and kinesthetic input informing about the movement of the body in space) plays a critical role in generating our sense of self, recent studies suggest the integration of interoceptive signals (i.e., neural mapping of multiple body states) is also crucial for bodily self-perception (Craig, 2009). Interoception involves the detection and processing of multiple signals from the body such as temperature, itch, pain, cardiac signals, respiration, hunger, thirst and touch – both discriminative and affective. Importantly, interoception has been associated with the generation of subjective feelings and self-perception (Damasio, 2010), in connection with insular cortex activation (Craig, 2002; Seth, 2013). In fact, it has been hypothesized that the anterior insula integrates sensorimotor signals and body schema networks (which involve different cerebral area such as parietal and frontal regions) creating a neural representation of the body into an embodied form of self-awareness (Namkung et al., 2017; Atzil et al., 2018). The insular cortex has a number of significant features. First, it is amongst the earliest cortical regions to develop and differentiate and is at an advanced

maturational stage from 27 weeks of gestation (Afif et al., 2007). This indicates that the insular cortex has an early involvement in the elaboration of somatosensory stimulation – again at an early maturational level from ~ 12 weeks gestational age, to which the fetus and the newborn are exposed. Second, insula neurons (as with cingulate neurons) have a specific cytoarchitecture: in contrast with pyramidal neurons which have a series of small basal dendrites that stretch from the cell body, these nerve cells, called spindle, or Von Economo neurons, are bipolar with only one bigger basal dendrite (Allman et al., 2005). Von Economo neurons are found in 4- month-old infants, increasing in number until four years of age and, interestingly, have also been identified in those mammals (such as big apes, some cetaceans and elephants) that use inter-individual contact to maintain affiliative behaviors. They are hypothesized to be implicated in body self-perception (Reiss and Marino, 2001; de Waal et al., 2005; Plotnik et al., 2006; Hakeem et al., 2009). Third, insula activity is central for affective and socio-cognitive information processing involving the self and others (Namkun et al., 2017). Lastly, this brain region is strongly involved in the regulation of homeostasis, in the elaboration of interoceptive inputs and in somatosensory perception with respect to pleasant tactile stimulations (Olausson et al, 2002; Björnsdotter et al., 2009). Moreover, while the anterior insular cortex has been implicated in the integration of autonomic and visceral signals into emotional and cognitive processes, the dorsal posterior insula supports primary cortical representations of ascending interoceptive pathways, reporting inputs (e.g., mechanical, thermal or chemical) from the skin, muscles, joints and internal organs (Craig, 2002). Importantly, the roles of the insular lobe for body-awareness in general (Craig, 2002, 2009), and of the right posterior insula in particular for the subjective experience of body-ownership has been well-established (Tsakiris, Schutz-Bosbach and Gallagher, 2007).

For the aim of this review it is critical to understand the development of insula functioning. Although the research is at the very beginning, recent neuroimaging studies targeting the insular

and somatosensory cortices in 2-month old babies report that slow stroking compared to fast stroking touch activated insular cortex, suggesting that this brain structure is functional at very early stages of postnatal life (Jönsson et al., 2018; Tuulari et al., 2019). Thus, since the first period of life the insula might integrate bodily and environmental information, providing a neural representation of bodily-self as a sentient entity (Namkun et al., 2017). In turn, this may constitute a basis for the elaboration of the distinction between the “inside” and the “outside” of the body, between “self” and “other” (Björnsdotter et al., 2009; Craig, 2009; Miguel et al., 2019). Nurturing touch could play a special role in this process. In fact, despite the exteroceptive origins of affectionate touch it has the capacity to simultaneously convey information about the “inner” body and the external world (Fotopoulou and Tsakiris, 2017).

The role of affectionate touch in infancy

When we are touched, or we touch something/somebody, the stimulus is processed in terms of its exteroceptive, discriminatory information content via fast conducting myelinated peripheral nerve pathways projecting to primary somatosensory cortical areas (McGlone and Reilly, 2010). However, an increasing body of evidence is establishing that there is a specialized sub-modality of touch sensing and processing its affective (i.e., rewarding, pleasant) properties (Essick et al., 2010; Morrison et al., 2010; Olausson et al., 2010). Recent neurophysiological studies have documented the existence of a dual-touch system consisting of two parallel neural pathways, one subserving discriminative touch and the other affective touch (McGlone et al., 2007; McGlone et al., 2014). The latter is encoded and processed by a recently characterized (in humans) population of mechanosensitive unmyelinated C-fibers that respond optimally to the gentle caressing touch, typical of nurturing care. These nerves, called C-tactile afferents (CT) project primarily to the dorsal posterior insula (Olausson et al., 2002; McGlone et al., 2014), and to brain networks involved in social perception processing (see below). CTs have only been found in the

hairy skin of the body and not in glabrous skin, such as the palmar skin of the hand. Despite this dual tactile transmission, the response to pleasant touch clearly assumes a particular valence in interpersonal relationships (the so-called 'social touch hypothesis', Olausson et al., 2002; Morrison et al., 2010). The preferred stimulus for CTs could not have been more appropriately evolved as they are tuned to respond to the specific velocities, forces and temperatures experienced by an infant during skin-to-skin nurturing care (Löken et al., 2009; Morrison, 2011; Ackerley et al., 2014; McGlone et al., 2014). CTs optimal firing response to gentle skin stroking, when recorded during microneurography studies (Löken et al. 2009), is between 1-10 cm/s, corresponding to self-reported pleasant perception of touch as measured psychophysically (Essick et al., 2010), leading to the hypothesis that CTs have evolved in all social mammals to encode the socio-affective and rewarding dimensions of touch (Morrison et al., 2011). Interestingly, a recent lesion study which has investigated deficits in the perceived affectivity of CT optimal touch has found that posterior and anterior right insula lesions reduce pleasantness sensitivity in perceiving CT-optimal touch (Kirsch et al., 2020). Cutaneous afferent C-fibers are a diverse population of unmyelinated, slowly conducting nerves that evolved primarily to provide a basic function of protection by detecting and transmitting information to the brain of negatively hedonic (i.e., nociceptive, pruriceptive) events from the skin of the body or viscera, or of positively hedonic (affiliative touch) events. C-fibers mature early in development: nociceptive stimuli evoke neural responses already at 25 weeks of gestation (Slater et al., 2006), and at 35-37 weeks, EEG brain responses are similar to those observed in adults (Fabrizi et al., 2011).

In adults, affective touch leads to widespread activation of a network of brain regions which comprise areas involved in social perception and social cognition, including the posterior insula, posterior superior temporal sulcus, medial prefrontal cortex, and dorsal anterior cingulate cortex (Bjornsdotter et al., 2014; Gordon et al., 2013; Olausson et al., 2002), amygdala and

orbitofrontal cortex (McGlone et al., 2007; Nees et al., 2019). Nevertheless, the emergence of this network in infancy is only partially understood. Research which has investigated infant brain response to affective touch during the first year of life leads to controversial findings. For example, using functional Near Infrared Spectroscopy (fNIRS) no increased responses for affective versus non-affective touch in posterior superior temporal sulcus and in prefrontal cortex regions were found in 5-month-old infants (Pirazzoli et al., 2019). On the other hand, while a few fNIRS studies indicate that the brain response to affective tactile stimuli emerges between 10 and 12 months of age (Miguel, Gonçalves, Cruz, and Sampaio, 2019; Miguel, Lisboa, Gonçalves and Sampaio, 2019; Miguel, Gonçalves and Sampaio, 2020), other neuroimaging research suggests that the affective touch system could be active very early in ontogeny (Jönsson et al., 2018; Tuulari et al., 2017). Notably, in children aged 5, maternal touch was predictive of resting activity in insula and in right posterior superior temporal sulcus, regions associated with the developing social brain and social perception (Brauer et al., 2016). Overall, these findings suggest that CT fibers could be crucial to convey socio-affective tactile information in the very early stages of development. Indeed, there are reasons to believe that affective touch is already being detected and processed during the early months of life. An infant's body is able to discriminate general tactile stimulation from pleasant tactile stimulation during development as they prefer tactile stimulations compatible with CT activation (Croy et al., 2019). A recent study reports that in young infants, sensitivity to pleasant touch is described by an inverted U-shaped function, similar to that observed in the adult, suggesting that a dual touch system might be potentially active in utero (Zieber et al., 2015), and the first months of life (Fairhurst et al., 2014). Particularly, 9-month-old infants displayed lower heart rate and greater behavioral engagement to body stimulation when they were stroked at a stroking velocity similar to the one that in adults activates CT fibers (Fairhurst et al., 2014). Furthermore, there is evidence that the optimal range for activating C-T fibers supports increasing

attention to contingent social information in 4-month-old infants (Della Longa et al., 2019). Moreover, compared to a non-stroking touch condition, during and after maternal stroking touch, the parasympatho-inhibitory regulation of infants increased while, intuitively, mothers stroked their infants at a CT afferent optimal velocity (Van Puyvelde et al., 2019). Thus, infants do not treat interpersonal touch as a purely mechanosensory event and, even more relevant, the infants' response to interpersonal touch is dependent on the source of the touch (Aguirre et al., 2019). In their study, Aguirre and colleagues (2019) found that infants' heart rate decreased more in response to stroking touch when their mother, rather than a stranger, was the source of the touch and, even more importantly, this effect was found only for CT-fiber-related tactile stimulation.

Importantly, unmyelinated CT afferents take a distinct ascending pathway from the periphery via spinal pathways to a different part of the thalamus than myelinated low threshold mechanoreceptors, projecting to dorsal posterior insular cortex (Olausson et al., 2002; Morrison et al. 2011; McGlone et al, 2014; Marshall et al., 2019). The signals conveyed via such affectionate touch provide, at the same time, bottom-up information about one's own body, as well as top-down information about the social contact and context. In other words, it simultaneously integrates information about 'the self' and about the external human environment. Thus, affective touch is here posited to be important not only in maintaining infant wellbeing and serving as a protective function by promoting affiliative behaviors (McGlone et al., 2014), but also in mediating the connection between self and other and developing bodily self-perception. In one study, using a preferential looking task, 5-month-old infants displayed a preference for the body-related visual-tactile synchrony when they were stroked at slow velocity, suggesting that affective CT-optimal touch might play a crucial role in the early development of bodily self-perception (Della Longa et al., 2020). Although CT affective touch might facilitate the process of discriminating the body boundaries of self (Fotopoulou and Tsakiris, 2017), early parent-infant embodied interactions

might also contribute to the experience of infant bodily-self through a key aspect of human caregiving, that is, interoceptive sensitivity, which in turn contributes to modulate self-other boundaries (Tajadura-Jiménez and Tsakiris, 2014).

Beyond a conventional view of maternal sensitivity

Implicit neurobehavioral sensitivity to interoceptive signals emerges in early infancy and it appears sensitive to emotional processing (Maister et al., 2017). Thus, it is highly plausible that, at least partially, in early infancy interoception is affected by interpersonal exchanges. A recent retrospective study found that young adults classified as demonstrating signs of avoidant attachment, based on attachment theory (Bowlby, 1969), reported lower interoceptive awareness, suggesting that the quality of early interpersonal relationships might be associated with later interoceptive functioning (Oldroyd et al., 2019). While research has mainly focused on the infant's interoceptive functioning it would not be surprising to find that a mother's interoceptive functioning could also play a role in the early exchanges with their infant. Indeed, while the mother may not always be aware of her own body signals, it is highly unlikely that embodied interactions with the infant would not impact their own bodily experiences, which imply an array of bodily tension, heart-beats, breathing and tactile pleasantness links to these sensations. Importantly, although interoceptive sensitivity does not imply full consciousness and the need for verbal expressions, the ability to represent one's own internal body state has been found to be associated with social attitudes (Ferri et al., 2013). Likewise, one could expect ventral-ventral parent-infant contact (e.g., skin-to-skin contact) and others embodied interactions involve activation of the mother's bodily sensations (see below), with possible modifications of interoceptive sensitivity which supports moment-by-moment adaptation to interactive demands and infant' needs. The focus on parental interoceptive functioning during early interactions sheds a different light on parental sensitivity. Over the past decades a mothers' sensitivity has been

considered a crucial aspect of the mother-infant relationship. Research in this field has devoted much attention to maternal sensitivity as the ability to respond promptly and in an active, warm, acceptant, appropriate and flexible way to the infant's signals (Kivijärvi et al., 2001). Furthermore, there is evidence that maternal sensitivity, at least partially, relies on perceptual processing of infants' body. For example, a recent study has explored the association between maternal sensitivity and the refinement of maternal visuo-perceptual processes in perceiving infants' body (Montirosso et al., 2016). Authors found that a more refined maternal perception of own infant's body cues was associated with higher maternal sensitivity.

Although traditional measures of maternal sensitivity implies higher-order social cognitive skills such as mentalization, recent studies highlight that parenting heavily relies on nonverbal, bodily based and biological co-regulation that have been referred to as *parental embodied mentalization* (PEM, i.e., the parental capacity to comprehend the infant's mental states from their body movement and adjust their own kinesthetic patterns accordingly (Shai and Belsky, 2011). Notably, infants of mothers displaying greater PEM during a mother–infant free-play at 6 months showed secure attachment at both 15 and 36 months; and at 54 months displayed greater social competence, fewer behavioral problems and greater cognitive functioning (Shai and Belsky, 2011). Interestingly, this prediction held even after accounting for traditional measures of maternal sensitivity. It should be noted that a high-order maternal sensitivity concept based on mentalization might be only partially paralleled and mimicked by the low-order and bodily based parenting level. Indeed, some recent findings suggested that maternal non-attuned mind-related comments were associated with touch behaviors that were not contingent with the infant's emotions (Crucianelli et al., 2018). Consistently, while traditional measures of maternal sensitivity did not significantly predict infant attachment security, affectionate touch during infant feeding at 3 months of age was associated with more secure attachment to their mother at 1 year of age

(Weiss et al., 2000). Interestingly, in their study the authors found that mother's felt security regarding their own tactile experience as a child, increased her infant's chances of having a secure attachment. A variety of maternal behaviors associated with providing a secure base provision, such as being available for connection, allowing infant physical proximity, providing emotional regulation through soothing physical contact assessed at 4.5 months significantly predicted infant attachment at 12 months, with an effect size 8 times larger than maternal sensitivity (Woodhouse et al., 2020). Collectively, these findings provide initial evidence for a conceptual shift in our understanding of maternal caregiving which goes beyond the conventional view of maternal sensitivity and is more closely associated with the embodied interaction view (Botero et al., 2019).

From this perspective, while appropriate parenting requires coordination of multiple systems, including both behavioral and physiological components, to read infant signals and respond sensitively (Barrett and Fleming, 2011; Welch, 2016), the mother's ability to perceive her own internal signals might further support the responsivity to her infant's cues. This view is corroborated by evidence suggesting that interoceptive sensitivity is not only crucial for the processing of emotional experience and self-regulation (Herbert and Pollatos, 2012), but most importantly it is associated with the ability to understand emotion in others (i.e., empathy; Fukushima et al., 2011). Moreover, a recent study found that activation of the parent's anterior insula (i.e., the cerebral hub which supports interoceptive sensitivity) in response to a video of self-interacting with his/her own infant predicted lower somatic problems in the child six years later, and this link was mediated by the parent's sensitive behavior at age four (Abraham et al., 2019). Remarkably, intuitive maternal stroking is not only tuned to the stimulation of CT fibers in infants aged 4 to 54 weeks, but stroking velocities were significantly correlated to maternal interoception, as measured by heart rate, suggesting that maternal affectionate touch behavior might be moderated by some aspects of maternal interoception (Bytowski et al., 2020).

Interestingly, mothers' interoceptive knowledge (i.e., parent knowledge of their own and other people's interoceptive sensations) about their own emotions has been found to be associated with social affective skills (i.e., emotion regulation, social initiative, cooperation, self-control) in middle childhood (MacCormack et al., 2019). Thus, if maternal sensitivity were, at least partially, associated with the accuracy with which mothers perceive their own bodily signals, then maternal interoceptive sensitivity might facilitate parental inferences about infant's physiological and emotional states and infant's ability to form accurate perceptions of bodily sensations. Mothers could be able to teach infants interoceptive cues which could help them differentiate bodily states (i.e., distress, hunger, tiredness, etc.). Furthermore, as suggested by studies which have documented a link between interoception and empathy (Fukushima et al., 2011), interoceptive sensitivity might support a mothers' ability to attune to her infant's bodily states, for managing these own bodily cues when infant experiences emotion and interacts with others, thereby modeling her emotion regulation and social skills.

Embodied reparation: A putative

process subserving experiences of embodied interactions

Based on existing evidence from developmental research, it stands to argue that in the early embodied interactions, the cycles of mutual regulation should imply both body attunement and body misattunement between mother and infant.

A number of studies suggest that during the first months of life, mothers and infants are able to reciprocally adjust their own physiological activities, above all when they are in close body contact, a sort of physiological body-to-body synchrony (i.e., bodily attunements or co-regulation; Neu et al., 2009; Atzil and Gendron, 2017). For example, a study documented thermal synchrony during skin-to-skin holding of twins (Ludington-Hoe et al., 2006). Notably, given that twins were held at the same time in skin-to-skin contact, each maternal breast temperature reacted to the

thermal needs of the infant on that breast. Maternal-infant heart rate variability attunement was reported during the first months of postnatal life (Feldman et al., 2011). Remarkably, during simple continuous skin-to-skin contact (i.e., the mothers were asked to hold their infants' hand or foot) 2-months-old infants adjust their heart rate to fluctuations of their mothers' cardiorespiratory activity (Van Puyvelde et al., 2015). Collectively, these findings indicate that mother-infant bodily attunements are established in various forms in the first months of life.

While it has been largely documented that bodily attunement is crucial for mother-infant interactions (Shai and Belsky, 2011), the potential role of misattunement conditions of psychobiological states have received less attention. Nevertheless, early adverse experiences (e.g., prematurity) or mother/infant characteristics (e.g., presence of at-risk conditions) might lead to bodily misattunement between partners. Some interesting results comes from studies using the Face-to-Face Still-Face paradigm (DiCorcia and Tronick, 2011), which allows for accurate measurement of infants' and mothers' reactions highlighting the impact of a misattunement process. For example, Ham and Tronick (2006) found that reciprocal regulatory effects of infant and mother heart rate variability were affected during the still-face episode in which the mother is temporarily unresponsive. A recent study has documented a complex pattern of coordination of hypothalamic-pituitary-adrenal (HPA) axis between infants and their mothers while facing a social stress associated maternal unavailability (Provenzi et al., 2019). Before the onset of the stressful condition the two partners presented coupled salivary cortisol concentration. Soon after the experimental manipulation of maternal behavior and the occurrence of a socio-emotional stress condition, non-significant cortisol coupling was found between mothers and infants suggesting that the dyad, like a system, was facing a perturbation and facing a challenge in their previously observed biological coordination. Importantly, this pattern of coordination of HPA axis was altered

between preterm infants and their mothers, suggesting that in at-risk development conditions, partners were less able to dynamically coordinate HPA axis regulation (Provenzi et al., 2019).

At this point, one could wonder how infants and their mothers might reach bio-behavioral synchrony. Although it remains largely unclear how infants and their caregivers synchronize, based on infant behavioral research, we propose that there is a putative process (i.e., *embodied reparation* see below) which could serve a similar function in facilitating body-to-body synchrony. In the Mutual Regulation Model (MRM), Tronick (2004) suggests that in typical interactions mothers and their infants fluctuate between attuned and misattuned states and recovery attunement states by a process called ‘reparation’. Indeed, mother–infant interactions during the first months of life are characterized by frequent misattunements (so called, interactive ruptures). Because misattunements are the rule rather than the exception (Tronick and Cohn, 1989), dyadic reparations needed to gain new dyadic states of attunements after normally occurring interactive ruptures (Tronick, 2004). Behavioral data suggest that, during face-to-face interactions, mother and infant are in misattunement states for almost 70% of time (Tronick and Cohn, 1989). Thus, dyadic reparation is critical for the development of infants’ self-regulatory capacities and scaffolding an implicit inner sense of self-efficacy in regulating the stress elicited by socially challenging conditions (DiCorcia and Tronick, 2011). For example, Provenzi et al. (2015) reported that 4-month-old infants who experience more frequent dyadic reparation within the interaction with their mother were more prone to the vagal reactivity (i.e., the vagal brake) in response to a social disruption condition (i.e., temporary maternal unavailability during the still-face paradigm), which is thought to reflect an adaptive parasympathetic functioning and effective stress regulation (Porges, 2007).

While, based on existing evidence, bodily attunements is through enhanced social connectedness, effective communication and interpersonal regulation (Atzil et al., 2018), there is a

lack of theoretical ground and conceptual framework guiding the body-to-body perspective of mother-infant interaction which takes into account early forms of bodily misattunement and reparation. Critically, it is necessary to recognize that the maintenance of a constant bodily attunement between two partners is rather improbable, but also if it were so it would prevent any possibility of bodily misattunement, thus leading the infant to never experience a reparatory step. As mentioned, bodily attunements and bodily misattunement processes, separately considered, could be not sufficient to delineate how mothers and infants reciprocally adjust their own physiological activities. Here, we propose that inevitable bodily misattunement may be followed by a reparatory step, the process which focuses on the phase of transforming bodily misattunement states into bodily attunements. The construct of *embodied reparation* describes the process by which each partner modifies their own bodily states thereby making possible the shifting from bodily misattunement to bodily attunements (Fogel, 2011). It is likely that this process occurs, dynamically and continuously, at multiple bio-behavioral levels including the micro-temporal level where the mother regulates the infant's neurophysiological processes.

Of course, the shift from bodily misattunement to bodily attunement could be more evident when the infant experiences moments of discomfort, or tension or distress during which the infant's interoceptive system is strongly dysregulated (Welch, 2016). Recent neuroscience models suggest that interoceptive experience might reflect predictions about the expected state of the body (Barrett and Simmons, 2015). In infants and in mothers moment-by-moment prediction anticipates inputs related to sensory information from exteroceptive and interoceptive information and which would be the best potential action to manage this information (Atzil et al., 2018). Any discrepancy from the expected state of the body is a prediction error which generates a physiological unbalance in the infant's body (for instance, infant experiences low skin temperature). While infants have self-regulation ability to cope with distress (e.g., non-nutritive

sucking), their resources are limited. As a consequence, the manifestation of such physiological changes emerges as bodily, behavioral and emotional signals which convey to the mother the infant's state. Although a mother uses several behaviors to comfort her infant (i.e., voice, eye contact) and to restore emotional connection and physiological co-regulation (Atzil et al., 2018), we speculate physical proximity (e.g., affectionate/nurturing touch) and maternal interoceptive sensitivity play a key role in *embodied reparation* process. Indeed, physical proximity and maternal interoceptive sensitivity would support a sort of data processing strategy in which discrepancies between psychobiological states of partners can be minimized, either by performing actions (i.e., affectionate touch) to modify the partner bodily states to fit the own bodily states or by changing the own bodily states to fit the partner bodily states (Seth and Tsakiris, 2018). By interoceptive sensitivity a mother can implicitly adapt to her infant's body state by, for example, adjusting her own skin temperature to that of the infant (Ludington-Hoe et al., 2006). In other words, *embodied reparation* allows the infant to reduce the discrepancies within the interoceptive system using the mother's body. Thus, interoceptive information of the infant's body is gradually associated with inputs from the mother (Atzil et al., 2018) suggesting that an efficient *embodied reparation* allowing the infant to recover a stable interoceptive experience, which, in turn, would support positive experience of the own bodily-self.

As for physical proximity, while no infant study has researched this topic, there is indirect evidence from an adult study in which affectionate touch during pain administration increased bodily attunement. Specifically, during pain administration, partner touch increased interpersonal respiration coupling (Goldstein et al., 2017) and interpersonal neural synchronization in brain networks associated with analgesia (Goldstein et al., 2018). Overall, this evidence indicates that social touch not only increases interpersonal physiological and brain-to-brain coupling, but

promotes affiliative bonding and a positive subjective self-experience (Krahé et al., 2016; Krahé et al., 2018).

Furthermore, although it is quite possible that each partner contributes to the *embodied reparation* process, the infant has limited and immature regulatory, behavioral, and attentional capacities to repair the misattunements, so that most *embodied reparations* could be managed by the parent. The mother's ability to quickly perceive and respond to the infant's bodily signals depends on a subtle balance between the awareness of her own body signals and those of the infant's (Abraham et al., 2019). As a consequence, maternal interoceptive sensitivity is an essential component to adjust own bodily sensations to support infants' regulatory capacities and to restore physiological co-regulation.

Importantly, early adverse experiences, such as premature birth, can have a negative impact on dyadic co-regulation, affecting the *embodied reparation* process and therefore have an impact on the development of the infant bodily-self. For instance, along with other stressful experiences, preterm infants experience physical restraint which limits the possibility of contact with the peripersonal space (e.g., clothes, sheets) as well as of self-contact (Durier et al., 2015). Moreover, whereas their tactile and proprioceptive systems receive minimal input, preterm infants are exposed to atypical procedural handling routines, altered stimulations and multiple stressful procedural interventions which are associated with long-term changes in somatosensory function or pain response (Walker et al., 2018). Preterm birth is also a stressful event for mothers and may affect the mother–preterm infant relationship (Korja et al., 2012). In fact, preterm birth is associated not only with a less-than-optimal maternal bonding (Provenzi et al., 2017), but also can affect the mother's ability to perceive her infants' bodily cues. For example, mothers of preterm infants are less efficient in perception of their infants' bodily cues (Butti et al., 2018). Collectively, these conditions could limit opportunities for successful repair of bodily misattunements between

mother and infant. As a consequence, the extent to which *embodied reparations* are disrupted, preterm infants could develop a distorted sense of interoception, which in turn could have an impact on the development of the bodily-self. Notably, a recent study has documented that preterm birth might interfere with the development of body representations. Specifically, school-aged children born preterm without neurological sequelae exhibited an impairment of body representation assessed by a visual body recognition task (Butti et al., 2020).

In short, bodily attunements, bodily misattunements and *embodied reparation* provide much more than the infant's behavior regulation. To some extent they sculpt the infant's bodily-self. Empirical assessment of the relationship between these processes and problematic interoception and bodily perception is warranted

Concluding remarks and future perspectives

Infants have the capacity to perceive their own body, for example using sensorimotor contingencies or by self-contact. However, these capacities are unlikely to depend only on a single process but rather are integrated by inter-corporeal connections. The development of an infant's bodily-self is an interactive phenomenon and interpersonal nurturing touch, as processed by CTs and by the brain areas they project to, is here posited to lead to increased interpersonal bodily attunement, which in turn supports the representation of the infant bodily-self (Filippetti et al., 2013; Bremner and Spence, 2017). Moreover, it has been hypothesized that an altered processing in CT stimulation (McGlone et al., 2014) and in social brain network including insula (Allman et al., 2005) might underpin neurodevelopmental conditions currently grouped under the classification of autism spectrum disorders which are characterized by an atypical sense of self (Lombardo et al., 2009). Thus, it seems conceivable that the infant is not only able to synchronize with external stimuli (Provasi et al., 2014), but, during embodied interactions with their mother, is able to encode maternal affectionate touch, specifically through CTs (Aguirre et al., 2019) and to

coordinate with maternal somatic inputs (i.e., cardiorespiratory patterns, body temperature, etc.) (e.g., Van Puyvelde et al., 2019). Importantly, it is plausible that affective touch and close physical proximity allow the dyad to attune both during pleasant exchanges and, even more importantly, after a bodily misattunement (i.e., *embodied reparation*). The renewed bodily attunement would regulate the infant's interoceptive states (e.g., decreases heart rate) and support bodily states which facilitate the interoceptive sensitivity and therefore the infant's bodily-self-perception.

Maternal interoceptive sensitivity may be another way for mothers to support bodily attunement into their embodied interactions with infants. There is now neuroimaging evidence suggesting that interoceptive awareness significantly enhances neural activity during empathy in bilateral anterior insula (Ernst et al., 2013). Thus, we speculate that maternal interoceptive sensitivity expands a mother's ability to perceive their infant's body, which in turn would affect the mother's social engagement and emotional responsiveness. Therefore, the focus on parental interoceptive sensitivity in early interactions offers the theoretical framework as well as one practical access of research to investigate the contribution of the mother's body in infant bodily-self-development.

Additionally, while relational repairs by a sensitive mother are crucial for socio-emotional development (Tronick, 2004), *embodied reparations* are here posited to be critical for building a representation of the infant bodily-self. In this perspective, less embodied reparations would impair the infant's ability to form accurate representations of bodily sensations. Thus, naturally occurring variations in *embodied reparations* during exchanges with the mother's body would contribute to both infant bodily self-perception and interactive embodied meaningful patterns. Therefore, individual differences in tactile biography may determine not only how an infant will respond to touch, but also, at least partially, how the infant will perceive its own bodily-self. In other words, specific patterns both of embodied interactions (e.g., *how*, *when*, *how much* and

where the mothers touch their infants) and *embodied reparations* (i.e., *how, when, how much* and *in which circumstances* dyad actively shifts from bodily misattunement to bodily attunement) could become more and more a part of the workings of the dyadic interactive processes which sculpt the sense of infant bodily-self. This framework has implications about the role of, specifically, CT touch both in clinical research in children with atypical development (Provenzi et al., 2020) and for promoting the use of affective touch in early parental interventions (Botero et al., 2019). For instance, video-feedback intervention in which mothers watch videos of their interaction with their at-risk infants, helps them not only to become more aware of their infants' signals and to interpret these signals more accurately, but also makes them aware of the impact of misattunement states on the interaction (Giusti et al., 2018).

The opinions suggested here have implications for research in developmental neuroscience and future research is necessary to examine the implications related to our hypothesis. A first point of interest involves affective touch as a factor supporting infant's bodily-self perception. How much affectionate touch would be appropriated? Do adverse experiences affect the affectionate touch processing? For example, currently we are studying the brain responses to gentle skin stroking, a type of tactile stimulus associated with affective touch, in the postcentral gyrus and posterior insular cortex - two cerebral regions known to process this kind of tactile stimulation – in preterm infants. Second, if maternal interoceptive sensitivity could have a critical role to adjust own bodily sensations to support infants' regulatory capacities and to restore physiological co-regulation, then we need further evidence in order to corroborate that mothers' interoceptive sensitivity is related to infant's bodily-self perception. Finally, despite the above insights, embodied interaction has remained generally understood to be a unidirectional process (i.e., from mother to infant), this assumption is however challenged by evidence that even the mother's body is affected by the inter-corporal connections with her own infant (Matthiesen et al.,

2001), and parents experience the kinetics and intensity of body sensations through the proximity of affective touch mechanisms. How does the infant's touch affect the mother's bodily-self perception? How does this inter-corporality process modify maternal interoceptive sensitivity? Furthermore, future research is needed to investigate mother and infant interoception during the first period of life. Given the evidence that infants have an implicit neurobehavioral sensitivity to interoceptive signals (Maister et al., 2017), it would be important to examine how individual differences in interoception can develop in connection with maternal interoceptive sensitivity. More in general, in order to investigate the role of *embodied reparations*, research involving behavioral, physiological and neural approach should be performed at various ages and conditions to establish the dynamics of the interplay between maternal touch, maternal interoceptive sensitivity and infant's bodily-self perception.

In recent years, there has been an increasing interest in the pre-reflective bodily foundations of the self, which are thought to be grounded in our embodied experience of being a separate entity (Damasio, 2000), providing a rudimental awareness of ourselves (Gallagher, 2000; Allen and Tsakiris, 2019). Therefore, the "self starts with body" (Baumeister, 1999) and the bodily experience is ubiquitous in everyday experience, above all for an infant. However, given the primacy of the embodied interactions at the very first stages of human life, this view should be integrated with a further consideration: an infant's bodily self-perception emerges from the dynamic interplay between signals arising from both (a) inside the body and (b) interpersonal affective exchanges with mother. The latter is a very critical point given that, as Ciaunica (2019) highlighted, before an infant "meets" mother's mind, the infant "meets" mother's body, a process that critically depends on affective touch, as mediated by CTs. Moreover, to some extent, the contrary is equally true, before mother "meets" infant's mind, mother "meets" infant's body. Thus, bodies come first and along with several other interactive processes, *embodied reparations*

make embodied meaning within and between individuals, which increase the coherence and complexity of each infant's and, likely parent's sense of bodily-self establishing reciprocal awareness.

Declarations of interest: none.

Funding: This study was partially supported by Italian Ministry of Health (Ricerca Finalizzata RF-2016-02361884 and Ricerca Corrente 2018-2019-2020) to RM.

Acknowledgments.

The first author is deeply indebted with Ed Tronick, whom scientific work on early mother-infant relationship has inspired the view discussed in the current review. We would like to thank Eleonora Mascheroni, Eleonora Visintin, Elisa Rosa, Niccolò Butti and Elena Guida for their valuable support at *0-3 Center for the at-Risk Infant*, Scientific Institute IRCCS Eugenio Medea, Bosisio Parini (Lecco), Italy. The authors are grateful to anonymous reviewers for helpful comments.

References

- Abraham, E., Hendler, T., Zagoory-Sharon, O., Feldman, R. (2019). Interoception sensitivity in the parental brain during the first months of parenting modulates children's somatic symptoms six years later: The role of oxytocin. *International Journal of Psychophysiology*, 136, 39-48.
- Ackerley, R., Wasling, H. B., Liljencrantz, J., Olausson, H., Johnson, R. D., Wessberg, J. (2014). Human C-tactile afferents are tuned to the temperature of a skin-stroking caress. *Journal of Neuroscience*, 34(8), 2879-2883.
- Afif, A., Bouvier, R., Buenerd, A., Trouillas, J., Mertens, P. (2007). Development of the human fetal insular cortex: Study of the gyration from 13 to 28 gestational weeks. *Brain Structure and Function*, 212(3-4), 335-346.
- Aguirre, M., Couderc, A., Epinat-Duclos, J., Mascaro, O. (2019). Infants discriminate the source of social touch at stroking speeds eliciting maximal firing rates in CT-fibers. *Developmental Cognitive Neuroscience*, 36, 100639.
- Allen, M., Tsakiris, M. (2019). The body as first prior: Interoceptive predictive processing and the primacy. In *The Interoceptive Mind: From Homeostasis to Awareness* (First Edition, pp. 27–45). Great Clarendon Street, Oxford, OX2 6DP: Oxford University Press.
- Allman, J.M., Watson, K., K., Tetreault N.,A., Hakeem A.Y. (2005). Intuition and autism: A possible role for Von Economo neurons. *Trends in Cognitive Sciences*. 9, 367-373.
- André, V., Henry, S., Lemasson, A., Hausberger, M., Durier, V. (2018). The human newborn's umwelt: Unexplored pathways and perspectives. *Psychonomic Bulletin & Review*, 25(1), 350-369.

- Atzil, S., Gendron, M. (2017). Bio-behavioral synchrony promotes the development of conceptualized emotions. *Current Opinion in Psychology*, 17, 162-169.
- Atzil, S., Gao, W., Fradkin, I., Barrett, L. F. (2018). Growing a social brain. *Nature Human Behaviour*, 2(9), 624-636.
- Barrett, J., Fleming, A. S. (2011). Annual research review: All mothers are not created equal: Neural and psychobiological perspectives on mothering and the importance of individual differences. *Journal of Child Psychology and Psychiatry*, 52(4), 368-397.
- Barrett, L. F., Simmons, W. K. (2015). Interoceptive predictions in the brain. *Nature Reviews Neuroscience*, 16(7), 419.
- Baumeister, R. (1999). The nature and structure of the self: An overview. In R. Baumeister (Ed.), *The self in social psychology* (pp. 1–21). Philadelphia, PA: Psychology Press.
- Björnsdotter, M., Löken, L., Olausson, H., Vallbo, Å., Wessberg, J. (2009). Somatotopic organization of gentle touch processing in the posterior insular cortex. *Journal of Neuroscience*, 29(29), 9314-9320.
- Björnsdotter, M., Gordon, I., Pelphrey, K. A., Olausson, H., & Kaiser, M. (2014). Development of brain mechanisms for processing affective touch. *Frontiers in Behavioral Neuroscience*, 8, 1-10. 10.3389/fnbeh.2014.00024
- Botero, M., Langley, H. A., and Venta, A. (2019). The untenable omission of touch in maternal sensitivity and attachment research. *Infant and Child Development*, 1-24 <https://doi.org/10.1002/icd.2159>
- Bowlby, J. (1969). *Attachment and loss: Vol 1. Attachment*. New York: Basic Books.
- Brauer, J., Xiao, Y., Poulain, T., Friederici, A. D., & Schirmer, A. (2016). Frequency of maternal touch predicts resting activity and connectivity of the developing social brain. *Cerebral Cortex*, 26(8), 3544-3552.
- Bremner, A.J., Spence, C. (2017). The development of tactile perception. *Advances in Child Development and Behavior*, 52, 227-268. doi.org/10.1016/bs.acdb.2016.12.002
- Burr, D., Gori, M. (2012). Multisensory integration develops late in humans. In *The neural bases of multisensory processes*. In: Murray MM, Wallace MT, editors. *The Neural Bases of Multisensory Processes*. Boca Raton (FL): CRC Press/Taylor and Francis.
- Butti N., Urgesi, C., Montiroso R. (2020). Premature birth affects visual body representation and body schema in preterm children. *Unpublished manuscript*.
- Butti, N., Montiroso, R., Borgatti, R., Urgesi, C. (2018). Maternal sensitivity is associated with configural processing of infant's cues in preterm and full-term mothers. *Early Human Development*, 125, 35-45.

- Bytowski, A., Ritschel, G., Bierling, A., Bendas, J., Weidner, K., & Croy, I. (2020). Maternal stroking is a fine-tuned mechanism relating to C-tactile afferent activation: An exploratory study. *Psychology & Neuroscience*. Advance online publication. <https://doi.org/10.1037/pne0000184>
- Cascio, C. J., Moana-Filho, E. J., Guest, S., Nebel, M. B., Weisner, J., Baranek, G. T., and Essick, G. K. (2012). Perceptual and neural response to affective tactile texture stimulation in adults with autism spectrum disorders. *Autism Research*, 5(4), 231-244.
- Caspers, S., Zilles, K., Laird, A. R., & Eickhoff, S. B. (2010). ALE meta-analysis of action observation and imitation in the human brain. *Neuroimage*, 50(3), 1148-1167.
- Ciaunica, A. (2019). The 'meeting of bodies': Empathy and basic forms of shared experiences. *Topoi. An International Review of Philosophy*, 38, 185-195. doi.org/10.1007/s11245-017-9500-x
- Ciaunica, A., Crucianelli, L. (2019). Minimal self-awareness: From within a developmental perspective. *Journal of Consciousness Studies*, 26(3-4), 207-226.
- Craig, A.D. (2002), How do you feel? Interoception: The sense of the physiological condition of the body. *Nature Reviews Neuroscience*, 3, 655-666.
- Craig, A. D. (2009). Emotional moments across time: a possible neural basis for time perception in the anterior insula. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 364(1525), 1933-1942.
- Croy, I., Sehlstedt, I., Wasling, H. B., Ackerley, R., & Olausson, H. (2019). Gentle touch perception: From early childhood to adolescence. *Developmental Cognitive Neuroscience*, 35, 81-86.
- Crucianelli, L., Filippetti, M. L. (2018). Developmental perspectives on interpersonal affective touch. *Topoi. An International Review of Philosophy*. 1-12. doi.org/10.1007/s11245-018-9565-1
- Crucianelli, L., Krahé, C., Jenkinson, P. M., Fotopoulou, A. K. (2018). Interoceptive ingredients of body ownership: Affective touch and cardiac awareness in the rubber hand illusion. *Cortex*, 104, 180-192.
- Crucianelli, L., Wheatley, L., Filippetti, M. L., Jenkinson, P. M., Kirk, E., Fotopoulou, A. K. (2019). The mindedness of maternal touch: An investigation of maternal mind-mindedness and mother-infant touch interactions. *Developmental cognitive neuroscience*, 35, 47-56.. <https://doi.org/10.1016/j.dcn.2018.01.010>
- Damasio, A. (2000). *The feeling of what happens: Body and emotion in the making of consciousness*. Fort Washington, PA: Harvest Books.
- Della Longa, L., Gliga, T., & Farroni, T. (2019). Tune to touch: Affective touch enhances learning of face identity in 4-month-old infants. *Developmental Cognitive Neuroscience*, 35, 42-46. [doi: 10.1016/j.dcn.2017.11.002](https://doi.org/10.1016/j.dcn.2017.11.002)

- Della Longa L., Filippetti M.L., Dragovic D. and Farroni T. (2020). Synchrony of caresses: Does affective touch help infants to detect body-related visual-tactile synchrony? *Frontiers in Psychology* 10:2944.doi: 10.3389/fpsyg.2019.02944
- de Waal, F.B.M., Dindo, M., Freeman, C.A., Hall, M. (2005). The monkey in the mirror: Hardly a stranger. *Proceedings of the National Academic Sciences USA*, 102, 11140-11147.
- DiCorcia, J. A., Tronick, E. D. (2011). Quotidian resilience: Exploring mechanisms that drive resilience from a perspective of everyday stress and coping. *Neuroscience and Biobehavioral Reviews*, 35(7), 1593-1602.
- DiMercurio, A. J., Connell, J. P., Clark, M., Corbetta, D. (2018). A naturalistic observation of spontaneous touches to the body and environment in the first 2 months of life. *Frontiers in Psychology*, 9, 2613. doi: 10.3389/fpsyg.2018.02613
- Dunn, B. D., Galton, H. C., Morgan, R., Evans, D., Oliver, C., Meyer, M., Cusack, R., Lawrence, A.D., Dalgleish, T. (2010). Listening to your heart: How interoception shapes emotion experience and intuitive decision making. *Psychological Science*, 21(12), 1835–1844.
- Durier, V., Henry, S., Martin, E., Dollion, N., Hausberger, M., Sizun, J. (2015). Unexpected behavioural consequences of preterm newborns' clothing. *Scientific Reports*, 5, 9177.
- Ehrsson, H. H., Holmes, N. P., & Passingham, R. E. (2005). Touching a rubber hand: feeling of body ownership is associated with activity in multisensory brain areas. *Journal of Neuroscience*, 25(45), 10564-10573.
- Ernst, J., Northoff, G., Böker, H., Seifritz, E., Grimm, S. (2013). Interoceptive awareness enhances neural activity during empathy. *Human Brain Mapping*, 34(7), 1615-1624.
- Essick, G. K., McGlone, F., Dancer, C., Fabricant, D., Ragin, Y., Phillips, N., Jones, T., Guest S. (2010). Quantitative assessment of pleasant touch. *Neuroscience and Biobehavioral Reviews*, 34(2), 192-203. doi.org/10.1016/j.neubiorev.2009.02.003
- Fabrizi, L., Slater, R., Worley, A., Meek, J., Boyd, S., Olhede, S., Fitzgerald, M. (2011). A shift in sensory processing that enables the developing human brain to discriminate touch from pain. *Current Biology*, 21(18), 1552-1558.
- Fairhurst, M. T., Löken, L., Grossmann, T. (2014). Physiological and behavioral responses reveal 9-month-old infants' sensitivity to pleasant touch. *Psychological science*, 25(5), 1124-1131.
- Feldman, R. (2012). Parent–infant synchrony: A biobehavioral model of mutual influences in the formation of affiliative bonds. *Monographs of the Society for Research in Child Development*, 77(2), 42-51.
- Feldman, R., Magori-Cohen, R., Galili, G., Singer, M., Louzoun, Y. (2011). Mother and infant coordinate heart rhythms through episodes of interaction synchrony. *Infant Behavior and Development*, 34(4), 569-577.

- Feldman, R., Rosenthal, Z., Eidelman, A. I. (2014). Maternal-preterm skin-to-skin contact enhances child physiologic organization and cognitive control across the first 10 years of life. *Biological Psychiatry*, 75(1), 56-64.
- Ferri, F., Ardizzi, M., Ambrosecchia, M., and Gallese, V. (2013). Closing the gap between the inside and the outside: Interoceptive sensitivity and social distances. *PLoS One*, 8(10), e75758.
- Filippetti, M. L., Johnson, M. H., Lloyd-Fox, S., Dragovic, D., Farroni, T. (2013). Body perception in newborns. *Current Biology*, 23(23), 2413-2416.
- Filippetti, M. L., Lloyd-Fox, S., Longo, M. R., Farroni, T., & Johnson, M. H. (2015). Neural mechanisms of body awareness in infants. *Cerebral Cortex*, 25(10), 3779-3787.
- Filippetti, M. L., Orioli, G., Johnson, M. H., Farroni, T. (2015). Newborn body perception: sensitivity to spatial congruency. *Infancy*, 20(4), 455-465.
- Fogel, A. (2011). Embodied awareness: Neither implicit nor explicit, and not necessarily nonverbal. *Child Development Perspectives*, 5(3), 183-186.
- Fotopoulou, A., Tsakiris, M. (2017). Mentalizing homeostasis: The social origins of interoceptive inference. *Neuropsychoanalysis*, 19(1), 3-28.
- Fukushima, H., Terasawa, Y., Umeda, S. (2011). Association between interoception and empathy: evidence from heartbeat-evoked brain potential. *International Journal of Psychophysiology*, 79(2), 259-265.
- Gallagher, S. (2000). Philosophical conceptions of the self: implications for cognitive science. *Trends in cognitive sciences*, 4(1), 14-21. doi.org/10.1016/S1364-6613(99)01417-5
- Gallese, V. (2007). Before and below 'theory of mind': Embodied simulation and the neural correlates of social cognition. *Philosophical Transactions of the Royal Society B*, 362(1480), 659-669.
- Garfinkel, S. N., Seth, A. K., Barrett, A. B., Suzuki, K., & Critchley, H. D. (2015). Knowing your own heart: distinguishing interoceptive accuracy from interoceptive awareness. *Biological psychology*, 104, 65-74.
- Gentsch, A., Crucianelli, L., Jenkinson, P., Fotopoulou, A. (2016). The touched self: Affective touch and body awareness in health and disease. In *Affective touch and the neurophysiology of CT afferents* (pp. 355-384). Springer New York.
- Georgie, Y. K., Schillaci, G., Hafner, V. V. (2019). An interdisciplinary overview of developmental indices and behavioral measures of the minimal self. *arXiv preprint arXiv:1907.00709*.
- Gergely, G., Watson, J. S. (1999). Early socio-emotional development: Contingency perception and the social-biofeedback model. *Early social cognition: Understanding others in the first months of life*, 60, 101-136.

- Giusti, L., Provenzi, L., Montirosso, R. (2018). The Face-to-Face Still-Face (FFSF) paradigm in clinical settings: Socio-emotional regulation assessment and parental support with infants with neurodevelopmental disabilities. *Frontiers in Psychology*, 9, 789.
- Goldstein, P., Weissman-Fogel, I., Dumas, G., Shamay-Tsoory, S. G. (2018). Brain-to-brain coupling during handholding is associated with pain reduction. *Proceedings of the national academy of sciences*, 115(11), E2528-E2537.
- Goldstein, P., Weissman-Fogel, I., Shamay-Tsoory, S.G. (2017). The role of touch in regulating inter-partner physiological coupling during empathy for pain. *Scientific reports*, 7(1), 3252.
- Gordon, I., Voos, A. C., Bennett, R. H., Bolling, D. Z., Pelphrey, K. A., & Kaiser, M. D. (2013). Brain mechanisms for processing affective touch. *Human Brain Mapping*, 34(4), 914-922.
- Hakeem, A.Y., Sherwood C.C., Bonar, C.J., Butti C., Hof, P.R., Allman, J.M. (2009). Von Economo Neurons in the elephant brain. *The Anatomical Record*. 292, 242-248.
- Ham, J., Tronick, E. (2006). Infant resilience to the stress of the still-face: Infant and maternal psychophysiology are related. *Annals of the New York Academy of Sciences*, 1094, 297–302. doi.org/10.1196/annals.1376.038
- Herbert, B. M., Pollatos, O. (2012). The body in the mind: on the relationship between interoception and embodiment. *Topics in cognitive science*, 4(4), 692-704.
- Herbert, B. M., Ulbrich, P., Schandry, R. (2007). Interoceptive sensitivity and physical effort: Implications for the self-control of physical load in everyday life. *Psychophysiology*, 44(2), 194–202.
- Ionta, S., Gassert, R., Blanke, O. (2011). Multi-sensory and sensorimotor foundation of bodily self-consciousness—an interdisciplinary approach. *Frontiers in Psychology*, 2, 383.
- Jean, A. D., Stack, D. M., & Fogel, A. (2009). A longitudinal investigation of maternal touching across the first 6 months of life: Age and context effects. *Infant Behavior and Development*, 32(3), 344-349.
- Jönsson, E. H., Kotilahti, K., Heiskala, J., Wasling, H. B., Olausson, H., Croy, I., Mustaniemi, H., Hiltunen, P., Tuulari, J.J., Scheinin, N.M., Karlsson, L., Karlsson H., Nissilä, I. (2018). Affective and non-affective touch evoke differential brain responses in 2-month-old infants. *NeuroImage*, 169, 162-171. doi.org/10.1016/j.neuroimage.2017.12.024
- Kirsch, L. P., Besharati, S., Papadaki, C., Crucianelli, L., Bertagnoli, S., Ward, N., Moro, V., Jenkinson, P.M. & Fotopoulou, A. (2020). Damage to the right insula disrupts the perception of affective touch. *Elife*, 9.
- Kivijärvi, M., Voeten, M. J., Niemelä, P., Räihä, H., Lertola, K., Piha, J. (2001). Maternal sensitivity behavior and infant behavior in early interaction. *Infant Mental Health Journal*, 22(6), 627-640.
- Korja, R., Latva, R., Lehtonen, L. (2012). The effects of preterm birth on mother–infant interaction and attachment during the infant's first two years. *Acta Obstetrica et Gynecologica Scandinavica*, 91(2), 164-173.

- Krahé, C., Drabek, M. M., Paloyelis, Y., & Fotopoulou, A. (2016). Affective touch and attachment style modulate pain: a laser-evoked potentials study. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 371(1708), 20160009.
- Krahé, C., von Mohr, M., Gentsch, A., Guy, L., Vari, C., Nolte, T., & Fotopoulou, A. (2018). Sensitivity to CT-optimal, affective touch depends on Adult Attachment Style. *Scientific Reports*, 8(1), 1-10.
- Leong, V., Byrne, E., Clackson, K., Georgieva, S., Lam, S., Wass, S. (2017). Speaker gaze increases information coupling between infant and adult brains. *Proceedings of the National Academy of Sciences*, 114(50), 13290-13295. doi: 10.1073/pnas.1702493114
- Lester, B. M., Conratt, E., LaGasse, L. L., Tronick, E. Z., Padbury, J. F., & Marsit, C. J. (2018). Epigenetic programming by maternal behavior in the human infant. *Pediatrics*, 142(4).
- Lewkowicz, D. J., & Bremner, A. J. (2020). The development of multisensory processes for perceiving the environment and the self. In: Sathian, K. & Ramachandran, V.S. (Eds). *Multisensory Perception: From Laboratory to Clinic*. Academic Press/Elsevier (pp. 89-112).
- Löken, L.S., Wessberg, J., McGlone, F., Olausson, H. (2009). Coding of pleasant touch by unmyelinated afferents in humans. *Nature Neuroscience*, 12(5), 547.
- Lombardo, M. V., Chakrabarti, B., Bullmore, E. T., Sadek, S. A., Pasco, G., Wheelwright, S. J., ... & Baron-Cohen, S. (2010). Atypical neural self-representation in autism. *Brain*, 133(2), 611-624.
- Ludington-Hoe, S. M., Lewis, T., Morgan, K., Cong, X., Anderson, L., Reese, S. (2006). Breast and infant temperatures with twins during shared kangaroo care. *Journal of Obstetric, Gynecologic and Neonatal Nursing*, 35(2), 223-231.
- MacCormack, J. K., Castro, V. L., Halberstadt, A. G., Rogers, M. L. (2019). Mothers' interoceptive knowledge predicts children's emotion regulation and social skills in middle childhood. *Social Development*, 00, 1-22.
- Maister, L., Tang, T., Tsakiris, M. (2017). Neurobehavioral evidence of interoceptive sensitivity in early infancy. *ELIFE*, 6, e25318. <https://doi.org/10.7554/eLife.25318.001>
- Markova, G., Nguyen, T., Hoehl, S. (2019). Neurobehavioral interpersonal synchrony in early development: The role of interactional rhythms. *Frontiers in psychology*, 10, 2078.
- Marshall, A. G., Sharma, M. L., Marley, K., Olausson, H., McGlone F. P. (2019) Spinal signaling of C-fiber mediated pleasant touch in humans. *eLife*;8:e51642 DOI: 10.7554/eLife.51642
- Marshall, P. J., Meltzoff, A. N. (2015). Body maps in the infant brain. *Trends in Cognitive Sciences*, 19(9), 499-505.
- Matthiesen, A.S., Ransjö-Arvidson, A.B., Nissen, E., Uvnäs-Moberg, K. (2001). Postpartum maternal oxytocin release by newborns: effects of infant hand massage and sucking. *Birth*, 28(1), 13-19.

- McGlone, F., Vallbo, A. B., Olausson, H., Loken, L., & Wessberg, J. (2007). Discriminative touch and emotional touch. *Canadian Journal of Experimental Psychology/Revue Canadienne de Psychologie Expérimentale*, 61(3), 173.
- McGlone, F., Reilly, D. (2010). The cutaneous sensory system. *Neuroscience and Biobehavioral Reviews*, 34(2), 148-159.
- McGlone F., Wessberg J., Olausson H. (2014) Discriminative and affective touch: Sensing and feeling. *Neuron*, 82 (4), 737-755, 10.1016/j.neuron.2014.05.001
- Meltzoff, A.N., Ramírez, R.R., Saby, J.N., Larson, E., Taulu, S., Marshall, P.J. (2018). Infant brain responses to felt and observed touch of hands and feet: An MEG study. *Developmental science*, 21(5), e12651.
- Meltzoff, A.N. (2017). Elements of a comprehensive theory of infant imitation. *Behavioral and Brain Sciences*, 40. DOI:10.1017/S0140525X1600193X
- Mercuri, M., Stack, D. M., Trojan, S., Giusti, L., Morandi, F., Mantis, I., Montirosso, R. (2019). Mothers' and fathers' early tactile contact behaviors during triadic and dyadic parent-infant interactions immediately after birth and at 3-months postpartum: Implications for early care behaviors and intervention. *Infant Behavior and Development*, 57, 101347.
- Miguel, H. O., Gonçalves, Ó. F., Cruz, S., & Sampaio, A. (2019). Infant brain response to affective and discriminative touch: A longitudinal study using fNIRS. *Social Neuroscience*, 14(5), 571-582.
- Miguel, H. O., Lisboa, I. C., Gonçalves, Ó. F., & Sampaio, A. (2019). Brain mechanisms for processing discriminative and affective touch in 7-month-old infants. *Developmental Cognitive Neuroscience*, 35, 20-27.
- Miguel, H. O., Gonçalves, Ó. F., & Sampaio, A. (2020). Behavioral response to tactile stimuli relates to brain response to affective touch in 12-month-old infants. *Developmental Psychobiology*, 62(1), 107-115.
- Montirosso, R., Casini, E., Borgatti, R., Urgesi, C. (2016). Relationship between maternal sensitivity during early interaction and maternal ability in perceiving infants' body and face. *Infancy*, 21(5), 582-602.
- Montirosso R., Provenzi L., Giorda R., Fumagalli M., Morandi F., Sirgiovanni I., Pozzoli U., Grunau R., Oberlander T., Mosca F., Borgatti R. (2016). SLC6A4 promoter region methylation and socio-emotional stress response in very preterm and full-term infants. *Epigenomics*, 8, 895-907.
- Montirosso, R., Piazza, C., Giusti, L., Provenzi, L., Ferrari, P. F., Reni, G., & Borgatti, R. (2019). Exploring the EEG mu rhythm associated with observation and execution of a goal-directed action in 14-month-old preterm infants. *Scientific Reports*, 9(1), 1-12.
- Morrison I., Löken L.S., Olausson H.(2010). The skin as a social organ. *Experimental Brain Research*, 204 (3), 305-314, 10.1007/s00221-009-2007-y

- Morrison I., Bjornsdotter M., Olausson H. (2011). Vicarious responses to social touch in posterior insular cortex are tuned to pleasant caressing speeds *Journal of Neuroscience*, 31 (26) (2011), pp. 9554-9562, 10.1523/JNEUROSCI.0397-11.2011
- Moszkowski, R. J., Stack, D. M., Girouard, N., Field, T. M., Hernandez-Reif, M., & Diego, M. (2009). Touching behaviors of infants of depressed mothers during normal and perturbed interactions. *Infant Behavior and Development*, 32(2), 183-194.
- Murgatroyd, C., Quinn, J. P., Sharp, H. M., Pickles, A., & Hill, J. (2015). Effects of prenatal and postnatal depression, and maternal stroking, at the glucocorticoid receptor gene. *Translational psychiatry*, 5(5), e560-e560.
- Namkung, H., Kim, S. H., Sawa, A. (2017). The insula: An underestimated brain area in clinical neuroscience, psychiatry, and neurology. *Trends in Neurosciences*, 40(4), 200-207.
- Nees, F., Usai, K., Löffler, M., & Flor, H. (2019). The evaluation and brain representation of pleasant touch in chronic and subacute back pain. *Neurobiology of Pain*, 5, 100025.
- Neu, M., Laudenslager, M. L., Robinson, J. (2009). Coregulation in salivary cortisol during maternal holding of premature infants. *Biological Research for Nursing*, 10(3), 226-240.
- Olausson, H., Lamarre, Y., Backlund, H., Morin, C., Wallin, B.G., Starck, G., Ekholm, S., Strigo I., Worsley, K., Vallbo A.B., Bushnell, M.C. (2002). Unmyelinated tactile afferents signal touch and project to insular cortex. *Nature Neuroscience* 5, 900-904.
- Olausson H., Wessberg J., Morrison I., McGlone F., Vallbo Å. (2010). The neurophysiology of unmyelinated tactile afferents. *Neuroscience and Biobehavioral Reviews*, 34 (2), 185-191, 10.1016/j.neubiorev.2008.09.011
- Oldroyd, K., Pasupathi, M., Wainry, C. (2019). Social antecedents to the development of interoception: Attachment related processes are associated with interoception. *Frontiers in Psychology*, 10. 10:712. doi: 10.3389/fpsyg.2019.00712
- Pirazzoli, L., Lloyd-Fox, S., Braukmann, R., Johnson, M. H., & Gliga, T. (2019). Hand or spoon? Exploring the neural basis of affective touch in 5-month-old infants. *Developmental Cognitive Neuroscience*, 35, 28-35.
- Plotnik J.M., De Waal, F.B.M., Reiss, D. (2006). Self-recognition in an Asian elephant. *Proceedings of the National Academic Sciences USA* 103, 17053-17057.
- Porges, S. W. (2007). The polyvagal perspective. *Biological Psychology*, 74(2), 116-143.
- Provasi, J., Anderson, D.I., Barbu-Roth, M. (2014). Rhythm perception, production, and synchronization during the perinatal period. *Frontiers in Psychology*, 5, 1048.
- Provenzi, L., Casini, E., De Simone, P., Reni, G., Borgatti, R., Montirosso, R. (2015). Mother–infant dyadic reparation and individual differences in vagal tone affect 4-month-old infants’ social stress regulation. *Journal of experimental child psychology*, 140, 158-170.

- Provenzi, L., Giusti, L., Fumagalli, M., Tasca, H., Ciceri, F., Menozzi, G., ... and Montirosso, R. (2016). Pain-related stress in the Neonatal Intensive Care Unit and salivary cortisol reactivity to socio-emotional stress in 3-month-old very preterm infants. *Psychoneuroendocrinology*, 72, 161-165.
- Provenzi, L., Fumagalli, M., Bernasconi, F., Sirgiovanni, I., Morandi, F., Borgatti, R., Montirosso, R. (2017). Very preterm and full-term infants' response to socio-emotional stress: The role of postnatal maternal bonding. *Infancy*, 22(5), 695-712.
- Provenzi, L., Giusti, L., Fumagalli, M., Frigerio, S., Morandi, F., Borgatti, R., Mosca, F., Montirosso, R. (2019). The dual nature of hypothalamic-pituitary-adrenal axis regulation in dyads of very preterm infants and their mothers. *Psychoneuroendocrinology*, 100, 172-179.
- Provenzi, L., Fumagalli, M., Scotto di Minico, G., Giorda, R., Morandi, F., Sirgiovanni, I., Schiavolin P., Mosca F., Borgatti R. and Montirosso, R. (2019). Pain-related increase in serotonin transporter gene methylation associates with emotional regulation in 4.5-year-old preterm-born children. *Acta Paediatrica*. 00, 1-9 doi: 10.1111/apa.15077
- Provenzi L., Rosa E., Visintin E., Mascheroni E., Guida E., Cavallini A., Montirosso R. (2020). Understanding the role and function of maternal touch in children with neurodevelopmental disabilities. *Infant Behavior and Development*. *Accepted for publication*.
- Reiss, D., Marino, L. (2001). Mirror self-recognition in the bottlenose dolphin: A case of cognitive convergence. *Proceedings of the National Academic Sciences USA*, 98, (2001), 5937-5942.
- Riva, G. (2017). The neuroscience of body memory: From the self through the space to the others. *Cortex*, 1-20.
- Rochat, P. (2003). Five levels of self-awareness as they unfold early in life. *Consciousness and cognition*, 12(4), 717-731.
- Seth, A. K. (2013). Interoceptive inference, emotion, and the embodied self. *Trends in cognitive sciences*, 17(11), 565-573.
- Seth, A. K., & Tsakiris, M. (2018). Being a beast machine: the somatic basis of selfhood. *Trends in Cognitive Sciences*, 22(11), 969-981.
- Shai, D., Belsky, J. (2017). Parental embodied mentalizing: How the nonverbal dance between parents and infants predicts children's socio-emotional functioning. *Attachment & Human Development*, 19(2), 191-219.
- Sharp, H., Hill, J., Hellier, J., & Pickles, A. (2015). Maternal antenatal anxiety, postnatal stroking and emotional problems in children: Outcomes predicted from pre- and postnatal programming hypotheses. *Psychological Medicine*, 45(2), 269–283. doi.org/10.1017/S0033291714001342
- Slater, R., Cantarella, A., Gallella, S., Worley, A., Boyd, S., Meek, J., and Fitzgerald, M. (2006). Cortical pain responses in human infants. *Journal of Neuroscience*, 26(14), 3662-3666.
- Slaughter, V., & Brownell, C. A. (Eds.). (2011). *Early development of body representations* (Vol. 13). Cambridge University Press.

- Stack, D. M., Jean, A. D. (2011). Communicating through touch: Touching during parent–infant interactions. *The handbook of touch: Neuroscience, Behavioral, and Health perspectives*, 273-298.
- Stein, B. E., Stanford, T. R., and Rowland, B. A. (2014). Development of multisensory integration from the perspective of the individual neuron. *Nature Reviews Neuroscience*, 15(8), 520-535.
- Tajadura-Jiménez, A., Tsakiris, M. (2014). Balancing the “inner” and the “outer” self: Interoceptive sensitivity modulates self–other boundaries. *Journal of Experimental Psychology: General*, 143(2), 736.
- Tanaka, Y., Kanakogi, Y., Kawasaki, M., & Myowa, M. (2018). The integration of audio– tactile information is modulated by multimodal social interaction with physical contact in infancy. *Developmental Cognitive Neuroscience*, 30, 31-40.
- Thomas, R., Misra, R., Akkunt, E., Ho, C., Spence, C., & Bremner, A. J. (2017). Sensitivity to auditory-tactile colocation in early infancy. *Developmental Science*. doi:10.1111/desc.12597
- Tronick, E. Z. (2004). Why is connection with others so critical? Dyadic meaning making, messiness and complexity governed selective processes which co-create and expand individuals’ states of consciousness. *Emotional development*, 293-315.
- Tronick, E.Z., Cohn, J. F. (1989). Infant-mother face-to-face interaction: Age and gender differences in coordination and the occurrence of miscoordination. *Child Development*, 85-92.
- Tronick, E.Z., Als, H., Adamson, L.B., Wise, S., Brazelton, T.B. (1978). The infant’s response to entrapment between contradictory messages in face-to-face interaction. *The American Academy of Child Psychiatry* 17, 1–13.
- Tsakiris, M., Schutz-Bosbach, S. and Gallagher, S. (2007). On agency and body-ownership: Phenomenological and neurocognitive reflections. *Consciousness and Cognition*, 16, 645–660.
- Tsakiris, M. (2010). My body in the brain: a neurocognitive model of body-ownership. *Neuropsychologia*, 48(3), 703-712.
- Tsakiris, M. (2017). The multisensory basis of the self: From body to identity to others. *The Quarterly Journal of Experimental Psychology*, 70:4, 597-609. doi: 10.1080/17470218.2016.118176
- Tuulari, J.J., Scheinin, N.M., Lehtola, S., Merisaari, H., Saunavaara, J., Parkkola, R., . Sehlstedt, I., Karlsson, L., Karlsson, H., Björnsdotter, M. (2019). Neural correlates of gentle skin stroking in early infancy. *Developmental Cognitive Neuroscience*, 35, 36-41. doi.org/10.1016/j.dcn.2017.10.004
- Tylka, T. L., Lumeng, J. C., & Eneli, I. U. (2015). Maternal intuitive eating as a moderator of the association between concern about child weight and restrictive child feeding. *Appetite*, 95, 158-165.
- Van Puyvelde, M., Loots, G., Meys, J., Neyt, X., Mairesse, O., Simcock, D., Pattyn, N. (2015). Whose clock makes yours tick? How maternal cardiorespiratory physiology influences newborns’ heart rate variability. *Biological Psychology*, 108, 132-141.

- Van Puyvelde, M., Collette, L., Gorissen, A. S., Pattyn, N., McGlone, F. (2019). Infants autonomic cardio-respiratory responses to nurturing stroking touch delivered by the mother or the father. *Frontiers in Physiology*, 10, 1117.
- Van Puyvelde, M., Gorissen, A. S., Pattyn, N., McGlone, F. (2019a). Does touch matter? The impact of stroking versus non-stroking maternal touch on cardio-respiratory processes in mothers and infants. *Physiology and Behavior*, 207, 55-63.
- Walker, S. M., Melbourne, A., O'Reilly, H., Beckmann, J., Eaton-Rosen, Z., Ourselin, S., Marlow, N. (2018). Somatosensory function and pain in extremely preterm young adults from the UK EPICure cohort: sex-dependent differences and impact of neonatal surgery. *British Journal of Anaesthesia*, 121(3), 623-635.
- Weiss, S. J., Wilson, P., Hertenstein, M. J., and Campos, R. (2000). The tactile context of a mother's caregiving: Implications for attachment of low birth weight infants. *Infant Behavior and Development*, 23(1), 91-111.
- Welch, M. G. (2016). Calming cycle theory: the role of visceral/autonomic learning in early mother and infant/child behaviour and development. *Acta Paediatrica*, 105(11), 1266-1274.
- Winberg, J. (2005), Mother and newborn baby: Mutual regulation of physiology and behavior- A selective review. *Developmental Psychobiology*, 47: 217-229. doi:10.1002/dev.20094
- Woodhouse, S. S., Scott, J. R., Hepworth, A. D., Cassidy, J. (2020). Secure base provision: A new approach to examining links between maternal caregiving and infant attachment. *Child Development*. 91, e249-e265. doi.org/10.1111/cdev.13224
- Zieber, N., Kangas, A., Hock, A., & Bhatt, R. S. (2015). Body structure perception in infancy. *Infancy*, 20(1), 1-17. doi:10.1111/infa.12064
- Zmyj, N., Jank, J., Schütz-Bosbach, S., Daum, M. M. (2011). Detection of visual–tactile contingency in the first year after birth. *Cognition*, 120(1), 82-89.