

Accepted Article

Measuring disturbance at a swift breeding colonies due to the visual aspects of a drone: a quasi-experiment study.

Geison P. MESQUITA^{a,c*}, José D. RODRÍGUEZ-TEIJEIRO^{b,c}, Serge A. WICH^{d,e}, Margarita MULERO-PÁZMÁNY^d

^aDepartment of Animal Biology Animal, Vegetal Biology and Ecology, Faculty of Bioscience, Autonomous University of Barcelona, Bellaterra, Spain; ^bDepartment of Evolutionary Biology, Ecology and Environmental Sciences, Faculty of Biology, University of Barcelona, and IRBio, Barcelona, Spain; ^cInstitut de Recerca de la Biodiversitat, University of Barcelona, Barcelona, Spain; ^dSchool of Biological and Environmental Sciences, Liverpool John Moores University, Liverpool, UK; ^eInstitute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, Amsterdam, The Netherlands.

*Address correspondence to Geison P. Mesquita. E-mail: geison.pires@e-campus.uab.cat

Handling editor: Zhi-Yun JIA

Received on 9 April 2020; accepted on 8 July 2020

Abstract

There is a growing body of research indicating that drones can disturb animals. However, it is usually unclear whether the disturbance is due to visual or auditory cues. Here, we examined the effect of drone flights on the behaviour of great dusky swifts *Cypseloides senex* and white-collared swifts *Streptoprocne zonaris* in two breeding sites where drone noise was obscured by environmental noise from waterfalls and any disturbance must be largely visual. We performed 12 experimental flights with a multirotor drone at different vertical, horizontal and diagonal distances from the colonies. From all flights, 17% caused <1% of birds to temporarily abandon the breeding site, 50% caused half to abandon and 33% caused more than half to abandon. We showed that the diagonal distance explained 98.9% of the variability of the disturbance percentage and while at distances greater than 50 m the disturbance percentage does not exceed 20%, at less than 40 m the disturbance percentage increase to above 60%. We recommend that flights with a multirotor drone during the breeding period should be conducted at a distance of > 50 m and that recreational flights should be discouraged or conducted at larger distances (e.g. 100 m) in nesting birds areas such as waterfalls, canyons and caves.

Keywords: Drones, Disturbance, *Cypseloides senex*, *Streptoprocne zonaris*, Multi-rotors, Unmanned Aircraft Systems

Multirotor drones are one of the most widely used drone platforms in the civilian environment and with the greatest commercial growth in recent years (Droneii 2019). The main growth factors for scientific, commercial and recreational drone use are associated with a diversity of models relatively easy-to-use, vertical take-off / landing and easy transport. The high maneuverability of multirotor drone and its ability to hover in the air make this type of drone the preferred option for filming and data collection in hard-to- access places (Bakó et al. 2014; Chabot et al. 2015). For these reasons, along with the affordability of commercial models, they are currently the most popular choice for recreational flyers (Rebolo-Ifrán et al. 2019), commercial services (Droneii 2019) and scientists (Chabot and Bird 2015; Jiménez and Mulero-Pázmány 2019).

Within the scientific environment, the integration of drones as data-collection platforms has significantly facilitated vertebrate studies, mainly focused on birds and mammals (Wich and Koh 2018) to address a wide variety of topics, such as species monitoring (Rey et al. 2017; Hodgson et al. 2018); behavioral analysis (Canal et al. 2016; Mulero-Pázmány et al. 2017; Cliff et al. 2018); management (Mulero-Pázmány et al. 2014); habitat mapping (Castellanos-Galindo 2019); and spatial ecology and wildlife diseases (Barasona et al. 2014; Mulero-Pázmány et al. 2015; Laguna et al. 2018). Some of the main advantages of using drones to study wildlife are the reduction of logistical difficulties; costs; risks; and disturbance on wildlife if compared to conventional methods such as manned aircraft surveys or researchers on the ground (Dulava et al. 2015; Christie et al. 2016). Because of this advantage and with the growing appearance of new commercial and recreational models the multirotor drone use has been increasing in research and conservation.

The increase in drone use has raised concerns about the potential disturbance these systems can cause on wildlife (Weston et al. 2020; Bennitt et al. 2019; Bevan et al. 2018). There are a number of factors associated with drone characteristics (drone size, motor type, and flight pattern) and animals (species, life-history stage and level of aggregation) that can be related to the level of disturbance caused by these systems (Mulero-Pázmány et al. 2017). The threshold of disturbance caused by a drone in a given species is often formed by a set of interconnected factors: the sound signature of the drone, the environmental noise level, the visual ability of the species and the association degree of the drone with a threatening stimulus of the species (Bevan et al. 2018). Although all these factors are connected, in the case of birds that in general have worse auditory sensitivity than humans (Dooling and Popper, 2007) and more acute visual perception, the visual stimuli generated by the drone can have a greater effect. Even though some studies that assessed drone disturbance in birds relating flight patterns and distances to the sound and visual aspects of the drone (McEvoy et al. 2016; Rümmler et al. 2016; Brisson-Curadeau et al. 2017; Reintsma et al. 2018), so far it has not been possible to analyze separately the disturbances caused by the visual stimuli of the sound stimuli coming from the drones.

Here, we describe an experiment in which we investigate responses from two species of swifts, great dusky swift *Cypseloides senex* and white-collared swift *Streptoprocne zonaris*, to drone flights in a scenario where noise is mainly masked by the background noise of waterfalls and the visual stimulus the main disturbance factor. We measure the disturbance caused by a multirotor drone at varying distances from swift colonies located in wet rocks walls next or behind waterfalls where the environmental noise is louder than the drone noise. Our aims were 1) bring a new perspective of visual disturbance analysis caused by multirotor drones disassociated from the drone noise; 2) facilitate establishing guidelines that allow minimizing disturbance to bird colonies that use places such as rocks walls next or behind waterfalls, canyons and caves around the world as resting and nesting sites, places with high probability of drone-bird interaction due to the increased recreational drone use and the tourist interest of such places.

Material and Methods

Study area and species

This study was conducted in Chapada das Mesas National Park, Maranhão, Brazil, in October, 2018. The park covers a total area of 1,600 km² within the Cerrado biome, that has various vegetation types, from “cerradão”, which is a type of seasonal forest with dense tree vegetation to “campos limpos” that are open fields as savannas with few trees (Marques and Amorim 2014). The two breeding areas of the study species were: Cachoeira do Prata (6°59'36"S, 47°9'55"W) and Cachoeira de São Romão (7°1'11"S, 47°2'26"W). Both are located in the North of the park along different stretches of the "Farinha" river, a tributary of Araguaia / Tocantins basin, and are approximately 14 km away from each other in a straight line. The breeding areas are the two most voluminous waterfalls present within the park. The Cachoeira do

Prata is formed by a set of falls that reach up to 18 m in height, and the Cachoeira de São Romão has falls of up to 25 m in height (Figure 1). The region has a humid tropical climate characterized by two well-defined seasons: dry, which runs from May to October and wet from November to April, with an annual temperature varying between 24° and 26°C and an annual rainfall varying between 1200 and 1600 mm (IMESC 2008). The waterfalls are accessible to tourists but the number of visitors is low because the access is currently limited to 50 km of dirt road that can only be accessed by 4 × 4 vehicle.

The two study species were the great dusky and white-collared swifts. These are globally considered of least concern according to the Red List (IUCN 2020) with stable population for the great dusky swift and declining population for the white-collared swift population. The great dusky swift distribution is restricted to Argentina, Bolivia, Brazil, and Paraguay (Stopiglia and Raposo 2007) and the white-collared swift is distributed from the USA to Argentina (Chantler 1999). In Brazil, data for both species are sparse, leading to an inaccurate distribution map. Both species are strongly associated to areas with wet rocks walls next or behind waterfalls, canyons and caves. These sites are used with great fidelity for breeding and nesting that occurs between October and November (Whitacre 1989; Stopiglia and Raposo 2007). The two species often share nesting sites (Pearman et al. 2010). In this study most of the individuals identified in the nesting sites were the great dusky swift and few individuals of the white-collared swift.

Drone and experimental flights

The drone model used was a DJI Mavic Pro quad-copter, black color, with diagonal size of 335 mm, 743 g weight, ± 77 dBA noise level, maximum flight speed of 65 km / h, and 20 min average flight autonomy, that carried a camera with a 1/2.3" (CMOS) and sensor with 12.35 effective megapixels. In each of the two swift breeding sites we performed six experimental flights at varying heights above the ground and distances to the breeding rocks walls (Table 1).

All the swift nests were located in the rock wall at $10 \text{ m} \pm 1 \text{ m}$ above the ground in the Cachoeiras do Prata and $15 \text{ m} \pm 1 \text{ m}$ in the Cachoeiras de São Romão (Figure 2). Flights were conducted between 15-18h local time. The drone was launched at a minimum distance of 100 m from the breeding site. During a pilot study conducted a week before the actual experiments we checked that at this distance the drone did not lead to any noticeable reaction from the birds. Between the launch sites and the breeding areas there was vegetation that prevented birds from viewing the drone's take-off. We approached the nesting sites horizontally at a speed between 14 and 21 km / h which in a previous study on birds did not seem to influence bird behaviour (Vas et al. 2015) and allows for good control of the drone. Once the drone reached the set point, which corresponds to the diagonal distance of each flight according to Table 1, it remained hovering stationary for a maximum time of 10 min or until we detected any swifts' behavioural reaction (flying away or mobbing). Once detected any reaction we kept the flight time no more than 5 min to minimize negative effects on the species. An experienced observer using a binocular (10×50) counted the number of birds that were present at the breeding site 5 min before the take-off of each flight and after the drone was landed. At both field sites the observer was positioned between the nesting rocks walls and the drone, with free view to both. Due to the difficulty of approaching the nesting rocks walls and to avoid possible disturbance to the colony, the observer was positioned at a horizontal distance of 15 to 20 m from the base of the rocks walls, hidden from the colony's line of sight. Because of the large number of individuals of the two-species agglomerated and the low luminosity at the waterfalls, we could not determine the number of individuals of each of the two species at the breeding sites and therefore recorded the total number of birds. We established a minimum interval of 30 minutes after landing of each flight or until the birds regrouped in the breeding sites, and a maximum of 2 daily flights, to avoid major disturbances in the species during the same day.

The visual analysis included an assessment of the spots size on the walls, which were agglomerations of the birds, and were used to define whether the birds had regrouped. This is, if the spot size returned to its original size, we assumed that the individuals had returned. For the visual analysis of spot sizes, we compare the spot sizes with rock wall features as atypical marks, deformations or some plants. Due to the high environmental noise caused by the waterfalls, in all the experimental flights in the two studied places it was not possible to hear the drone noise by the observer who was positioned between the drone and the rock walls at a horizontal distance of 15 to 20 m from the base of the rocks walls.

Statistical analysis

As drone disturbance we considered the change in swifts' behaviour (flying away or mobbing). We calculated this disturbance for each experimental flight as the percentage of birds present in the breeding colony 5 minutes before drone exposure minus the percentage of birds present after drone landing. Following Chabot et al. 2015, we classified the drone disturbance level in three categories based on the percentage of birds reacting: 1) noticeable disturbance, when the percentage does not exceed 1%; 2) moderate disturbance, when the percentage does not exceed 50%; and 3) high disturbance, when the percentage is greater than 50%. For vertical distance we considered the difference in height between the nest and the drone on each flight. The horizontal was measured from the projection of the drone to the ground to the colony and the diagonal distance (hereafter distance) was obtained through the Pythagorean theorem. We also calculated the return time of the individuals to the breeding sites after the drone had landed on each flight, and the average time for each of the three categories of disturbance.

A previous descriptive scatter plot showed the possibility of a non-linear association between variables in the two ran models. The first model with diagonal distance as a predictor variable and the disturbance percentage as a dependent variable, and the second model with the disturbance percentage as a predictor variable and the return time as a dependent variable. To choose the best models we initially consider the nature of the variables and Akaike's information criterion (AIC). For model validation we tested for normality test (Shapiro-Wilk), heteroscedasticity (Breusch – Pagan) and set the significance level at 0.05. All analyses and charts were made using “car” (Fox 2016), “drc” (Ritz et al. 2015), “investr” (Greenwell and Schubert 2014) packages in R 3.6.2 with RStudio 1.2.5033 (R Core Team 2019).

Results

Twelve drone flights were performed at different distances from two swift breeding colonies. A maximum disturbance of 93.3% was recorded when the drone flew at 25.5 m distance from a bird's colony, and a minimum of 0.7% disturbance when the flight was conducted at 64.0 m distance (Table 2). During the six flights that produced moderate disturbance initially, a few swifts, ranging from 5 to 40 individuals, showed a mobbing behaviour against the drone. However, the majority of other individuals who showed reactions just left the breeding sites and began to perform circular flights at a distance $20 \text{ m} \pm 5 \text{ m}$ above the drone. Flights performed at less than 29 m produced high disturbance, causing the departure of most of the colony of the breeding sites with just an average of 15.8% of the individuals remaining. In flights with high disturbance we also recorded a larger number of individuals performing mobbing behaviour towards the drone. In each of these flights, we landed as fast as possible.

The nonlinear Gompertz model is the one that presents a lower AIC, 80.16, and the distance from the drone to the colony explained 98.9% of the variability of disturbance percentage. Thus, while at distances greater than 50 m the

percentage of disturbances does not exceed 20%, at less than 40 m the disturbance percentage increase to above 60% (Figure 3). The relationship between the disturbance percentage and the return time, that is, the time it takes for the swifts to return to the colonies is better fitted to a nonlinear power model that explains 97.3% of the variability of return time, and it was the one that presents a lower AIC, 54.5 (Figure 4). On the four flights classified as high disturbance it took an average of 23.5 ± 2.4 minutes for all individuals in the colony to return to the breeding sites after the drone had landed. On flights classified as moderate disturbance this time was reduced to 12 ± 2.9 minutes, whereas on flights with just noticeable disturbance the individuals returned almost immediately after the drone landing.

Discussion

For the first time, we measured the drone visual disturbance separate from the drone noise disturbance in birds breeding colonies from a quasi-experiment where the drone's noise is masked by environment noise, and we found that the response of birds to drone use follows a sigmoidal distribution with the diagonal distance from the drone to the colonies. Although our results are similar to studies that indicate that drone disturbance on birds increases as flight height decreases under different conditions and with different bird species (Rümmeler et al. 2016, Mulero-Pázmány et al. 2017, Vliet et al. 2019), we found that the recommended minimum distance must be greater than 50 m to avoid moderate and high disturbance in breeding sites, which is different from other studies, that were 15 m by common gulls and other species in the bird reserve island Langenwerder in the Baltic Sea (Grenzdörffer 2013) and at least 20 m with drones to survey cliff-nesting seabirds as murre (Brisson-Curadeau et al. 2017). However, unlike all the studies mentioned above, our results show that this reaction to the drone at a greater distance from the colony could be due to the idiosyncrasy of these species but it could also be a consequence of the fact that the drone, without any apparent sound, is more similar to a natural situation of approach of a winged predator to the colony and trigger the defensive reaction earlier. The drone's sound could initially prevent the colony's reaction by being an artificial stimulus not associated with a winged predator, and only when the drone is close enough then triggers this defensive reaction.

The median bird hearing thresholds from 49 bird species suggest that the birds hear best at frequency between about 2 and 3 kHz, while humans generally have better auditory sensitivity with lower auditory thresholds and with wider bandwidth than typical birds (Dooling and Popper, 2007). Therefore, if an observer was unable to hear the drone at 15 m, suppressed or muffled by waterfalls in this experiment, it is assumed that the swifts could not hear the drone at 25 m in the flight closest to the colony. This suggests that the drone noise may lose importance for the disturbance, while the visual aspects such as the shape or the flight pattern can be determinant for the swift's behaviour change. Indeed, the drone visual stimulation was one of the possible causes of disturbances in colonies of greater crested tern *Thalasseus bergii* in a study that suggested that the noise emitted by multirotor drones may not be audible to colonies of this species (Bevan et al. 2018). However, the drone shape of our study eschews the classic "hawk / goose" rule (Schleidt et al. 2011) because a multirotor does not look like any potential swift predator. The new multirotor shape was one of the explanations for the lack of flight response in waterfowl at low flight altitudes in other studies (McEvoy et al. 2016). In contrast, we found that swifts showed mobbing behaviour in flights near the nesting sites and may have recognized the multirotor drone as a potential predator. In the case of the great dusky swift and white-collared swift, the only known aerial predator is the peregrine falcon *Falco peregrinus* which has been observed near the others colony sites awaiting to catch swifts as they enter or leave the colony to feed and collect nest materials (Whitacre 1989). So even though the multirotor does not have a "hawk" shape it is possible that the mobbing behaviour of the swifts facing the drone can be elicited due to the drone being perceived as an unknown potential predator.

The time that swifts took to return to the colony after multirotor flights considered of high disturbance was about 2 times longer than flights considered of moderate disturbance and about 20 times longer than flights considered of low disturbance. This time between departure and return to the original location after the disturbance is also considered a way to measure an animal's response to a disturbance (Vliet et al. 2019). These types of responses can have a negative impact on the reproductive process in the case of birds in their breeding season, since it causes the individual to spend more energy, alters the incubation cycle and the care of altricial nestlings and exposes them to possible predators. This negative impact caused by the return time to the nests was different from others bird studies that measured this time after drone disturbance in breeding colonies: ranging from 1 min to common terns *Sterna hirundo* (Reintsma et al. 2018), 1 to 3 min for Iceland gulls *Larus glaucoideus* and 5 to 10 min for thick-billed murres *Uria lomvia* (Brisson-Curadeau et al. 2017), while our experiment demonstrated much longer return time, whether on high disturbance flights, ranging from 20 to 25 min, or moderate disturbance flights, 9 to 16 min. This variability in return time suggests the need to carry out specific tests to know this effect in different species. Our experiment shows that this delay time in returning to the nesting site can cause very negative impacts on the reproductive process if the presence of these drones is intense over time.

Understanding the minimum operating distance at which drones can cause disturbance, which factors can cause them, and for which species each distance can be tolerated is critical, whether for the preparation of flight missions in scientific studies or to regulate the growing recreational use of drones in such environments. Despite the great diversity of responses to the drone use from different bird species due to the different types of ecological contexts in which they are found, almost always the greater the frequency and intensity of the disturbance, the greater the negative impacts on breeding bird populations. In this sense, the drone use, which is expanding in sites as bird nesting areas, such as this study, should be considered as a possible source of negative effects in certain colony bird. Therefore, we suggest the flight distance with multirotor drone to avoid high disturbance in the great dusky and white-collared swifts during the breeding period in nesting areas should be done > 50 m. We also recommend that recreational flights are generally discouraged or conducted at larger distances (e.g. 100 m) in areas where swifts occur such as waterfalls, canyons and caves. This study serves as a basis both for the elaboration of new protocols for the use of drones with birds by researchers in conservation studies and for possible regulations for the recreational use of drones in protected areas or not with the presence of these species.

Acknowledgments

We are grateful to Neotropical Grassland Conservancy for donating the drone used in this study; Fundação de Amparo a Pesquisa e Desenvolvimento Científico (FAPEMA) for funding part of the study; and the Chico Mendes Institute for Biodiversity Conservation (ICMBio) for providing us with permission and access to the park, in particular the chief of the park Mr. Deijacy Rego for the availability and assistance in the field. This study has been carried out within the framework of biodiversity PhD program at the Autonomous University of Barcelona.

Ethics statement

This project was the authorized N. 64630-1 (scientific purpose) by System of Authorization and Information on Biodiversity – SISBIO in Brazil (art. 28 of IN 03/2014) from Chico Mendes Institute for Biodiversity Conservation – ICMBio, and the flight drone was register certificate N. PP-019272726 by the National Civil Aviation Agency – ANAC.

References

- ANAC – Agência Nacional de Aviação Civil, 2020. Drones Cadastrados. <https://www.anac.gov.br/aceso-a-informacao/dados-abertos/areas-de-atuacao/aeronaves/drones-cadastrados> (Accessed 27 May 2020).
- Bakó G, Tolnai M, Takács Á, 2014. Introduction and Testing of a monitoring and colony-mapping method for waterbird populations that uses high-speed and ultra-detailed aerial remote sensing. *Sensors* 14:12828–12846.
- Barasona JA, Mulero-Pázmány M, Acevedo P, Negro JJ, Vicente J et al., 2014. Unmanned aircraft systems for studying spatial abundance of ungulates: Relevance to spatial epidemiology. *PLoS ONE* 12:1–17.
- Bates D, Maechler M, Bolker B, Walker S, 2015. Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* 67:1–48.
- Bennitt E, Bartlam-Brooks HLA, Hubel TY, Wilson AM, 2019. Terrestrial mammalian wildlife responses to unmanned aerial systems approaches. *Scientific Reports* 9:2142.
- Bevan E, Whiting S, Tucker T, Guinea M, Raith A et al., 2018. Measuring behavioral responses of sea turtles, saltwater crocodiles, and crested terns to drone disturbance to define ethical operating thresholds. *PloS ONE* 13:e0194460.
- Brisson-Curadeau É, Bird D, Burke C, Fifield DA, Pace P et al., 2017. Seabird species vary in behavioural response to drone census. *Scientific Reports* 7:17884.
- Canal D, Mulero-Pázmány M, Negro JJ, Sergio F, 2016. Decoration increases the conspicuousness of raptor nests. *PLoS ONE*, 11:e0157440.
- Castellanos-Galindo GA, Casella E, Mejía-Rentería JC, Rovere A, 2019. Habitat mapping of remote coasts: Evaluating the usefulness of lightweight unmanned aerial vehicles for conservation and monitoring. *Biological Conservation* 239:108282
- Chabot D, Craik SR, Bird DM, 2015. Population census of a large common tern colony with a small unmanned aircraft. *PLoS ONE* 10:e0122588.
- Chabot D, Bird DM, 2015. Wildlife research and management methods in the 21st century: Where do unmanned aircraft fit in? *Journal of Unmanned Vehicle Systems* 3:137–155.
- Chantler P, 1999. Family Apodidae (Swifts). In: *Handbook of the Birds of the World: Barn-Owls to Hummingbirds*. Barcelona: Lynx Edicions, 387–466.
- Christie KS, Gilbert SL, Brown CL, Hatfield M, Hanson L, 2016. Unmanned aircraft systems in wildlife research: Current and future applications of a transformative technology. *Frontiers in Ecology and the Environment* 14:241–251.
- Cliff OM, Saunders DL, Fitch R, 2018. Robotic ecology: Tracking small dynamic animals with an autonomous aerial vehicle. *Science Robotics* 3:eaat8409.
- Dooling RJ, Popper AN, 2007. *The Effects of Highway Noise on Birds*. California Department of Transportation Division of Environmental Analysis.
- Droneii, 2019. Report: Drone Manufacturer Ranking 2019. <https://www.droneii.com/project/drone-manufacturer-ranking-2019> (Accessed 13 December 2019).
- Dulava S, Bean WT, Richmond OMW, 2015. Environmental reviews and case studies: applications of unmanned aircraft systems (UAS) for waterbird surveys. *Environmental Practice* 17:201–210.
- Fox J, 2016. *Applied Regression Analysis and Generalized Linear Models*. 3rd edn. Sage.
- Greenwell BM, Schubert KCM, 2014. investr: an r package for inverse estimation. *The R Journal* 6:90–100.
- Grenzdörffer GJ, 2013. UAS-based automatic bird count of a common gull colony. *ISPRS - International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences* XL-1/W2:169–174.
- Hodgson JC, Mott R, Baylis SM, Pham TT, Wotherspoon S et al., 2018. Drones count wildlife more accurately and precisely than humans. *Methods in Ecology and Evolution* 9:1160–1167.

- IMESC, 2008. Perfil do Maranhão 2006/2007. <http://imesc.ma.gov.br/porta/Post/view/outras-publicacoes/38> (Accessed 8 January 2019).
- IUCN, 2020. *The IUCN Red List of Threatened Species*. Version 2020-1. <https://www.iucnredlist.org>. (Accessed 15 April 2020).
- Jiménez López J, Mulero-Pázmány M, 2019. Drones for conservation in protected areas: present and future. *Drones* 3:1–23.
- Laguna E, Barasona JA, Triguero-Ocaña R, Mulero-Pázmány M, Negro JJ et al., 2018. The relevance of host overcrowding in wildlife epidemiology: A new spatially explicit aggregation index. *Ecological Indicators* 84:695–700.
- Lyons M, Brandis K, Callaghan C, Mccann J, Mills C et al., 2018. Bird interactions with drones from individuals to large colonies. *Australian Field Ornithology* 35:51–56.
- Marques AR, Amorim MCCT, 2014. Saberes geográficos integrados aos estudos territoriais sob a ótica da implantação do Parque Nacional da Chapada das Mesas, Sertão de Carolina/MA. *Geografia em Questão* 7:100–117.
- McEvoy JF, Hall GP, McDonald PG, 2016. Evaluation of unmanned aerial vehicle shape, flight path and camera type for waterfowl surveys: disturbance effects and species recognition. *PeerJ* 4:e1831.
- Montgomerie RD, Weatherhead PJ, 1988. Risks and rewards of nest defense by parent birds. *Quarterly Review of Biology* 63:167–187.
- Mulero-Pázmány M, Stolper R, Essen LD, Negro JJ, Sassen T, 2014. Remotely piloted aircraft systems as a rhinoceros anti-poaching tool in Africa. *PLoS ONE* 9:1–10.
- Mulero-Pázmány M, Barasona JÁ, Acevedo P, Vicente J, Negro JJ, 2015. Unmanned aircraft systems complement biologging in spatial ecology studies. *Ecology and Evolution*, 5: 4808–4818.
- Mulero-Pázmány M, Jenni-Eiermann S, Strebel N, Sattler T, Negro JJ et al., 2017. Unmanned aircraft systems as a new source of disturbance for wildlife: A systematic review *PLoS ONE* 12:e0178448.
- Pearman M, Areta JJ, Roesler I, Bodrati A, 2010. Confirmation of the sooty swift *Cypseloides fumigatus* in Argentina with notes on its nest placement, seasonality, and distribution. *Ornitologia Neotropical* 21:351–359.
- R Core Team, 2019. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>. (Accessed 22 December 2019).
- Rebolo-Ifrán N, Grilli MG, Lambertucci SA, 2019. Drones as a threat to wildlife: youtube complements science in providing evidence about their effect. *Environmental Conservation* 46:205–210.
- Reintsma KM, McGowan PC, Callahan C, Collier T, Gray D et al., 2018. Preliminary evaluation of behavioral response of nesting waterbirds to small unmanned aircraft flight. *Waterbirds* 41:326–331.
- Rey N, Volpi M, Joost S, Tuia D, 2017. Detecting animals in African Savanna with UAVs and the crowds. *Remote Sensing of Environmental* 200:341–351.
- Rümmler M-C, Mustafa O, Maercker J, Peter H-U, Esefeld J, 2016. Measuring the influence of unmanned aerial vehicles on Adélie penguins. *Polar Biology* 39:1329–1334.
- Schleidt W, Shalter MD, Moura-Neto H, 2011. The hawk/goose story: the classical ethological experiments of Lorenz and Tinbergen, revisited. *Journal of Comparative Psychology* 125:121–133.
- Stopiglia R, Raposo MA, 2007. Distribuição e biologia do andorinhão-preto-da-cascata *Cypseloides fumigatus* e do andorinhão-velho-da-cascata *C. senex* no Brasil: uma síntese. *Cotinga* 28:49–57.
- Svagej WS, Trivellini MM, Quintana F, 2011. Parental investment theory and nest defense by Imperial Shags: effects of offspring number, offspring age, laying date and parent sex. *Ethology* 118:251–259.

- Vas E, Lescroel A, Duriez O, Boguszewski G, Grémillet D, 2015. Approaching birds with drones: first experiments and ethical guidelines. *Biology Letters* 11: 20140754.
- Vliet RE van der, Jeninga L, Burg A van den, 2019. RPAS over Natura 2000 areas: Disturbance responses of wildlife and opportunities for research RPAS over Natura 2000 areas: Disturbance responses of wildlife and opportunities for research. RPAS Civil Operators & Operations Forum 8th Annual International Conference, December.
- Weston MA, O'Brien C, Kostoglou K, Symonds MRE, 2020. Escape responses of terrestrial and aquatic birds to drones: towards a code of practice to minimise disturbance. *Journal of Applied Ecology* 1365-2664.
- Whitacre DF, 1989. Conditional use of nest structures by white-naped and white-collared swift. *The Condor* 91:813-825.
- Wich SA, Koh LP, 2018. *Conservation Drones: Mapping and Monitoring Biodiversity*. Oxford: Oxford University Press.
- Wickham H, 2016. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

Table 1. Experimental flights parameters. Distances are in meters.

Flight	Date	Time	Study Site	Height nests	Flight Altitude	Vertical Distance	Horizontal Distance	Diagonal Distance
1	22/10/2018	16:00	Cachoeira do Prata	10	50	40	50	64.03
2	22/10/2018	17:30	Cachoeira do Prata	10	25	15	50	52.20
3	23/10/2018	16:00	Cachoeira de São Romão	15	50	35	50	61.03
4	23/10/2018	17:30	Cachoeira de São Romão	15	25	10	50	50.99
5	24/10/2018	16:00	Cachoeira do Prata	10	10	0	50	50.00
6	24/10/2018	17:30	Cachoeira do Prata	10	50	40	25	47.17
7	25/10/2018	16:00	Cachoeira de São Romão	15	10	-5	50	50.25
8	25/10/2018	17:30	Cachoeira de São Romão	15	50	35	25	43.01
9	26/10/2018	16:00	Cachoeira do Prata	10	25	15	25	29.15
10	26/10/2018	17:30	Cachoeira de São Romão	15	25	10	25	26.93
11	27/10/2018	16:00	Cachoeira do Prata	10.00	10	0	25	25.00
12	27/10/2018	17:30	Cachoeira de São Romão	15.00	10.00	-5	25	25.50

Table 2. Percentage disturbed and classification of experimental flights. 1 (noticeable disturbance), 2 (moderate disturbance) and 3 (high disturbance).

Classification	Flight	Date	Time	Study Site	Diagonal Distance (m)	Total Swifts	Disturbed (%)	Return Time (min)
1	1	22/10/2018	16:00	Cachoeira do Prata	64.03	3000	0.7	1
1	3	23/10/2018	16:00	Cachoeira de São Romão	61.03	1000	1.0	1
2	2	22/10/2018	17:30	Cachoeira do Prata	52.20	1000	5.0	9
2	4	23/10/2018	17:30	Cachoeira de São Romão	50.99	3000	10.0	9
2	5	24/10/2018	16:00	Cachoeira do Prata	50.00	1000	15.0	12
2	6	24/10/2018	17:30	Cachoeira do Prata	47.17	2500	32.0	15
2	7	25/10/2018	16:00	Cachoeira de São Romão	50.25	2500	20.0	12
2	8	25/10/2018	17:30	Cachoeira de São Romão	43.01	1500	46.7	16
3	9	26/10/2018	16:00	Cachoeira do Prata	29.15	1000	70.0	20
3	10	26/10/2018	17:30	Cachoeira de São Romão	26.93	3000	83.3	22
3	11	27/10/2018	16:00	Cachoeira do Prata	25.00	1000	90.0	25
3	12	27/10/2018	17:30	Cachoeira de São Romão	25.50	3000	93.3	25




Figure 1. Localization of the studied swift breeding sites in Chapada das Mesas National Park at Brazil. Cachoeira do Prata (white square), Cachoeira de São Romão (white circle).




Figure 2. Design of experimental flights. Breeding group from “Cachoeira do Prata” and “Cachoeira de São Romão”. Classification (circle, noticeable disturbance; triangle, moderate disturbance; square, high disturbance), Diagonal distance (meters) and disturbance (%) for each drone flight.




Figure 3. Nonlinear gompertz regression between diagonal distance and % disturbed of swifts. In blue 95% confidence band and green prediction band.




Figure 4. Nonlinear power regression between % disturbed od swifts and return time. In blue 95% confidence band and green prediction band.